

Safety evaluation of the food enzyme cellobiose phosphorylase from the genetically modified *Escherichia coli* strain LE1B109-pPB130

EFSA Panel on Food Contact Materials, Enzymes and Processing Aids (CEP) |

Claude Lambré | José Manuel Barat Baviera | Claudia Bolognesi | Pier Sandro Cocconcelli | Riccardo Crebelli | David Michael Gott | Konrad Grob | Evgenia Lampi | Marcel Mengelers | Alicja Mortensen | Gilles Rivière | Inger-Lise Steffensen | Christina Tlustos | Henk Van Loveren | Laurence Vernis | Holger Zorn | Yrjö Roos | Ana Gomes | Magdalena Andryszkiewicz | Erik Boinowitz | Cristina Fernández-Fraguas | Yi Liu | Andrew Chesson

Correspondence: fip@efsa.europa.eu

Abstract

The food enzyme cellobiose phosphorylase (cellobiose: phosphate α -D-glucosyltransferase; EC 2.4.1.20) is produced with the genetically modified *Escherichia coli* strain LE1B109-pPB130 by c-LEcta GmbH. The genetic modifications do not give rise to safety concerns. The food enzyme is considered free from viable cells of the production organism and its DNA. It is intended to be used in combination with a sucrose phosphorylase in the production of the specialty carbohydrate cellobiose. Since residual amounts of total organic solids are removed by downstream purification steps, the Panel considered that toxicological studies other than assessment of allergenicity were unnecessary and a dietary exposure was not estimated. A search for similarity of the amino acid sequence of the food enzyme to known allergens was made and no match was found. The Panel considered that, under the intended conditions of use, the risk of allergic reactions upon dietary exposure cannot be excluded, but the likelihood is low. Based on the data provided, the Panel concluded that this food enzyme does not give rise to safety concerns under the intended conditions of use.

KEYWORDS

cellobiose phosphorylase, cellobiose:phosphate α -D-glucosyltransferase, *E. coli*, EC 2.4.1.20, EFSA-Q-2021-00297, genetically modified microorganism, food enzyme

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.

© 2024 European Food Safety Authority. *EFSA Journal* published by Wiley-VCH GmbH on behalf of European Food Safety Authority.

CONTENTS

Abstract.....	1
1. Introduction	3
1.1. Background and Terms of Reference as provided by the requestor.....	3
1.1.1. Background as provided by the European Commission.....	3
1.1.2. Terms of Reference.....	3
2. Data and methodologies.....	3
2.1. Data.....	3
2.2. Methodologies.....	4
3. Assessment.....	4
3.1. Source of the food enzyme.....	4
3.1.1. Characteristics of the parental and recipient microorganism.....	4
3.1.2. Characteristics of introduced sequences.....	5
3.1.3. Description of the genetic modification.....	5
3.1.4. Safety aspects of the genetic modification.....	5
3.2. Production of the food enzyme	5
3.3. Characteristics of the food enzyme	5
3.3.1. Properties of the food enzyme.....	5
3.3.2. Chemical parameters.....	6
3.3.3. Purity.....	6
3.3.4. Viable cells and DNA of the production strain.....	7
3.4. Toxicological data	7
3.4.1. Allergenicity	7
3.5. Dietary exposure.....	7
3.5.1. Intended use of the food enzyme.....	7
3.5.2. Dietary exposure estimation.....	8
3.6. Margin of exposure	8
4. Conclusions.....	8
5. Documentation as provided to EFSA	8
Abbreviations	8
Conflict of interest	8
Requestor.....	8
Question number.....	8
Copyright for non-EFSA content.....	8
Panel members	8
Note.....	9
References.....	9

1 | INTRODUCTION

Article 3 of the Regulation (EC) No 1332/2008¹ provides definition for ‘food enzyme’ and ‘food enzyme preparation’.

‘Food enzyme’ means a product obtained from plants, animals or microorganisms or products thereof including a product obtained by a fermentation process using microorganisms: (i) containing one or more enzymes capable of catalysing a specific biochemical reaction; and (ii) added to food for a technological purpose at any stage of the manufacturing, processing, preparation, treatment, packaging, transport or storage of foods.

‘Food enzyme preparation’ means a formulation consisting of one or more food enzymes in which substances such as food additives and/or other food ingredients are incorporated to facilitate their storage, sale, standardisation, dilution or dissolution.

Before January 2009, food enzymes other than those used as food additives were not regulated or were regulated as processing aids under the legislation of the Member States. On 20 January 2009, Regulation (EC) No 1332/2008 on food enzymes came into force. This Regulation applies to enzymes that are added to food to perform a technological function in the manufacture, processing, preparation, treatment, packaging, transport or storage of such food, including enzymes used as processing aids. Regulation (EC) No 1331/2008² established the European Union (EU) procedures for the safety assessment and the authorisation procedure of food additives, food enzymes and food flavourings. The use of a food enzyme shall be authorised only if it is demonstrated that:

- it does not pose a safety concern to the health of the consumer at the level of use proposed;
- there is a reasonable technological need;
- its use does not mislead the consumer.

All food enzymes currently on the European Union market and intended to remain on that market, as well as all new food enzymes, shall be subjected to a safety evaluation by the European Food Safety Authority (EFSA) and approval via an EU Community list.

1.1 | Background and Terms of Reference as provided by the requestor

1.1.1 | Background as provided by the European Commission

Only food enzymes included in the Union list may be placed on the market as such and used in foods, in accordance with the specifications and conditions of use provided for in Article 7(2) of Regulation (EC) No 1332/2008 on food enzymes.

An application has been introduced by the applicant “c-LEcta GmbH” for the authorisation of the food enzyme Cellobiose phosphorylase from a genetically modified strain of *E. coli* K12 (strain LE1B109-pPB130).

Following the requirements of Article 12.1 of Regulation (EC) No 234/2011³ implementing Regulation (EC) No 1331/2008, the Commission has verified that the application falls within the scope of the food enzyme Regulation and contains all the elements required under Chapter II of that Regulation.

1.1.2 | Terms of Reference

In accordance with Article 29(1)(a) of Regulation (EC) No 178/2002, the European Commission requests the European Food Safety Authority to carry out the safety assessment on the following food enzyme: Cellobiose phosphorylase from a genetically modified strain of *E. coli* K12 (strain LE1B109_pPB130) in accordance with Regulation (EC) No 1331/2008 establishing a common authorisation procedure for food additives, food enzymes and flavourings.

2 | DATA AND METHODOLOGIES

2.1 | Data

The applicant has submitted a dossier in support of the application for authorisation of the food enzyme cellobiose phosphorylase from a genetically modified *Escherichia coli* strain LE1B109_pPB130.

¹Regulation (EC) No 1332/2008 of the European Parliament and of the Council of 16 December 2008 on Food Enzymes and Amending Council Directive 83/417/EEC, Council Regulation (EC) No 1493/1999, Directive 2000/13/EC, Council Directive 2001/112/EC and Regulation (EC) No 258/97. OJ L 354, 31.12.2008, pp. 7–15.

²Regulation (EC) No 1331/2008 of the European Parliament and of the Council of 16 December 2008 establishing a common authorisation procedure for food additives, food enzymes and food flavourings. OJ L 354, 31.12.2008, pp. 1–6.

³Commission Regulation (EU) No 234/2011 of 10 March 2011 implementing Regulation (EC) No 1331/2008 of the European Parliament and of the Council establishing a common authorisation procedure for food additives, food enzymes and food flavourings. OJ L 64, 11.03.2011, pp. 15–24.

Additional information was requested from the applicant during the assessment process on 21 June 2022 and received on 23 January 2023 (see [‘Documentation provided to EFSA’](#)).

2.2 | Methodologies

The assessment was conducted in line with the principles described in the EFSA ‘Guidance on transparency in the scientific aspects of risk assessment’ (EFSA, 2009a) and following the relevant existing guidance documents of EFSA Scientific Committee.

The ‘Guidance on the submission of a dossier on food enzymes for safety evaluation’ (EFSA, 2009b) as well as the ‘Statement on characterisation of microorganisms used for the production of food enzymes’ (EFSA CEP Panel, 2019) have been followed for the evaluation of the application. Additional information was requested in accordance with the updated ‘Scientific Guidance for the submission of dossiers on food enzymes’ (EFSA CEP Panel, 2021) and the guidance on the ‘Food manufacturing processes and technical data used in the exposure assessment of food enzymes’ (EFSA CEP Panel, 2023).

3 | ASSESSMENT

IUBMB nomenclature	Cellobiose phosphorylase
Systematic name	Cellobiose: phosphate α -D-glucosyltransferase
Synonyms	Cellobiose:orthophosphate α -D-glucosyltransferase
IUBMB no	2.4.1.20
CAS no	9030-20-0
EINECS no	–

Abbreviations: CAS, Chemical Abstracts Service; EINECS, European Inventory of Existing Commercial Chemical Substances; IUBMB, International Union of Biochemistry and Molecular Biology.

Cellobiose phosphorylases catalyse the formation of a β -(1,4) glycosidic bond between α -D-glucose-1-phosphate and glucose, forming cellobiose and releasing phosphate. The enzyme under application is intended to be used in combination with a sucrose phosphorylase in the production of the specialty carbohydrate cellobiose.

3.1 | Source of the food enzyme

The cellobiose phosphorylase is produced with the genetically modified bacterium *Escherichia coli* strain LE1B109_pPB130, which is deposited at the German Collection of Microorganisms and Cell Cultures (DSMZ, Germany), with the deposit number [REDACTED].⁴

The production strain was identified as *E. coli* using [REDACTED]

[REDACTED].⁵ No acquired antimicrobial resistance genes or virulence factors of concern were identified in its genome.

3.1.1 | Characteristics of the parental and recipient microorganism

The parental microorganism is [REDACTED], a strain that has been well-characterised and whose safety (non-pathogenicity) has been documented (Gorbach, 1978). *E. coli* K-12 was shown to be ineffective in colonising the human gut. Its genome has been fully sequenced (Hayashi et al., 2006). [REDACTED]

⁴Technical dossier/Confidential Dossier/Annex V_b.

⁵Technical dossier/Confidential Dossier/Annex V_c.

⁶Technical dossier/Confidential Dossier/Annex V_c.

3.1.2 | Characteristics of introduced sequences

The sequence encoding the cellobiose phosphorylase [REDACTED]

3.1.3 | Description of the genetic modification

The purpose of the genetic modification was to enable the production strain to synthesise the cellobiose phosphorylase [REDACTED].

3.1.4 | Safety aspects of the genetic modification

The technical dossier contains all necessary information on the recipient microorganism, the donor organism and the genetic modification process.

The production strain *E. coli* LE1B109_pPB130 differs from the recipient strain in its capacity to produce the cellobiose phosphorylase [REDACTED].

No issues of concern arising from the genetic modification were identified by the Panel.

3.2 | Production of the food enzyme

The food enzyme is manufactured according to the Food Hygiene Regulation (EC) No 852/2004,⁹ with food safety procedures based on Hazard Analysis and Critical Control Points, and in accordance with current Good Manufacturing Practice.¹⁰

The production strain is grown as a pure culture using a typical industrial medium in a submerged, fed-batch fermentation system with conventional process controls in place. [REDACTED] is added to the fermentation medium to induce the production of the cellobiose phosphorylase. After completion of the fermentation, the solid biomass is removed from the fermentation broth by centrifugation and the cells are mechanically lysed to release the intracellular enzyme. The lysate containing the enzyme is treated with [REDACTED] and further purified and concentrated, including an ultrafiltration step in which enzyme protein is retained, while most of the low molecular mass material passes the filtration membrane and is discarded.¹¹ The applicant provided information on the identity of the substances used to control the fermentation and in the subsequent downstream processing of the food enzyme.¹²

The Panel considered that sufficient information has been provided on the manufacturing process and the quality assurance system implemented by the applicant.

3.3 | Characteristics of the food enzyme

3.3.1 | Properties of the food enzyme

The cellobiose phosphorylase is a single polypeptide chain of [REDACTED] amino acids.¹³ The molecular mass of the mature protein, calculated from the amino acid sequence, is 91.4 kDa.¹⁴ The food enzyme was analysed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). A consistent protein pattern was observed across all batches. The gels

⁷Technical dossier/Confidential Dossier/Annex V_c.

⁸Technical dossier/Confidential Dossier/Annex V_c.

⁹Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of food additives. OJ L 226, 25.6.2004, pp. 3–21.

¹⁰Technical Dossier/Risk assessment data/p. 20 and additional information January 2023/Reply to question 3.

¹¹Technical Dossier/Risk assessment data/p. 22–30 and Annex VI.

¹²Technical Dossier/Annex VII.

¹³Technical Dossier/Risk assessment data/p. 11.

¹⁴Technical Dossier/Risk assessment data/p. 11.

showed the target protein migrating between the marker proteins of 66 and 116 kDa, consistent with the expected molecular mass of the enzyme.¹⁵ No other enzymatic activities were reported.¹⁶

The in-house determination of cellobiose phosphorylase activity is based on the hydrolysis of cellobiose (reaction conditions: pH 7.0, 30°C, 5 min), by measuring the release of α -D-glucose-1-phosphate by means of a coupled reaction that forms NADPH, which is then detected spectrophotometrically at 340 nm. The enzyme activity is expressed in U/mL. One Unit (U) corresponds to the formation of 1 μ mol glucose-1-phosphate per minute under the conditions of the assay.¹⁷

The food enzyme has a temperature optimum around 44°C (pH 7.0) and a pH optimum around pH 7.0 (30°C).¹⁸ Thermostability was tested after a pre-incubation of the food enzyme for 15 min at different temperatures (pH 7.0). The cellobiose phosphorylase activity decreased sharply above 50°C, showing 20% residual activity at 55°C.¹⁹

3.3.2 | Chemical parameters

Data on the chemical parameters of the food enzyme preparation were provided for three batches used for commercialisation (Table 1).²⁰ The mean total organic solids (TOS) was 5.8% and the mean enzyme activity/TOS ratio was 11.9 U/mg TOS.

TABLE 1 Composition of the food enzyme preparation.

Parameters	Unit	Batches		
		1	2	3
Sucrose phosphorylase activity	U/mL batch ^a	459	736	892
Protein	%	3.6	4.6	4.7
Ash	%	0.8	0.8	0.8
Water	%	44.3	43.3	42.7
Excipient (glycerol)	%	50.0	50.0	50.0
Total organic solids (TOS)^b	%	4.9	5.9	6.5
Activity/TOS ratio	U/mg TOS	9.4	12.5	13.7

^aU: Unit (see Section 3.3.1).

^bTOS calculated as 100% – % water – % ash – % excipient.

3.3.3 | Purity

The lead content in the three commercial batches was below 5 mg/kg,²¹ which complies with the specification for lead as laid down in the general specifications for enzymes used in food processing (FAO/WHO, 2006). In addition, the concentrations of cadmium, mercury and arsenic were below the limit of detection (LoD) of the employed methods.^{22,23}

The food enzyme preparation complies with the microbiological criteria for total coliforms, *Escherichia coli* and *Salmonella* as laid down in the general specifications for enzymes used in food processing (FAO/WHO, 2006).²⁴ No antimicrobial activity was detected in any of the tested batches.²⁵

██████ was used as an inducer in the fermentation process. Its concentration in the three batches of the food enzyme was below the LoD of the employed method. The Panel noted that this method had a LoD of ████████.²⁶ Considering the absence of TOS in the final food (Section 3.5.1), any residual ████████ would also have been removed during the downstream purification steps.

The Panel considered that the information provided on the purity of the food enzyme is sufficient.

¹⁵Technical Dossier/Risk assessment data/p. 11, 13.

¹⁶Technical Dossier/Risk assessment data/p. 17.

¹⁷Technical Dossier/Annex II a.

¹⁸Technical Dossier/Risk assessment data/p. 16–17.

¹⁹Technical Dossier/Risk assessment data/p. 16–17.

²⁰Technical Dossier/Risk assessment data/p. 12 and Annexes III a-c.

²¹Technical Dossier/Annexes III a-c.

²²LoDs: Pb=0.02 mg/kg; As=0.05 mg/kg; Cd=0.01 mg/kg; Hg=0.01 mg/kg.

²³Technical Dossier/Annexes III a-c.

²⁴Technical Dossier Annexes III a-c.

²⁵Technical Dossier/Annexes III a-c.

²⁶Technical dossier/Additional data January 2023/Answer to question 1 and Annexes II Section 1.

3.3.4 | Viable cells and DNA of the production strain

The absence of viable cells of the production strain in the food enzyme was demonstrated in three independent batches analysed in quintuplicate. One millilitre of product was incubated in non-selective enrichment medium at [REDACTED]. Subcultures were then plated onto selective agar and incubated at [REDACTED]. No colonies were produced.²⁷ A positive control was included.

The absence of recombinant DNA in the food enzyme was demonstrated by polymerase chain reaction analysis of three batches in triplicate.²⁸ No DNA was detected with primers that would have amplified a [REDACTED] in the production strain, with a LoD of [REDACTED].²⁹

3.4 | Toxicological data

In the course of the food manufacturing process, the food enzyme is removed by the purification steps applied (see Section 3.5). Consequently, in the absence of exposure, the Panel considered that no toxicological studies other than assessment of allergenicity were needed for the assessment of this food enzyme.

3.4.1 | Allergenicity

The allergenicity assessment considers only the food enzyme and not any carrier or other excipient that may be used in the final formulation.

The potential allergenicity of the cellobiose phosphorylase produced with the *E. coli* strain LE1B109_pPB130 was assessed by comparing its amino acid sequence with those of known allergens according to the 'Scientific opinion on the assessment of allergenicity of GM plants and microorganisms and derived food and feed of the Scientific Panel on Genetically Modified Organisms' (EFSA GMO Panel, 2010). Using higher than 35% identity in a sliding window of 80 amino acids as the criterion, no match was found.³⁰

No information was available on oral and respiratory sensitisation or elicitation reactions of this cellobiose phosphorylase.³¹

Some phosphorylases have shown food allergenic potential (Rangkakulnuwat et al., 2023). However, a literature search performed by the applicant did not identify any publication on allergic reactions to cellobiose phosphorylase.

The Panel considered that a risk of allergic reactions upon dietary exposure to this food enzyme cannot be excluded, but the likelihood is low.

3.5 | Dietary exposure

3.5.1 | Intended use of the food enzyme

The food enzyme is intended to be used in the production of the specialty carbohydrate cellobiose at a recommended use level of 27,200 Unit/kg glucose, corresponding to 2,893.62 mg TOS/kg glucose.³²

In the manufacturing of cellobiose, the food enzyme is used in combination with a sucrose phosphorylase and is added to sucrose, glucose and phosphate. The successive action of sucrose phosphorylase and cellobiose phosphorylase converts the sucrose and glucose into cellobiose. The reaction product is then passed through a membrane ([REDACTED]) to retain the enzyme in the reactor. The cellobiose-containing permeate is subjected to electro dialysis to remove charged constituents. Cellobiose is then obtained by crystallisation.³³

To establish the extent of the food enzyme–TOS removal, the applicant measured the amount of proteins present in the cellobiose product by SDS-PAGE analysis with silver staining, showing a removal of > 99%.³⁴ These data were considered by the Panel as sufficient to confirm the absence of TOS in the final product.

²⁷Technical dossier/Additional data January 2023/Answer to question 4 and Annex II Section 4.

²⁸Technical dossier/Annex V_e. and additional data January 2023/Answer to question 5 and Annex II Section 5.

²⁹Technical dossier/Additional data January 2023/Answer to question 5 and Annex II Section 5.

³⁰Technical Dossier/Risk assessment data/p. 34–36/Annex IX.

³¹Technical Dossier/Additional information January 2023/Answer to question 6.

³²Technical dossier/Additional information January 2023/Answer 7.

³³Technical dossier/Annex VIII.

³⁴Technical dossier/Additional information January 2023/Answer 8 and Annex II Section 8.

3.5.2 | Dietary exposure estimation

The Panel accepted the evidence provided as sufficient to conclude that the residual amounts of food enzyme–TOS in the cellobiose final products is negligible. Consequently, a dietary exposure was not calculated.

3.6 | Margin of exposure

Since no toxicological assessment was considered necessary by the Panel, a margin of exposure was not calculated.

4 | CONCLUSIONS

Based on the data provided and the removal of TOS during the intended food manufacturing process, the Panel concluded that the food enzyme cellobiose phosphorylase produced with the genetically modified *Escherichia coli* strain LE1B109_pPB130 does not give rise to safety concerns under the intended conditions of use.

The CEP Panel considered the food enzyme free from viable cells of the production organism and recombinant DNA.

5 | DOCUMENTATION AS PROVIDED TO EFSA

Cellobiose phosphorylase from the GM *E. coli* strain LE1B109_pPB130. Submitted by c-LEcta GmbH. The dossier was submitted on 09 June 2021.

Additional information. January 2023. Submitted by c-LEcta GmbH.

ABBREVIATIONS

CAS	Chemical Abstracts Service
CEP	EFSA Panel on Food Contact Materials, Enzymes and Processing Aids
EC	European Commission
EINECS	European Inventory of Existing Commercial Chemical Substances
FAO	Food and Agriculture Organization of the United Nations
IPTG	isopropyl β -D-1-thiogalactopyranoside
IUBMB	International Union of Biochemistry and Molecular Biology
kDa	kiloDalton
LoD	limit of detection
SDS-PAGE	sodium dodecyl sulfate-polyacrylamide gel electrophoresis
TOS	total organic solids
WGS	whole genome sequencing
WHO	World Health Organization

CONFLICT OF INTEREST

If you wish to access the declaration of interests of any expert contributing to an EFSA scientific assessment, please contact interestmanagement@efsa.europa.eu.

REQUESTOR

European Commission

QUESTION NUMBER

EFSA-Q-2021-00297

COPYRIGHT FOR NON-EFSA CONTENT

EFSA may include images or other content for which it does not hold copyright. In such cases, EFSA indicates the copyright holder and users should seek permission to reproduce the content from the original source.

PANEL MEMBERS

José Manuel Barat Baviera, Claudia Bolognesi, Andrew Chesson, Pier Sandro Cocconcelli, Riccardo Crebelli, David Michael Gott, Konrad Grob, Claude Lambré, Evgenia Lampi, Marcel Mengelers, Alicja Mortensen, Gilles Rivière, Vittorio Silano (until 21 December 2020[†]), Inger-Lise Steffensen, Christina Tlustos, Henk Van Loveren, Laurence Vernis and Holger Zorn.

[†]Deceased.

NOTE

The full opinion will be published in accordance with Article 12 of Regulation (EC) No 1331/2008 once the decision on confidentiality will be received from the European Commission.

REFERENCES

- EFSA (European Food Safety Authority). (2009a). Guidance of the Scientific Committee on transparency in the scientific aspects of risk assessments carried out by EFSA. Part 2: General principles. *EFSA Journal*, 7(5), 1051. <https://doi.org/10.2903/j.efsa.2009.1051>
- EFSA (European Food Safety Authority). (2009b). Guidance of EFSA prepared by the scientific panel of food contact material, enzymes, Flavourings and processing aids on the submission of a dossier on food enzymes. *EFSA Journal*, 7(8), 1305. <https://doi.org/10.2903/j.efsa.2009.1305>
- EFSA CEP Panel (EFSA Panel on Food Contact Materials, Enzymes and Processing Aids). (2019). Statement on the characterisation of microorganisms used for the production of food enzymes. *EFSA Journal*, 17(6), 5741. <https://doi.org/10.2903/j.efsa.2019.5741>
- EFSA CEP Panel (EFSA Panel on Food Contact Materials, Enzymes and Processing Aids), Lambré, C., Barat Baviera, J. M., Bolognesi, C., Cocconcelli, P. S., Crebelli, R., Gott, D. M., Grob, K., Lampi, E., Mengelers, M., Mortensen, A., Rivière, G., Steffensen, I.-L., Tlustos, C., Van Loveren, H., Vernis, L., Zorn, H., Glandorf, B., Herman, L., ... Chesson, A. (2021). Scientific Guidance for the submission of dossiers on Food Enzymes. *EFSA Journal*, 19(10), 6851. <https://doi.org/10.2903/j.efsa.2021.6851>
- EFSA CEP Panel (EFSA Panel on Food Contact Materials, Enzymes, Processing Aids), Lambré, C., Barat Baviera, J. M., Bolognesi, C., Cocconcelli, P. S., Crebelli, R., Gott, D. M., Grob, K., Lampi, E., Mengelers, M., Mortensen, A., Rivière, G., Steffensen, I.-L., Tlustos, C., van Loveren, H., Vernis, L., Zorn, H., Roos, Y., Aperi, K., ... Chesson, A. (2023). Food manufacturing processes and technical data used in the exposure assessment of food enzymes. *EFSA Journal*, 21(7), 8094. <https://doi.org/10.2903/j.efsa.2023.8094>
- EFSA GMO Panel (EFSA Panel on Genetically Modified Organisms). (2010). Scientific Opinion on the assessment of allergenicity of GM plants and microorganisms and derived food and feed. *EFSA Journal*, 8(7), 1700. <https://doi.org/10.2903/j.efsa.2010.1700>
- FAO/WHO (Food and Agriculture Organization of the United Nations/World Health Organization). (2006). General specifications and considerations for enzyme preparations used in food processing in Compendium of food additive specifications. 67th meeting. *FAO JECFA Monographs*, 3, 63–67. <https://www.fao.org/3/a-a0675e.pdf>
- Gorbach, S. L. (1978). Risk assessment of recombinant DNA experimentation with *Escherichia coli* K12. *The Journal of Infectious Diseases*, 137, 613–714. <https://doi.org/10.1093/infdis/137.5.704>
- Hayashi, K., Morooka, N., Yamamoto, Y., Fujita, K., Isono, K., Choi, S., Ohtsubo, E., Baba, T., Wanner, B. L., Mori, H., & Horiuchi, T. (2006). Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and W3110. *Molecular Systems Biology*, 2, 2006.0007. <https://doi.org/10.1038/msb4100049>
- Rangkakulnuwat, P., Aluksanasuwan, S., Somsuan, K., Chiangjong, W., Walker, A., Thongklang, N., & Lao-Araya, M. (2023). Trehalose phosphorylase as a novel potential allergen in a case of allergic reaction due to oyster mushroom (*Pleurotus ostreatus*) ingestion. *Journal of Allergy and Clinical Immunology: Global*, 2, 100095.

How to cite this article: EFSA CEP Panel (EFSA Panel on Food Contact Materials, Enzymes and Processing Aids), Lambré, C., Barat Baviera, J. M., Bolognesi, C., Cocconcelli, P. S., Crebelli, R., Gott, D. M., Grob, K., Lampi, E., Mengelers, M., Mortensen, A., Rivière, G., Steffensen, I.-L., Tlustos, C., Van Loveren, H., Vernis, L., Zorn, H., Roos, Y., Gomes, A., ... Chesson, A. (2024). Safety evaluation of the food enzyme cellobiose phosphorylase from the genetically modified *Escherichia coli* strain LE1B109-pPB130. *EFSA Journal*, 22(5), e8774. <https://doi.org/10.2903/j.efsa.2024.8774>