# CORRECTION

# **Open Access**



Correction to: Skin rash following Administration of Apalutamide in Japanese patients with Advanced Prostate Cancer: an integrated analysis of the phase 3 SPARTAN and TITAN studies and a phase 1 open-label study

Hiroji Uemura<sup>1</sup>, Yosuke Koroki<sup>2\*</sup><sup>®</sup>, Yuki Iwaki<sup>3</sup>, Keiichiro Imanaka<sup>4</sup>, Takeshi Kambara<sup>5</sup>, Angela Lopez-Gitlitz<sup>6</sup>, Andressa Smith<sup>6</sup> and Hirotsugu Uemura<sup>7</sup>

## Correction to: BMC Urol (2020) 20:139

https://doi.org/10.1186/s12894-020-00689-0

In the original publication of this article [1] there were several errors in Table 1 related to the values for Disease status (nmCRPC and mCSPC).

In this correction article the correct and incorrect values are shown.

**Table 1** The correct and incorrect values

Incorrect					Correct Disease status, n				
Disease sta									
nmCRPC	55	-	-	55	nmCRPC	34	-	-	34
mCSPC	-	51	-	51	mCSPC	-	28	-	28

Furthermore, the **Time-to-event analyses** section has several errors with the decimal values/rounding of numbers, the incorrect and correct information is shown below

The original article can be found online at https://doi.org/10.1186/s1289 4-020-00689-0.

\*Correspondence: ykoroki@its.jnj.com

<sup>2</sup> Medical Affairs, Janssen Pharmaceutical K.K., Tokyo, Japan

Full list of author information is available at the end of the article



Incorrect	Correct				
66 days	66.0 days				
45 days	45.0 days				
52 days	52.0 days				
38 days	38.0 days				
82 days	82.0 days				
In the global population of the SPARTAN study, skin rash of any grade resolved for <b>81%</b> of the patients within 59.5 days, while the median time to resolution of skin rash of any grade in the TITAN study was 100 days (Sup- plementary Table 2)	In the global population of the SPARTAN study, skin rash of any grade resolved for <b>80.6%</b> of the patients within 59.5 days, while the median time to resolution of skin rash of any grade in the TITAN study was 100.0 days (Supplemen- tary Table 2)				
100 days	100.0 days				
35 days	35.0 days				
37 days	37.0 days				
66 days	66.0 days				

#### Author details

<sup>1</sup> Department of Urology and Renal Transplantation, Yokohama City University Medical Center, Yokohama, Japan. <sup>2</sup> Medical Affairs, Janssen Pharmaceutical K.K., Tokyo, Japan. <sup>3</sup> Clinical Pharmacology, Janssen Pharmaceutical K.K, Tokyo, Japan. <sup>4</sup> Clinical Science, Janssen Pharmaceutical K.K, Tokyo, Japan. <sup>5</sup> Department of Dermatology, Yokohama City University Medical Center, Yokohama,

© The Author(s) 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Japan.<sup>6</sup> Janssen Global Research & Development, Spring House, PA, USA. <sup>7</sup> Department of Urology, Faculty of Medicine, Kindai University, Osaka, Japan.

## Published online: 22 October 2020

#### Reference

 Uemura H, Koroki Y, Iwaki Y, et al. Skin rash following Administration of Apalutamide in Japanese patients with Advanced Prostate Cancer: an integrated analysis of the phase 3 SPARTAN and TITAN studies and a phase 1 open-label study. BMC Urol. 2020;20:139. https://doi. org/10.1186/s12894-020-00689-0.

## **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

#### At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

