

Asymptomatic Essential Thrombocythemia in a Child: A Rare Case Report

majid vafaie*, kaveh jaseb, majid ghanavat, mohamad pedram, tooran rahiminia.

Research Center of Thalassemia & Hemoglobinopathy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Corresponding Author: majid vafaie

Research Center of Thalassemia & Hemoglobinopathy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Email: baran_t_r@yahoo.com

Received: 12, Feb, 2013

Accepted: 2, Mar, 2013

ABSTRACT

Essential thrombocythemia is a rare myeloproliferative disorder in pediatrics. This myeloproliferative disorder is characterized by thrombocytosis and hyperplasia of megakaryocytes in the bone marrow. Other cell lines are not involved. JAK2V617F mutation has been identified in approximately half the patients with this disorder. We describe a 12-year-old boy with essential thrombocythemia. The patient had a persistent thrombocytosis over 600×10^9 /L and the time of diagnosis, his platelet count ranged between 900×10^9 and 2150×10^9 /L. Megakaryocytes in the bone marrow were increased in number. The chromosomal analysis was normal and bcr/abl rearrangement was negative. He remained asymptomatic throughout the follow-up period.

KEY WORDS: Thrombocythemia, Myeloproliferative disorder

INTRODUCTION

Essential thrombocythemia (primary thrombocytosis) is rare in children occurring in approximately 1/1000,000 children.¹⁻⁶ ET is a myeloproliferative disorder characterized by thrombocytosis and hyperplasia of the megakaryocytes in the bone marrow.^{1, 2} Other cell lines are not involved.¹ Reactive (secondary) thrombocytosis is a more common cause of this elevated platelet count among children and the common causes of this entity such as infections, iron deficiency anemia, surgery, cancer, major trauma and post splenectomy need to be excluded prior to considering the diagnosis ET.^{1, 2, 3} ET is primarily a diagnosis of exclusion^{2, 6, 8, 9} and is considered in the presence of a persistent thrombocytosis of greater than 600×10^9 /L in the absence of alternative cause.² These patients may be predisposed to thrombosis or hemorrhage.³ JAK2V617F mutation has been identified in approximately half the patients with this disorder.^{1, 2, 4, 7, 9} It occurs in more than 90% of patients with

polycythemia vera and primary myeloproliferative disease^{1, 2} and 30% of cases of ET.^{4, 7} Familial thrombocythemia, a rare condition,⁶ is similar to ET but differs from ET in that the platelet count is usually significantly lower than ET associated with a lower incidence of hepatomegaly, and with fewer associated thrombotic complications.^{1, 2, 4, 5, 6} No familial case converted to leukemia or myelofibrosis.^{1, 2, 6}

CASE REPORT

We describe a 10 years old male with an incidental finding of an elevated platelet count. This patient had been admitted due to upper respiratory tract infection. His post medical history was unremarkable with a normal birth history and was taking no medications. Immunizations status was up to date. The platelet count at the time of referral was 950×10^9 /L. The physical examination was normal with the exception of mild splenomegaly (just palpable splenomegaly). He had lower leg pain but deep vein thrombosis ruled out with color

duplex ultrasonography. His parent platelet count was normal and there was no family history of thrombocytosis. His white blood cell count and differential count were unremarkable, and platelet size was not enlarged. Hb, ESR and C-reactive protein were all within normal ranges. Bone marrow examination was normocellular with relatively normal numbers and maturation of erythroid and myeloid lineages but show clusters of large megakaryocytes with increased hyperlobulated nuclei (Fig-1). Myeloblasts are about 5% of all nucleated cells. Bone marrow biopsy show remarkable bony trabeculae and marrow spaces contain hematopoietic elements with predominance of large hyperlobulated megakaryocytes. Reticulin staining was negative for myelofibrosis. The overall histological features suggest chronic myeloproliferative disorder without myelofibrosis suggestive of ET. Cytogenetic showed normal 46, XY karyotype. No bcr/abl chimaeric transcript was demonstrated by reverse transcription polymerase chain reaction. JAK2-V617F mutation was negative. Evaluation of the MPL W515L/K gene was negative in the patient. He received low dose aspirin (80mg/day). Her platelet count ranging from 303×10^9 to 2131×10^9 /L. He was admitted to hospital because of anxiety and right lower leg pain. At this time his platelet count was 2131×10^9 /L and we started hydroxyurea 1000mg and anagrelide 2mg for him. The mean platelet count was 1029×10^9 /L. The platelet count has shown a trend of gradual fall over the past year and the latest platelet count was 580×10^9 /L. He is asymptomatic now.




Figure 1 Bone marrow aspiration showed megakaryocytic hyperplasia (magnification $\times 40$)

DISCUSSION

Thrombocytosis is relatively common in young children occurring in 3% to 13% of children.^{5, 8} Extreme thrombocytosis (platelet $>1000,000$) is uncommon, occurring in less than 2% of children.⁸ Almost all reported case of thrombocytosis in children are secondary to infection, iron deficiency, surgery, cancer, postsplenectomy, chronic inflammation, collagen vascular and renal disease, langerhans cell histiocytosis, hemolytic anemia, Kawasaki disease drugs.^{1, 2, 3, 4, 8} Essential thrombocythemia extremely rare in childhood,¹⁻⁸ with an estimated incidence of 0.09 case per million for children younger than 14 years 60 times lower than in adults.⁴ Based on the criteria of the PVSG approximately 75 children have been reported with essential thrombocythemia between 1966 and 2000.² Edwin B. Robins described a 2 years old male with an incidental finding of an elevated platelet count discovered on health maintenance screening. The platelet count at the time of referral was 900×10^9 /L, and his physical examination was remarkable for bony abnormalities of the right upper extremity and hand.⁴ Maria Luigia Randi et al., reported five children with essential thrombocythemia. Moreover only one case of them was infant.⁵ Abeer Abd El-Moneim et al., also reported 12 cases of children with essential thrombocythemia.⁹ Dror et al., from the Hospital for sick children in Toronto reviewed the clinical course of 36 children with ET. Forty eight children with ET were described in the medical literature prior to 2006, with the age ranging from 6 weeks to 18 years.⁶ ET in children is not always a benign entity, in adults with ET, thrombosis is the most common and hazardous complication, occurring in about 40% of patients. In contrast, most children with ET seen to have few major thrombotic complication.⁵ Out of the 42 children with ET reviewed by Dror Four children experienced severe bleeding episodes, six children had mild hemorrhage, four children suffered from severe arterial thrombosis, and in two cases venous thrombosis were documented. Two children had headache, erythromelalgia and limb paresthesia. Splenomegaly was found in 44%, hepatomegaly in 22% of the patients.⁶ Children with thrombocytosis may develop unspecific symptoms that are potentially related to thrombocytosis. The clinical course is heterogeneous.^{8, 9} Some patients

may recover spontaneously.⁹ Due to the lower incidence of ET in children optimal first line therapy remains unclear in this age group.⁶ Furthermore treatment of asymptomatic patients remains problematic. Hydroxyurea has been used successfully in patients with ET.^{4, 6, 9, 10} Anagrelide has a high specificity toward megakaryocytes and effectively controls extreme thrombocytosis in adults and children.^{1, 2, 3, 4, 6} The use of anagrelide to treat children with ET has been described since 1991 and long term use of anagrelide has been studied in adults with a 93% response rate in a prospective cohort of 335 adults, and in children.⁴ Indication for treatment included thrombotic events, major bleeding, minor bleeding, or merely high platelet counts.⁶ Antiplatelet agents such as low-dose acetylsalicylic acid or dipyridamol have proved efficacious in preventing recurrence of thrombosis in the acral, coronary and cerebral arterial circulations in patients with ET, and have been recommended as initial therapy in patients with minor benign problems.⁶

REFERENCES

1. Cooper TM, Hasle, Smith FO, In Pizzo PA, Poplack DG: Principle and practice of pediatric oncology. Philadelphia :Lippincott Williams &Wilkins;2011.pp.566-610
2. Lanzkowsky P.Disorders of platelets; Manual of pediatric hematology &oncology, 5th edition .Amsterdam: Elsevier; 2011. P.321-377
3. Wilson DB.Acquired platelet defects .In :Nathan D ,Oski FA ,Hematology of infancy and childhood .Philadelphia :WB Sanders;7th edition .2009 . p.1553-1590
4. Robins EB.Niazi M, Essential thrombocythemia in a child with elevated thrombopoietin concentrations and skeletal anomalies. *Pediatric Blood Cancer* 2008; 50:859-861
5. Randi ML;Putti MC,Fabris L,Zanesco L and Girolami A. Features of essential thrombocythemia in childhood :A study of five children .*British Journal of Hematology* , 2000,108,86-89
6. Dror Y,Blanchelette VS .Essential thrombocythemia in children :Review.*British Journal of Hematology* ,1999,107,691-698
7. Randi ML,Putti MC ,Scapin M,Pacquola E,Tucci F,Micaillizzi C and et al. Pediatric patients with essential thrombocythemia are mostly polyclonal and V617FJAK2 negative :Brief report .*Blood* .2006,108:3600-3602
8. Mantadkis A,Tsalkidis A ,Chatzimichael A.Thrombocytosis in childhood :Review article .*Indian pediatrics* .volume 45,August 17 ,2008,P:669-677
9. El-Moneim AA,Kartz CP ,Boll S, Rister M, Pahl HL, Niemeyer CM. Essential versus reactive thrombocythemia in children :Retrospective analysis of 12 cases.*Pediatr Blood Cancer* ,2007;49:52-55
10. Cortelazzo S, Finazzi G,Ruger M,Vestri O,Galli M ,Rodeghiero F and et al .Hydroxyurea for patients with essential thrombocythemia and a high risk of thrombosis .*The New England Journal of medicine*.April 27,1995