

Retention of aberrant cortisol secretion in a patient with bilateral macronodular adrenal hyperplasia after unilateral adrenalectomy

This article was published in the following Dove Medical Press journal:
Therapeutics and Clinical Risk Management

Sho Tanaka¹
Midori Fujishiro²
Yoshihiro Nakamura¹
Yoshinari Hatanaka¹
Masanori Abe¹

¹Division of Nephrology, Hypertension and Endocrinology, Department of Internal Medicine, Nihon University School of Medicine, Tokyo, Japan; ²Division of Diabetes and Metabolic Diseases, Nihon University School of Medicine, Tokyo, Japan

Abstract: Aberrant cortisol secretion responses after exogenous stimuli such as upright posture, eating a mixed meal or receiving agents influencing aberrant G-protein-coupled receptors in adrenal glands, are often observed in patients with bilateral macronodular adrenal hyperplasia (BMAH). However, little is known about whether this aberrant response is retained after unilateral adrenalectomy. Here, we describe a 61-year-old postmenopausal Japanese woman with unsatisfactorily controlled hypertension who was referred to us for further investigation due to her pre-obesity characteristics (body mass index 28.4 kg/m²). Cushing's signs and serum cortisol at 16.2 µg/dL with undetectable adrenocorticotropic hormone indicated adrenal Cushing's syndrome. Adrenal imaging revealed bilaterally enlarged adrenal glands with 131-I adosterol uptake; hence, BMAH was diagnosed. Preoperatively, in vivo screening for aberrant adrenal receptors revealed an aberrant response of cortisol secretion on metoclopramide challenge. The patient underwent unilateral adrenalectomy; thereafter, glucocorticoid replacement therapy was reduced to hydrocortisone 15 mg/day at postoperative day 6. Fasting morning serum cortisol level measured at postoperative day 8 was 2.96 µg/dL, suggesting adrenal insufficiency. However, following metoclopramide administration serum cortisol level rose to 19.7 µg/dL, indicating potential efficient adrenal function. Aberrant cortisol secretory capacity was thus preserved in BMAH, even in a state of adrenal insufficiency after unilateral adrenalectomy. Caution should be exercised when assessing the hypothalamus-pituitary-adrenal axis, because in this patient, a high cortisol level did not guarantee appropriate adrenal function when the patient was challenged by exogenous stimuli.

Keywords: adrenal glands, Cushing's syndrome, glucocorticoids, hydrocortisone, metoclopramide, receptors, G-protein-coupled

Introduction

Bilateral macronodular adrenal hyperplasia (BMAH) is a subtype of adrenal Cushing's syndrome, and is characterized by bilaterally enlarged adrenal glands consisting of non-pigmented macronodules. The molecular mechanisms of cortisol secretion in BMAH, which does not depend on circulating adrenocorticotropic hormone (ACTH) released by the pituitary gland, have become clear in the last two decades. It is widely recognized that cAMP signaling evoked by activation of aberrant G-protein-coupled receptors (GPCRs) in adrenal glands largely contributes to steroid genesis.¹ Clinically, this is confirmed by aberrant responses of serum cortisol level on challenge by endogenous stimuli such as upright posture, eating a mixed meal or exposure to certain chemicals.¹ Previous investigations have demonstrated that metoclopramide, an agent frequently used for alleviating gastrointestinal symptoms, stimulates adrenal serotonin type 4

Correspondence: Sho Tanaka
Division of Nephrology, Hypertension and Endocrinology, Department of Internal Medicine, Nihon University School of Medicine, 30-1 Kamicho, Oyaguchi, Itabashi-ku, Tokyo 173-8610, Japan
Tel +81 33 972 8111
Fax +81 33 972 8311
Email tanaka.sho@nihon-u.ac.jp

receptors in some cases of BMAH, leading to excess cortisol in vivo and in vitro.^{2–8} There are several different treatment modalities for patients with BMAH, including unilateral adrenalectomy.⁹ In patients who underwent this operation, assessing residual cortisol secretion capacity is clinically important in order to avoid unnecessary steroid replacement therapy while reducing the risk of adrenal insufficiency or crisis. While more than 20 cases of BMAH with aberrant responses against metoclopramide have been reported in the literature, the impact of metoclopramide on cortisol secretion from the remaining adrenal gland in patients after unilateral adrenalectomy remains unknown.^{2,3,5,7,8} Herein, we report a Japanese woman with BMAH in whom aberrant serum cortisol levels on metoclopramide use were seen despite adrenal insufficiency after surgery.

Case description

A 61-year-old postmenopausal Japanese woman was referred to us for further investigation of her pre-obese state. She had had hypertension, dyslipidemia and osteoporosis for 10 years. Hypertension was treated with telmisartan 80 mg/day, and amlodipine 7.5 mg/day but was not satisfactorily controlled. Home-monitored blood pressure level remained 130–160/70–90 mmHg. Her family history included hypertension in an elderly sister but was otherwise unremarkable. The patient habitually smoked a pack of cigarettes per day and consumed 30 g alcohol daily.

On physical examination, pre-obesity was observed: height 147 cm, weight 61.3 kg and BMI 28.4 kg/m². Additionally, centripetal obesity, facial plethora, skin atrophy with easy bruising, proximal lower limb weakness and mild moon face were apparent. Randomly measured serum cortisol at 16.2 µg/dL (reference range, 6.24–18.0) with undetectable ACTH at <2 pg/mL (reference range, 7.2–63.3) indicated adrenal Cushing's syndrome. Computed tomography and single photon emission computed tomography using ¹³¹I adosterol revealed bilateral nodules in the adrenal glands. Maximum nodule size in the left-side gland was 34 mm with strong uptake, and 31 mm with relatively weak uptake in the right-side gland (Figure 1). Circadian variation of serum cortisol and ACTH levels was abnormal; Cortisol levels were 11.2, 11.4, 14.5 and 10.8 µg/dL at 0.00am, 6.00am, 0.00pm and 6.00pm, respectively, and ACTH was undetectable throughout the day. Urinary free cortisol excretion was high at 380 µg/day (reference range, 11.2–80.3). Reduction of serum cortisol level was not seen in overnight dexamethasone suppression tests; morning serum cortisol levels following 1 or 8 mg dexamethasone administration were 11.3 and 15.3 µg/




Figure 1 Imaging of adrenal glands.

Notes: The arrows indicate the nodule in the right (yellow) and left (red) adrenal gland. (A) Image of CT. (B) Merged image of CT and ¹³¹I adosterol SPECT. (C) Image of ¹³¹I adosterol SPECT.

Abbreviations: CT, computed tomography; SPECT, single photon emission computed tomography.

dL, respectively. Responses of serum cortisol after eating a mixed meal, on intravenously administered glucagon 1 mg, or orally administered metoclopramide 10 mg were investigated. Additionally, responses against ACTH and gonadotropin-releasing hormone (GnRH) were assessed using intravenous synacthen 250 µg and gonadorelin 0.1 mg, respectively. Responses on transitioning to an upright posture could not be investigated due to the patient's refusal because of knee pain and lower limb muscle weakness. Results are shown in Figure 2. Serum cortisol levels increased only in response to




Figure 2 Aberrant responses of cortisol secretion preoperatively. **Notes:** Vertical axis indicates serum cortisol level. Horizontal axis indicates time course after exogenous stimuli. **Abbreviations:** ACTH, adrenocorticotropic hormone; GnRH, gonadotropin-releasing hormone.

ACTH and metoclopramide administration. The increased cortisol level following metoclopramide administration was 239% from a baseline of 11.2 to a peak of 26.8 µg/dL.

Based on the above data, the patient was diagnosed as having BMAH and underwent left adrenalectomy. Surgery was successful, and the dose of postoperative glucocorticoid replacement therapy was uneventfully reduced to hydrocortisone 15 mg/day (10 mg after breakfast and 5 mg after dinner) at day 6. On the morning of postoperative day 8, prior to use of hydrocortisone in the morning, the fasting serum cortisol response against oral metoclopramide 10 mg was investigated once again. A baseline morning cortisol level of 2.96 µg/dL strongly suggested adrenal insufficiency, but the peak level following administration of metoclopramide was at 19.7 µg/dL (representing an increase of 666%, Figure 3). On receiving hydrocortisone 15 mg/day, the patient showed no signs of adrenal insufficiency thereafter, and was therefore uneventfully discharged at postoperative day 10.

Serum cortisol levels prior to hydrocortisone in the morning were monitored on an outpatient basis on postoperative day 72 and 109, when the patient was receiving hydrocortisone 12.5 mg/day (10 mg after breakfast and 2.5 mg after dinner). Markedly decreased morning serum cortisol levels were repeatedly confirmed, and adrenal insufficiency was strongly suggested even at day 109; thus, steroid coverage could not be withdrawn (Figure 4). Undetectable ACTH




Figure 3 Cortisol secretion on metoclopramide challenge at postoperative day 8. **Notes:** Vertical axis indicates serum cortisol level. Horizontal axis indicates time course after metoclopramide administration. Numerical values are also shown with trajectory.

at <2 pg/mL was sustained throughout the observation period, which also suggested an unrestored hypothalamus-pituitary-adrenal (HPA) axis. Meanwhile, the patient’s weight was successfully reduced to 53.0 kg with a BMI of 24.5 kg/m², and hypertension is currently controlled at 120–140/70–90 mmHg with amlodipine 5 mg/day alone.

Discussion

BMAH is a rare subtype of adrenal Cushing’s syndrome, and genetic involvement is suggested because of bilateral




Figure 4 Time course of morning serum cortisol before and after left adrenalectomy. **Notes:** Vertical axis indicates morning serum cortisol level. Horizontal axis indicates time course before and after left adrenalectomy (week 0). Three data points collected before surgery, and data collected at postoperative days 8, 72 and 109 are shown. Regimen of steroid coverage at day 8 was hydrocortisone 15 mg/day (10 and 5 mg after breakfast and dinner, respectively). Regimen of steroid coverage at days 72 and 109 was hydrocortisone 12.5 mg/day, 10 and 2.5 mg after breakfast and dinner, respectively. Numerical values are also shown with trajectory.

involvement and reports of familial cases. A major genetic characteristic is the presence of inactivating mutations of *ARMC5*, a potential tumor suppressor gene; biallelic alterations consisting of an additive pathogenic somatic mutation and a germline mutation inactivating *ARMC5* result in nodule formation in the adrenal glands of BMAH patients.¹⁰ Clinically, Cushing's syndrome of BMAH type is characterized by marked bilateral enlargement of the adrenal glands and aberrant cortisol secretion responses on challenge with exogenous stimuli.¹ Previously, several investigators reported that this aberrant response is associated with an upright posture, eating a mixed meal, ACTH, GnRH, thyrotropin-releasing hormone, glucagon, vasopressin and serotonin type 4 receptor agonists including metoclopramide which was first described by Lacroix et al in 1999.²⁻⁸ The present case we describe here is a patient with BMAH associated with an aberrant response of cortisol secretion on metoclopramide challenge. Clinical profiles such as age, nodule sizes and urinary cortisol excretion in the present case seem similar to the previously reported BMAH cases with aberrant responses to metoclopramide administration (Table 1).^{2,3,5,7,8} However, this patient highlights two clinically important issues. First, metoclopramide-stimulated aberrant cortisol secretion responses by the remaining adrenal gland are retained even under conditions of adrenal insufficiency in the early postoperative period. Second, the impact of metoclopramide use in such cases has a substantial ability to conceal adrenal insufficiency as assessed by morning, or randomly timed, serum cortisol level determinations.

In the present case, adrenal insufficiency after left adrenalectomy was apparent, and did not recover over the observation period; thus, morning serum cortisol levels were 2.96, 4.13 and 2.12 $\mu\text{g/dL}$ on postoperative day 8, 72 and 109. This is sufficiently low to predict adrenal insufficiency because a recent retrospective cohort study reported that morning serum cortisol level was correlated with peak serum cortisol level on an insulin hypoglycemia test, with morning cortisol level below 4 $\mu\text{g/dL}$ or above 17 $\mu\text{g/dL}$ being highly predictive of insufficient and sufficient adrenal function, respectively.¹¹ Furthermore, another retrospective cohort study reported an association between randomly sampled serum cortisol levels and the results of ACTH testing. In that study, a cortisol level below 5.1 $\mu\text{g/dL}$ predicted a 30 minutes serum cortisol level below 20 $\mu\text{g/dL}$ during the ACTH test, indicating insufficient adrenal function, with 100% specificity and 35% sensitivity.¹² In contrast, a value exceeding >15.2 $\mu\text{g/dL}$ guaranteed a 30 minutes value exceeding 20 $\mu\text{g/dL}$ during the ACTH test, indicating adequate adrenal function.¹² Therefore, it seems

highly likely that our patient had adrenal insufficiency sustained until at least postoperative day 109.

The postoperative course of adrenal function observed in this present case is not a chance event, because a previous retrospective study reported that postoperative adrenal insufficiency following unilateral adrenalectomy in BMAH patients was observed in 40% of patients (6 of 15 enrolled).¹³ The report also remarked that adrenal insufficiency and the following latent adrenal insufficiency persisted for two years or more.¹³ It was also mentioned that the 250 μg synacthen test was unreliable for diagnosis of adrenal insufficiency in such situations because the cortisol secretory response to a challenge of 250 μg synacthen could occur even under conditions of adrenal insufficiency after surgery.¹³ Similarly, this present case indicates that the remaining gland still did have substantial cortisol secretory capacity when stimulated via an aberrant GPCRs, even though the patient was ordinarily adrenal-insufficient.

The 666% increase in the serum cortisol level following metoclopramide stimulation observed after surgery is of interest. Although the baseline serum cortisol level of 2.96 $\mu\text{g/dL}$ was strongly suggestive of adrenal insufficiency, the peak value of 19.7 $\mu\text{g/dL}$ would indicate sufficient adrenal function when measured either in the morning or randomly.^{11,12} In the clinical setting, morning or random serum cortisol measurement is frequently used to assess HPA axis functionality because it is easy to perform. However, the marked fluctuation of serum cortisol level observed in this present case suggests that external stimuli associated with aberrant cortisol secretion can conceal underlying adrenal insufficiency and mislead clinicians assessing the HPA axis after unilateral adrenalectomy in BMAH patients. Because metoclopramide is frequently used, also after surgery, caution is required to avoid the risk of inappropriate withdrawal of steroid coverage. Moreover, habitual use of metoclopramide might suppress the hypothalamus and pituitary via negative feedback due to cortisol excess, and lead to delayed recovery of the HPA axis. Similarly, this risk should be considered in all patients with BMAH receiving unilateral adrenalectomy, because not only metoclopramide but also upright posture or eating a mixed meal, always happening in daily life, were also reported to induce cortisol secretion.^{3,7,8} To conclude, detailed surveillance of aberrant cortisol secretion responses on challenge with exogenous stimuli influencing aberrant GPCRs is clinically important in BMAH patients.

This report has certain limitations. First, it is an observational case report of a single patient, and there are several

Table I Review of BMAH cases with paradoxical cortisol secretion response on metoclopramide challenge

Case	Author	Sex	Age (years)	BMI (kg/m ²)	Function	Maleifestation	Nodule size (mm)		U-Cor (µg/day)	Degree of increase (%)
							Right	Left		
1	Libé R	Male	53	31.2	CS	Cushing's sign	35	45	293	133
2		Female	34	30.0	CS	Cushing's sign	45	30	373	180
3		Female	70	25.0	CS	Incidentaloma	50	30	122	204
4		Female	45	21.3	SCS	Incidentaloma	60	30	44	438
5		Female	37	30.5	SCS	Incidentaloma	33	15	26	185
6		Male	59	25.6	SCS	Hypertension	30	60	31	426
7		Female	66	27.8	SCS	Incidentaloma	14	30	29	209
8		Female	35	27.4	SCS	Incidentaloma	26	27	18	180
9		Female	67	28.6	SCS	Incidentaloma	14	40	56	209
10		Female	54	28.2	SCS	Incidentaloma	18	44	36	137
11		Female	53	26.7	SCS	Incidentaloma	35	27	83	310
12		Male	74	23.3	SCS	Incidentaloma	20	25	85	160
13		Male	51	20.6	SCS	Incidentaloma	27	40	63	206
14		Male	52	27.0	SCS	Incidentaloma	55	46	50	138
15		Male	51	24.0	SCS	Incidentaloma	40	37	47	183
16		Female	43	22.3	SCS	Incidentaloma	45	15	46	154
17		Female	55	33.6	SCS	Hypertension	30	24	26	183
18	Bourdeau I	Female	51	NA	SCS	Hypertension, weight gain	44	50	NA	145
19		Male	61	NA	SCS	NA	61	73	NA	191
20		Male	53	25.3	SCS	Hypertension	35	40	NA	300
21	Vezzosi D	Female	56	NA	CS	Hypertension, centripetal obesity, facial plethora, dorsal fat pad supraclavicular fat pad, glucose intolerance, mild hirsutism	NA	NA	34	270
22		Female	54	NA	CS	Hypertension, centripetal obesity, facial plethora, easy bruising, diabetes mellitus	NA	NA	50	301
23	Feelders RA	Female	60	24.4	CS	Centripetal obesity, facial plethora, muscle weakness, easy bruising depression, skin atrophy	35	50	199–228	289
24		Female	40	32.8	CS	Hypertension, centripetal obesity, facial plethora, weight gain, muscle weakness, easy bruising, dorsal fat pad, supraclavicular fat pad, depression	50	30	326–652	195
25	Lacroix A	Female	63	28.9	CS	Hypertension, weight gain, muscle weakness, numbness, hot flushes, decrease in concentration and memory	40	40	279	257

Note: Clinical features of cases with BMAH reported to be associated with paradoxical response of cortisol secretion on metoclopramide challenge are shown together with the percentage increase of cortisol level.

Abbreviations: BMAH, bilateral macronodular adrenal hyperplasia; BMI, body mass index; CS, Cushing's syndrome; NA, not applicable; SCS, subclinical Cushing's syndrome; U-Cor, urinary cortisol excretion.

unavailable data, such as results of a vasopressin test, a postoperative synacthen test and postoperative urinary free cortisol excretion. Further cases with fully-assessed pre- and postoperative aberrant responses to exogenous stimuli must be accumulated to assess cortisol secretory

capacity after surgery in BMAH. Second, cortisol levels following metoclopramide stimulation after surgery were possibly modified by hydrocortisone use, despite the small doses applied (15 mg/day), so underestimates cannot be excluded.

Conclusion

The aberrant cortisol secretion response after exogenous stimulation influencing aberrant G-protein-coupled receptors was found to be retained in a patient with BMAH, even in a state of adrenal insufficiency after unilateral adrenalectomy. Thus, a high serum cortisol level when the patient receives such exogenous stimuli does not guarantee efficient adrenal function. Caution is thus required for assessing the actual status of the HPA axis.

Abbreviations

ACTH, adrenocorticotrophic hormone; BMAH, bilateral macronodular hyperplasia; BMI, body mass index; GnRH, gonadotropin-releasing hormone; GPCRs, G-protein-coupled receptors; HPA, hypothalamus-pituitary-adrenal.

Ethics and consent for publication

The patient described in this study provided permission to publish data and accompanying images, and written informed consent was obtained. A formal ethical review by an institutional review board was not required because this is a case report.

Acknowledgment

This report was not funded by any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions

ST designed the study, contributed to collect, analyze and interpret data, and wrote the initial draft of the manuscript. MF contributed to interpret the data, and assisted in the preparation of the manuscript. All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. El Ghorayeb N, Bourdeau I, Lacroix A. Multiple aberrant hormone receptors in Cushing's syndrome. *Eur J Endocrinol.* 2015;173(4): M45–M60.
2. Lacroix A, Hamet P, Boutin JM. Leuprolide acetate therapy in luteinizing hormone – dependent Cushing's syndrome. *N Engl J Med.* 1999; 341(21):1577–1581.
3. Bourdeau I, D'Amour P, Hamet P, Boutin JM, Lacroix A. Aberrant membrane hormone receptors in incidentally discovered bilateral macronodular adrenal hyperplasia with subclinical Cushing's syndrome. *J Clin Endocrinol Metab.* 2001;86(11):5534–5540.
4. Cartier D, Lihmann I, Parmentier F, et al. Overexpression of serotonin4 receptors in cisapride-responsive adrenocorticotropin-independent bilateral macronodular adrenal hyperplasia causing Cushing's syndrome. *J Clin Endocrinol Metab.* 2003;88(1):248–254.
5. Feelders RA, Lamberts SW, Hofland LJ, et al. Luteinizing hormone (LH)-responsive Cushing's syndrome: the demonstration of LH receptor messenger ribonucleic acid in hyperplastic adrenal cells, which respond to chorionic gonadotropin and serotonin agonists in vitro. *J Clin Endocrinol Metab.* 2003;88(1):230–237.
6. Louiset E, Contesse V, Groussin L, et al. Expression of serotonin7 receptor and coupling of ectopic receptors to protein kinase A and ionic currents in adrenocorticotropin-independent macronodular adrenal hyperplasia causing Cushing's syndrome. *J Clin Endocrinol Metab.* 2006;91(11):4578–4586.
7. Vezzosi D, Cartier D, Régnier C, et al. Familial adrenocorticotropin-independent macronodular adrenal hyperplasia with aberrant serotonin and vasopressin adrenal receptors. *Eur J Endocrinol.* 2007;156(1): 21–31.
8. Libé R, Coste J, Guignat L, et al. Aberrant cortisol regulations in bilateral macronodular adrenal hyperplasia: a frequent finding in a prospective study of 32 patients with overt or subclinical Cushing's syndrome. *Eur J Endocrinol.* 2010;163(1):129–138.
9. Lacroix A. ACTH-independent macronodular adrenal hyperplasia. *Best Pract Res Clin Endocrinol Metab.* 2009;23(2):245–259.
10. Assié G, Libé R, Espiard S, et al. Armc5 mutations in macronodular adrenal hyperplasia with Cushing's syndrome. *N Engl J Med.* 2013; 369(22):2105–2114.
11. Erturk E, Jaffe CA, Barkan AL. Evaluation of the integrity of the hypothalamic-pituitary-adrenal axis by insulin hypoglycemia test. *J Clin Endocrinol Metab.* 1998;83(7):2350–2354.
12. Kadiyala R, Kamath C, Baglioni P, Geen J, Okosieme OE. Can a random serum cortisol reduce the need for short synacthen tests in acute medical admissions? *Ann Clin Biochem.* 2010;47(Pt 4):378–380.
13. Debillon E, Velayoudom-Cephise FL, Salenave S, et al. Unilateral adrenalectomy as a first-line treatment of Cushing's syndrome in patients with primary bilateral macronodular adrenal hyperplasia. *J Clin Endocrinol Metab.* 2015;100(12):4417–4424.

Therapeutics and Clinical Risk Management

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS,

Submit your manuscript here: <http://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>

Dovepress

EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.