

Suppression of NF- κ B activity by mutant I κ B α : A molecular target for radiosensitization of adenoid cystic carcinoma

ZHE LIU^{1*}, SHENGYUN HUANG^{1*}, SHIZHOU ZHANG¹, JIAWEN SI¹, QIANG WANG¹,
QIANGXIU WANG², WENLI MU³, JUNQING HAN⁴ and DONGSHENG ZHANG¹

Departments of ¹Oral and Maxillofacial Surgery and ²Pathology; ³Medical Research Center and ⁴Cancer Center, Provincial Hospital Affiliated to Shandong University, Jinan 250021, P.R. China

Received October 21, 2012; Accepted January 14, 2013

DOI: 10.3892/ol.2013.1153

Abstract. The constitutive activation of the nuclear factor κ B (NF- κ B) signaling pathway is involved in oncogenesis, invasive growth, metastasis and induced resistance to radiation and chemotherapy. Selective inhibition of the NF- κ B signaling pathway, either by a mutant inhibitor or pharmacological agents, improves the therapeutic efficiency of irradiation. In the present study, the changes in NF- κ B expression and the rate of apoptosis were investigated following irradiation of cells of an adenoid cystic carcinoma cell line (ACC-M) in which NF- κ B expression had been inhibited by transient transfection with a mutant I κ B α plasmid. ACC-M cells were transiently transfected with the mutant I κ B α plasmid using Lipofectamine and the expression of this mutant I κ B α gene was verified. The presence of the mutant I κ B α gene alone did not result in a reduction in cell proliferation. Furthermore, a significant inhibition of translocation and synthesis of NF- κ B protein in the transfected cells was observed after irradiation. NF- κ B protein was activated by different doses of irradiation in a dose- and time-dependent manner with concordant changes in the radiosensitivity of ACC-M cells. We conclude that the mutant I κ B α gene selectively inhibited the NF- κ B pathway, which may be a promising method to improve the radiosensitivity of adenoid cystic carcinomas.

Introduction

Adenoid cystic carcinoma (ACC) is one of the most common subtypes of malignant tumors occurring in the salivary glands. It is characterized by a high rate of recurrence, a strong

tendency of perineural invasion and an early development of hematogenous metastasis. ACC patients possessing prognostic factors, including lymph node positivity, a solid histological subtype or perineural invasion with involvement of a named nerve, suffer poor prognostic survival prospects (1). The rates of perineural invasion and distant metastases after 10 years, regardless of aggressive surgery, have been demonstrated to be 50 and 39%, respectively (2). Postoperative radiotherapy is thus recommended for its additive improvement in the control of local and regional recurrence compared with surgery alone (1,3). However, considering the relatively low radiosensitivity and the aggressive growth pattern of ACC, methods to improve the therapeutic efficiency have potential for further investigation.

Nuclear factor κ B (NF- κ B) was first discovered in 1986 as a eukaryotic transcription factor that bound to the κ immunoglobulin light chain enhancer in the nuclei of the B lymphoid cell lineage, and is presently considered to exist in the majority of cell types (4,5). Homo- and heterodimeric complexes of its five members, p50, p52, p65/RelA, c-Rel and Rel-B, are sequestered inactive in the cytoplasm preformed and bound to the inhibitory subunits of the I κ B family (I κ B α , I κ B β , I κ B ϵ , Bcl-3, p100 and p105) (6). In the canonical NF- κ B activation pathway, the activation of NF- κ B depends on stimuli including bacterial and viral products, proinflammatory cytokines and ionizing radiation, all of which intrigue phosphorylation of I κ B α by I κ B kinase (I κ K)-induced ubiquitination and subsequent proteolysis by the 26 S proteasome. The phosphorylation of I κ B α at specific serine (Ser) residues liberates the captive cytoplasmic NF- κ B for translocation to the nucleus and initiation of target genetic transcription (7). The active NF- κ B transcription factor regulates the expression of >150 downstream genes. The majority of proteins encoded by NF- κ B target genes participate in a wide variety of physiological processes, including embryonic development, lymphoid differentiation, immune and inflammatory responses and apoptotic resistance to radio- and chemotherapy, as well as being involved in oncogenesis and tumor proliferation (8).

The aberrant activation of NF- κ B is characteristic of numerous solid tumor types, including breast epithelial tumors, pancreatic adenocarcinoma and bladder, prostate and non-small cell lung cancer (9). High levels of NF- κ B activity contribute to a negative prognosis by playing a key

Correspondence to: Professor Dongsheng Zhang, Department of Oral and Maxillofacial Surgery, Provincial Hospital Affiliated to Shandong University, 324 Jingwu Road, Shandong, Jinan 250021, P.R. China

E-mail: ds63zhang@sdu.edu.cn

*Contributed equally

Key words: nuclear factor κ B, radiosensitization, apoptosis, transfection, ACC-M cells, pathway analysis

role in the induction of anti-apoptosis, the acceleration of cell cycle progression, increased resistance to radiation and chemotherapy, as well as in elevated aggressive growth and metastatic frequency in malignant cancer (10). It is widely accepted that constitutively activating NF- κ B exerts a negative impact on the radiosensitivity of different cancer cell lines. Sandur *et al* demonstrated that suppression of the NF- κ B pathway via curcumin-induced inhibition sensitized colorectal cancer cells to radiation by downregulating the phosphorylation and degradation of I κ B α , inhibition of I κ K activity and inhibition of Akt phosphorylation (10). By transfecting a gene encoding mutated I κ B α that is not able to be phosphorylated and thereby inhibits the activation of the NF- κ B pathway, similar findings have been demonstrated in that suppressive NF- κ B radiosensitizes prostate cancer cells (6). However, the role of the NF- κ B pathway in regulating the radiosensitivity remains controversial in different diseases. Jung *et al* demonstrated that by transfecting a phosphorylation-defective I κ B α gene into cells derived from a patient with ataxia telangiectasia (AT) group D, the constitutive activation of NF- κ B was suppressed and the previous radiation hypersensitivity in AT was restored to the normal level (11). Additionally, to the best of our knowledge, the impact of this mutant I κ B α gene on the expression of NF- κ B and subsequent changes in radiosensitivity in ACC cells had not yet been studied.

In the present study, we investigated the role of NF- κ B in regulating the radiosensitivity of adenoid cystic carcinoma cells (ACC-M) *in vitro*. To achieve a conclusion, we transiently transfected a plasmid encoding a Flag-tagged phosphorylation defective mutant of I κ B α (S32, 36A) (pBabe-SR-I κ B α) into ACC-M cells, thereby suppressing the activity of the NF- κ B pathway. While analyzing the results, we observed that suppressing the NF- κ B pathway by transfecting ACC-M cells with mutant I κ B α , as compared with cells with a control pBabe plasmid transfection, leads to increased radiosensitivity. Meanwhile, intragroup data analysis of the SR-I κ B α group demonstrated that different doses of irradiation induced the expression of NF- κ B in a dose- and time-dependent manner, with corresponding changes in the radiosensitivity of ACC-M cells.

Materials and methods

Chemicals and reagents. The pBabe-SR-I κ B α and control pBabe plasmids were provided by Professor Wantao Chen (Department of Oral and Maxillofacial Surgery, Ninth People's Hospital, College of Stomatology, Shanghai Jiao Tong University, Shanghai, China). Lipofectamine was purchased from Invitrogen Life Technologies (Carlsbad, CA, USA).

An MTT and an Annexin V-FLUOS staining kit were purchased from Roche Diagnostics (Indianapolis, IN, USA). Rabbit anti-human NF- κ B, goat anti-rabbit secondary antibodies, anti- β -actin and horseradish peroxidase (HRP)-conjugated goat anti-rabbit immunoglobulin were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA). Anti-I κ B α antibody was purchased from Cell Signaling Technology, Inc. (Denver, MA, USA).

Cell culture. The human salivary adenoid cystic carcinoma cell line (ACC-M) was provided by Professor Wantao Chen.

ACC-M cells were cultured in RPMI-1640 medium supplemented with 10% filtered fetal bovine serum (FBS; Hyclone, Logan, UT, USA) and 100 U/ml penicillin/streptomycin. The cultures were incubated at 37°C in a humidified 5% CO₂ incubator.

Plasmid transfection. The site-specific, signal-induced degradation of I κ B α depends on phosphorylation at Ser 32 and 36. Therefore, the pBabe-SR-I κ B α plasmid that consisted of a double point mutation (Ser to Lactamine) was thus resistant to phosphorylation (12). The mutant and control plasmids were transiently transfected into ACC-M cells with the use of Lipofectamine, according to the manufacturer's instructions. In brief, ACC-M cells were removed by trypsin/EDTA treatment and seeded at a density of 2x10⁵ cells/ml in 6-cm culture dishes. Cells were grown to 90% confluence and subjected to 24-h synchronization in serum-free medium. ACC-M cells were transfected with 4 μ g of the pBabe-SR-I κ B α or control pBabe plasmid per dish with the use of Lipofectamine. Following incubation for 6 h, the transfection medium was replaced by fresh medium for an additional 48-h incubation to allow for gene expression to occur.

MTT. The effect of transfection on cellular proliferation was assessed using MTT. In brief, the non-transfected, transfected pBabe-SR-I κ B α or control plasmid cells were seeded in 96-well plates in fresh medium (5000 cells/well) and incubation was continued for an additional 24, 48 or 72 h following transfection. Thereafter, MTT solution (5 mg/ml) was added to each well. Following incubation for 4 h at 37°C, the blue dye taken up by the cells was dissolved in dimethyl sulfoxide (100 μ g/ml), and then the optical density was measured at 570 nm using a 96-well multiscanner. All assays were run in triplicate.

Cells in the different groups mentioned previously were irradiated at room temperature with a medical linear accelerator (Primus-H; Siemens, Munich, Germany) at different doses. Cells in the transfected pBabe-SR-I κ B α plasmid group were harvested at different time points (1, 3, 6, 10, 24 and 48 h) following exposure to graded irradiation doses (0, 2, 4, 6, 8 and 10 Gy). Cells in the pBabe and non-transfected groups were harvested 3 h following exposure to graded irradiation doses. The 3 h time point was selected due to the fact that, as described by Sandur *et al*, the maximum expression of NF- κ B occurs 3 h following irradiation (10). Treated cells in the indicated groups were prepared for sequential immunocytochemistry, as well as western blot and flow cytometric analyses.

Immunocytochemistry. Changes in NF- κ B nuclear translocation were examined by immunocytochemistry. In brief, cells were plated onto 10x10-mm glass slides, and subjected to transfection and irradiation treatment. Thereafter, cells were fixed with 4% paraformaldehyde and permeabilized with 0.2% Triton X-100. Following washing in phosphate-buffered saline (PBS), slides were blocked with 5% normal goat serum for 1 h and then incubated with rabbit anti-NF- κ B antibody at a dilution of 1:100. Following overnight incubation at 4°C and rewarming to 37°C, the slides were washed with PBS and incubated for 20 min with goat anti-rabbit secondary antibody at a dilution of 1:100. The slides were then washed in PBS

and visualized using diaminobenzidine. Negative controls for each group were processed in the same manner, using a non-immunized rabbit IgG (at a dilution of 1:100) in place of the primary antibody. Immunocytochemical staining for NF- κ B in the nucleus was quantitatively analyzed using the Image-Pro Plus image analytical system (Media Cybernetics; Silver Spring, MD, USA) with the gray scale method.

Western blot. The changes in the expression of NF- κ B and mutant I κ B α following transfection and irradiation treatment were evaluated by western blot analysis. In brief, whole cell extracts were prepared in RIPA buffer (50 mM Tris-HCl, pH 7.4; 150 mM NaCl; 0.5 sodium deoxycholate; 0.1% SDS; 1% NP40 and 1 mM phenylmethylsulfonyl fluoride; Shenneng Bocai Biotechnology Co., Ltd.; Shanghai, China) from the treated cells grown in 6-cm dishes. The protein concentration was quantified using the BCA protein measurement kit (Shenneng Bocai Biotechnology Co., Ltd.). Extracts (40 μ g) were subjected to SDS-PAGE and then electrophoretically transferred to nitrocellulose membranes, which were then blocked with 5% (w/v) dried skimmed milk-TBST (10 mM Tris-HCl, pH 8.0; 150 mM NaCl; 0.05% Tween-20) for 1 h at room temperature. Membranes were probed with anti-NF- κ B antibody (at a dilution of 1:1000) or anti-I κ B α antibody (at a dilution of 1:1000), incubated overnight in 5% milk-TBST, washed three times with 1X TBST and exposed to HRP-conjugated secondary antibody (at 1:10000 dilution) in 5% dried skimmed milk for 1 h. All blots were reprobed with anti- β -actin (at 1:1000 dilution) in 5% dried skimmed milk, followed by HRP-conjugated secondary antibody (at a dilution of 1:1000) in 5% dried skimmed milk. Protein bands were visualized by the Alpha Imager 2200 system (Alpha Innotech, San Leandro, CA, USA).

Flow cytometric analysis. The effects on apoptosis in cells following the treatment indicated previously were determined by flow cytometric analysis. In brief, cells were trypsinized, then harvested by centrifugation at 1000 x g for 3 min. Thereafter, the cell collections were washed twice with cold PBS, resuspended in Annexin V and propidium iodide (PI), and incubated for 15 min at 4°C. Fluorescence was measured at 488 nm in a flow cytometer (Coulter, Luton, UK). Early apoptotic cells are indicated by Annexin V-positive and PI-negative staining, whereas late apoptotic cells demonstrate both Annexin V- and PI-positive staining.

Statistical analysis. SPSS/PC (version 17.0; SPSS, Inc., Chicago, IL, USA) was used for statistical analysis. Data are expressed as means \pm standard deviation and were analyzed by a one-way ANOVA with an LSD test. The Pearson's correlation coefficient analysis was used to assess the correlation between radiation doses, NF- κ B expression and apoptotic rate. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Expression of I κ B α protein is affected following transfection in ACC-M cells. To examine the changes in NF- κ B following irradiation in ACC-M cells, we transiently transfected the cells by use of a pBabe-SR-I κ B α plasmid prior to irradiation,




Figure 1. Expression of I κ B α protein following pBabe-SR-I κ B α plasmid transfection, demonstrated by western blot analysis; (A) the non-transfected, (B) pBabe and (C) pBabe-SR-I κ B α groups.




Figure 2. Effects of pBabe-SR-I κ B α gene transfection on the growth of adenoid cystic carcinoma cell line (ACC-M) cells were measured by MTT assay.

and thus suppressed the activity of the NF- κ B pathway. The expression of the Flag-tagged I κ B α following transfection was confirmed by western blot analysis. As demonstrated in Fig. 1, in the pBabe and non-transfected groups, cells exhibited a major band corresponding to endogenous I κ B α protein. However, the ACC-M cells that had been transfected with the pBabe-SR-I κ B α plasmid expressed endogenous I κ B α protein, as well as the mutant I κ B α protein with a higher molecular weight.

In vitro growth properties of transfected ACC-M cells. To determine whether the mutant I κ B α gene exerted its antiproliferative activity on the transfected cells, cells were subjected to an MTT analysis. As demonstrated in Fig. 2, no visible transfection-related inhibition was observed between the pBabe and non-transfected groups, while the growth property of cells in the pBabe-SR-I κ B α group was slightly reduced to 98.07, 97.79 and 97.5% after 24, 48 and 72 h, respectively. However, no significant difference was observed between the pBabe-SR-I κ B α group and the pBabe or non-transfected groups at the 95% probability level. Accordingly, it is likely that the expression of mutant I κ B α protein does not necessarily lead to the *in vitro* growth inhibition of non-stimulated ACC-M cells.

Translocation of NF- κ B following irradiation in ACC-M cells. NF- κ B staining was located within the cell cytoplasm in resting cells. Following exposure to different doses of irradiation, the previously suppressed NF- κ B pathway (achieved by transfecting the ACC-M cells with mutant I κ B α) was activated and thereafter liberated NF- κ B translocated to the nucleus. The staining of NF- κ B in the nucleus was categorized as positive expression. Compared with the pBabe and non-transfected groups at 3 h following exposure to equal doses of irradiation, the NF- κ B pathway was notably inhibited in the

pBabe-SR-I κ B α group, as demonstrated by a relatively lightly stained nucleus and the fact that the immunoprecipitates were mainly distributed in the cytoplasm. The majority of the NF- κ B protein in the cells of the pBabe-SR-I κ B α group was found to be clustered in the cytoplasm around the nuclear envelope. With exposure to graded doses of irradiation, the NF- κ B protein was observed to pass through the nuclear envelope and gradually accumulate in the nucleus. The NF- κ B protein was first observed in the center of the nucleus at 6 h (data not shown).

The quantitative analysis of the immunocytochemistry images supported our findings. Compared with the pBabe and non-transfected groups at 3 h following exposure to equal doses of irradiation, the average nuclear gray scale of cells in the pBabe-SR-I κ B α plasmid group was significantly increased ($P < 0.05$; data not shown). As demonstrated in Fig. 3, further analysis of the SR-I κ B α group revealed that the average nuclear gray scale of the various dose groups was lower than that of the 0 Gy group with only marginally detectable expression of NF- κ B protein, and the data reached the deepest valley 6-10 h following irradiation. At the same time point, we disclosed that at 3 h following irradiation, the average nuclear gray scale of the ACC-M cells decreased as the doses of irradiation increased ($P < 0.05$).

Expression of NF- κ B protein is affected following irradiation in ACC-M cells. To explore the quantitative changes in NF- κ B protein levels in ACC-M cells, cell lysates were subjected to western blot analysis following transfection and subsequent irradiation. Cells in the pBabe-SR-I κ B α plasmid group demonstrated downregulated levels of NF- κ B protein ($P < 0.05$), which suggested transfection of mutant I κ B α simultaneously blocked the synthesis and translocation of NF- κ B protein (data not shown). Further analysis on the time course and dose-response characteristics of NF- κ B protein in the pBabe-SR-I κ B α plasmid group are demonstrated in Fig. 4. The levels of NF- κ B protein in the various dosage groups were significantly higher than that of the 0 Gy group. The NF- κ B protein levels peaked at 6-10 h and began gradually declining at 24 h in each group with equal irradiation exposure. At the 3 h time point, the levels of NF- κ B protein increased as the dose of irradiation increased ($P < 0.05$). Overall, the western blot analysis results are consistent with our immunocytochemistry data.

Changes in the apoptotic rate following irradiation in ACC-M cells. It has been demonstrated that irradiation results in deleterious DNA damages and consequentially launches cell apoptosis. However, the simultaneously triggered cellular defense mechanism prevents impaired cells from the radiation-induced apoptosis (13,14). NF- κ B has been demonstrated to be activated by irradiation, and to therefore elucidate the elevated irradiation resistance (6). As demonstrated in Fig. 5, treatment with mutant I κ B α transfection significantly enhanced the radiosensitivity of ACC-M cells, as assessed by flow cytometric analysis ($P < 0.05$). A significant increase in the number of both Annexin V-positive and PI-negative (early apoptosis) and Annexin V- and PI-positive (late apoptosis) ACC-M cells was detected in the pBabe-SR-I κ B α plasmid group with graded doses of irradiation, compared with the




Figure 3. Average nuclear gray scale at different time points following exposure to graded doses of irradiation in the pBabe-SR-I κ B α group.




Figure 4. (A) Quantitative changes in NF- κ B expression at different time points following exposure to graded doses of irradiation, as demonstrated by western blot analysis in the pBabe-SR-I κ B α group. (B) The relative level of NF- κ B expression in 40 μ g of whole cell extract as determined by western blot analysis. Data are presented as the mean integrated density value (IDV) of three separate experiments (\pm standard deviation).




Figure 5. Rate of apoptosis in the different transfection groups, as measured by flow cytometry.




Figure 6. Apoptosis of adenoid cystic carcinoma cell line (ACC-M) cells following irradiation in the pBabe-SR-I κ B α group, which comprised the following groups: (A) Control; (B) 2 Gy, 3 h; (C) 4 Gy, 24 h; (D) 6 Gy, 24 h; (E) 8 Gy, 24 h; and (F) 10 Gy, 3 h.




Figure 7. Apoptosis of adenoid cystic carcinoma cell line (ACC-M) cells following exposure to graded doses of irradiation in the pBabe-SR-I κ B α group at different time points, demonstrated by flow cytometry.

0 Gy group (Fig. 6). Furthermore, cells with equal irradiation exposure exhibited a similar survival tendency in that the apoptotic rate reached the minimal value at 6-10 h, and the maximal value at 24 h. The apoptotic rates of cells with mutant I κ B α transfection, 3 h following irradiation treatment, were 50.83, 48.61, 44.32, 42.53 and 40.42%, between the different dosage groups (2, 4, 6, 8 and 10 Gy, respectively). These results suggested that the increasing expression of NF- κ B was attributed to the decreasing number of total (early and late) apoptotic cells at 3 h (Fig. 7).

The linear correlation estimation was applied to demonstrate that the apoptotic rates correlated well with the quantitative changes in NF- κ B protein expression. The apoptotic rate of the cells was significantly negatively correlated with the expression of NF- κ B protein. The overexpression of NF- κ B protein 6-10 h following irradiation in the pBabe-SR-I κ B α plasmid group led to the lowest rate of apoptosis. Meanwhile, graded doses of irradiation amplified the activation of the NF- κ B pathway, which in reverse decreased the apoptotic rates at 3 h, and the increased expression of NF- κ B at 3 h decreased the apoptotic rates.

Discussion

The primary aim of our study was to determine whether the transfected mutant I κ B α gene was able to inhibit the activity of the NF- κ B pathway, and to investigate the negative

contribution of the radio-activated NF- κ B pathway on radiosensitivity. We transiently transfected a mutant I κ B α gene into ACC-M cells, obtaining simultaneous blockage of the synthesis and translocation of the NF- κ B protein. The mutant I κ B α gene promoted radiosensitivity via suppression of the NF- κ B pathway. Additionally, exposure to radiation invoked the activity of the NF- κ B pathway that was initially inhibited by transfection, in a dose- and time-dependent manner. The highly active NF- κ B protein in reverse exhibited a radio-resistant effect in the ACC-M cells and improved survival following irradiation.

NF- κ B plays a pivotal role in various physiological processes, including immune and inflammatory responses, and is also considered to regulate apoptosis, cell proliferation and resistance to chemotherapy- and radiotherapy-induced cytotoxicity (15). NF- κ B is sequestered inactive in the cytoplasm by interaction with an inhibitor of I κ B α in resting cells. It is now widely accepted that the degradation of I κ B α plays a crucial role in the activation of the NF- κ B pathway (12). When cells were stimulated by different exogenous stimuli, the Ser 32 and 36 residues of I κ B α were phosphorylated by IKK and I κ B α underwent subsequent ubiquitylation and degradation, thus releasing the NF- κ B protein for nuclear translocation (16). Fujioka *et al* transfected a retroviral vector encoding phosphorylation-defective I κ B α by inducing point mutations at the potential phosphoacceptor sites of Ser 32 and 36, and observed decreased NF- κ B DNA binding activity in the non-metastatic human pancreatic tumor cell line (17). However, to the best of our knowledge, applying a mutant form of I κ B α to specifically inhibit the NF- κ B pathway in ACC-M cells has not been previously studied. As we have demonstrated in ACC-M cells, the synthesis and translocation of the NF- κ B protein were highly specifically suppressed by transfecting the cells with the use of a pBabe-SR-I κ B α plasmid, and this blocking effect persisted following irradiation and hence when the NF- κ B pathway was activated.

Given the involvement of the NF- κ B pathway in oncogenesis and induction of resistance to conventional therapy, such as chemotherapy and irradiation, blockage of the pathway by specific inhibitors is beneficial and may be a promising antitumor therapy (18). NF- κ B is suppressed by multiple strategies, including the application of a

phosphorylation-defective form of I κ B α , proteasome inhibitors or pharmacological agents (10,17,18). van Hogerlinden *et al* applied a super-repressor form of I κ B α with mutant phosphorylation sites that selectively inhibited the NF- κ B pathway, yielding elevated apoptosis in the basal and suprabasal keratinocytes *in vivo* (19). In addition, the super-repressor considerably sensitized the keratinocytes to UV-induced apoptosis. Duffey *et al* also demonstrated that stable expression of a mutant form of I κ B α in squamous carcinoma cells resulted in an augmented apoptotic rate, and this blocking effect was not relieved by TNF- α induced activation (20). However, in the present study, no significant effects on cell proliferation were observed in ACC-M cells with pBabe-SR-I κ B α plasmid transfection. By contrast, cells with prior mutant I κ B α plasmid transfection were susceptible to radio-induced apoptosis. Accordingly, we estimate that inhibition of the NF- κ B pathway in non-stimulated ACC-M cells does not necessarily result in increased apoptosis, whereas inhibition of the NF- κ B pathway combined with irradiation yielded a significant radiosensitizing outcome.

Cytoplasmic NF- κ B is activated by various stimuli, including IL-1, TNF- α and viral infection, as well as irradiation and chemotherapeutic agents (8). Li *et al* demonstrated that maslinic acid, a triterpene derivative obtained from olive pomace, inhibited endogenous NF- κ B activity in a time-dependent manner and the TNF α -induced NF- κ B activation in a dose-dependent manner in pancreatic cancer cells, and thus prevented the expression of downstream target genes, such as Bcl-2, Survivin, COX-2 and IAP-1 (21). Sandur *et al* also demonstrated that maximal expression of NF- κ B occurred 3 h following exposure to 10 Gy irradiation in colorectal cancer cells (10). However, there are a limited number of studies concerning the time-dose expression pattern of radio-induced activation in previously inhibited NF- κ B pathway. In our study, a similar expression tendency was observed in the pBabe-SR-I κ B α plasmid group; the maximal expression appeared at 6-10 h following irradiation. At 3 h, graded doses of irradiation corresponded well with the increased activity of the NF- κ B pathway.

Accumulating studies indicate that the constitutive activation of NF- κ B is closely correlated with oncogenesis and tumor invasion. Aberrant activation of the NF- κ B pathway has been identified in various types of cancer, including both solid and hematopoietic malignancies (22). Zhang *et al* revealed that an activated NF- κ B pathway was significantly correlated with advanced clinical stage, high frequencies of tumor angiogenesis and perineural invasion, aggressive locoregional recurrence and distant metastasis in ACC, all of which predict an adverse prognostic prospect (23). Sustained activation of the NF- κ B pathway has been demonstrated to promote survival in tumor cells with radiotherapy by permitting cells to repair DNA damage and to escape elimination by apoptosis, which is induced by the destruction of DNA double-strand (24). In our study, the peak expression levels of NF- κ B protein at 6-10 h following irradiation were paralleled by minimal apoptotic rates in the pBabe-SR-I κ B α plasmid group. Amplified expression levels of NF- κ B activated by graded doses of irradiation at 3 h reflected attenuated apoptosis. These findings to an extent explain the radio-resistant role of NF- κ B in stimulated ACC-M cells.

In the present study, we demonstrated that ACC-M cells exhibited markedly increased radiosensitivity following selective inhibition of the NF- κ B pathway by transfection with a mutant I κ B α plasmid. However, the NF- κ B pathway also plays a crucial role in various physiological processes, such as regulation of innate immunity by activating downstream pro-inflammatory cytokines, including IL-1, TNF- α and RANTES. One of the pitfalls that may be neglected is the likelihood of immunodeficiency and great susceptibility to infections following specific inhibition of the NF- κ B pathway. The correlation between downregulation of the NF- κ B pathway and human diseases is reported in *Incontinentia pigmenti* and anhidrotic ectodermal dysplasia (25). van Hogerlinden *et al* observed interferences in normal epidermal homeostasis and hair follicle development in selectively inhibited genetic mice (19). The transgenic mouse model exhibited dysplasia of the epidermis characterized by flaky skin, hair loss and hyperkeratotic lesions. The increased susceptibility to the spontaneous occurrence of well-differentiated skin squamous cell carcinoma was also demonstrated in the transgenic mice. Accordingly, the appropriate application of NF- κ B pathway inhibitors while avoiding their side-effects requires further investigation.

In conclusion, the results of our study demonstrated that by transfecting a mutant form of the I κ B α gene, the synthesis and translocation of the NF- κ B protein were specifically inhibited, and a concomitant decrease in survival rate following irradiation was observed in ACC-M cells. Furthermore, the NF- κ B expression after irradiation represents a time-dose dependent manner of the transfected cells, followed by a corresponding change in radiosensitivity. Accordingly, selective inhibition of the NF- κ B pathway by novel target agents reveals a promising method for improvements in the radiosensitivity of ACC. However, the potential side-effects that may be activated by systemic suppression of immune system require further investigation.

References

1. Fordice J, Kershaw C, El-Naggar A and Goepfert H: Adenoid cystic carcinoma of the head and neck: predictors of morbidity and mortality. *Arch Otolaryngol Head Neck Surg* 125: 149-152, 1999.
2. Terhaard CH, Lubsen H, Van der Tweel I, *et al*: Salivary gland carcinoma: independent prognostic factors for locoregional control, distant metastases and overall survival: results of the Dutch head and neck oncology cooperative group. *Head Neck* 26: 681-693, 2004.
3. Polat B, Fassnacht M, Pfreundner L, *et al*: Radiotherapy in adrenocortical carcinoma. *Cancer* 115: 2816-2823, 2009.
4. Karin M and Ben-Neriah Y: Phosphorylation meets ubiquitination: the control of NF-kappa B activity. *Annu Rev Immunol* 18: 621-663, 2000.
5. Ghosh S, May MJ and Kopp EB: NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol* 16: 225-260, 1998.
6. Pajonk F, Pajonk K and McBride WH: Inhibition of NF- κ B, clonogenicity, and radiosensitivity of human cancer cells. *J Natl Cancer Inst* 91: 1956-1960, 1999.
7. Greten FR and Karin M: The IKK/NF- κ B activation pathway-A target for prevention and treatment of cancer. *Cancer Lett* 206: 193-199, 2003.
8. Pahl HL: Activators and target genes of Rel/ NF- κ B transcription factors. *Oncogene* 18: 6853-6866, 1999.
9. Schwartz SA, Hernandez A and Mark Evers B: The role of NF-kappaB/ IkappaB proteins in cancer: implications for novel treatment strategies. *Surg Oncol* 8: 143-153, 1999.

10. Sandur SK, Deorukhkar A, Pandey MK, *et al*: Curcumin modulates the radiosensitivity of colorectal cancer cells by suppressing constitutive and inducible NF- κ B activity. *Int J Radiat Oncol Biol Phys* 75: 534-542, 2009.
11. Jung M, Zhang Y, Lee S and Dritschilo A: Correction of radiation sensitivity in ataxia telangiectasia cells by truncated I κ B- α . *Science* 268: 1619-1621, 1995.
12. Brown K, Gerstberger S, Carlson L, Franzoso G and Siebenlist U: Control of I κ B- α proteolysis by site-specific signal induced phosphorylation. *Science* 267: 1485-1488, 1995.
13. Cataldi A, Zauli G, Di Pietro R, Castorina S and Rana R: Involvement of the pathway phosphatidylinositol-3-kinase/AKT-1 in the establishment of the survival response to ionizing radiation. *Cell Signal* 13: 369-375, 2001.
14. Weichselbaum RR, Hallahan D, Fuks Z and Kufe D: Radiation induction of immediate early genes: effectors of the radiation-stress response. *Int J Radiat Oncol Biol Phys* 30: 229-234, 1994.
15. Magné N, Toillon RA, Bottero V, *et al*: NF- κ B modulation and ionizing radiation: mechanisms and future directions for cancer treatment. *Cancer Lett* 231: 158-168, 2006.
16. Tergaonkar V, Bottero V, Ikawa M, Li Q and Verma IM: I κ B kinase-independent I κ B α degradation pathway: functional NF- κ B activity and implications for cancer therapy. *Mol Cell Biol* 23: 8070-8083, 2003.
17. Fujioka S, Scwab GM, Schmidt C, *et al*: Inhibition of constitutive NF- κ B activity by I κ B α M suppresses tumorigenesis. *Oncogene* 22: 1365-1370, 2003.
18. Darnell JE Jr: Transcription factors as targets for cancer therapy. *Nat Rev Cancer* 2: 740-749, 2002.
19. van Hogerlinden M, Rozell BL, Ahrlund-Richter L and Toftgård R: Squamous cell carcinomas and increased apoptosis in skin with inhibited Rel/nuclear factor-kappaB signaling. *Cancer Res* 59: 3299-3303, 1999.
20. Duffey DC, Crowl-Bancroft CV, Chen Z, *et al*: Inhibition of transcription factor nuclear factor-kappa B by a mutant inhibitor-kappa B α attenuates resistance of human head and neck squamous cell carcinoma to TNF-alpha caspase-mediated cell death. *Br J Cancer* 83: 1367-1374, 2000.
21. Li C, Yang Z, Zhai C, *et al*: Maslinic acid potentiates the anti-tumor activity of tumor necrosis factor alpha by inhibiting NF-kappaB signaling pathway. *Mol Cancer* 9: 73, 2010.
22. Baud V and Karin M: Is NF-kappaB a good target for cancer therapy? Hopes and pitfalls. *Nat Rev Drug Discov* 8: 33-40, 2009.
23. Zhang J, Peng B and Chen X: Expressions of nuclear factor kappaB, inducible nitric oxide synthase, and vascular endothelial growth factor in adenoid cystic carcinoma of salivary glands: correlations with the angiogenesis and clinical outcome. *Clin Cancer Res* 11: 7334-7343, 2005.
24. Magné N, Toillon RA, Bottero V, *et al*: NF-nB modulation and ionizing radiation: mechanisms and future directions for cancer treatment. *Cancer Lett* 231: 158-168, 2006.
25. Courtois G and Gilmore TD: Mutations in the NF- κ B signaling pathway: implications for human disease. *Oncogene* 25: 6831-6843, 2006.