

RESEARCH ARTICLE

Prognostic Role of BRAF Mutation in Stage II/III Colorectal Cancer Receiving Curative Resection and Adjuvant Chemotherapy: A Meta-Analysis Based on Randomized Clinical Trials

Lizhen Zhu^{1☯‡}, Caixia Dong^{1☯‡}, Ying Cao¹, Xuefeng Fang¹, Chenhan Zhong¹, Dan Li¹, Ying Yuan^{1,2*}

1 Department of Medical Oncology, The Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, Zhejiang Province, China, **2** Cancer Institute, Key Laboratory of Cancer Prevention and Intervention, Chinese National Ministry of Education, Key Laboratory of Molecular Biology in Medical Sciences, The Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, Zhejiang Province, China

☯ These authors contributed equally to this work.

‡ These authors are co-first authors on this work.

* yuanying1999@zju.edu.cn



CrossMark
click for updates

OPEN ACCESS

Citation: Zhu L, Dong C, Cao Y, Fang X, Zhong C, Li D, et al. (2016) Prognostic Role of BRAF Mutation in Stage II/III Colorectal Cancer Receiving Curative Resection and Adjuvant Chemotherapy: A Meta-Analysis Based on Randomized Clinical Trials. PLoS ONE 11(5): e0154795. doi:10.1371/journal.pone.0154795

Editor: Nikolas K. Haass, University of Queensland Diamantina Institute, AUSTRALIA

Received: December 31, 2015

Accepted: April 19, 2016

Published: May 3, 2016

Copyright: © 2016 Zhu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: The work was supported by four research funds: Key Projects in the National Science & Technology Pillar Program during the Twelfth Five-year Plan Period (2014BAI09B07), <http://www.most.gov.cn>; Training Program of the Major Research Plan of the National Natural Science Foundation of China (91229104), <http://www.most.gov.cn>; The National High Technology Research and Development Program of China (863 Program) (2012AA02A506),

Abstract

Background and Objective

Studies examining the prognostic value of the BRAF mutation on relapse-free survival (RFS), disease-free survival (DFS) and overall survival (OS) in stage II/III colorectal cancer (CRC) patients receiving curative resection and adjuvant chemotherapy so far showed discrepant results. Therefore, a meta-analysis of relevant studies was performed for clarification.

Methods

Randomized trials of stage II/III colorectal cancer treated with curative resection followed by adjuvant chemotherapy were selected to conduct a meta-analysis. The necessary descriptive and statistical information such as hazard ratios (HRs) and 95% confidence intervals (CIs) were derived from published survival data.

Results

Seven phase III randomized clinical trials (RCTs) including 1,035 BRAF mutation stage II/III CRC patients receiving curative resection and adjuvant chemotherapy were analyzed. Overall, BRAF mutation resulted in poorer OS (HR = 1.42, 95% CI: 1.25–1.60; P < 0.00001), and poorer DFS (HR = 1.26, 95% CI: 1.07–1.48, P = 0.006) compared with BRAF wild-type CRC. The prognostic role on RFS could not be elucidated in the meta-analysis because of limited data.

<http://www.most.gov.cn>; The National High Technology Research and Development Program of China (863 Program) (2012AA02A204), <http://www.most.gov.cn>. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All authors read and approved the final manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Conclusions

BRAF mutation was significantly related with shorter DFS and OS among stage II/III CRC patients receiving adjuvant chemotherapy after curative resection. Its prognostic role for RFS needs to be further analyzed when more data is available.

Introduction

Curative resection followed by adjuvant chemotherapy remarkably improved survival in patients with high-risk stage II and stage III colorectal cancer (CRC). However, it has soon become clear that not all patients benefit equally and a significant proportion of patients even has worse prognosis than others. Retrospective researches and randomized clinical trials (RCTs) carried out in these patients confirmed that BRAF mutation is one of the strong negative prognostic factors [1, 2].

A negative prognostic impact of the BRAF mutation in metastatic colorectal cancer had already been corroborated by several meta-analyses [3, 4]. Irrespective of the stage of cancer, one meta-analysis has shown that mutated BRAF determines poor overall survival (OS) in colorectal cancer when compared with wild type BRAF [Hazard ratio (HR) = 2.25, 95% confidence interval (CI): 1.82–2.83] [5]. However, in the subgroup of stage II/III CRC patients treated with curative resection and adjuvant chemotherapy, the prognostic value of BRAF mutation is still uncertain. Although several RCTs including PETACC-3 [6], or NSABP C-07 and C-08 [7], recognize BRAF mutation as an unfavorable prognostic factor for OS in this subgroup of CRC, but it was not confirmed by others, take CALGB 89803 [8] and MOSAIC [9] as examples. Furthermore, in respect of disease-free survival (DFS), results from the RCTs are inconclusive: for example CALGB 89803 found no difference between BRAF-mut and BRAF-wt CRC whereas N0147 [2] observed worse DFS in the BRAF-mut subpopulation compared to the BRAF-wt subpopulation. Interestingly, the prognostic value of BRAF mutation was not the same according to different cancer sites in N0147 trial.

Therefore a systematic review and meta-analysis of RCTs was performed to quantitatively evaluate whether BRAF mutation plays a negative prognostic role in stage II/III CRC receiving curative resection and adjuvant chemotherapy.

Methods

Selection criteria and search strategy

Three individuals carried out the literature search. All randomized trials (1) of stage II/III CRC undergoing curative resection, followed by adjuvant chemotherapy, and (2) with sufficient quantitative data of the prognosis according to BRAF mutation status were eligible. We considered sufficient quantitative data as data including the necessary descriptive and statistical information to perform meta-analysis, such as HR, CI, observed or logrank expected events and Kaplan-Meier curve; there was no need to included all of these data, but enough data to perform meta-analysis was required. The insufficient quantitative data means the article lacked necessary information to perform meta-analysis.

This meta-analysis was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (S1 Table) [10]. We searched PubMed, EMBASE, Web of Science and the Cochrane library using the following terms: (colon cancer or colon carcinoma or colorectal cancer or colorectal carcinoma or rectal cancer or rectal

carcinoma) and (BRAF or B-RAF or B RAF) and (((stage and (II or III)) or resect* or surg* or opera* or adjuvant*). Database searches were restricted to articles published in English until 19 November 2015. Exclusion criteria were described as follows: (1) mixed prognostic data of stage I or IV with stage II/III colorectal cancer; (2) neoadjuvant chemotherapy or radiation prior to resection; (3) data partly or totally included in other eligible studies; (4) insufficient prognostic data according to BRAF mutations.

Data extraction and quality assessment

The following data were collected from each study: author, year of publication, treatment protocol, BRAF detection, mutational data, and survival outcomes, namely, relapse-free survival (RFS), DFS and OS. Overall survival was defined from the time random assignment to the date of death, and disease free survival was calculated from the time study enrollment to tumor recurrence, occurrence of a new primary colorectal tumor, or death from any cause, while RFS was defined from the time study enrollment to tumor recurrence or occurrence of a new primary colon tumor, patients that died without these two issues were censored. Specifically, the hazard ratios and associated 95% confidence intervals of survival outcomes for BRAF-mut patients compared to those with BRAF-wt were extracted. Pooled estimates of the prognosis for survival according to the BRAF status were weighted and pooled using the generic inverse-variance. An Egger's funnel plot was performed to assess the publication bias.

Statistical analysis

The comparative effects were initially analyzed by the traditional pairwise meta-analysis method using Cochrane Collaboration review manager software (RevMan v.5.3.0). The relative risk for dichotomous outcomes and the standardized mean difference for continuous outcomes pooled across studies were estimated by using the DerSimonian and Laird random-effects model [11]. The level of significance was set at 5%. If the result of analysis showed $p > 0.05$, the studies were considered homogeneous and a fixed-effect model for meta-analysis was chosen. Otherwise, a random-effect model was performed. Inconsistency was quantified using the I^2 statistic, $I^2 < 25\%$ reflects a small level of inconsistency while $I^2 > 50\%$ implies significant inconsistency. And we excluded one study each time to perform the sensitivity analysis to examine the robustness of the estimates. For all analyses, $P < 0.05$ was considered statistically significant.

Results

1. Literature retrieval and study characteristics

The literature search identified 2631 citations, of which seven phase III RCTs with 14,699 patients were considered eligible and were included in this meta-analysis. (Fig 1) Among these, 8721 patients had been assessed for the BRAF mutation, and a total of 1,035 BRAF-mut CRC patients, 11.9% of the tested population, were incorporated in these studies. Since we defined the OS, DFS and RFS as mentioned above, the RFS defined in the research of Roth et al [6] was transferred into DFS in this meta-analysis. Sinicrope et al had observed that BRAF-mut had significant negative prognostic value for DFS and OS in proximal colon cancer. However, the prognostic value of BRAF mutation was not significant for DFS and OS in colon cancer in the same report. [2] Therefore, analysis about proximal side and distal side from that study were considered as separate ones—named Sinicrope 1 (proximal colon cancer) and Sinicrope 2 (distal colon cancer)—in the meta-analysis. Although most studies included were enrolled in colon cancer, in the study of Pentheroudakis 27.4% cancers' primary site was rectum, and all the




Fig 1. Selection of studies included in the analysis.

doi:10.1371/journal.pone.0154795.g001

studies included both sides cancer. The characteristics and biomarker analyses of included studies are outlined in [Table 1](#). Particularly, in these RCTs, BRAF mutations are more prominent in DNA mismatch repair (MMR) defective and microsatellite instability (MSI) high tumors, for examples, Roth et al found 46% BRAF-mut patients were with MSI-high; Gavin et al found 29% BRAF-mut patients were with dMMR, while 9% BRAF-wt patients with dMMR; in French's study [12], 45% BRAF-mut patients were with dMMR, while 6% BRAF-wt patients with dMMR, while only French's study [12] performed stratified analysis of BRAF mutation's prognostic role according to MMR status, and only Gavin et al's study [7] provided stratified analysis of MMR status' prognostic role according to BRAF mutations. Therefore, there is no sufficient data to do further stratified. None of the included studies analyzed KARS mutation's prognostic role in BRAF wild type group.

2. Quality of individual studies

Among the eligible studies, seven RCTs were double blinded, and all described the randomization processes that they had used. All included a power calculation to determine the optimal sample size. The funnel plots of publication bias of OS meta-analysis provided a qualitative estimation of publication bias of the studies, and no evidence of bias was found. ([Fig 2](#))

Table 1. Characteristics of included phase III RCTs.

Author, year [ref]	Treatment	Total n	Pts with tissue available for BRAF mutations	Median follow-up time	BRAF mut type	BRAF assessment method	BRAF mut n, tested % (n)	OS	DFS	RFS
French, 2008 [12]	Curative surgery + FU/LV	878	490	NA	V600E	CSGE, PCR	15.7% (77)	mut VS wt HR = 1.2, 95% CI = (0.8–1.8), p = 0.31	mut VS wt HR = 1.0, 95% CI = (0.6–1.6), p = 0.97	NA
Gavin, 2012 [7]	Curative surgery + FLOX vs FU/LV, curative surgery + mFOLFOX6 vs mFOLFOX6 + Bevacizumab	4802	2226	NA	V600E, D594V,	sequenom, PCR	V600E 13.15% (293), D594V 0.80% (18)	mut VS wt HR = 1.46; 95% CI, (1.20–1.79); P = 0.0002	NA	mut VS wt HR, 1.02; 95% CI, 0.82– 1.28; P = .0.86
Roth, 2012 [6]	Curative surgery + FOLFIRI vs 5-FU/LV	2094	1404	69 months	V600E	PCR, sequencing	8% (103)	mut vs wt HR = 1.56, 95% CI = (1.02 to 2.39), p = 0.04	mut vs wt HR = 1.17, 95% CI = (0.79 to 1.73) p = 0.44	NA
Ogino, 2013 [8]	Curative surgery + FU/LV vs IFL	1264	580	7.6 years	V600E	PCR, pyrosequencing	16% (94)	mut VS wt HR = 1.46, 95% CI = (1.00– 2.15)	mut VS wt HR = 1.35, 95% CI = (0.94– 1.93)	mut VS wt HR 1.44, CI (0.99– 2.11)
Sinicrope 1, 2015 [2]	Curative surgery + mFOLFOX6 ± cetuximab	1511	1423	4.9 years	V600E	PCR, sequencing	20% (287)	mut vs wt HR = 1.58, 95% CI = (1.17– 2.12), p = 0.003	mut vs wt HR = 1.35, 95% CI = (1.03– 1.76), p = 0.028	NA
Sinicrope 2, 2015 [2]	Curative surgery + mFOLFOX6 ± cetuximab	1463	1375	4.9 years	V600E	PCR, sequencing	4% (54)	mut vs wt HR = 1.79, 95% CI = (0.99– 3.24), p = 0.053	mut vs wt HR = 1.25, 95% CI = (0.75– 2.11), p = 0.394	NA
André, 2015 [9]	Curative surgery + LV5FU2 vs FOLFOX4	2246	902	10 years	V600E	PCR, pyrosequencing	10.42% (94)	mut VS wt HR = 0.99, 95% CI = 0.67 to 1.47, p = 0.965	NA	NA
Pantheroudakis, 2015 [13]	Curative surgery + FOLFOX vs XELOX	441	321	74.7 months	V600E	PCR, sequencing	4.7% (15)	mut VS wt HR 1.75, 95% CI (0.7– 4.35) p = 0.23	mut VS wt HR 1.28, 95% CI (0.52– 3.17) p = 0.59	NA

PCR = polymerase chain reaction; RCT = randomized clinical trial; FOLFOX = bolus + continuous infusion 5-fluorouracil + bolus folinic acid days 1, 2 + oxaliplatin; FU = 5-fluorouracil; LV = folinic acid; FLOX = bolus infusion 5-fluorouracil + bolus folinic acid + oxaliplatin; IFL = bolus irinotecan + bolus infusion 5-fluorouracil + bolus folinic acid; FOLFIRI = 5-fluorouracil + folinic acid + irinotecan; BRAF = B-Raf proto-oncogene; DFS = disease-free survival; OS = overall survival; RFS = relapse-free survival; NA = not available.




Fig 2. Funnel plots for publication bias of OS (a) and DFS (b) meta-analysis: no evidence of bias was found.

doi:10.1371/journal.pone.0154795.g002

3. Prognostic role of BRAF mutation on OS in stage II/III CRC

Among the eligible studies, eight subgroups from seven RCTs were included to assess the predictive role of BRAF mutation on OS in CRC with adjuvant chemotherapy compared to BRAF wild. No heterogeneity was found among the trials ($P = 0.61$, $I^2 = 0\%$), and a fixed-effects model was thus chosen for the analysis. The prognostic role of BRAF mutation on OS is shown in Fig 3. Based on the meta-analysis, CRC with BRAF mutation have poorer overall survival after surgery and adjuvant chemotherapy, and the HR for OS with BRAF mutation is 1.42 (95% CI, 1.25–1.60; $P < 0.00001$), compared to BRAF wild CRC. (Fig 3)

4. Prognostic role of BRAF mutation on DFS in stage II/III CRC

There were six subgroups from five RCTs with enough data to assess the efficacy of BRAF mutation on DFS compared to BRAF wild. No heterogeneity was found among the trials ($P = 0.92$, $I^2 = 0\%$), and a fixed-effects model was thus chosen for the analysis. Based on the




Fig 3. Forest plots showing hazard ratio for a negative prognostic role of BRAF mutation on over all survival in stage II/III CRC.

doi:10.1371/journal.pone.0154795.g003

analysis of four RCTs, CRC with BRAF mutation had shorter DFS than those with BRAF wild (HR = 1.26, 95% CI: 1.07–1.48, P = 0.006). (Fig 4)

5. Prognostic role of BRAF mutation on RFS in stage II/III CRC

Among the seven RCTs, only Gavin et al [7] and Ogino et al [8] analyzed the prognostic role of BRAF mutation on RFS, reporting HRs of 1.02 (0.82–1.28) and 1.44 (0.99–2.11), respectively. Therefore, we did not perform the meta-analysis about the prognostic role of BRAF mutation on RFS due to the limited data.

6. Sensitivity analysis

One study was removed at a time to conduct the sensitivity analysis. The pooled HRs and CIs were not significantly changed for OS, indicating the stability of the result. However, it is noteworthy that in the sensitivity analysis for DFS the outcome seemed similar except that when the study Sinicrope 1 excluded, the HR (95%CI) become 1.21 (0.99, 1.48), suggesting the lack of robustness of the estimate and a cautious interpretation about the result that BRAF mutation plays a unfavorable role on DFS in stage II/III CRC is needed. (Table 2)




Fig 4. Forest plots showing hazard ratio for a negative prognostic role of BRAF mutation on disease free survival in stage II/III CRC.

doi:10.1371/journal.pone.0154795.g004

Table 2. Sensitivity analysis by omitting each of the included studies in different outcomes.

Outcomes	Omitted Study	HR (95%CI)	P	I ²	Ph
OS	André 2015	1.47 (1.29, 1.67)	< 0.00001	0	0.93
	French 2008	1.44 (1.27, 1.64)	< 0.00001	0	0.59
	Gavin 2012	1.40 (1.20, 1.63)	< 0.0001	0	0.50
	Ogino 2013	1.41 (1.24, 1.61)	< 0.00001	0	0.49
	Pentheroudakis 2015	1.41 (1.25, 1.59)	< 0.00001	0	0.52
	Roth 2012	1.40 (1.24, 1.59)	< 0.00001	0	0.52
	Sinicrope 1 2015	1.39 (1.21, 1.58)	< 0.00001	0	0.56
	Sinicrope 2 2015	1.40 (1.24, 1.59)	< 0.00001	0	0.57
	DFS	French 2008	1.30 (1.09, 1.54)	0.004	0
Roth 2012		1.28 (1.07, 1.53)	0.008	0	0.88
Ogino 2013		1.23 (1.02, 1.49)	0.03	0	0.88
Sinicrope 1 2015		1.21 (0.99, 1.48)	0.07	0	0.91
Sinicrope 2 2015		1.26 (1.06, 1.50)	0.009	0	0.84
Pentheroudakis 2015		1.26 (1.06, 1.49)	0.007	0	0.84

OS = overall survival; DFS = disease free survival; HR = hazard ratio; Ph = P for heterogeneity

doi:10.1371/journal.pone.0154795.t002

Discussion

The cellular RAS/RAF/Mitogen-activated protein kinase/ERK kinase (MEK)/extracellular signal regulated kinase (ERK) pathway controls cell proliferation, apoptosis and differentiation by transferring signals from cell surface receptors to transcription factors. Activated by RAS, the RAF phosphorylates and activates downstream MEK, which then activates ERK subsequently and ultimately results in the activation of transcription factors. Mutations of these signals can cause abnormal cell growth, invasion and metastasis. Among these, RAF consists of serine–threonine kinases ARAF, BRAF and CRAF. While ARAF and CRAF mutations are detected only rarely, BRAF mutations are found at a higher frequency in several cancers. Meanwhile, 90% BRAF mutations are V600E mutation, which is reported to occur in melanoma (40–60%), papillary thyroid carcinoma (45%), low grade serous ovarian carcinoma (35%), colorectal adenocarcinoma (5–15%) and other cancers [14]. Besides, Zheng et al found that about 20.6% of BRAF-mut tumors in CRC are non-V600E mutations, including N581S, Y472C, D594G and D594N mutations, and the latter two are related to concomitant activation of KRAS or NRAS mutations [15]. Patients harboring BRAF 594 or 596 mutated CRC have a longer OS than BRAF wild type and BRAF V600E mutated CRC (62.0 versus 35.9 months versus 12.6 months; HR = 0.55, P = 0.081 and HR 0.36, P = 0.002) [16].

In CRC, while it is nearly mutually exclusive with KRAS mutation, BRAF mutation is associated with specific high-risk clinicopathologic characteristics, such as lymph-node metastasis, high grade, T4 tumors, mucinous histology, MSI or MMR and right-side location [8, 9]. Recently, Loupakis et al found that CRC patients with right-side primary, female gender and mucinous histology have 81% chance to bear a BRAF V600E-mutant [17, 18]. Since these characteristics themselves are related to worse survival, it may be possible that the unfavorable prognostic value of BRAF mutation and these high-risk factors are interactive. For example, BRAF V600E mutation is detected in less than 12% MMR-proficient tumors, while in MMR-defective tumors the incidence ranges from 35 to 60% [7, 12]. It was reported that the prognostic effect of BRAF and MMR status is additive, giving rise to worst prognosis for patients with MMR-proficient and BRAF-mutant tumors [7]. Similar results were observed in patients

according to MSI or microsatellite stability (MSS). On the other hand, compared to the right-sided BRAF-mut CRC, the left-sided BRAF-mut CRC had a worse prognosis. Hence, the MSS/left side BRAF-mut population had the worst survival [17]. However, it should be reminded that these clinical and pathologic features are typically all bundled together.

Several meta-analyses have demonstrated that advanced CRC with BRAF-mut has a much poorer prognosis compared to those with BRAF-wt [19, 20]. Conversely, the prognostic role of BRAF mutation has not been clarified in stage II and stage III CRC patients undergoing adjuvant chemotherapy after curative resection. The actual meta-analysis that quantitatively and systematically collected the evidence from seven RCTs is the first one to demonstrate that the presence of BRAF mutation plays a negative prognostic role for OS in the patients. As for DFS, in the sensitivity analysis when the study Sinicope 1 was removed, the unfavorable prognostic value of BRAF mutation was not statistically confirmed, with HR = 1.21, 95% CI: 0.99–1.48, $P = 0.07$, indicating the instability of the result. Hence, whether BRAF mutation is related with shorter DFS in stage II/III CRC needs a more cautious interpretation with the following two facts: 1) among the six studies included in the analysis about BRAF mutation's prognostic value on DFS, the study Sinicope 1 contains the largest number of BRAF-mut patients and has the narrowest 95% confidence interval (1.03–1.76), therefore the result of this study affect the meta-analysis' outcome remarkably, and deleting it could change the result significantly. 2) all patients in the study Sinicope 1 are proximal colon cancer, which not only is associated with high prevalence of BRAF mutation, but also, more importantly, indicates poor prognosis itself. The proximal colon location and BRAF mutation can synergistically play adverse role on the prognosis, and these patients may have the shorter DFS, thus the DFS may be better after removing the study Sinicope 1. Nevertheless, when the study was removed, I^2 of the heterogeneity remained to be 0%, meaning that the results of this meta-analysis is reliable in terms of the whole BRAF mutation population regardless of cancer locations. Consequently, the BRAF mutation has an unfavorable prognostic value on DFS in stage II/III CRC without regard to cancer locations and other situations. Certainly, it would be better to perform stratified analysis based on cancer locations to check whether BRAF mutation in proximal colon cancer is confirmed a adverse prognostic factor, while its prognostic value is not sure in distal colon cancer. Unfortunately we cannot work it out since only the study Sinicope performed stratified analysis of BRAF mutation's prognostic role according to cancer locations. Furthermore, a higher hazard ratio for OS than DFS in BRAF-mut patients may imply that population has a shorter survival time after recurrence in comparison to BRAF-wt patients, as suggested by Gavin et al [7].

Despite its negative prognostic value confirmed in this meta-analysis, there is no evidence of a predictive value of BRAF mutation in the terms of therapy outcomes. In stage II/III CRC, BRAF-mut patients receiving adjuvant chemotherapy plus cetuximab had no survival advantage over those receiving adjuvant chemotherapy only [21]. Moreover, the study of Ogino et al suggested no significant predictive value of BRAF mutation for irinotecan-based adjuvant chemotherapy either [22]. Similarly, meta-analyses have neither demonstrated statistical differences of anti-EGFR monoclonal antibodies' (mAbs) effect on progressive-free survival (PFS) or OS between BRAF-mut and BRAF-wt metastatic colorectal cancer, nor benefit from the addition of anti-EGFR mAbs to standard chemotherapy in BRAF-mut patients compared with chemotherapy alone or best support care [23, 24]. Fortunately, compared to FOLFIRI plus bevacizumab, FOLFOXIRI plus bevacizumab seems to generate better PFS and OS in BRAF-mut advanced CRC [25].

The following issues should also be taken into account when interpreting the results of this meta-analysis. First, because of the relatively low prevalence of BRAF mutation in CRC, the proportion of BRAF-mut patients is limited. Moreover, even though all the prognostic data were derived from phase III randomized clinical trials, some studies included in the meta-

analysis analyzed BRAF status retrospectively, leading to about 6–60% of patients lacking information about BRAF mutational status causing that it might miss a positive prognostic value during the evaluation. Second, since this analysis has been performed based on published and not on individual patient data, specific subgroup analyses could not be conducted, such as further stratified analysis according to MSI/MMR status in BRAF-mut patients. Third, the chemotherapies received in some trials are not used as standard management in stage II or III colorectal cancer. Besides, two RCTs contain irinotecan as treatment while five contain oxaliplatin, and bevacizumab is included in one study include compared to cetuximab in two study. However it is infeasible to do further stratified according to the regimens since these studies mixed the survival data from different therapies together. Finally, due to the uncommonness of most BRAF mutation types, this meta-analysis almost exclusively evaluated the prognostic value of the BRAF V600E mutation. For example, about 0.80% ($n = 18$) BRAF mutation in NSABP C-07 and C-08 clinical trials were BRAF D594G mutations, but such rare or undetected BRAF mutation types may not inevitably carry a similar prognostic value as BRAF V600E mutation. Nonetheless, this study is based on the most exhaustive overview of traceable randomized trials conducted so far in stage II/III CRC with curative resection and adjuvant chemotherapy. Furthermore, this is the first meta-analysis that suggests a negative prognostic role of the BRAF V600E mutation for this group of CRC patients.

In summary, this meta-analysis shows that BRAF mutation is significantly related with worse DFS and OS among stage II/III CRC patients receiving adjuvant chemotherapy after curative resection, while its prognostic role for RFS needs to be further analyzed when more data is available. Future studies, especially well-designed large randomized controlled trials conducted in BRAF-mut stage II/III CRC patients are urgently needed to find better treatments for that population.

Supporting Information

S1 Table. PRISMA checklist.
(DOC)

Acknowledgments

We gratefully acknowledge financial support from following four research funds: Key Projects in the National Science & Technology Pillar Program during the Twelfth Five-year Plan Period (2014BAI09B07); Training Program of the Major Research Plan of the National Natural Science Foundation of China (91229104); The National High Technology Research and Development Program of China (863 Program) (2012AA02A506); The National High Technology Research and Development Program of China (863 Program) (2012AA02A204). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. All authors read and approved the final manuscript.

Author Contributions

Conceived and designed the experiments: YY LZZ. Performed the experiments: CXD YC XFF. Analyzed the data: LZZ CHZ DL. Contributed reagents/materials/analysis tools: CXD LZZ. Wrote the paper: YY LZZ CXD.

References

1. Ooki A, Akagi K, Yatsuoka T, Asayama M, Hara H, Takahashi A, et al. Combined microsatellite instability and BRAF gene status as biomarkers for adjuvant chemotherapy in stage III colorectal cancer. *J Surg Oncol*. 2014 Dec; 110(8):982–8. doi: [10.1002/jso.23755](https://doi.org/10.1002/jso.23755) PMID: [25154726](https://pubmed.ncbi.nlm.nih.gov/25154726/)

2. Sinicrope FA, Mahoney MR, Yoon HH, Smyrk TC, Thibodeau SN, Goldberg RM, et al. Analysis of Molecular Markers by Anatomic Tumor Site in Stage III Colon Carcinomas from Adjuvant Chemotherapy Trial NCCTG N0147 (Alliance). *Clin Cancer Res*. 2015 Dec 1; 21(23):5294–304. doi: [10.1158/1078-0432.CCR-15-0527](https://doi.org/10.1158/1078-0432.CCR-15-0527) PMID: [26187617](https://pubmed.ncbi.nlm.nih.gov/26187617/)
3. Xu Q, Xu AT, Zhu MM, Tong JL, Xu XT, Ran ZH. Predictive and prognostic roles of BRAF mutation in patients with metastatic colorectal cancer treated with anti-epidermal growth factor receptor monoclonal antibodies: a meta-analysis. *J Dig Dis*. 2013 Aug; 14(8):409–16. doi: [10.1111/1751-2980.12063](https://doi.org/10.1111/1751-2980.12063) PMID: [23615046](https://pubmed.ncbi.nlm.nih.gov/23615046/)
4. Yang ZY, Wu XY, Huang YF, Di MY, Zheng DY, Chen JZ, et al. Promising biomarkers for predicting the outcomes of patients with KRAS wild-type metastatic colorectal cancer treated with anti-epidermal growth factor receptor monoclonal antibodies: a systematic review with meta-analysis. *Int J Cancer*. 2013 Oct 15; 133(8):1914–25. doi: [10.1002/ijc.28153](https://doi.org/10.1002/ijc.28153) PMID: [23494461](https://pubmed.ncbi.nlm.nih.gov/23494461/)
5. Safaee Ardekani G, Jafarnejad SM, Tan L, Saeedi A, Li G. The prognostic value of BRAF mutation in colorectal cancer and melanoma: a systematic review and meta-analysis. *PloS One*. 2012; 7(10): e47054. doi: [10.1371/journal.pone.0047054](https://doi.org/10.1371/journal.pone.0047054) PMID: [23056577](https://pubmed.ncbi.nlm.nih.gov/23056577/)
6. Roth AD, Delorenzi M, Tejpar S, Yan P, Klingbiel D, Fiocca R, et al. Integrated analysis of molecular and clinical prognostic factors in stage II/III colon cancer. *J Natl Cancer Inst*. 2012 Nov 7; 104(21):1635–46. doi: [10.1093/jnci/djs427](https://doi.org/10.1093/jnci/djs427) PMID: [23104212](https://pubmed.ncbi.nlm.nih.gov/23104212/)
7. Gavin PG, Colangelo LH, Fumagalli D, Tanaka N, Remillard MY, Yothers G, et al. Mutation profiling and microsatellite instability in stage II and III colon cancer: an assessment of their prognostic and oxaliplatin predictive value. *Clin Cancer Res*. 2012 Dec 1; 18(23):6531–41. doi: [10.1158/1078-0432.CCR-12-0605](https://doi.org/10.1158/1078-0432.CCR-12-0605) PMID: [23045248](https://pubmed.ncbi.nlm.nih.gov/23045248/)
8. Ogino S, Liao X, Imamura Y, Yamauchi M, McCleary NJ, Ng K, et al. Predictive and prognostic analysis of PIK3CA mutation in stage III colon cancer intergroup trial. *J Natl Cancer Inst*. 2013 Dec 4; 105(23):1789–98. doi: [10.1093/jnci/djt298](https://doi.org/10.1093/jnci/djt298) PMID: [24231454](https://pubmed.ncbi.nlm.nih.gov/24231454/)
9. André T, de Gramont A, Vernerey D, Chibaudel B, Bonnetain F, Tijeras-Raballand A, et al. Adjuvant Fluorouracil, Leucovorin, and Oxaliplatin in Stage II to III Colon Cancer: Updated 10-Year Survival and Outcomes According to BRAF Mutation and Mismatch Repair Status of the MOSAIC Study. *J Clin Oncol*. 2015 Dec 10; 33(35):4176–87. doi: [10.1200/JCO.2015.63.4238](https://doi.org/10.1200/JCO.2015.63.4238) PMID: [26527776](https://pubmed.ncbi.nlm.nih.gov/26527776/)
10. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009; 6: e1000097. doi: [10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097) PMID: [19621072](https://pubmed.ncbi.nlm.nih.gov/19621072/)
11. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986 Sep; 7(3):177–88. PMID: [3802833](https://pubmed.ncbi.nlm.nih.gov/3802833/)
12. French AJ, Sargent DJ, Burgart LJ, Foster NR, Kabat BF, Goldberg R, et al. Prognostic significance of defective mismatch repair and BRAF V600E in patients with colon cancer. *Clin Cancer Res*. 2008; 14(11):3408–15. doi: [10.1158/1078-0432.CCR-07-1489](https://doi.org/10.1158/1078-0432.CCR-07-1489) PMID: [18519771](https://pubmed.ncbi.nlm.nih.gov/18519771/)
13. Pentheroudakis G, Raptou G, Kotoula V, Wirtz RM, Vrettou E, Karavasilis V, et al. Immune response gene expression in colorectal cancer carries distinct prognostic implications according to tissue, stage and site: a prospective retrospective translational study in the context of a hellenic cooperative oncology group randomised trial. *PLoS One*. 2015 May 13; 10(5):e0124612. doi: [10.1371/journal.pone.0124612](https://doi.org/10.1371/journal.pone.0124612) PMID: [25970543](https://pubmed.ncbi.nlm.nih.gov/25970543/)
14. Pakneshan S, Salajegheh A, Smith RA, Lam AK. Clinicopathological relevance of BRAF mutations in human cancer. *Pathology*. 2013; 45(4):346–56. doi: [10.1097/PAT.0b013e328360b61d](https://doi.org/10.1097/PAT.0b013e328360b61d) PMID: [23594689](https://pubmed.ncbi.nlm.nih.gov/23594689/)
15. Zheng G, Tseng LH, Chen G, Haley L, Illei P, Gocke CD, et al. Clinical detection and categorization of uncommon and concomitant mutations involving BRAF. *BMC Cancer* 2015 Oct 24; 15:779. doi: [10.1186/s12885-015-1811-y](https://doi.org/10.1186/s12885-015-1811-y) PMID: [26498038](https://pubmed.ncbi.nlm.nih.gov/26498038/)
16. Cremolini C., Di Bartolomeo M., Amatu A., Antoniotti C., Moretto R., Berenato R., et al. BRAF codons 594 and 596 mutations identify a new molecular subtype of metastatic colorectal cancer at favorable prognosis. *Ann Oncol*. 2015 Oct; 26(10):2092–7. doi: [10.1093/annonc/mdv290](https://doi.org/10.1093/annonc/mdv290) PMID: [26153495](https://pubmed.ncbi.nlm.nih.gov/26153495/)
17. Chen D, Huang JF, Liu K, Zhang LQ, Yang Z, Chuai ZR, et al. BRAFV600E Mutation and Its Association with Clinicopathological Features of Colorectal Cancer: A Systematic Review and Meta-Analysis. *PLoS One*. 2014 Mar 3; 9(3):e90607. doi: [10.1371/journal.pone.0090607](https://doi.org/10.1371/journal.pone.0090607) PMID: [24594804](https://pubmed.ncbi.nlm.nih.gov/24594804/)
18. Loupakis F, Moretto R, Aprile G, Muntoni M, Cremolini C, Iacono D, et al. Clinico-pathological nomogram for predicting BRAF mutational status of metastatic colorectal cancer. *Br J Cancer*. 2016 Jan 12; 114(1):30–6. doi: [10.1038/bjc.2015.399](https://doi.org/10.1038/bjc.2015.399) PMID: [26575603](https://pubmed.ncbi.nlm.nih.gov/26575603/)
19. Therkildsen C, Bergmann TK, Henrichsen-Schnack T, Ladelund S, Nilbert M. The predictive value of KRAS, NRAS, BRAF, PIK3CA and PTEN for anti-EGFR treatment in metastatic colorectal cancer: A

systematic review and meta-analysis. *Acta Oncol.* 2014; 53(7):852–64. doi: [10.3109/0284186X.2014.895036](https://doi.org/10.3109/0284186X.2014.895036) PMID: [24666267](https://pubmed.ncbi.nlm.nih.gov/24666267/)

20. Cui D, Cao D, Yang Y, Qiu M, Huang Y, Yi C. Effect of BRAF V600E mutation on tumor response of anti-EGFR monoclonal antibodies for first-line metastatic colorectal cancer treatment: a meta-analysis of randomized studies. *Mol Biol Rep.* 2014; 41(3):1291–8. doi: [10.1007/s11033-013-2974-8](https://doi.org/10.1007/s11033-013-2974-8) PMID: [24390240](https://pubmed.ncbi.nlm.nih.gov/24390240/)
21. Alberts SR, Sargent DJ, Nair S, Mahoney MR, Mooney M, Thibodeau SN, et al. Effect of oxaliplatin, fluorouracil, and leucovorin with or without cetuximab on survival among patients with resected stage III colon cancer: a randomized trial. *Jama.* 2012; 307(13):1383–93. doi: [10.1001/jama.2012.385](https://doi.org/10.1001/jama.2012.385) PMID: [22474202](https://pubmed.ncbi.nlm.nih.gov/22474202/)
22. Ogino S, Shima K, Meyerhardt JA, McCleary NJ, Ng K, Hollis D, et al. Predictive and prognostic roles of BRAF mutation in stage III colon cancer: results from intergroup trial CALGB 89803. *Clin Cancer Res.* 2012; 18(3):890–900. doi: [10.1158/1078-0432.CCR-11-2246](https://doi.org/10.1158/1078-0432.CCR-11-2246) PMID: [22147942](https://pubmed.ncbi.nlm.nih.gov/22147942/)
23. Rowland A, Dias MM, Wiese MD, Kichenadasse G, McKinnon RA, Karapetis CS, et al. Meta-analysis of BRAF mutation as a predictive biomarker of benefit from anti-EGFR monoclonal antibody therapy for RAS wild-type metastatic colorectal cancer. *Br J Cancer.* 2015 Jun 9; 112(12):1888–94. doi: [10.1038/bjc.2015.173](https://doi.org/10.1038/bjc.2015.173) PMID: [25989278](https://pubmed.ncbi.nlm.nih.gov/25989278/)
24. Pietrantonio F, Petrelli F, Coinu A, Di Bartolomeo M, Borgonovo K, Maggi C, et al. Predictive role of BRAF mutations in patients with advanced colorectal cancer receiving cetuximab and panitumumab: a meta-analysis. *Eur J Cancer.* 2015; 51(5):587–94. doi: [10.1016/j.ejca.2015.01.054](https://doi.org/10.1016/j.ejca.2015.01.054) PMID: [25673558](https://pubmed.ncbi.nlm.nih.gov/25673558/)
25. Cremolini C, Loupakis F, Antoniotti C, Lupi C, Sensi E, Lonardi S, et al. FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. *Lancet Oncol.* 2015 Oct; 16(13):1306–15. doi: [10.1016/S1470-2045\(15\)00122-9](https://doi.org/10.1016/S1470-2045(15)00122-9) PMID: [26338525](https://pubmed.ncbi.nlm.nih.gov/26338525/)