

RESEARCH

Open Access



Preferential utilization and colonization of keratin baits by different myco-keratinophiles

Sandeep Kotwal* and Geeta Sumbali

Abstract

Myc-keratinophilic species have a predilection for different keratinous substrates but show variability in their affinity towards them. Keeping this in view, a survey was conducted in the Khardung and Khardung La soils of Ladakh (India) and 28 myco-keratinophilic species belonging to 15 fungal genera (*Sarocladium*, *Aspergillus*, *Beauveria*, *Chrysosporium*, *Cladosporium*, *Alternaria*, *Epicoccum*, *Fusarium*, *Gibberella*, *Clonostachys*, *Paecilomyces*, *Purpureocillium*, *Metarhizium*, *Penicillium* and *Sagenomella*) were isolated by using keratin bait technique. These isolated species were tested for their preferential utilization ability and colonization on different baits by morphological assessment. Different types of keratin baits used were feathers, human hair, human nails and wool. Overall assessment revealed that feathers were colonized and utilized by all the species (100 %), followed in decreasing order by nails (89.29 %), hair (85.71 %) and sheep wool (67.86 %). So, it is concluded that feather baiting technique, could be more useful in trapping keratinophilic fungi than the hair baiting technique which is till date regarded as the best method for the isolation of myco-keratinophiles. On the basis of succession on keratinous baits, the recovered keratinophilic species were also categorized into four categories: early successional species (pioneer colonizers), late successional species (final colonizers), persistent species and no-pattern species.

Keywords: Feathers, Keratinophilic fungi, Keratin baits, Preferential utilization

Background

Keratinophilic fungi are considered as an ecologically important group of highly specialized fungi, which are adapted to the utilization of keratin as the main or sole source of nutrition (Kunert 2000). This unique fungal group is attracting lot of attention throughout the world because of their ability to degrade hard keratin, which is otherwise resistant to degradation by most of the other microorganisms (Filipello 2000). In nature, they exist as self sufficient saprophytes as long as environmental conditions are favourable but they may become parasitic by accident and then pathogenic. Soils rich in keratinous material are found to be most conducive for the growth and occurrence of keratinophilic fungi (Otcenasek 1978; Mercantini et al. 1980). But occurrence of keratinophilic fungi also depends upon various genomic and climatic factors such as organic matter, soil humidity, pH, temperature, soil texture, depth of soil profile and other

microorganisms (Srivastava et al. 1990). For isolation of keratinophilic fungi from soil, keratin or hair baiting technique given by Vanbreuseghem (1952) is widely used. As per this procedure, different keratinous substrates are used as a bait to lure keratinophilic species. However, different species vary in their preference for colonization and utilization of these baits. Keeping this in view, an experiment was carried out to find out the best keratin baits for trapping these fungi and to know the successional pattern of the isolated species.

Methods

Keratin rich substrates found abundantly in nature were used as baits. These included feathers, human hair, human nails and sheep wool. Collected baits were thoroughly washed with water, air dried at room temperature and then cut into small pieces. Finally, they were washed with 70 % alcohol, air dried and sterilized by autoclaving at 15 lbs./sq inch for 20 min. Petridishes were half-filled with soil and sterilized in an oven at 180 °C for 4–5 h on three successive days. Thereafter, 7 days old culture of the test

*Correspondence: s.kotwal@yahoo.com
Department of Botany, University of Jammu, Jammu 180 006, India

fungus growing in a test tube was scrapped in 5 ml sterile water and added to the sterile soil contained in the Petridish inoculated with test culture. These were incubated at 28 ± 2 °C for 20 days and at the end of this period, degree of colonization and preferential utilization of the baits by keratinophilic fungal species was recorded. Morphological appearance of colonized and invaded keratin baits was examined under light microscopy using direct samples.

Results and discussion

After incubation period, the keratin baits were visually examined for the growth of keratinophilic fungal species and the results are presented in Table 1. Perusal of data shows that keratinophiles differ in their substrate preferences for colonization. Out of the four different keratinous baits used, feathers were colonized by all the

recovered keratinophilic species (28), human nails by 25 fungal species, human hair by 24 fungal species and sheep wool by 19 fungal species (Table 1). Earlier, Sundaram (1987) also reported sheep wool as a poor bait in comparison to bird feathers and human hair.

In the present investigation, substrate specificity was found to be variable within the species of the same genus also. Among the recovered *Chrysosporium* species, *C. merdarium* and *C. queenslandicum* showed luxuriant growth and maximum degradation of all the keratinous baits (Fig. 1), whereas *Chrysosporium* anamorph of *Gymnoascus demonbreunii* and *C. inops* showed moderate growth on bird feathers and human nails but slow growth on human hair and sheep wool. The ability of *Chrysosporium* species to colonize all the keratinous substrates confirmed the cosmopolitan nature of this genus.

Table 1 Preferential colonization of keratin baits by myco-keratinophiles

Myco-keratinophilic species	Keratin baits used			
	Feathers	Human hair	Human nails	Sheep wool
<i>Sarocladium bacillisporum</i>	++	+	++	+
<i>S. implicatum</i>	+++	+++	+++	++
<i>Aspergillus flavus</i>	+++	++	+++	+
<i>A. parasiticus</i>	+++	++	++	+
<i>A. sydowii</i>	++	++	+++	–
<i>A. ustus</i>	++	++	+++	+
<i>A. wentii</i>	+	+	–	–
<i>Beauveria bassiana</i>	+++	++	+++	+
<i>Chrysosporium inops</i>	++	+	++	+
<i>C. merdarium</i>	+++	+++	+++	++
<i>C. queenslandicum</i>	+++	+++	+++	++
<i>Chrysosporium</i> anamorph of <i>Gymnoascus demonbreunii</i>	++	+	++	+
<i>Cladosporium cladosporioides</i>	++	+	++	+
<i>Alternaria chlamydosporigena</i>	++	–	++	+
<i>Aspergillus stellatus</i>	++	+	–	–
<i>Epicoccum nigrum</i>	+++	++	+	–
<i>Fusarium oxysporum</i>	+++	+++	++	+
<i>F. sporotrichioides</i>	+++	++	+++	+
<i>F. trichothecioides</i>	++	–	+	–
<i>F. incarnatum</i>	++	–	++	–
<i>Gibberella fujikuroi</i>	++	+	++	+
<i>Clonostachys rosea</i>	++	++	++	+
<i>Purpureocillium lilacinum</i>	+++	+++	++	+
<i>Metarhizium marquandii</i>	+++	++	+++	+
<i>Paecilomyces divaricatus</i>	+++	++	++	+
<i>Penicillium brevicompactum</i>	+++	++	+	–
<i>P. griseofulvum</i>	++	+	–	–
<i>Sagenomella alba</i>	++	–	+	–
Total number of fungal species colonizing individual baits	28	24	25	19

+++ , excellent growth; ++ , medium growth; + , slow growth; – , no growth




Fig. 1 Differential colonizing ability of **a** *Sarocladium bacillisporum*, **b** *Sarocladium implicatum*, **c** *Aspergillus parasiticus*, **d** *Beauveria bassiana*, **e** *Chrysosporium merdarium*, **f** *Chrysosporium queenslandicum*, **g** *Alternaria chlamydosporigena*, **h** *Epicoccum nigrum*, **i** *Fusarium oxysporum*, **j** *Fusarium sporotrichioides*, **k** *Clonostachys rosea*, **l** *Metarhizium marquandii*

Since, all the *Chrysosporium* species were observed to be growing either luxuriantly, moderately or slowly on various keratinous residues, it is probable that they possess specific enzymatic system for degradation of keratin and it is, therefore, important to recognize their potential as opportunistic pathogens. Moreover, these fungi by virtue of their ability to colonize and degrade various keratinous substrates are able to add carbon, nitrogen and sulphur content to the soil.

Among *Sarocladium* species, *S. implicatum* showed excellent growth on feathers, human hair and human nails but medium growth on sheep wool (Table 1), whereas *S. bacillisporum* showed medium growth on feathers and human nails but slow growth on human hair and sheep wool (Fig. 1).

Different species of *Aspergillus* also depicted keratin specificity. Among the recovered species of *Aspergillus*, *A. flavus* and *A. parasiticus* showed excellent growth on feathers and human nails but showed moderate growth on human hair and slow growth on sheep wool (Fig. 1). On the other hand, *A. sydowii* and *A. ustus* grew luxuriantly on human nails and showed moderate growth on feathers and human hair (Fig. 2). However, *A. ustus* showed less growth on sheep wool, whereas *A. sydowii* showed no preference for this bait. *A. wentii* and *A. stellatus*, which showed low frequency in Khardung soils could not grow on nails and wool but were slow colonizers of feathers and hair.

Among *Fusarium* species, *F. oxysporum* and *F. sporotrichioides* were excellent colonizers of feathers, nails and hair but were poor colonizers of sheep wool (Fig. 1), whereas two, other species, *F. incarnatum* and *F. trichothecioides* showed moderate growth on feathers and nails and no preference for human hair and sheep wool. *Gibberella fujikuroi* showed moderate to low growth on all the investigated keratin baits.

Purpureocillium lilacinum, *Metarhizium marquandii* and *Paecilomyces divaricatus*, also showed more preference for feathers than for human nails and hair and were poor colonizers of sheep wool (Table 1). Similarly, *Penicillium* species also showed differential colonizing ability on varied baits. *P. brevicompactum* could grow luxuriantly on feathers, moderately on human hair, sparsely on human nails but failed to grow on sheep wool. *P. griseofulvum* grew moderately on feathers, sparsely on human hair but could not utilize human nails and sheep wool (Table 1). *Clonostachys rosea* usually showed moderate preference for most of the keratinous baits that were tested (Fig. 1).

Other isolated keratinophilic species, viz., *Beauveria bassiana*, *Cladosporium cladosporioides*, *Alternaria chlamydosporigena*, *Epicoccum nigrum* and *Sagenomella alba* also showed luxuriant growth on feathers and its maximum degradation (Figs. 1, 2). Among these, *Alternaria chlamydosporigena* and *Sagenomella alba* were not able

to grow on human hair, whereas *Epicoccum nigrum* and *Sagenomella alba* were not able to colonize sheep wool.

The ability of most of the recovered keratinophilic fungi to grow and hydrolyze feather keratin more efficiently than other baits suggests their use in isolating this group of fungi from the soil rather than using the earlier hair baiting technique of Vanbreuseghem (1952). Earlier Pugh (1971), Jain and Agrawal (1980) and Kaul and Sumbali (1994) also observed preferential utilization and maximum degradation of feathers by most of the isolated keratinophiles. Further, moderate to low growth of some of the recovered myco-keratinophiles on different baits suggests that they have low keratinase producing ability or it may be due to the biochemical differences existing in the keratin of various ectodermal appendages. Similar observations have been recorded by Kunert (2000), who found that in case of hard keratin (α -keratin), the rate of hydrolysis corresponds roughly to hardness, that is, cystine content and disulphide bonds. Therefore, since feathers and nails have lesser cystine content and thus few disulphide bonds, they are more easily cleaved by keratinophilic fungi than human hair and sheep wool.

Excellent colonization of all the keratinous substrates was shown mostly by *Chrysosporium* species and *Sarocladium implicatum*, indicating that they possess specific enzymatic system for degradation of keratin. Therefore, it is important to recognize their potential as pathogens. Substrate specificity was also found to be variable within the species of the same genus.

On the basis of their succession on keratinous baits the, keratinophilic mycobiota were categorized into four categories.

- (1) Early successional species (pioneer colonizers): Species which appear only at the beginning (within 15 days after soil was baited with keratin-baits) included *Sarocladium implicatum*, *Aspergillus parasiticus*, *A. sydowii*, *A. ustus*, *A. stellatus*, *Beauveria bassiana*, *Fusarium oxysporum*, *Clonostachys rosea* and *Penicillium griseofulvum*.
- (2) Late successional species (final colonizers): Species which appear after 15 days of the incubation period included *Chrysosporium merdarium*, *C. queenslandicum*, *C. inops* and *Chrysosporium* anamorph of *Gymnoascus demonbreunii*.
- (3) Persistent species: Species which are present persistently on the keratinous baits e.g., *Aspergillus flavus*, *Fusarium sporotrichioides*, *Purpureocillium lilacinum*, *Metarhizium marquandii*, *Paecilomyces divaricatus*, and *Penicillium brevicompactum*.
- (4) No-pattern species:




Fig. 2 showing colonization of various keratinous baits by some keratinophilic fungal species. **a** Conidiophores of *Aspergillus ustus* arising from a hair shaft. **b** Sporogenous cells of *Beauveria bassiana* arising from the colonized sheep wool. **c** Degradation of feather by the growth of *Chrysosporium merdarium*. **d** Conidiophores and conidia of *Epicoccum nigrum* arising from the slightly damaged hair shaft. **e** Hair shaft colonized by cleistothecia of *Aspergillus stellatus*. **f** Mycelium and phialides of *Fusarium sporotrichioides* arising from the colonized hair shaft. **g** Thick growth of conidiophores of *Clonostachys rosea* arising from the slightly damaged hair shaft. **h** Typical conidiophores of *Purpureocillium lilacinum* arising from the colonized hair shaft

Species that did not seem to have a clear successional pattern. These included *Sarocladium bacillisporum*, *Aspergillus wentii*, *Cladosporium cladosporioides*,

Alternaria chlamydosporigena, *Epicoccum nigrum*, *Fusarium trichothecioides*, *F. incarnatum* and *Sagenomella alba*.

Similar results were obtained by Ali-Shtayeh and Jamous (2000) while working on the succession of human hair by keratinophilic mycobiota of the soil. As observed during this investigation, Ali-Shtayeh and Jamous (2000) also found *Beauveria bassiana* and *Penicillium griseofulvum* as the early pioneer colonizers. In addition, as noted in the present investigation, even De Vries (1952) observed that the final fungal colonizers of keratinous baits were the typical keratinophilic hyphomycetes such as species of *Chrysosporium*.

Overall assessment of the recovered myco-keratinophiles for their growth on four different keratinous baits revealed that feathers were colonized and utilized by all the species (100 %), followed in decreasing order by nails (89.29 %), hair (85.71 %) and sheep wool (67.86 %). Similar observations have been recorded by Kunert (2000) who found that the rate and completeness of the degradation is dependent on the kind of substrate and correspond roughly to its hardness, that is, cystine content. Therefore, human, dog, horse and cattle hairs are attacked more slowly than the feathers of birds.

Conclusions

In view of these observations, it is concluded that hair baiting technique as given by Vanbreuseghem (1952) and recorded by many other workers (Sundaram 1987; Gugnani et al. 2012; Pakshir et al. 2013; Sarkar et al. 2014; Soleymani et al. 2015; Sharma and Choudhary 2015) as the best method for the isolation of myco-keratinophiles is not very true. Instead, as observed during the present investigation, feather baits, which could allow the growth of all the recovered keratinophiles are more useful in trapping this unique group.

It was also recorded that the isolated myco-keratinophiles showed a second preference for nails, which indicates that they possess an ability of efficiently hydrolyzing the nail keratin. Therefore, they may pose a potential threat for onychomycosis, particularly among the farmers, gardeners, children and old people.

Authors' contributions

Both the authors were involved in designing the study, experimentation, data collection, data analysis, manuscript preparation, typing, reading and final approval. Both authors read and approved the final manuscript.

Acknowledgements

The authors are thankful to National Fungal Culture Collection of India (NFCCI) Agharkar Research Institute (ARI), Pune, for confirming the identity of some of the keratinophilic fungal species.

Competing interests

The authors declare that they have no competing interests.

Received: 17 December 2015 Accepted: 19 July 2016

Published online: 28 July 2016

References

- Ali-Shtayeh MS, Jamous RMF (2000) Keratinophilic fungi and related dermatophytes in polluted soil and water habitats. In: Kushwaha RKS, Guarro J (eds) *Biology of dermatophytes and other keratinophilic fungi*. Revista Iberoamericana de Micología, Elsevier, Apdo, pp 51–59
- De Vries GA (1952) Contribution to the knowledge of the genus *Cladosporium*. Dissertation, University of Utrecht
- Filipello MV (2000) Keratinophilic fungi: their role in nature and degradation of keratinic substrates. In: Kushwaha RKS, Guarro J (eds) *Biology of dermatophytes and other keratinophilic fungi*. Revista Iberoamericana de Micología, Elsevier, Apdo, pp 86–92
- Gugnani HC, Sharma S, Gupta B, Gaddam S (2012) Prevalence of keratinophilic fungi in soils of St. Kitts and Nevis. *J Inf Dev Ctries* 6(4):347–351
- Jain PC, Agrawal SC (1980) A note on the keratin decomposing capability of some fungi. *Trans Mycol Soc Jpn* 21:513–517
- Kaul S, Sumbali G (1994) Prevalence of soil borne keratin degrading fungi in dairy farm soils of Jammu (J&K). *Geobios* 21:54–59
- Kunert J (2000) Physiology of keratinophilic fungi. In: Kushwaha RKS, Guarro J (eds) *Biology of dermatophytes and other keratinophilic fungi*. Revista Iberoamericana de Micología, Spain, pp 77–85
- Mercantini R, Marsella R, Caprilli F, Dovigiallo G (1980) Isolation of dermatophytes and correlated species from the soil of public gardens and parks in Rome. *Sabouraudia* 18:123–128
- Otcenasek M (1978) Ecology of the dermatophytes. *Mycopathologia* 65:67–72
- Pakshir K, Ghiasi MR, Zomorodian K, Gharavi AR (2013) Isolation and molecular identification of keratinophilic fungi from public parks soil in Shiraz, Iran. *BioMed Res Int* 2013:619576-1–619576-5
- Pugh GJF (1971) Factors which influence the early colonization of organic matter by fungi. In: IVth international colloquium for soil. Zoology Instiut National de la Recherche Agrenomique Paris, pp 319–327
- Sarkar AK, Ravi V, Gupta AK (2014) Incidence of keratinophilic fungi in areas of Raipur city, Chhattisgarh region, India. *Afr J Microbiol Res* 8(3):264–269
- Sharma R, Choudhary N (2015) Isolation of keratinophilic fungi from soils samples of agricultural fields of Saharanpur (U.P.), India. *Int J Curr Microbiol Appl Sci* 4(7):229–237
- Soleymani A, Sefidgar SAA, Hoseini M, Sharifi H (2015) Species diversity of keratinophilic fungi in various soil type of Babol Medical University's Hospitals' Yard. *Int J Appl Sci Technol* 5(3):55–59
- Srivastava OP, Jain M, Shukla PK (1990) A critical review on keratinophilic fungi in perspectives. In: Hasiji SK, Bilgrami HS (eds) *Prof. G.P. Agarwal Festschrift: Mycological research II*. Today and Tomorrow's Printers and Publishers, New Delhi, pp 269–292
- Sundaram BM (1987) Fungal flora of rice field soils. *Proc Natl Acad Sci* 87B:90–99
- Vanbreuseghem R (1952) Technique biologique pour one isolament des dermatophytes du sol (biological technique for the isolation of dermatophytes from the soil). *Ann Soc Belgs Med Trop* 32:173–178

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com