



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Key Highlights of the Canadian Thoracic Society's Position Statement on the Optimization of COPD Management During the Coronavirus Disease 2019 Pandemic



Mohit Bhutani, MD
Edmonton, AB, Canada
Paul Hernandez, MD
Halifax, NS, Canada
Jean Bourbeau, MD
Montreal, QC, Canada
Gail Dechman, PhD
Halifax, NS, Canada
Erika Penz, MD
Saskatoon, SK, Canada
Raymond Acheron, MN
Edmonton, AB, Canada
Marla Beauchamp, PhD
Joshua Wald, MD
Hamilton, ON, Canada
Michael Stickland, PhD
Edmonton, AB, Canada
Sharla-Rae Olsen, MD
Prince George, BC, Canada
Donna Goodridge, PhD
Saskatoon, SK, Canada

Patients living with COPD represent a vulnerable population during the coronavirus disease 2019 (COVID-19). Physicians and patients have many questions regarding the acute and chronic management of COPD during the pandemic. This commentary summarizes the Canadian Thoracic Society's (CTS) position statement on managing COPD during the COVID-19 pandemic¹ in an easy FAQ format. The full COPD position statement, and other valuable clinical tools including links to online patient support programs for self-management and exercise/pulmonary rehabilitation, can be found online (<https://cts-sct.ca/covid-19/>).

General Recommendations for All Patients With COPD

Patients with COPD should stay at home as much as possible, including working from home, if feasible. If you must leave the home, we suggest that all patients follow current local, national, and global public health advisories with respect to the indications for physical distancing and isolation. Patients should wash their hands with soap and water frequently for 20 s or use alcohol-based hand sanitizer containing at least 60% alcohol. Consider having at least a 30-day supply of all medications on hand to reduce the need for leaving the home, or select delivery options at your pharmacy, or have trusted individuals pick up your medication.

Longitudinal experience with seasonal influenza and preliminary data in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection (subsequently discussed) suggests that patients with chronic lung disease are at risk for severe complications of SARS-CoV-2 infection (COVID-19). Physical distancing is an important public health measure to flatten the curve of community spread of the virus. The workplace is a social environment which may expose patients to others in their community, particularly if physical distancing in the workplace setting is difficult to implement. There is also increasing concern regarding the challenges of implementing physical distancing measures in congregated living situations such as retirement homes and long-term care facilities that need to be urgently addressed by health-care providers and health systems to limit the spread of the virus in this vulnerable population. Until we fully understand the risks associated with

ABBREVIATIONS: COVID-19 = coronavirus disease 2019; CTS = Canadian Thoracic Society; ICS = inhaled corticosteroid; LABA = long-acting beta-2-agonist; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2

AFFILIATIONS: From the Division of Pulmonary Medicine, Department of Medicine (Dr Bhutani), University of Alberta; the Department of Medicine (Dr Hernandez), Dalhousie University; the Research Institute of the McGill University Health Centre (Dr Bourbeau), McGill University; the School of Physiotherapy (Dr Dechman), Dalhousie University; the Respiratory Research Centre (Dr Penz), University of Saskatchewan; the Faculty of Nursing (Mr Acheron), University of Alberta; the School of Rehabilitation Science (Dr Beauchamp), McMaster University;

SARS-CoV-2 infection in patients with chronic lung disease, we have placed a high value on limiting exposure based on prior experience with influenza.

Patients living with COPD, particularly those with severe disease and/or if associated with advanced frailty, should complete or update their advance care plans and indicate if they would accept transfer to hospital or admission to critical care (eg, for mechanical ventilation) in the event of a severe COVID-19-related illness. If patients lack the capacity to complete this document, we recommend that a discussion with the patient's most responsible designate occur to establish goals of care.

the Department of Medicine (Dr Wald), McMaster University; the Division of Pulmonary Medicine (Dr Stickland), Department of Medicine, University of Alberta; the Department of Medicine (Dr Olsen), Heritage Medical Centre; and the Respiratory Research Centre (Dr Goodridge), University of Saskatchewan.

FINANCIAL/NONFINANCIAL DISCLOSURES: The authors have reported to *CHEST* the following: M. Bhutani reports personal fees and grants outside the submitted work from AstraZeneca Canada Ltd, Boehringer Ingelheim Canada Ltd, GlaxoSmithKline Canada Ltd, Novartis, Sanofi-Genzyme, the Canadian Institute Health Research, CHEST, The Lung Association of Alberta, The University of Alberta Hospital Foundation, and Alberta Innovates Health Solutions. P. H. reports grants from the Canadian Institute Health Research, the Lung Association of Nova Scotia, the Nova Scotia Health Authority Research Fund, Astra Zeneca Canada Ltd, Boehringer Ingelheim Canada Ltd, Cyclomedica, Grifols, Respivant, and Vertex; and personal fees from Actelion, Astra Zeneca Canada Ltd, Boehringer Ingelheim Canada Ltd, GlaxoSmithKline Canada Ltd, Novartis, Sanofi-Aventis, and Trudell, outside the submitted work. J. B. reports grants from the Canadian Respiratory Research Network, the Canadian Institute Health Research, and the Respiratory Health Network of the Fonds de la recherche en santé du Québec; grants and personal fees from Astra Zeneca Canada Ltd, Boehringer Ingelheim Canada Ltd, GlaxoSmithKline Canada Ltd, Grifols, Novartis, and Trudell; and personal fees from the Canadian Thoracic Society and CHEST, outside the submitted work. E. P. reports personal fees from Boehringer Ingelheim Canada Ltd, Astra Zeneca Canada Ltd, GlaxoSmithKline Canada Ltd, and Novartis; and grants from the Canadian Institute Health Research, the Saskatchewan Research Foundation, the Respiratory Research Centre, and Astra Zeneca Canada Ltd, outside the submitted work. J. W. reports personal fees from GlaxoSmithKline Canada Ltd and a grant from Fisher & Paykel, outside the submitted work. M. S. reports investigator-initiated funding from the Canadian Institute Health Research, the Natural Sciences and Engineering Research Council of Canada, the Canadian Respiratory Research Network, the Canadian Foundation of Innovation, Alberta Innovates Health Solutions, the Lung Association of Alberta, and the University of Alberta Hospital Foundation/Alberta Boehringer Ingelheim Collaboration. D. G. reports grants from the Canadian Institute Health Research, the Centre for Aging and Brain Health Innovation, the Saskatchewan Health Research Foundation, the Royal University Hospital Foundation, the Lung Association of Saskatchewan, the Canadian Network for the Prevention of Elder Abuse, the Respiratory Research Centre, and the College of Medicine, University of Saskatchewan, outside the submitted work. None declared (G. D., R. A., M. Beauchamp, S.-R., O.).

CORRESPONDENCE TO: Mohit Bhutani, MD, Division of Pulmonary Medicine, Department of Medicine, University of Alberta, 3rd Floor, Clinical Sciences Bldg, 11350-83 Ave, Edmonton, AB, T6G 2G3, Canada; e-mail: mohit.bhutani@ualberta.ca

Copyright © 2020 American College of Chest Physicians. Published by Elsevier Inc. All rights reserved.

DOI: <https://doi.org/10.1016/j.chest.2020.05.530>

Are Patients With COPD at Higher Risk of Acquiring SARS-CoV-2 Infection?

Viral respiratory tract infections are a common cause of COPD exacerbations.² However, according to current available data, patients with COPD are not at an increased risk of acquiring SARS-CoV-2 infection. Published literature on the clinical characteristics of patients admitted to hospital for a SARS-CoV-2 infection suggest that patients with COPD are not overrepresented when compared with the general population.³⁻⁶

Will Patients With COPD Have More Severe Symptoms or Disease Course Because of COVID-19?

Patients with COPD hospitalized because of COVID-19 are more likely to require ICU support and have higher mortality when compared with other patient groups.^{7,8} A meta-analysis by Lippi and Henry⁹ shows that patients with COPD with COVID-19 have over a fivefold risk of having severe disease.

What Should Patients With COPD Do With Their Current Inhaled Therapies During the COVID-19 Pandemic?

We recommend that usual maintenance and exacerbation management for COPD be continued according to current CTS treatment guidelines.¹⁰ Based on what we know about viral respiratory infections in patients with COPD, optimal pharmacologic treatment is the best way to prevent exacerbations and/or reduce the severity of exacerbations. Maintenance inhaled therapies have been shown to improve lung function, symptoms, and quality of life and decrease the risk of future exacerbations, including those precipitated by viral infections.

Is There Any Risk of Using Inhaled Corticosteroids for COPD Treatment During the COVID-19 Pandemic?

There is no evidence that inhaled corticosteroids (ICSs) increase the risk of acquiring SARS-CoV-2 infection or complicate/worsen this infection, such as increasing the need for hospitalization, intubation for mechanical ventilation, or death. This includes patients using their ICSs in combination inhalers with long-acting bronchodilators (ie, long-acting muscarinic antagonist, long-acting beta-2-agonist [LABA]), such as ICS/LABA or ICS/LABA/long-acting muscarinic antagonist combination inhalers. Patients should

continue their maintenance and exacerbation management for COPD according to CTS treatment guidelines.^{10,11}

Are Systemic Corticosteroids Safe to Use in Acute Exacerbations of COPD Caused by SARS-CoV-2 Infections?

This question needs careful consideration because we need to separate the use of systemic corticosteroids for treatment of acute exacerbations of COPD from the use of systemic corticosteroids in a more general setting of COVID-19.

In the absence of evidence of harm and an expectation of a low risk of harm, we prioritized the high value of current evidence-based recommendations to treat acute exacerbations of COPD with prednisone to reduce the need for urgent health service utilization. The high value to reduce acute care utilization supersedes the low risk of concern that prednisone may prolong viral replication.¹

It remains unclear as to whether systemic steroids (such as prednisone) are helpful or harmful in the treatment of COVID-19. Most of what we know comes from studies on SARS-CoV-2 acute lung injury. Russel et al¹² reviewed observational data and concluded there was no benefit to using prednisone to treat SARS-CoV-2-related acute lung injury. However, this literature is evolving and the evidence of the benefit/risk of systemic steroids may change.

Is a Nebulizer Safe to Use During the Pandemic?

We do not recommend the use of a nebulizer during the pandemic because there is an increased risk of aerosol spread of virus particles.¹³ Instead, patients should use metered dose inhalers with spacing devices, soft mist inhalers, or dry powder inhalers to administer all COPD medications at home, and inside health-care facilities and nursing homes. Patients who are already using nebulizers at home should continue to do so until they can be switched to alternative delivery methods. However, they should consider nebulizing their medicines in a separate room from others in the house, and implement other infection control recommendations.

Is Self-Management Education, Pulmonary Rehabilitation, and Exercise for Patients With COPD Still Available During the Pandemic?

In-person programs are closed until further notice; however, self-management and pulmonary

rehabilitation counseling can still be done by telephone or tele-health technologies in some institutions.

Patients should remain physically active (eg, daily walks while physical distancing, functional resistance exercises for strength training) and continue their treatment plan (regular medications and self-management using their action plan with additional treatment in the event of an exacerbation).

The CTS COVID-19 webpage (<https://cts-sct.ca/covid-19/>) provides links to online resources that can help facilitate the teaching and implementation of self-management and rehabilitation strategies.

Should Patients With COPD Continue to Use Oxygen at Home?

Yes. Patients who currently are on home oxygen should continue to use it as prescribed. They should follow the manufacturer's instructions for cleaning and maintenance of their equipment. If the patient has had to increase the flow rate, they should inform the physician, and/or case manager. If in extreme distress, they should call emergency medical services.

The COVID-19 pandemic is a rapidly evolving situation. Health-care professionals are advised to monitor the CTS website for additional COPD resources (action plans and tutorial videos for adults for the proper use of inhalers, etc). Updates on COVID-19 and other lung diseases (eg, asthma) and a link to recommendations regarding the clinical management of patients in the event of a salbutamol metered dose inhaler shortage can also be found on this webpage.

References

1. Bhutani M, Hernandez P, Bourbeau J, et al. Addressing therapeutic questions to help Canadian health care professionals optimize COPD management for their patients during the COVID-19 pandemic [published online ahead of print April 30, 2020]. *Can Respir, Crit Care, and Sleep Med*. doi:10.1080/24745332.2020.1754712.
2. Wedzicha JA, Seemungal TA. COPD exacerbations: defining their cause and prevention. *Lancet*. 2007;370(9589):786-796.
3. Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China [published online ahead of print February 19, 2020]. *Allergy*. <http://doi.org/10.1111/all.14238>
4. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-1720.
5. Korean Society of Infectious Diseases; Korean Society of Pediatric Infectious Diseases; Korean Society of Epidemiology; Korean Society for Antimicrobial Therapy; Korean Society for Healthcare-associated Infection Control and Prevention; Korea Centers for Disease Control and Prevention. Report on the epidemiological features of coronavirus disease 2019 (COVID-19) outbreak in the Republic of Korea from January 19 to March 2, 2020. *J Korean Med Sci*. 2020;35(10):e112.

6. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of covid-19 in New York City. *N Engl J Med*. 2020;382:2372-2374.
7. Guan WJ, Liang WH, Zhao Y, et al. Comorbidity and its impact on 1590 patients with covid-19 in China: a nationwide analysis. *Eur Respir J*. 2020;55(5):2000547.
8. Zhao Q, Meng M, Kumar R, et al. The impact of COPD and smoking history on the severity of covid-19: a systemic review and meta-analysis [published online ahead of print April 15, 2020]. *J Med Virol*. <https://doi.org/10.1002/jmv.25889>.
9. Lippi G, Henry BM. Chronic obstructive pulmonary disease is associated with severe coronavirus disease 2019 (COVID-19). *Respir Med*. 2020;167:105941.
10. Bourbeau J, Bhutani M, Hernandez P, et al. Canadian Thoracic Society Clinical Practice Guideline on pharmacotherapy in patients with COPD - 2019 update of evidence. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*. 2019;3(4):210-232.
11. O'Donnell DE, Hernandez P, Kaplan A, et al. Canadian Thoracic Society recommendations for management of COPD - 2008 update - highlights for primary care. *Can Respir J*. 2008;15(suppl A):1A-8A.
12. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet*. 2020;395(10223):473-475.
13. van Doremalen N, Morris DH, Holbrook MG, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020;382(16):1564-1567.