

# Phagocytosis of Microglia in the Central Nervous System Diseases

Ruying Fu · Qingyu Shen · Pengfei Xu · Jin Jun Luo · Yamei Tang

Received: 30 August 2013 / Accepted: 15 December 2013 / Published online: 7 January 2014  
© The Author(s) 2014. This article is published with open access at Springerlink.com

**Abstract** Microglia, the resident macrophages of the central nervous system, rapidly activate in nearly all kinds of neurological diseases. These activated microglia become highly motile, secreting inflammatory cytokines, migrating to the lesion area, and phagocytosing cell debris or damaged neurons. During the past decades, the secretory property and chemotaxis of microglia have been well-studied, while relatively less attention has been paid to microglial phagocytosis. So far there is no obvious concordance with whether it is beneficial or detrimental in tissue repair. This review focuses on phagocytic phenotype of microglia in neurological diseases such as Alzheimer's disease, multiple sclerosis, Parkinson's disease, traumatic brain injury, ischemic and other brain diseases. Microglial morphological characteristics, involved receptors and signaling pathways, distribution variation along with time and space changes, and environmental factors that affecting phagocytic function in each disease are reviewed. Moreover, a comparison of contributions between macrophages from peripheral circulation and the

resident microglia to these pathogenic processes will also be discussed.

**Keywords** Microglia · Phagocytosis · Neurological diseases · Macrophages

## Introduction

Microglia, which constitute 10~20 % of glia cells, are the resident macrophages of the brain and spinal cord and act as the main immune defense in the central nervous system (CNS). Since first described by del Rio-Hortega, microglia have attracted much attention in exploring their multifaceted meticulous functions, especially related to neuronal inflammation. Microglia are inactive under physiological conditions, with small cell body and highly ramified branching processes. In response to injury or pathogen invasion however, microglia transform into active phagocytic microglia [1], migrate, and accumulate at the site of injury through a process known as chemotaxis [2]. Being different to their resting phenotype, activated microglia are identified by their retracted processes and “amoeboid” morphology, release of both pro- and anti-inflammatory molecules, and high capacity for phagocytic removal of apoptotic cells and debris [3, 4]. Activated microglia are observed in nearly all kinds of neurological diseases, including neurodegenerative diseases such as Alzheimer's disease (AD) [5], Parkinson's disease (PD) [6], and amyotrophic lateral sclerosis (ALS) [7]; infectious and inflammatory diseases such as multiple sclerosis (MS) [8, 9]; stroke [10]; and traumatic [11] and radiation-induced brain injury [12].

In recent years, numerous papers have focused on microglial function of chemotaxis and secretion of pro- and anti-inflammatory cytokines, growth factors, chemokines, and neurotrophins; however, studies regarding microglial phagocytosis are rare. Recently, “phagoptosis” has been proposed as

---

Ruying Fu and Qingyu Shen contributed equally to the study.

R. Fu · Q. Shen · P. Xu · Y. Tang (✉)  
Department of Neurology, Sun Yat-Sen Memorial Hospital, Sun Yat-Sen University, Number 107, Yan Jiang Xi Road, Guangzhou 510120, Guangdong Province, China  
e-mail: yameitang@hotmail.com

Q. Shen  
Department of Neurology, Zengcheng People's Hospital, Sun Yat-Sen University, Guangzhou, China

J. J. Luo  
Department of Neurology, School of Medicine, Temple University, Philadelphia, PA, USA

Y. Tang  
Key Laboratory of Malignant Tumor Gene Regulation and Target Therapy of Guangdong Higher Education Institutes, Sun Yat-Sen University, Guangzhou, China

a form of cell death, caused by phagocytosis of viable cells [13]. It is also known as “primary phagocytosis.” It is provoked by exposure of “eat-me” signals and/or loss of “don’t-eat-me” signals by viable cells, causing their phagocytosis by phagocytes and resulting in self-destruction [13]. Phagocytosis is normally exerted by phagocytic cells, such as microglia and macrophages [13]. Whether microglial phagocytosis plays a beneficial or detrimental role in brain diseases remains controversial, though most researchers are in favor of the former claim since efficient clearance of tissue debris is critical in reconstruction and reorganization of neuronal networking after an injury in the brain [14–16]. The crucial beneficial role of microglial phagocytosis in axon regeneration and in restoration of the microenvironment has also been shown during the recovery of an acute brain injury.

In this article, we will review the characteristics of phagocytic microglia in some brain diseases and discuss quantity variation along with time and localization, receptors and following signaling pathways that mediated microglial phagocytosis, and correlative influential factors. In addition, the relative contribution made by resident microglia and hematogenous macrophages during disease processes is also reviewed.

### Receptors Involved in Microglial Phagocytosis


The activity of microglial phagocytosis relies on specific receptors expressed on the cell surface and downstream signaling pathways that contribute to the reorganization of actin

protein and engulfment of harmful microparticles (Fig. 1; Table 1). Microglial phagocytosis may need different types of receptors to initiate function [17]. In general, there are two distinctive types of receptors, one with a high affinity to bind to foreign microbial pathogens, such as Toll-like receptors (TLRs), and another recognizing apoptotic cellular substances, such as triggering receptor expressed on myeloid cells 2 (TREM-2). Besides these two types, some receptors including Fc receptors, complement receptors [18], scavenger receptors (SR), pyrimidinergic receptor P2Y, G-protein coupled, 6 (P2RY6), macrophage antigen complex 2 (MAC-2), mannose receptor [19], and low-density lipoprotein receptor-related protein (LRP) receptor also participate in microglial clearance of misfolded, apoptotic cells and dead neurons in both acute and chronic brain injury [20].

### TLRs

TLRs are a class of proteins that play a key role in the innate immune system and the digestive system. TLRs are single, membrane-spanning, non-catalytic receptors usually expressed in sentinel cells, such as macrophages and dendritic cells, that recognize structurally conserved molecules derived from microbes. Once these microbes have breached physical barriers such as the skin or intestinal tract mucosa, they are recognized by TLRs, which activate immune cell responses. TLR1–9, which belong to interleukin (IL)-1R super-family, expressed exclusively on antigen presenting cells including microglia [21], macrophages, antigen presenting dendritic cells, and

**Fig. 1** Signaling pathways involved in microglial phagocytosis. *A* Extracellular nucleotides, such as UDP and UTP, trigger microglial phagocytosis through P2Y<sub>6</sub>R/PLC/InsP<sub>3</sub> pathway. *B* Apoptotic debris induces phagocytosis via TREM-2/DAP12/ERK/PKC pathway. *C* Endogenous or ectogenic detriments, such as LPS, viral nucleotides,  $\alpha$ -synuclein, and f-A $\beta$ , provoke phagocytosis by microglia via TLRs through activation of MyD88-dependent IRAK4/p38/scavenger receptors pathway or MyD88-independent actin-Cdc42/Rac signaling pathway



**Table 1** Receptors involved in microglial phagocytosis

Receptors	Stimulus	Cell types expressed	Diseases/animal models	References
TLRs	$\alpha$ -Synuclein Fibrillar A $\beta$ Bacterial strains LPS Amyloid protein	Microglia Astrocytes Oligodendrocytes Neurons Microglia Macrophages Monocyte-derived dendritic cells Osteoclasts	alpha-Synucleinopathies, Alzheimer's disease CNS infections, spinal cord injury, and neuronal injury Alzheimer's disease Nasu-Hakola disease Multiple sclerosis/EAE	[23, 120–122] [34, 123]
Fc receptors (Fc $\gamma$ R)	$\alpha$ -Synuclein	Neurons Astrocytes Microglia	Parkinson's disease Multiple sclerosis Alzheimer's disease	[124, 125]
Complement receptors (CR3 (CD11b/CD18), CR4 (CD11c/CD18))	Degenerated myelin A $\beta$ Apoptotic cells	Oligodendrocytes Microglia Macrophages	Alzheimer's disease Wallerian degeneration Multiple sclerosis	[126, 127]
Scavenger receptors (SR-A, SR-BI, CD36, RAGE)	Degenerated myelin fA $\beta$ , thrombospondin-1 Antionic polysaccharide Polynucleotides Chemically modified proteins Apoptotic cells Bacteria	Microglia, astrocytes, mato cells, cerebral microvascular endothelial cells, cerebral arterial smooth muscle cells, retinal pigment epithelial cells	Multiple sclerosis/EAE Amyotrophic lateral sclerosis Alzheimer's disease	[128–131]
P2RY6	ATP and other nucleotides	Microglia	Neuropathic pain	[37, 132]
Galectin-3/MAC-2	Degenerated myelin	Microglia Macrophages Schwann cells Neurons Brain capillary endothelium	Traumatic brain injury Spinal cord injury Multiple sclerosis/EAE Alzheimer's disease	[133, 134] [135–137]
LRP receptors (low-density lipoprotein receptor-related protein-1) Phosphatidylserine receptors (phosphatidyl-serine-specific receptor)	Apoptotic and necrotic neurons	Macrophages Fibroblasts Epithelial cells Dendritic cells Astrocytes Perivascular Meningeal Choroid plexus macrophages	Alzheimer's disease	[138, 139]
Mannose receptors	Soluble immune components, cell debris, cytotoxic substances, myelin glycoproteins, degrading enzymes, microbial ligands		Ischemia	[140, 141]

cerebral parenchyma cells which contain neurons, oligodendrites, and astrocytes. TLRs not only trigger the recognition of pathogen-associated molecular patterns, such as LPS or viral nucleotides, but also recognize danger-associated molecular patterns, such as deposited amyloid  $\beta$  ( $A\beta$ ) fibril and  $\alpha$ -synuclein [22, 23]. TLRs are also implicated in a variety of cerebral disorders, including bacterial or viral infections; neurodegenerative disorders such as AD; inflammatory demyelinating disorders such as MS; spinal cord injury (SCI); and in development or physiological processes such as neurogenesis, learning, and memory [22–25]. TLRs and TLR-dependent signaling pathways are involved in antibacterial immunity and restricting viral infection in CNS infection. Of note, TLR2 and TLR4 mediate brain injury and subsequent inflammation after ischemic stroke [25–28]. TLR4-, TLR2-, and TLR9-dependent signaling pathways are involved in mediating microglial phagocytosis of neurotoxic  $A\beta$  deposit in AD brain and exert a protective role in nerve regeneration [28–30]. It has been reported that TLRs regulate phagocytosis through myeloid differentiation factor 88(MyD88)-dependent and MyD88-independent signaling pathways. The MyD88-dependent pathway is triggered by TLRs through activation of IL-1 receptor-associated kinase (IRAK)-4 and p38, resulting in up-regulation of scavenger receptors [31]. On the other hand, TLRs also regulate phagocytosis by MyD88-independent actin-Cdc42/Rac pathway [32].

#### TREM-2

TREM-2 is a kind of pattern receptor specific for polyanionic and locates mainly on the cell surface of osteoclasts in bones and in microglia of the CNS [8, 33]. In addition to up-regulating the synthesis of chemokines and mediating protective phagocytosis of apoptotic cell debris, activation of TREM-2 receptors suppresses secretion of pro-inflammatory factors such as cytokines and ROS [8, 33]. Clinical observation showed that administration of specific agonist or antibody of TREM during the effector phase of MS led to a more severe immune response and resulted in more extensive demyelination [34]. TREM-2 on microglia via binding with DNAX-activation protein 12 (DAP12), an ITAM-containing adaptor protein, triggers the reorganization of F-actin and phosphorylation of ERK/MAPK, mediating the clearance of apoptotic neurons [34, 35]. Nasu–Hakola disease, a systemic bone cystic disorder with progressive presenile dementia followed by extensive sclerosis in the front-temporal lobe and the basal ganglia, occurs due to genetic mutation of TREM-2 and DAP12 resulting in aberrant TREM-2/DAP12 signaling pathway [36].

#### P2Y6

P2Y6 receptor, a member of the G-protein-coupled receptor family, is actively responsive to UDP and partially responsive

to UTP and ADP. The study of P2Y6 receptor has gained increasing attention during the past several years since the elegant demonstration that P2Y6 receptor triggers the UDP-evoked microglial phagocytosis [37]. In other words, UDP, which is released from injured neurons after trauma or ischemia, acts as “eat me” signal and mediates the P2Y6-dependent phagocytosis. P2Y6, when combined with UDP, activates phospholipase C (PLC) which in turn causes the synthesis of inositol 1,4,5-trisphosphate (InsP3) and triggers the booted release of  $Ca^{2+}$  from InsP3-receptor-sensitive stores [37]. In addition to triggering the intracellular  $Ca^{2+}$  over-loading, P2Y6-receptor-dependent signaling pathway also triggers actin cytoskeleton polarization to shape filopodia-like protrusions, thus facilitates the engulfment of cell debris [37].

### Microglial Phagocytosis in CNS Diseases

#### Microglial Phagocytosis in AD

AD is a progressive neurodegenerative disease characterized by progressive memory loss, change in personality, and dementia. The distinctive pathological hallmarks of AD are the presence of extracellular plaques of  $A\beta$  peptide and intracellular neurofibrillar tangles of tau protein. Activated microglia, reactive astrocytes, and profound neuronal loss are also evident [38]. Human studies disclosed that the accumulation of  $A\beta$  deposition in the brain correlated well with cognitive impairment and neuronal loss [39]. Microglial cells play a crucial role in removing  $A\beta$  in different ways. Using 3D reconstruction of microglia and amyloid in an animal model of AD, Stalder et al. demonstrated that clusters of activated microglia contained lysosomes or vacuoles in close vicinity of the dense core plaques [40]. An in vitro study of pulse-chase experiments with Cy3-labeled  $\alpha_2$ -macroglobulin ( $\alpha_2M$ ) and Cy3-labeled  $A\beta$  microaggregates in the cultured microglia showed that rapid uptake of both  $\alpha_2M$  and  $A\beta$  microaggregates took place within 15 min, and those vacuoles could reach the acidic endosomes and lysosomes within 1 h. After 4 h, most internalized  $\alpha_2M$  were degraded or released from cells. Interestingly, microglial cells became engorged with undigested  $A\beta$  after continuous incubation with  $A\beta$  aggregation for 4 days, suggesting a slow rate of degradation of the fibrils [41]. Bamberger et al. reported the identification of the multicomponent receptor complex of fibrillar  $A\beta$  (f $A\beta$ ) on microglial cell surface [42]. The principle constituents of the complex are CD36, CD47 (also termed integrin-associated protein), and the  $\alpha 6\beta 1$ -integrin [43]. This receptor complex mediates the adhesion of  $A\beta$  fibrils (f $A\beta$ ) to microglia and elicits phagocytosis. The bond of f $A\beta$  to the ensemble of receptors leads to activation of tyrosine kinase-mediated signal transduction cascades, which then results in a respiratory burst and production of IL-1 $\beta$  [42]. Koenigsnecht and

Landreth provided data arguing that the cell surface receptor complex-mediated engagement of  $\text{fA}\beta$  was driven principally by a novel  $\beta_1$ -integrin-dependent mechanism, but not by classical phagocytic mechanisms mediated either by Ig receptors or the complement receptor 3 [43]. Mo/Hu APPswe PS1dE9 mice are capable of producing numerous  $\text{A}\beta$  deposits in the brain. C3H/HeJ mice are mutated with destructive *TLR4* gene (*Tlr<sup>Lps-d</sup>*), which causes inactivation of TLR4 by LPS. Mo/Hu APPswe PS1dE9 mice with *TlrLps-d/TlrLps-d* genotype were developed from Mo/Hu APPswe PS1dE9 mice mating with C3H/HeJ mice. These animal models exhibited an increase of cerebral  $\text{A}\beta$  load, which suggested that TLRs, particularly TLR-2 and TLR-4, were involved in the phagocytosis of  $\text{A}\beta$  deposit in brain parenchyma and thus exerted a protective role in AD patients [29]. FcR-mediated phagocytosis and complement activation also play a critical role in removal of plaques from the AD brain [30]. Additionally, monocyte chemoattractant protein-1 (MCP-1/CCL2), coupled with its binding receptor, CC-chemokine receptor 2, was crucial in neuroinflammatory response that affected disease process in a mouse model of AD [44–46]. CCL2-deficient AD mice (APP/PS1 mice) showed decreased microglial phagocytosis for both monomeric and oligomeric  $\text{A}\beta_{42}$  and accelerated  $\text{A}\beta$  deposits and oligomers [46]. Interestingly, transgenic overexpression of CCL-2 in APP/CCL2 mice dramatically facilitated  $\text{A}\beta$  uptake and subsequent intracellular  $\text{A}\beta$  oligomerization and resulted in progression of neurocognitive decline [45]. The exact role of CCL-2 in AD remains further study. Koenigsnecht-Talboo and Landreth have studied that the pro-inflammatory environment of the AD brain impaired the microglial capacity of removal of  $\text{A}\beta$ , whereas anti-inflammatory factors enhanced  $\text{A}\beta$  clearance [47]. Microglia treated with pro-inflammatory cytokines such as LPS, IL-1 $\beta$ , TNF- $\alpha$ , IFN- $\gamma$ , MCP-1, and CD40L suppressed  $\text{fA}\beta$ -stimulated microglial phagocytic activity in vitro [47]. This effect was even significant during activation of the CR3 or  $\text{fA}\beta$  receptor complex, but did not affect IgG- or FcR-mediated phagocytosis [47]. On the other hand, when BV2 cells were incubated with anti-inflammatory cytokines in a pro-inflammatory milieu, these phagocytes exhibited a stronger competence to uptake fibrillar  $\text{A}\beta$  [47]. Anti-inflammatory cytokines, such as IL-4, IL-10, cyclooxygenase (COX) inhibitors, ibuprofen, or an E prostanoid receptor antagonist, blocked NF $\kappa$ B-dependent stimulation of COX-2 expression and prostaglandin E2 (PGE2) production or its cascade pathways to eliminate the inhibitory activity of pro-inflammatory cytokines and consequently restored  $\text{fA}\beta$ -stimulated phagocytic response [42, 43, 47, 48]. These observations are consistent with previous demonstration that deletion of PGE2 EP2 receptor in animal model of familial AD resulted in a marked reduction of  $\text{A}\beta$  plaque burden and enhanced phagocytosis [49].

Although microglia exhibit “activated” phenotype, they are presumably unable to efficiently uptake  $\text{A}\beta$  plaque to prevent

the neurodegenerative progression of AD patients [42]. Treatment with antibodies against  $\text{A}\beta$  peptide to PDAPP transgenic mouse model of AD dramatically reduced  $\text{A}\beta$  deposit by facilitating clearance of preexisting amyloid rather than by simply preventing new plaques formation [50]. Recent studies showed that treatment with anti- $\text{A}\beta$  antibodies may be effective in reducing plaque burden and reversing memory deficits in mouse model for AD [51, 52]. Therefore, modulating phagocytosis and degradation of  $\text{fA}\beta$  by microglia might be a potential treatment of AD.

#### Microglial Phagocytosis in Trauma

Traumatic brain injury (TBI) and SCI are the most common forms of CNS traumatic injury. Much attention has been given to their pathological mechanisms. SCI occurs from a primary mechanical insult on the spinal cord, resulting in damage of neurons and axons [53], followed by a second-wave injury. Synthesis of free radicals and nitric oxide, as well as glutamate receptor-associated excitotoxicity and release of protease, was observed in the process of secondary injury. Microglia, together with infiltrated peripheral macrophages and T cells, have long been proposed to be involved in the delayed secondary injury. Microglia, as the principal immune effectors in the brain, undergo a conversion into a reactive pro-inflammatory phenotype in neurotrauma rapidly [54]. ED1-positive microglia began to be detected at 6 h after SCI and mostly located close to the lesion epicenter [55]. Within 48 h post-injury, the ED-1 positive microglia showed the tendency of a higher density and a broader macrophages/microglia infiltrating extension [55]. It is striking that a higher density of macrophages/microglia in the gray matter than white matter continued to be observed [55]. Since activated microglia and infiltrated peripheral macrophages exhibit the similar morphology, gene expression, as well as surface and endocellular markers, there is still no discriminating cellular markers to distinguish these two cell types in the CNS so far.

Accumulating evidence has shown that microglia play a pivotal role in the process of spinal cord regeneration and degeneration during the so-called secondary injury [11]. Microglia, when exposed to myelin in vitro, rapidly transformed from ramified to activated amoeboid morphology. These activated microglia synthesized and released a wide range of pro-inflammatory cytokines, chemokines, anti-inflammatory cytokines, neurotrophic factors, and phagocytosed the cellular debris.

In a dog model for SCI, a significant myelinophagia occurred 5 days after trauma, which was concomitant to the phagocytic activation of microglia [56]. An enhanced phagocytic capacity of single microglial cell was observed, particularly in the early 5 days, but the number of phagocytes did not change [56]. In this study, the capacity of microglial phagocytosis was measured by flow cytometric assay using FITC-

labeled *Staphylococcus aureus*. The bacteria were added into microglial cell suspension to determine the percentage of microglia with phagocytic phenotype. The enhanced phagocytic property was proposed to be required to remove the injured neurons, axons, and myelin sheaths. However, in the later stage, number of microglia with phagocytic phenotype increased, accompanied by less activity of phagocytosis of individual cell [56]. The rapid clearance of degenerated myelin is beneficial to trigger efficient remyelination.

Following neuronal axon injury, a classical response in the CNS is to remove or “strip off” synapses from the affected cell body, which is mediated by glia cells. The early studies by Cullheim and Thams elegantly demonstrated that major histocompatibility complex (MHC) class I molecules and MHC class I receptors played a key role in synaptic plasticity and nerve-regeneration after axotomy [57]. Synaptic stripping preferred to remove excitotoxic glutamatergic rather than the inhibitory glycinergic and GABAergic nerve terminals, suggesting a beneficial effect for the repair. Functional MHC class I took part in normal “stripping” by specifically keeping the inhibitory influence on injured motoneurons.  $\beta$ 2-Microglobulin, a subunit of MHC class I and transporter associated with antigen processing 1 knockdown or knockout mice, displayed remarkable elimination of synapses, resulting in less synaptic terminals remaining on the surface of affected neurons [57]. This observation was supported by a recent study by Sabha and colleagues showing a robust up-regulation of MHC I class after axotomy in different mice strains [58]. C57BL/6J mice with a low level of MHC I expression showed poor axon regeneration and slow synaptic stripping from motor neuron surface following peripheral nerve axotomy [58]. Additionally, A/J mice after axotomy exhibited significant elevation of MHC I expression and stronger regenerative potential, accompanied with more intense of synaptic elimination [58]. Zanon and colleagues indicated that IFN- $\beta$  treatment notably increased MHC class I expression after sciatic nerve crush in mice [59].

A recent study by Gitik et al. demonstrated that CD47 (also known as integrin-associated protein), expressed on myelin, when reacting to immune inhibitory receptor signal regulatory protein- $\alpha$  (SIRP- $\alpha$ ) on microglia, notably down-regulated myelin phagocytosis by microglia in vitro [60]. This finding suggested that CD47 on one hand exerts a neuroprotective function by protecting normal intact myelin from activated phagocytes, while on the other hand impedes normal phagocytosis therefore is unfavorable to the repair of lesioned neurons. Previous experiments have demonstrated that complement receptor-3 (CR3/MAC-1) signaling pathway was involved in the clearance of degenerated myelin [61]. CR3/MAC-1, as a member of integrin superfamily, is composed by  $\alpha_M$  and  $\beta_2$  subunit [62] and mediates myelin phagocytosis via binding to both complement (iC3b) and noncomplement ligand (e.g., fibrinogen and I-CAM) [61]. A study by

Rotshenker and colleagues demonstrated that CR3/MAC-1 activated guanine nucleotide exchange factors that facilitated the conversion of inactive K-Ras-GDP to active K-Ras-GTP, which then led to activation of PI3K and myelin phagocytosis [63]. Treating microglia with TNF- $\alpha$  suppressed myelin phagocytosis mediated by CR3/MAC-1 [64]. Scavenger receptor AI/II (SRAI/II) also participates in myelin clearance after trauma. However, the relative potential contribution of CR3/MAC-1 to myelin phagocytosis is two- or three-folds more than that of SRAI/II [65]. Microglia immunophenotypical characterization in dogs with SCI revealed a remarkable upregulation of mediators for phagocytosis, including ICAM-I, CD14, CD44, and CD45, and the increasing of expression intensity of these surface molecules significantly enhanced the removal of tissue debris [56].

Myelin structures, such as Nogo-A, myelin-associated glycoprotein, oligodendrocytes myelin glycoprotein, the transmembrane semaphoring 4D (Sema4D/CD100), and ephrinB3, translocate to cell surface on injured myelin and act as inhibitors of axonal regeneration [66]. Therefore, efficient microglial phagocytosis of damaged myelin and cell debris is beneficial for the survival of injured neurons and myelin regeneration in acute SCI and TBI [56, 67]. The enhanced microglial phagocytosis of myelin debris was demonstrated dramatically to decrease dead cells and myelin debris and result in prominent functional recovery [61].

#### Microglial Phagocytosis in Ischemic/Stroke

Stroke is the most common disease in the CNS among the elderly. Rapid infiltration of polymorphonuclear neutrophils (PMNs) and peripheral monocytes/macrophages and abundant activation of resident microglia are observed in the injured brain post-ischemia lesion [68, 69]. In an in vivo study, Schilling et al. demonstrated that both resident microglia and hematogenous macrophages made contributions to debris clearance post-cerebral infarction, while microglia played a more major role than did macrophages [70]. Activated microglia were observed to rapidly migrated into the infarction area and elicited phagocytic response at day 1 after ischemia [71, 72], but no peripheral infiltration was seen [70]. The total number of activated microglia showed an increased tendency in the following several days and reached maximum at day 10 after transient focal cerebral ischemia [70], whereas blood-borne macrophages began to occur in the damaged lesion at day 4, increased to the peak number in the following 3 days, and began to decrease until 2 weeks after ischemic insult [70]. Strikingly, phagocytic microglia only accounted for one quarter of the total activated microglia. This phenotype of microglia rapidly had their maximum number as early as day 1 and stayed at the same level within the following days until the hematogenous macrophages infiltrated [70]. These findings lead to the hypothesis that participation of microglia in the

debris clearance takes place in the early stage while the assistance from blood-borne macrophages infiltration in the infarction zone occurs later.

In vivo study showed that PMN rapidly infiltrated and accumulated in ischemic-injured brain as early as day 1 after cerebral ischemia [69, 73]. Application of PMNs onto post-oxygen–glucose deprivation organotypic hippocampal slice cultures (OGD-OHCs) led to a remarkable exacerbation of neuronal damage, characterized by severe loss of axons and dendrites, as well as the appearance of apoptotic or necrotic neurons containing subcellular material at day 1 [69]. Interestingly, the co-application of PMNs and microglia onto the OGD-OHCs significantly reduced the PMN-induced damage to neurons, which was presumably due to direct engulfment of those viable, motile, and nonapoptotic PMNs by microglia [69]. Uptake of PMNs by microglia significantly reduced the release of pro-inflammatory mediators, such as cytokines and ROS, and was associated with the upregulation of TGF- $\beta$  in microglia, both of which exerted a beneficial role in the injured brain [69]. The exact mechanisms and signals involved in recognition and uptake of PMN by microglia are not yet defined. The capacity of microglia to engulf PMN can be inhibited by TNF- $\alpha$ . Furthermore, microglia also elicited phagocytic response through interaction with cell surface receptors, for example, osteopontin (OPN), an adhesive glycoprotein [74]. It has previously been accepted that OPN acts only as a chemoattractant after ischemic insult, but recently, it has been reported that OPN also takes part in the phagocytosis of the cell debris [75]. Elevated expression of OPN protein was seen mainly along the membranes lining after ischemic stroke; thus, researchers concluded that OPN elicited phagocytic response of fragmented debris selectively [74].

Both microglia and hematogenous macrophages contribute to clearance of cell debris generated in the context of ischemia. The efficient phagocytosis of debris is beneficial for axon regeneration. In addition, phagocytosis by microglia/macrophages exerts favorable effect through engulfment of infiltrated PMN. Depletion of PMN subsequently converts the microenvironment from pro-inflammatory to anti-inflammatory, which reduce neuronal injury in the process of disease. In general, microglial phagocytosis might exert beneficial effect in ischemic brain.

#### Microglial Phagocytosis in PD

PD is characterized by progressive dopaminergic neuronal loss in substantia nigra and nerve terminals in the striatum, resulted in bradykinesia, static tremor, instability of gait and posture, and muscle stiffness. Overexpression of extracellular  $\alpha$ -synuclein is a well-known etiological pathology of PD [76]. Microglia treated with monomeric  $\alpha$ -synuclein exhibited an enhanced phagocytic activity in vitro, in a both time- and dose-dependent manner [77]. In this study, microglial

phagocytosis was evaluated by the activity of microglia in ingesting extracellular fluorescent microspheres. The exact mechanism underlying in monomeric  $\alpha$ -synuclein induced phagocytosis by microglia is still an open question. Studies have confirmed the involvement of CR3,  $\alpha 6\beta 1$ , and CD47, which are important components of receptor complex in clearance of A $\beta$  [77]. In comparison, aggregated  $\alpha$ -synuclein inhibited phagocytosis of cell debris and dead neurons not only by antagonizing monomeric-facilitated clearance but also through decreasing the basal microglial phagocytic capability [77]. It is also reported that microglia were capable of phagocytosing and degenerating neuromelanin (NM) released from degenerated dopaminergic neurons. Microglia co-cultured with NM revealed a rapid transform from a ramified to an amoeboid morphology and were engaged in attaching to and engulfing NM particles [78]. In a 6-hydroxydopamine (6-OHDA) model of PD, phagocytic microglia (CD68 positive) were found to surround the intact tyrosine hydroxylase-positive dopaminergic substantia nigra pars compacta (SNc) neurons to remove damaged NM particles in the SNc ipsilateral to the lesion side at day 3. At day 7, an increasing number of microglia were confined to the medial area of SNc to contact with intact dopaminergic neurons, as well as apoptotic cells. The number of microglia increased in a time-dependent manner and reached significance at day 9 and had their maximum at day 15 post-6-OHDA lesioning. Microglia were found to localize in the medial and lateral part of ipsilateral SNc. A vast proportion of phagocytic microglia (CD68 positive) were seen to adhere to and engulf degenerated dopaminergic neurons and axons [79]. Using proteomic technology, Liu et al. have shown that a variety of types of membrane proteins were potentially involved in the internalization of  $\alpha$ -synuclein [80]. Clathrin was demonstrated to play a critical role in the endocytosis of aggregated  $\alpha$ -synuclein, probably in a receptor-ligand sequestration-related manner [80], but the exact mechanism needs further study. Recently, TLR4 signaling pathway is demonstrated to mediate  $\alpha$ -synuclein phagocytosis and exert a beneficial role in deferring disease progression both in vivo and in vitro [81]. In in vivo study of transgenic murine model of  $\alpha$ -synucleinopathies (ASP), mice overexpressed human  $\alpha$ -synuclein (hAS) with TLR4 deficiency (AS/TLR4<sup>-/-</sup>) exhibited severer neuronal loss, motor disability, and predominant reduced phagocytic activity than those with normal TLR4 expression (AS/TLR4<sup>+/+</sup>). Counterstaining of anti-hAS and CD11b showed abundant hAS-positive structure appeared in the cytoplasm of CD11b-positive microglia in AS/TLR4<sup>+/+</sup> mice; on the contrary, hAS-positive structure was found only outside microglia in AS/TLR4<sup>-/-</sup> mice. By further study with immunogold labeling for AS in the brain of transgenic mice, microglia in AS/TLR4<sup>+/+</sup> mice showed abundant gold particles in phagocytic cytoplasmic organelles, while fewer gold particles were found in microglia in AS/TLR4<sup>-/-</sup> mice [81].

Numerous studies also displayed that C1q-mediated pathway [82] scavenger receptors [83] and MAC-1 [78] are also involved in microglial endocytosis of  $\alpha$ -synuclein. Overexpression of human wild type and mutant  $\alpha$ -synuclein(A30P and A30T) in BV2 cells resulted in downregulation of phagocytosing bioparticles and a marked low lysosomal associated protein 1 expression, accompanied with elevated COX-2 and proinflammatory cytokines such as PGE2 [84].

The rate of internalization and subsequent intracellular degradation of extracellular  $\alpha$ -synuclein aggregates were compared in the major brain cell types of neurons, microglia, and astrocytes. The finding demonstrated that all these three types of cells were capable of clearing  $\alpha$ -synuclein. Among them, microglia showed to be the most effective [85]. Whether microglial phagocytosis of  $\alpha$ -synuclein favors or harms the process of PD is still under debate. Zhang et al. argued that internalization of  $\alpha$ -synuclein took a central role in dopaminergic neurotoxicity through activation of NADPH oxidase and subsequently oxidative stress [86]. However, as mentioned above, impaired microglial phagocytic capacity by ablation of TLR4 in ASP mouse model led to aggregation of extracellular  $\alpha$ -synuclein and accelerated neurodegeneration. In brief, the role of microglial phagocytosis on PD remains further investigation.

#### Microglial Phagocytosis in ALS

ALS is the most common progressive neurodegenerative disorder that selectively affects motoneurons in the CNS. Its remarkable characteristic is the simultaneous death of lower and upper motor neuron, leading to progressive muscle weakness and atrophy. Patients suffering from ALS usually end up with death from respiratory paralysis within 2 to 5 years of onset. Although ALS is overwhelmingly a sporadic disorder, genetic studies have established that mutations in the Cu/Zn superoxide dismutase 1 (SOD1) gene are the most well-known cause of familial ALS [87, 88]. Studies have shown that microglia has an important function in propagation of the disease process both in sporadic and familial ALS [89, 90] and in the transgenic animals overexpressing human mutant SOD1 (hmSOD1) [91]. By analysis of autopsy cases of ALS, increased numbers of macrophages were observed in the regions with motor neuron loss, such as lower motor neuron XII, upper motor neuron beta cells, spinocerebellar inferior olivary nuclei and red nuclei, somatosensory caudate nuclei and thalamus, cerebral cortex amygdaloid, and the ventral horn of the spinal cord [90]. This observation is supported by a recent experimental study which argued that activated microglia aggregated in the anterior horn of the lumbar spinal cord, particularly around impaired motor neurons [92]. This study also demonstrated that activated microglia attached to somata of motoneurons and exhibited

phagocytic properties as early as presymptomatic stage [92]. Interestingly, activated microglia were visualized not only in regions where there was severe motor neuron loss but also in areas of mild motor neuronal damage [90]. However, the exact role of microglia in motor neuron degeneration remains unclear. In transgenic hmSOD1 (G93A) rat, it has been shown that CD11b-positive macrophages/microglia accumulated both in the ventral horn and more distally at the peripheral nerve during pre-symptom stage. As the disease progressed, microglia aggregated in the ventral horn and became more prevalent. These focal aggregates of microglia were observed often near but never fully encapsulating NeuN-labeled neurons. However, after onset of the symptoms, microglia labeled with CD68 and MHC class II (markers of activated microglia with phagocytic activity) began to be observed in the ventral horn. Therefore, microglia may cause damage to nearby motor neurons through secretion of proinflammatory factors at the early stage of ALS, while play a neuroprotective role through phagocytosing the degenerated debris following clinical onset [91].

TLR-dependent signaling pathways that modulate microglial phagocytosis play a key role in Wallerian degeneration (WD) and ALS. Injection of TLR-2 and TLR-4 enhanced myelin debris clearance and repairment of locomotor function after sciatic nerve lesion [93]. Blockade of TLRs and downstream signaling MyD88 decreased axons debris removal and delayed neuron regeneration [93]. MCP-1, macrophage inflammatory protein-1 $\alpha$ , and interleukin-1 $\beta$  were reported to be involved in the progress of WD. Injection of these proinflammatory cytokines into the sciatic nerve led to rapid myelin damage and elevated macrophage/microglial phagocytic activity in WD [94]. Efficient removal of degenerated neurons or cell debris is necessary to rebuild beneficial environment for neuronal regeneration in degenerated disease. In response to neuronal cell death, microglia become activated and aggregate around the lesioned area, exhibiting potent phagocytic activity [95]. Enhancement of myelin phagocytosis by microglial/macrophages favors neuron regeneration and restoration of locomotor function.

#### Microglial Phagocytosis in MS/EAE

MS is a chronic neurological disorder of the CNS which leads to nontraumatic disability among young adults. The pathological hallmarks of MS are extensive demyelination and formation of sclerosis plaques in cerebral cortex and spinal cord, which contribute to irreversible neurological injury. Experimental allergic encephalomyelitis (EAE) and Theiler's murine encephalomyelitis virus infection are the two most commonly used animal models of human MS. Immune effector cells including reactive microglia and hematogenous macrophages, as well as T cells, are observed aggregated in the lesion area in rat model of EAE [96, 97]. The exact mechanisms of



demyelination remained unclear. Glial cell proliferation during the process of MS may be the major effector in the development of a demyelinating plaque due to the composition of the inflammation infiltrating, as well as the dynamics of remyelination [98]. Early remyelinating lesions contained a mixture of infiltrated macrophages and microglia. Thin myelin sheaths were observed to surround the damaged axons [20, 34, 98], while in late remyelinating lesions only a few monomorphic populations of phagocytic macrophages presented in these lesions [98]. These studies suggested that insufficient myelin clearance in the CNS after MS onset may be involved in the failure of axonal regeneration. The relative contribution to disease progression of brain resident macrophages versus that of blood-borne macrophages remains controversial. Most studies favor the viewpoint that resident macrophages play a decisive role in disease progression and selective depletion of perivascular and meningeal macrophages using clodronate liposome injection in rat model of acute EAE slows disease progression [96]. Recent studies using bone marrow chimeras claimed that the number of microglia were several folds over blood-derived macrophages, though they shared equal phagocytic capacity [97]. These findings support the hypothesis that microglia are more efficient in phagocytosing myelin debris compared with peripheral macrophages. In an *in vitro* Wallerian degeneration model by cutting axons of the cortical explants, co-cultured microglia exposed to degenerated axon debris were first seen to engulf debris within 15 h [99], whereas perivascular and meningeal macrophages infiltrating into the lesion area occurred at day 9 reached their maximum number at day 15 and decreased to normal level at day 24 post-EAE induction in rats [96]. The different types of macrophage activation in MS/EAE are closely correlated with the stage of the demyelinating activity and with the type of MS tissue [99]. In early-active MS lesion, massive activated macrophages aggregated at the plaque border and engaged in phagocytosing myelin degradation products; however, in late-active demyelination lesion, a small number of amoeboid-like macrophages were observed diffusely infiltrated in the lesion [98, 99]. The brain resident microglia are the main macrophages engaged in phagocytosis in the early stage of demyelination, while in the late stage a number of infiltrated blood-borne macrophages contribute to axon debris clearance. A variety of types of receptors are involved in MS, including Fc receptors, complement receptors, macrophage scavenger receptors, the galectin-3/MAC-2 receptors,  $\alpha$ 2-macroglobulin/low-density lipoprotein receptor, and mannose receptors [8]. In these receptor-linked pathways, TREM-2 signaling pathway has been the most well-studied. TREM-2 mediated phagocytosis of apoptotic neurons without release of pro-inflammation molecules in animal model of MS [35, 100]. Han et al. recently demonstrated that downregulation of CD47 promoted phagocytosis of myelin in a SIRP- $\alpha$ -dependent mechanism by using CD47<sup>-/-</sup> EAE mice [101]. The capacity

of macrophages/microglia to phagocytose degenerated myelin can be altered by environmental inflammatory mediators, such as IFN- $\gamma$ , TNF- $\alpha$ , IL-4, IL-10, and so on [20]. This notion was confirmed by an *in vitro* study that investigated the effect of certain cytokines on phagocytosis of <sup>14</sup>C-labeled myelin by cultured macrophages or microglia. In this study, TNF- $\alpha$  was shown to increase the phagocytic activity of microglia, but have no effect on macrophages [20]. TNF- $\gamma$  was demonstrated to reduce removal of myelin by macrophage, on the contrary enhance debris clearance driven by microglia. IL-4 and IL-10 exerted a role of up-regulating phagocytosis in macrophages/microglia, while accompanied by a reduction of inflammatory response [20]. In demyelinated diseases, efficient removal of degenerated myelin is necessary to rebuild beneficial environment for remyelination and axon regeneration. Therefore, robust microglial phagocytosis might be critical in the recovery of MS/EAE.

### Microglial Phagocytosis in Brain Tumor

Nearly 18,000 patients die of primary tumors of the brain, and another 130,000 die from metastases neoplasm. Most primary brain tumors are of glial-cell origin, including glioblastoma multiforme, astrocytomas, oligodendrogliomas, ependymomas, and so on. Among which, malignant gliomas are the most common primary brain tumors and are highly aggressive with limited treatment efficiency. Macrophages/microglia were observed to present in tumors, tumor periphery, and contralateral tumor-free hemispheres, particularly around tumor margins [102, 103]. Badie and Scharfner revealed that infiltrated microglia comprise 13.2 to 34.0 % of tumor mass, and the extent of microglia infiltration depends on the tumor type rather than its size [102]. Glioma-infiltrating microglia/macrophages show protumorigenic activity by up-regulation of metalloprotease-II [104], but cannot secrete cytokines such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$ , which is distinctive from the inflammatory phenotype [105]. In glioma microenvironment, normal human astrocytes (NHA), glioma cells, and microglia all show capability of phagocytosing glioma cells and specifically apoptotic tumor cells [106]. The phagocytosis index (the index to assess the phagocytic activity of phagocytes through counting of engulfed apoptotic cells) value of microglia showed about four-fold higher than that of NHA or glioma cells, indicating that microglia were the most efficient phagocytes in brain tumor [106]. IL-12 and LPS administration obviously enhance microglial phagocytotic activity through the TRAIL pathway [107]. *In vitro* study revealed that glioma cancer stem cells (gCSCs) dramatically inhibited phagocytosis by human microglia [108]. Further study regarding underlying mechanisms showed reversal of phagocytosis inhibition after STAT3 blockade by WP1066 and STAT3 siRNA in gCSCs, suggesting involvement of p-STAT3 pathway in inhibition of phagocytosis by gCSCs

[108]. Microglia play a key role in phagocytosing tumor cells. This activity can be modulated by cancer cells or cancer environment. Whether microglial phagocytosis serves as cancer defense or cancer contribution still needs further investigation.

#### Microglial Phagocytosis in Other Brain Diseases

Studies have shown that immune response is pivotal in prion disease [9, 109–112]. In prion disease, microglia become activated and take a decisive role in the progression of this disease [9]. Microglia in prion disease displayed low level of secretory profiles and elevated expression of phagocytotic machinery such as receptors for advanced glycation end products, TREM-2, and the scavenger receptors SAR2, CD68, and SRB(CD36) [109]. It is striking that intracerebral injection of LPS significantly stimulated synthesis and secretion of proinflammatory cytokines, but without alteration of PrPs clearance [109], which is thoroughly different from the generally accepted notion that microglial phagocytic state is positively in line with its activation degree [113]. Numerous findings demonstrated that microglia also participated in radiation induced neuronal damage [114–119]. Microglia with phagocytic morphology could be seen accumulated in the lesion cavity/spinal cord interface in irradiated spinal cord hemisection. Interestingly, microglial population rapidly decreased in the irradiated, lesioned hemisection when compared to nonirradiated animal [114, 118]. However, the exact mechanism underlying phagocytic process and the role of clearance by microglia in radiation induced brain injury needs further investigation.

#### Conclusion

Microglia, as the resident macrophage in the CNS, play a critical role in the pathological and physiological processes of CNS diseases, sustaining the homeostasis of the local brain parenchyma. Even a minor disturbance can provoke the activation of microglia and following inflammation-induced neuronal damage. Most investigators favor that microglial phagocytosis exerts a beneficial effect in repair and regeneration. Variety types of receptors expressed on the cell surface of microglia are involved in mediating phagocytic process. These activated microglia exhibit an array of phenotypes and are governed by their surrounding environment. Activated homogenous macrophages infiltrate the blood–brain barrier and contribute to debris clearance in brain disease, such as ischemia, MS, trauma, and so forth. The relative contribution of resident microglia and peripheral macrophages remains controversial. The exact mechanism and functional aspect of phagocytosis by microglia needs further study, and modulation of microglial phagocytosis might be a potential therapeutic strategy for neurological diseases.

**Acknowledgments** This work was supported by the National Natural Science Foundation of China (no. 81272576 and no. 81072242), the Fundamental Research Funds for the Central Universities, and the Funds for Pearl River Science & Technology Star of Guangzhou City (2012J2200088) to Yamei Tang.

**Conflict of Interest** The authors declare that there are no conflicts of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

#### References

1. Stence N, Waite M, Dailey ME (2001) Dynamics of microglial activation: a confocal time-lapse analysis in hippocampal slices. *Glia* 33(3):256–266
2. Eugenin EA et al (2001) Microglia at brain stab wounds express connexin 43 and in vitro form functional gap junctions after treatment with interferon-gamma and tumor necrosis factor-alpha. *Proc Natl Acad Sci U S A* 98(7):4190–4195
3. Gehrmann J, Matsumoto Y, Kreutzberg GW (1995) Microglia: intrinsic immunoeffector cell of the brain. *Brain Res Brain Res Rev* 20(3):269–287
4. Neumann H, Kotter MR, Franklin RJ (2009) Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain* 132(Pt 2):288–295
5. Prokop S, Miller KR, Heppner FL (2013) Microglia actions in Alzheimer's disease. *Acta Neuropathol* 126(4):461–477
6. Long-Smith CM, Sullivan AM, Nolan YM (2009) The influence of microglia on the pathogenesis of Parkinson's disease. *Prog Neurobiol* 89(3):277–287
7. Sargsyan SA, Monk PN, Shaw PJ (2005) Microglia as potential contributors to motor neuron injury in amyotrophic lateral sclerosis. *Glia* 51(4):241–253
8. Napoli I, Neumann H (2010) Protective effects of microglia in multiple sclerosis. *Exp Neurol* 225(1):24–28
9. Brown DR (2001) Microglia and prion disease. *Microsc Res Tech* 54(2):71–80
10. Yenari MA, Kauppinen TM, Swanson RA (2010) Microglial activation in stroke: therapeutic targets. *Neurotherapeutics* 7(4):378–391
11. Loane DJ, Bymes KR (2010) Role of microglia in neurotrauma. *Neurotherapeutics* 7(4):366–377
12. Chiang CS, McBride WH, Withers HR (1993) Radiation-induced astrocytic and microglial responses in mouse brain. *Radiother Oncol* 29(1):60–68
13. Brown GC, Neher JJ (2012) Eaten alive! Cell death by primary phagocytosis: 'phagoptosis'. *Trends Biochem Sci* 37(8):325–332
14. Sierra A et al (2010) Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell* 7(4):483–495
15. Weldon DT et al (1998) Fibrillar beta-amyloid induces microglial phagocytosis, expression of inducible nitric oxide synthase, and loss of a select population of neurons in the rat CNS in vivo. *J Neurosci* 18(6):2161–2173
16. Ito U et al (2007) Fate of disseminated dead neurons in the cortical ischemic penumbra: ultrastructure indicating a novel scavenger mechanism of microglia and astrocytes. *Stroke* 38(9):2577–2583
17. Aderem A, Underhill DM (1999) Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol* 17:593–623

18. Veerhuis R, Nielsen HM, Tenner AJ (2011) Complement in the brain. *Mol Immunol* 48(14):1592–1603
19. Lucin KM, Wyss-Coray T (2009) Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron* 64(1):110–122
20. Smith ME (1999) Phagocytosis of myelin in demyelinating disease: a review. *Neurochem Res* 24(2):261–268
21. Collin M, McGovern N, Haniffa M (2013) Human dendritic cell subsets. *Immunology* 140(1):22–30
22. Marsh BJ, Williams-Karnesky RL, Stenzel-Poore MP (2009) Toll-like receptor signaling in endogenous neuroprotection and stroke. *Neuroscience* 158(3):1007–1020
23. Hanke ML, Kielian T (2011) Toll-like receptors in health and disease in the brain: mechanisms and therapeutic potential. *Clin Sci (Lond)* 121(9):367–387
24. Schiff D, Rosenblum MK (1998) Herpes simplex encephalitis (HSE) and the immunocompromised: a clinical and autopsy study of HSE in the settings of cancer and human immunodeficiency virus-type 1 infection. *Hum Pathol* 29(3):215–222
25. Rivest S (2003) Molecular insights on the cerebral innate immune system. *Brain Behav Immun* 17(1):13–19
26. Ziegler G et al (2007) TLR2 has a detrimental role in mouse transient focal cerebral ischemia. *Biochem Biophys Res Commun* 359(3):574–579
27. Lehnardt S et al (2007) Toll-like receptor 2 mediates CNS injury in focal cerebral ischemia. *J Neuroimmunol* 190(1–2):28–33
28. Caso JR et al (2008) Toll-like receptor 4 is involved in subacute stress-induced neuroinflammation and in the worsening of experimental stroke. *Stroke* 39(4):1314–1320
29. Tahara K et al (2006) Role of toll-like receptor signalling in A $\beta$  uptake and clearance. *Brain* 129(Pt 11):3006–3019
30. Lee CY, Landreth GE (2010) The role of microglia in amyloid clearance from the AD brain. *J Neural Transm* 117(8):949–960
31. Doyle SE et al (2004) Toll-like receptors induce a phagocytic gene program through p38. *J Exp Med* 199(1):81–90
32. Kong L, Ge BX (2008) MyD88-independent activation of a novel actin-Cdc42/Rac pathway is required for Toll-like receptor-stimulated phagocytosis. *Cell Res* 18(7):745–755
33. Klesney-Tait J, Turnbull IR, Colonna M (2006) The TREM receptor family and signal integration. *Nat Immunol* 7(12):1266–1273
34. Piccio L et al (2007) Blockade of TREM-2 exacerbates experimental autoimmune encephalomyelitis. *Eur J Immunol* 37(5):1290–1301
35. Takahashi K, Rochford CD, Neumann H (2005) Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *J Exp Med* 201(4):647–657
36. Satoh J et al (2012) Phosphorylated Syk expression is enhanced in Nasu-Hakola disease brains. *Neuropathology* 32(2):149–157
37. Koizumi S et al (2007) UDP acting at P2Y6 receptors is a mediator of microglial phagocytosis. *Nature* 446(7139):1091–1095
38. Selkoe DJ (1991) Amyloid protein and Alzheimer's disease. *Sci Am* 265(5):68–71, 74–6, 78
39. Nathalie P, Jean-Noel O (2008) Processing of amyloid precursor protein and amyloid peptide neurotoxicity. *Curr Alzheimer Res* 5(2):92–99
40. Stalder M et al (2001) 3D-Reconstruction of microglia and amyloid in APP23 transgenic mice: no evidence of intracellular amyloid. *Neurobiol Aging* 22(3):427–434
41. Paresce DM, Chung H, Maxfield FR (1997) Slow degradation of aggregates of the Alzheimer's disease amyloid  $\beta$ -protein by microglial cells. *J Biol Chem* 272(46):29390–29397
42. Bamberger ME et al (2003) A cell surface receptor complex for fibrillar  $\beta$ -amyloid mediates microglial activation. *J Neurosci* 23(7):2665–2674
43. Koenigsnecht J, Landreth G (2004) Microglial phagocytosis of fibrillar  $\beta$ -amyloid through a  $\beta$ 1 integrin-dependent mechanism. *J Neurosci* 24(44):9838–9846
44. Bose S, Cho J (2013) Role of chemokine CCL2 and its receptor CCR2 in neurodegenerative diseases. *Arch Pharm Res* 36(9):1039–1050
45. Kiyota T et al (2009) CCL2 accelerates microglia-mediated A $\beta$  oligomer formation and progression of neurocognitive dysfunction. *PLoS One* 4(7):e6197
46. Kiyota T et al (2013) CCL2 affects  $\beta$ -amyloidosis and progressive neurocognitive dysfunction in a mouse model of Alzheimer's disease. *Neurobiol Aging* 34(4):1060–1068
47. Koenigsnecht-Talboo J, Landreth GE (2005) Microglial phagocytosis induced by fibrillar  $\beta$ -amyloid and IgGs are differentially regulated by proinflammatory cytokines. *J Neurosci* 25(36):8240–8249
48. Inoue Y et al (2004) Novel regulatory mechanisms of CD40-induced prostanoind synthesis by IL-4 and IL-10 in human monocytes. *J Immunol* 172(4):2147–2154
49. Liang X et al (2005) Deletion of the prostaglandin E2 EP2 receptor reduces oxidative damage and amyloid burden in a model of Alzheimer's disease. *J Neurosci* 25(44):10180–10187
50. Bard F et al (2000) Peripherally administered antibodies against amyloid  $\beta$ -peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. *Nat Med* 6(8):916–919
51. DeMattos RB et al (2001) Peripheral anti-A $\beta$  antibody alters CNS and plasma A $\beta$  clearance and decreases brain A $\beta$  burden in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 98(15):8850–8855
52. Wilcock DM et al (2003) Intracranially administered anti-A $\beta$  antibodies reduce  $\beta$ -amyloid deposition by mechanisms both independent of and associated with microglial activation. *J Neurosci* 23(9):3745–3751
53. Brockhaus J, Moller T, Kettenmann H (1996) Phagocytosing amoeboid microglial cells studied in a mouse corpus callosum slice preparation. *Glia* 16(1):81–90
54. Hawthorne AL, Popovich PG (2011) Emerging concepts in myeloid cell biology after spinal cord injury. *Neurotherapeutics* 8(2):252–261
55. Carlson SL et al (1998) Acute inflammatory response in spinal cord following impact injury. *Exp Neurol* 151(1):77–88
56. Boekhoff TM et al (2012) Microglial contribution to secondary injury evaluated in a large animal model of human spinal cord trauma. *J Neurotrauma* 29(5):1000–1011
57. Cullheim S, Thams S (2007) The microglial networks of the brain and their role in neuronal network plasticity after lesion. *Brain Res Rev* 55(1):89–96
58. Sabha MJ et al (2008) MHC I expression and synaptic plasticity in different mice strains after axotomy. *Synapse* 62(2):137–148
59. Zanon RG et al (2010) Interferon (IFN)  $\beta$  treatment induces major histocompatibility complex (MHC) class I expression in the spinal cord and enhances axonal growth and motor function recovery following sciatic nerve crush in mice. *Neuropathol Appl Neurobiol* 36(6):515–534
60. Gitik M et al (2011) Myelin down-regulates myelin phagocytosis by microglia and macrophages through interactions between CD47 on myelin and SIRP $\alpha$  (signal regulatory protein- $\alpha$ ) on phagocytes. *J Neuroinflammation* 8:24
61. Rotshenker S (2003) Microglia and macrophage activation and the regulation of complement-receptor-3 (CR3/MAC-1)-mediated myelin phagocytosis in injury and disease. *J Mol Neurosci* 21(1):65–72
62. Hughes PE, Pfaff M (1998) Integrin affinity modulation. *Trends Cell Biol* 8(9):359–364
63. Rotshenker S et al (2008) Galectin-3/MAC-2, Ras and PI3K activate complement receptor-3 and scavenger receptor-AI/II mediated myelin phagocytosis in microglia. *Glia* 56(15):1607–1613
64. Bruck W, Bruck Y, Friede RL (1992) TNF- $\alpha$  suppresses CR3-mediated myelin removal by macrophages. *J Neuroimmunol* 38(1–2):9–17

65. Reichert F, Rotshenker S (2003) Complement-receptor-3 and scavenger-receptor-AI/II mediated myelin phagocytosis in microglia and macrophages. *Neurobiol Dis* 12(1):65–72
66. Yiu G, He Z (2006) Glial inhibition of CNS axon regeneration. *Nat Rev Neurosci* 7(8):617–627
67. Redondo-Castro E et al (2013) Phagocytic microglial phenotype induced by glibenclamide improves functional recovery but worsens hyperalgesia after spinal cord injury in adult rats. *Eur J Neurosci* 38:3786–3798
68. Dimagl U, Iadecola C, Moskowitz MA (1999) Pathobiology of ischaemic stroke: an integrated view. *Trends Neurosci* 22(9):391–397
69. Neumann J et al (2008) Microglia cells protect neurons by direct engulfment of invading neutrophil granulocytes: a new mechanism of CNS immune privilege. *J Neurosci* 28(23):5965–5975
70. Schilling M et al (2005) Predominant phagocytic activity of resident microglia over hematogenous macrophages following transient focal cerebral ischemia: an investigation using green fluorescent protein transgenic bone marrow chimeric mice. *Exp Neurol* 196(2):290–297
71. Kato H et al (1996) Progressive expression of immunomolecules on activated microglia and invading leukocytes following focal cerebral ischemia in the rat. *Brain Res* 734(1–2):203–212
72. Morioka T, Kolehua AN, Streit WJ (1993) Characterization of microglial reaction after middle cerebral artery occlusion in rat brain. *J Comp Neurol* 327(1):123–132
73. Dinkel K, Dhabhar FS, Sapolsky RM (2004) Neurotoxic effects of polymorphonuclear granulocytes on hippocampal primary cultures. *Proc Natl Acad Sci U S A* 101(1):331–336
74. Shin YJ et al (2011) Osteopontin: correlation with phagocytosis by brain macrophages in a rat model of stroke. *Glia* 59(3):413–423
75. Pedraza CE et al (2008) Osteopontin functions as an opsonin and facilitates phagocytosis by macrophages of hydroxyapatite-coated microspheres: implications for bone wound healing. *Bone* 43(4):708–716
76. Cookson MR (2005) The biochemistry of Parkinson's disease. *Annu Rev Biochem* 74:29–52
77. Park JY et al (2008) Microglial phagocytosis is enhanced by monomeric alpha-synuclein, not aggregated alpha-synuclein: implications for Parkinson's disease. *Glia* 56(11):1215–1223
78. Zhang W et al (2011) Neuromelanin activates microglia and induces degeneration of dopaminergic neurons: implications for progression of Parkinson's disease. *Neurotox Res* 19(1):63–72
79. Marinova-Mutafchieva L et al (2009) Relationship between microglial activation and dopaminergic neuronal loss in the substantia nigra: a time course study in a 6-hydroxydopamine model of Parkinson's disease. *J Neurochem* 110(3):966–975
80. Liu J et al (2007) Identification of proteins involved in microglial endocytosis of alpha-synuclein. *J Proteome Res* 6(9):3614–3627
81. Stefanova N et al (2011) Toll-like receptor 4 promotes alpha-synuclein clearance and survival of nigral dopaminergic neurons. *Am J Pathol* 179(2):954–963
82. Depboylu C et al (2011) Possible involvement of complement factor C1q in the clearance of extracellular neuromelanin from the substantia nigra in Parkinson disease. *J Neuropathol Exp Neurol* 70(2):125–132
83. Michelakakis H et al (2012) Evidence of an association between the scavenger receptor class B member 2 gene and Parkinson's disease. *Mov Disord* 27(3):400–405
84. Lee HJ et al (2008) Clearance and deposition of extracellular alpha-synuclein aggregates in microglia. *Biochem Biophys Res Commun* 372(3):423–428
85. Rojanathammanee L, Murphy EJ, Combs CK (2011) Expression of mutant alpha-synuclein modulates microglial phenotype in vitro. *J Neuroinflammation* 8:44
86. Zhang W et al (2005) Aggregated alpha-synuclein activates microglia: a process leading to disease progression in Parkinson's disease. *FASEB J* 19(6):533–542
87. Rosen DR et al (1993) Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 362(6415):59–62
88. Sakuma R et al (1995) A clinical variance in familial amyotrophic lateral sclerosis with a point mutation in Cu/Zn superoxide dismutase gene. *Eur J Neurol* 2(4):369–374
89. Kawamata T et al (1992) Immunologic reactions in amyotrophic lateral sclerosis brain and spinal cord tissue. *Am J Pathol* 140(3):691–707
90. Ince PG et al (1996) Familial amyotrophic lateral sclerosis with a mutation in exon 4 of the Cu/Zn superoxide dismutase gene: pathological and immunocytochemical changes. *Acta Neuropathol* 92(4):395–403
91. Graber DJ, Hickey WF, Harris BT (2010) Progressive changes in microglia and macrophages in spinal cord and peripheral nerve in the transgenic rat model of amyotrophic lateral sclerosis. *J Neuroinflammation* 7:8
92. Sanagi T et al (2010) Appearance of phagocytic microglia adjacent to motoneurons in spinal cord tissue from a presymptomatic transgenic rat model of amyotrophic lateral sclerosis. *J Neurosci Res* 88(12):2736–2746
93. Boivin A et al (2007) Toll-like receptor signaling is critical for Wallerian degeneration and functional recovery after peripheral nerve injury. *J Neurosci* 27(46):12565–12576
94. Perrin FE et al (2005) Involvement of monocyte chemoattractant protein-1, macrophage inflammatory protein-1alpha and interleukin-1beta in Wallerian degeneration. *Brain* 128(Pt 4):854–866
95. Henkel JS et al (2006) The chemokine MCP-1 and the dendritic and myeloid cells it attracts are increased in the mSOD1 mouse model of ALS. *Mol Cell Neurosci* 31(3):427–437
96. Polfliet MM et al (2002) The role of perivascular and meningeal macrophages in experimental allergic encephalomyelitis. *J Neuroimmunol* 122(1–2):1–8
97. Rinner WA et al (1995) Resident microglia and hematogenous macrophages as phagocytes in adoptively transferred experimental autoimmune encephalomyelitis: an investigation using rat radiation bone marrow chimeras. *Glia* 14(4):257–266
98. Schonrock LM et al (1998) Identification of glial cell proliferation in early multiple sclerosis lesions. *Neuropathol Appl Neurobiol* 24(4):320–330
99. Tanaka T, Ueno M, Yamashita T (2009) Engulfment of axon debris by microglia requires p38 MAPK activity. *J Biol Chem* 284(32):21626–21636
100. Bruck W et al (1996) Macrophages in multiple sclerosis. *Immunobiology* 195(4–5):588–600
101. Han MH et al (2012) Janus-like opposing roles of CD47 in autoimmune brain inflammation in humans and mice. *J Exp Med* 209(7):1325–1334
102. Badie B, Schartner JM (2000) Flow cytometric characterization of tumor-associated macrophages in experimental gliomas. *Neurosurgery* 46(4):957–961, discussion 961–2
103. Sasaki A et al (2013) Characterization of microglia/macrophages in gliomas developed in S-100beta-v-erbB transgenic rats. *Neuropathology* 33(5):505–514
104. Markovic DS et al (2005) Microglia stimulate the invasiveness of glioma cells by increasing the activity of metalloprotease-2. *J Neuropathol Exp Neurol* 64(9):754–762
105. Hanisch UK, Kettenmann H (2007) Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 10(11):1387–1394
106. Chang GH, Barbaro NM, Pieper RO (2000) Phosphatidylserine-dependent phagocytosis of apoptotic glioma cells by normal human microglia, astrocytes, and glioma cells. *Neuro Oncol* 2(3):174–183
107. Chiu TL, Peng CW, Wang MJ (2011) Enhanced anti-glioblastoma activity of microglia by AAV2-mediated IL-12 through TRAIL and phagocytosis in vitro. *Oncol Rep* 25(5):1373–1380

108. Wu A et al (2010) Glioma cancer stem cells induce immunosuppressive macrophages/microglia. *Neuro Oncol* 12(11):1113–1125
109. Hughes MM et al (2010) Microglia in the degenerating brain are capable of phagocytosis of beads and of apoptotic cells, but do not efficiently remove PrPSc, even upon LPS stimulation. *Glia* 58(16):2017–2030
110. Kranich J et al (2010) Engulfment of cerebral apoptotic bodies controls the course of prion disease in a mouse strain-dependent manner. *J Exp Med* 207(10):2271–2281
111. Polazzi E, Monti B (2010) Microglia and neuroprotection: from in vitro studies to therapeutic applications. *Prog Neurobiol* 92(3):293–315
112. Siskova Z et al (2009) Degenerating synaptic boutons in prion disease: microglia activation without synaptic stripping. *Am J Pathol* 175(4):1610–1621
113. Raivich G et al (1999) Neuroglial activation repertoire in the injured brain: graded response, molecular mechanisms and cues to physiological function. *Brain Res Brain Res Rev* 30(1):77–105
114. Kalm M et al (2009) Irradiation-induced loss of microglia in the young brain. *J Neuroimmunol* 206(1–2):70–75
115. Hellstrom NA et al (2009) Differential recovery of neural stem cells in the subventricular zone and dentate gyrus after ionizing radiation. *Stem Cells* 27(3):634–641
116. Espejel S, Romero R, Alvarez-Buylla A (2009) Radiation damage increases Purkinje neuron heterokaryons in neonatal cerebellum. *Ann Neurol* 66(1):100–109
117. Jenrow KA et al (2011) Combined atorvastatin and ramipril mitigate radiation-induced impairment of dentate gyrus neurogenesis. *J Neurooncol* 101(3):449–456
118. Gilmore SA et al (2003) Radiation-induced modulation of the microglial population in the normal and injured mature spinal cord. *Exp Neurol* 182(1):169–179
119. Peng Y et al (2013) Blockade of Kv1.3 channels ameliorates radiation-induced brain injury. *Neuro Oncol* (in press)
120. Song M et al (2011) TLR4 mutation reduces microglial activation, increases A $\beta$  deposits and exacerbates cognitive deficits in a mouse model of Alzheimer's disease. *J Neuroinflammation* 8:92
121. Fellner L et al (2013) Toll-like receptor 4 is required for alpha-synuclein dependent activation of microglia and astroglia. *Glia* 61(3):349–360
122. Ribes S et al (2010) Fibronectin stimulates *Escherichia coli* phagocytosis by microglial cells. *Glia* 58(3):367–376
123. Jones B (2013) Alzheimer disease: TREM2 linked to late-onset AD. *Nat Rev Neurol* 9(1):5
124. Okun E, Mattson MP, Arumugam TV (2010) Involvement of Fc receptors in disorders of the central nervous system. *Neuromolecular Med* 12(2):164–178
125. Cao S, Standaert DG, Harns AS (2012) The gamma chain subunit of Fc receptors is required for alpha-synuclein-induced pro-inflammatory signaling in microglia. *J Neuroinflammation* 9:259
126. Rogers J, Lue LF (2001) Microglial chemotaxis, activation, and phagocytosis of amyloid beta-peptide as linked phenomena in Alzheimer's disease. *Neurochem Int* 39(5–6):333–340
127. Hadas S et al (2012) Complement receptor-3 negatively regulates the phagocytosis of degenerated myelin through tyrosine kinase Syk and cofilin. *J Neuroinflammation* 9:166
128. Hendrickx DA et al (2013) Selective upregulation of scavenger receptors in and around demyelinating areas in multiple sclerosis. *J Neuropathol Exp Neurol* 72(2):106–118
129. Alarcon R et al (2005) Expression of scavenger receptors in glial cells. Comparing the adhesion of astrocytes and microglia from neonatal rats to surface-bound beta-amyloid. *J Biol Chem* 280(34):30406–30415
130. Husemann J et al (2002) Scavenger receptors in neurobiology and neuropathology: their role on microglia and other cells of the nervous system. *Glia* 40(2):195–205
131. Roberts K et al (2013) Extracellular aggregated Cu/Zn superoxide dismutase activates microglia to give a cytotoxic phenotype. *Glia* 61(3):409–419
132. Kobayashi K et al (2012) Multiple P2Y subtypes in spinal microglia are involved in neuropathic pain after peripheral nerve injury. *Glia* 60(10):1529–1539
133. Rotshenker S (2009) The role of galectin-3/MAC-2 in the activation of the innate-immune function of phagocytosis in microglia in injury and disease. *J Mol Neurosci* 39(1–2):99–103
134. Reichert F, Rotshenker S (1999) Galectin-3/MAC-2 in experimental allergic encephalomyelitis. *Exp Neurol* 160(2):508–514
135. Fricker M, Oliva-Martin MJ, Brown GC (2012) Primary phagocytosis of viable neurons by microglia activated with LPS or A $\beta$  is dependent on calreticulin/LRP phagocytic signalling. *J Neuroinflammation* 9:196
136. Deane R, Sagare A, Zlokovic BV (2008) The role of the cell surface LRP and soluble LRP in blood–brain barrier A $\beta$  clearance in Alzheimer's disease. *Curr Pharm Des* 14(16):1601–1605
137. Marzolo MP et al (2000) Expression of alpha(2)-macroglobulin receptor/low density lipoprotein receptor-related protein (LRP) in rat microglial cells. *J Neurosci Res* 60(3):401–411
138. De Simone R, Ajmone-Cat MA, Minghetti L (2004) Atypical anti-inflammatory activation of microglia induced by apoptotic neurons: possible role of phosphatidylserine-phosphatidylserine receptor interaction. *Mol Neurobiol* 29(2):197–212
139. Hirt UA, Leist M (2003) Rapid, noninflammatory and PS-dependent phagocytic clearance of necrotic cells. *Cell Death Differ* 10(10):1156–1164
140. Galea I et al (2005) Mannose receptor expression specifically reveals perivascular macrophages in normal, injured, and diseased mouse brain. *Glia* 49(3):375–384
141. Giraldo-Guimaraes A et al (2012) Bone marrow mononuclear cells and mannose receptor expression in focal cortical ischemia. *Brain Res* 1452:173–184