

## ORIGINAL ARTICLE

# The Frequency and Severity of Gastrointestinal Symptoms in Patients with Early Parkinson's Disease

Hye-Young Sung,<sup>a</sup> Jeong-Wook Park,<sup>b</sup> Joong-Seok Kim<sup>b</sup>

<sup>a</sup>Division of Gastroenterology, Digestive Disease Research Institute, Department of Internal Medicine, Wonkwang University Sanbon Hospital, Gunpo, Korea

<sup>b</sup>Department of Neurology, College of Medicine, The Catholic University of Korea, Seoul, Korea

## ABSTRACT

**Objective** Although gastrointestinal dysfunctions occur in the majority of patients with Parkinson's disease (PD), they are often unrecognized because many patients remain relatively asymptomatic in the early stage. We investigated the frequency of gastrointestinal symptoms in patients with PD using newly developed gastrointestinal symptom questionnaires.

**Methods** Early PD patients with a symptom duration not exceeding 3 years were included in this study. All PD patients were evaluated using a questionnaire, which consisted of three relevant domains: oropharyngoesophageal (10 items); gastric (3 items); and intestinal-anorectal (7 items). The frequency of symptoms was calculated as a proportion with an item score  $\geq 2$ .

**Results** Of the 54 patients enrolled, 48 patients (88.9%) responded that bowel symptoms developed before the onset of Parkinsonian motor symptoms, and four patients reported that the onset of two types of symptoms (i.e., bowel and neurological) occurred approximately simultaneously, with only months between them. The frequencies of gastrointestinal symptoms are as follows: speech disturbance (40.7%), drooling (24.1%), sense of getting stuck (31.5%), choking (27.8%), globus pharyngis (16.7%), repetitive deglutition (29.6%), pain during swallowing (5.6%), food regurgitation (3.7%), acid reflux (7.4%), nausea/vomiting (11.1%), early satiety (16.7%), postprandial fullness (14.8%), epigastric soreness (9.3%), abdominal pain (3.7%), constipation (46.3%), excessive strain during defecation (33.3%), fecal incontinence (7.4%), tenesmus (20.4%), loose stool or diarrhea (3.7%), and difficulty in relaxing anal sphincter (11.1%). Two patients were scored at zero.

**Conclusions** Our findings confirm that gastrointestinal dysfunction occurs in early PD in relatively high frequency.

**Key Words** Parkinson's disease; Gastrointestinal symptoms; Frequency; Severity.

Gastrointestinal (GI) symptoms are relatively common in patients suffering from Parkinson's disease (PD), and these symptoms can be a significant burden on patients and their caregivers. Recent advances in neuropathology address the role of autonomic nuclei in the brainstem and the enteric nervous system in the control of gastrointestinal function, explaining early involvement of this function in PD.<sup>1,2</sup>

The under-recognition and low levels of awareness of the presence of gastrointestinal dysfunctions by patients, caregiver, and health professionals result in underreporting and a low fre-

quency of complaints. Early detection and effective intervention can help prevent the serious consequences in the quality of life. Several questionnaires have previously been developed for screening GI symptoms;<sup>3-6</sup> however, many of these questionnaires relied on specific, subjective complaints and did not include many types of symptoms.

We developed new gastrointestinal symptom questionnaires and evaluated the prevalence of gastrointestinal symptoms in early PD patients.

Received: February 15, 2014 Revised: February 20, 2014 Accepted: February 20, 2014

Corresponding author: Joong-Seok Kim, MD, PhD, Department of Neurology, College of Medicine, The Catholic University of Korea, Seoul St. Mary's Hospital, 222 Banpo-daero, Seocho-gu, Seoul 137-701, Korea / Tel: +82-2-2258-6078 / Fax: +82-2-599-9686 / E-mail: neuronet@catholic.ac.kr

## METHODS

### Patients

Consecutive early PD patients were enrolled from the Movement Disorders Clinic at Seoul St. Mary's Hospital, Seoul, Korea. A clinical diagnosis of PD was made according to the UK brain bank criteria.<sup>7</sup> None of the patients had ever taken medication for PD. The evaluation procedure included a detailed medical and drug history, a physical and neurological examination and a brief neuropsychological assessment. Excluded from the study were patients with the following: 1) neurological abnormalities related to atypical PD or secondary Parkinsonism, 2) patients with diabetic neuropathy, and 3) patients taking medications known to influence gastrointestinal motility, such as levodopa, dopamine agonists, anticholinergics, or GI dysmotility remedies.

This study was approved by the local ethics committee, and all subjects gave their consent to participate.

### Clinical assessments

All patients were subsequently evaluated using the Unified Parkinson's Disease Rating Scale (UPDRS), the Hoehn and Yahr (H&Y) stage, and the Korean version of the Schwab and England activities of daily living. The patients were also asked to complete a self-administered scale to evaluate the presence and frequency or severity of bowel symptoms during the last 3 months. The self-administered questionnaire consisted of the following 20 items (Table 1): 1) speech disturbance, 2) drooling, 3) sense of getting stuck, 4) choking, 5) globus pharyngis, 6) repetitive deglutition, 7) pain during swallowing, 8) food regurgitation, 9) acid reflux, 10) nausea/vomiting, 11) early satiety, 12) postprandial fullness, 13) epigastric soreness, 14) abdominal pain, 15) constipation, 16) excessive strain during defecation, 17) fecal incontinence, 18) tenesmus, 19) loose stool or diarrhea, and 20) difficulty in relaxing anal sphincter. Each symptom was scored with respect to either the frequency [1 = rare (< 1/week), 2 = often (1/week), 3 = frequent (several times

**Table 1.** Score distribution of the gastrointestinal symptom scale

Measure	Mean	SD	Min.	Max.
Speech disturbance	2.17	3.08	0	15
Drooling	1.54	2.51	0	10
Sense of getting stuck	1.91	2.02	0	6
Choking	1.76	2.05	0	8
Globus (foreign body sensation)	1.02	2.10	0	10
Repetitive deglutition	1.89	2.60	0	10
Pain during swallowing	0.72	1.25	0	5
Food regurgitation	0.50	2.14	0	15
Acid reflux	0.74	2.42	0	15
Nausea/vomiting	0.94	2.09	0	15
Oropharyngo-esophageal subdomain	13.18	11.17	0	48
Early satiety	1.48	3.16	0	15
Postprandial fullness	0.91	1.50	0	5
Epigastric soreness	0.87	1.74	0	8
Gastric subdomain	3.26	4.74	0	23
Abdominal pain	0.93	1.93	0	12
Constipation	2.83	2.70	0	10
Excessive strain during defecation	2.35	2.69	0	10
Fecal incontinence	0.41	1.12	0	6
Tenesmus	1.65	2.45	0	15
Loose stool or diarrhea	0.22	0.92	0	6
Difficulty in relaxing anal sphincter	0.59	1.28	0	5
Intestinal-anorectal subdomain	8.98	8.00	0	33
Total gastrointestinal symptom score	25.43	18.62	0	97

SD: standard deviation, Min.: minimum, Max.: maximum.

per week), and 4 = always (daily or all the time)] or severity (0 = no or minimum, 1 = mild symptoms present but causing little distress or disturbance, 2 = moderate symptoms causing some distress or disturbance, and 3 = severe symptoms representing a major source of distress or disturbance) of the symptom. The overall score was calculated by multiple severity and frequency scores.

### Data analysis


Patients were categorized into the following two subgroups: 1) patients with no or mild problems (a score for each item of  $\leq 1$ ) and 2) patients with moderate-to-severe problems (a score for each item of  $\geq 2$ ), and the frequencies of each symptom were calculated using this categorical variable. Items were also grouped into three relevant domains: oropharyngoesophageal (10 items), gastric (3 items), and intestinal-anorectal (7 items). The symptom scores were also compared with Parkinsonian motor symp-

oms. Statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS) version 15.0 (SPSS Inc., Chicago, IL, USA), and the statistical significance level was established at  $p < 0.05$ .

### RESULTS

Among the 54 patients enrolled, 22 were men and 32 were women. The mean age at examination was  $67.1 \pm 10.3$  years (range, 33–86 years) and the mean motor symptom duration was  $14.7 \pm 9.3$  months (range, 1–36 months). The severity of Parkinsonian motor symptoms was as follows: mean total UPDRS score was  $25.1 \pm 18.6$  (part 1,  $2.3 \pm 2.3$ ; part 2,  $8.3 \pm 6.9$ ; part 3,  $14.4 \pm 10.6$ ), and the mean H&Y stage was  $1.6 \pm 0.4$ . Fourteen patients were scored as H&Y stage 1, 20 patients as stage 1.5, and 20 patients as stage 2.

The frequencies of gastrointestinal symptoms are



**Figure 1.** Gastrointestinal symptom frequencies in early Parkinson's disease patients.


as follows (Table 1, Figure 1): speech disturbance (40.7%), drooling (24.1%), sense of getting stuck (31.5%), choking (27.8%), globus (16.7%), fragmented deglutition (29.6%), pain during swallowing (5.6%), food regurgitation (3.7%), acid reflux (7.4%), nausea/vomiting (11.1%), early satiety (16.7%), postprandial fullness (14.8%), epigastric soreness (9.3%), abdominal pain (3.7%), constipation (46.3%), excessive strain during defecation (33.3%), fecal incontinence (7.4%), tenesmus (20.4%), loose stool or diarrhea (3.7%), and difficulty in relaxing anal sphincter (11.1%). Only two patients scored 0 on the self-administered scale. Forty-eight patients (88.9%) responded that the bowel symptoms developed before the onset of Parkinsonian motor symptoms and worsened after the onset, and four patients reported that the onset of two types of symptoms (i.e., bowel and neurological) occurred approximately simultaneously, with only months between them.

The gastrointestinal symptoms were positively correlated with the UPDRS scores and, in particular, the activity of daily living (ADL) portion of the scores (Figure 2). However, neither the specific motor subtype nor the disease duration was associated with the severity of bowel symptoms.

## DISCUSSION

The results of this study suggest that bowel symp-

toms are present in a majority of early, non-treated PD patients, even before the onset of clinical motor symptoms. The most frequent symptoms found in our patients were difficulty with deglutition and constipation. Dysphagia in patients with PD may be associated with abnormalities found during the oral to esophageal stages of swallowing.<sup>8</sup> Dysphagia can result from the motor symptoms of PD involving the oropharyngeal muscles or can occur from defective coordination of the oropharyngeal and esophageal musculature resulting from brainstem involvement.<sup>9</sup> About half of the patients with early PD in our study reported a variety of complaints associated with dysphagia. In particular, repetitive deglutition was predominantly related to oropharyngoesophageal dysfunction. Recently, we clearly showed that a majority of early PD patients underwent repetitive swallow attempts (double or triple swallow pattern) for a single bolus.<sup>8</sup> These abnormal swallow patterns were shown to be directly associated with abnormal peristalsis in the esophageal manometry. Repetitive deglutition might be a reactive response necessary to clear remnant bolus material on the pharynx. Such a repetitive deglutition was closely related to esophageal dysfunction though the vagal inhibitory dysfunction. Under normal physiological conditions, while there are repetitive swallow attempts, esophageal peristalsis only responds to the last stimuli, called deglutitive inhibition. Normal deglutitive in-



**Figure 2.** Correlation between gastrointestinal symptom score and Unified Parkinson's Disease Rating Scale (UPDRS); (A) total and (B) activity of daily living (ADL; part 2).

hibition indicates intact vagal inhibitory function. Deglutition-evoked peristaltic sequences are mediated via vagal efferent nerves, and peristalsis occurs with active inhibition followed by sequential esophageal contractions that are centrally mediated.<sup>10</sup> Therefore, repetitive deglutition may be the most relevant dysfunction, due to its association with esophageal dysmotility, and the above described abnormal deglutitive inhibition provides further evidence of neuronal degeneration of the brain stem swallowing center or abnormalities of the peripheral vagus nerve in patients with early PD.<sup>8</sup>

Early-stage PD patients in our study showed dysfunctions of the proximal part of the gastrointestinal system, encompassing the stomach and small intestine. Prior reports have suggested that impaired gastric motility is associated with different medications used for the treatment of PD, especially in those patients with response fluctuation. Although major pathological changes in the enteric nervous system were described, some previous studies indicate that the presence of gastroparesis in only 7.5% of PD patients was related to PD itself.<sup>11,12</sup> In this study, we show that the proportions of gastric symptoms ranged from 3.7% to 16.7%. Gastric symptoms are a well-recognized manifestation of PD, which can cause not only nausea and other symptoms, such as gastroparesis, but can also affect drug absorption.<sup>13</sup> In addition, antiparkinson medications may further exacerbate these symptoms. Therefore, it is important that clinicians help patients with PD find and overcome these disabling symptoms.

The most common gastrointestinal symptoms in our study were related to constipation and difficulty in defecation. Constipation and defecatory dysfunction in patients with PD may be associated with abnormalities occurring during colonic inertia to the anal outlet.<sup>14</sup> Defecatory dysfunction is a consequence of uncoordinated action of the muscles involved with defecation. Relaxation of the puborectalis muscle allows for the opening of the anorectal angle and perineal descent, facilitating fecal expulsion. This functional outlet obstruction may cause both excessive straining and a sense of incomplete emptying, or sometimes, may induce painful defecation.<sup>15-17</sup> Another mechanism appears to involve the colonic musculature, which induces a slow passage of feces through the colon.<sup>14</sup> Several studies have demonstrated considerably prolonged mean

colonic transit time in PD patients.<sup>18</sup>

Gastrointestinal symptoms were correlated with the UPDRS score, particularly the ADL score. This suggests that the severity of motor symptoms might be closely related to enteric involvement, which is distinct from the degree of nigrostriatal involvement, and other neurotransmitter systems can be selectively affected.

In summary, most of the included patients with early PD had several instances of gastrointestinal symptoms. Many patients reported that these symptoms developed before the clinical motor manifestation of PD. These findings suggest selective involvement of either the brain stem and/or the enteric nervous system at an early stage of PD. The more severe and frequent gastrointestinal symptoms were associated with severe motor symptoms but did not correlate with the duration of PD. Comprehensive screening of gastrointestinal symptoms in patients with early-stage PD can facilitate prompt recognition and effective therapeutic interventions for significant bowel dysfunctions.

#### Conflicts of Interest

The authors have no financial conflicts of interest.

#### REFERENCES

1. Braak H, de Vos RA, Bohl J, Del Tredici K. Gastric alpha-synuclein immunoreactive inclusions in Meissner's and Auerbach's plexuses in cases staged for Parkinson's disease-related brain pathology. *Neurosci Lett* 2006;396:67-72.
2. Hawkes CH, Del Tredici K, Braak H. Parkinson's disease: a dual-hit hypothesis. *Neuropathol Appl Neurobiol* 2007;33:599-614.
3. Edwards LL, Pfeiffer RF, Quigley EM, Hofman R, Balluff M. Gastrointestinal symptoms in Parkinson's disease. *Mov Disord* 1991;6:151-156.
4. Byrne KG, Pfeiffer R, Quigley EM. Gastrointestinal dysfunction in Parkinson's disease. A report of clinical experience at a single center. *J Clin Gastroenterol* 1994;19:11-16.
5. Edwards LL, Quigley EM, Pfeiffer RF. Gastrointestinal dysfunction in Parkinson's disease: frequency and pathophysiology. *Neurology* 1992;42:726-732.
6. Cersosimo MG, Raina GB, Pecci C, Pellene A, Calandra CR, Gutiérrez C, et al. Gastrointestinal manifestations in Parkinson's disease: prevalence and occurrence before motor symptoms. *J Neurol* 2013;260:1332-1338.
7. Gibb WR, Lees AJ. The relevance of the Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1988;51:745-752.
8. Sung HY, Kim JS, Lee KS, Kim YI, Song IU, Chung SW, et al. The prevalence and patterns of pharyngoesophageal dysmotility in patients with early stage Parkinson's disease. *Mov Disord* 2010;25:2361-2368.
9. Cersosimo MG, Benarroch EE. Neural control of the gastrointestinal tract: implications for Parkinson disease. *Mov*

- Disord 2008;23:1065-1075.
10. Vanek AW, Diamant NE. Responses of the human esophagus to paired swallows. *Gastroenterology* 1987;92:643-650.
  11. Djaldetti R, Baron J, Ziv I, Melamed E. Gastric emptying in Parkinson's disease: patients with and without response fluctuations. *Neurology* 1996;46:1051-1054.
  12. Soykan I, Sivri B, Sarosiek I, Kiernan B, McCallum RW. Demography, clinical characteristics, psychological and abuse profiles, treatment, and long-term follow-up of patients with gastroparesis. *Dig Dis Sci* 1998;43:2398-2404.
  13. Gurevich T, Korczyn AD, Giladi N. Gastric dysfunction. In: Pfeiffer RF, Bodis-Wollner I. *Parkinson's disease and non-motor dysfunction*. New Jersey: Humana Press, 2005:105-113.
  14. Kim JS, Sung HY, Lee KS, Kim YI, Kim HT. Anorectal dysfunctions in Parkinson's disease. *J Neurol Sci* 2011;310:144-151.
  15. Mathers SE, Kempster PA, Swash M, Lees AJ. Constipation and paradoxical puborectalis contraction in anismus and Parkinson's disease: a dystonic phenomenon? *J Neurol Neurosurg Psychiatry* 1988;51:1503-1507.
  16. Mathers SE, Kempster PA, Law PJ, Frankel JP, Bartram CI, Lees AJ, et al. Anal sphincter dysfunction in Parkinson's disease. *Arch Neurol* 1989;46:1061-1064.
  17. Sung HY, Choi MG, Kim YI, Lee KS, Kim JS. Anorectal manometric dysfunctions in newly diagnosed, early-stage Parkinson's disease. *J Clin Neurol* 2012;8:184-189.
  18. Sakakibara R, Odaka T, Uchiyama T, Asahina M, Yamaguchi K, Yamaguchi T, et al. Colonic transit time and recto-anal videomanometry in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2003;74:268-272.