

Diaquabis[2,6-bis(4*H*-1,2,4-triazol-4-yl)-pyridine- κ N²]bis(selenocyanato- κ N)-cobalt(II)

Yuan-Yuan Liu* and Pan Yang

Tianjin Key Laboratory of Structure and Performance for Functional Molecule, Tianjin Normal University, Tianjin 300071, People's Republic of China

Correspondence e-mail: liuyuan1973@yahoo.com.cn


Received 12 July 2012; accepted 17 July 2012

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.026; wR factor = 0.065; data-to-parameter ratio = 12.3.

In the title compound, $[\text{Co}(\text{NCSe})_2(\text{C}_9\text{H}_7\text{N}_7)_2(\text{H}_2\text{O})_2]$, the Co^{2+} cation is coordinated by two selenocyanate anions, two 2,6-bis(4*H*-1,2,4-triazol-4-yl)pyridine ligands and two water molecules within a slightly distorted N_4O_2 octahedron. The asymmetric unit consists of one Co^{2+} cation, which is located on a center of inversion, as well as one selenocyanate anion, one 2,6-bis(4*H*-1,2,4-triazol-4-yl)pyridine ligand and one water molecule in general positions. Intermolecular $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds join the complex molecules into layers parallel to the bc plane. The layers are linked by $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{Se}$ hydrogen bonds into a three-dimensional supramolecular architecture.

Related literature

For general background to this work, see: Liu *et al.* (2007). Previous research on compounds with Co(II) as cation have found a slow relaxation of the magnetization, see: Boeckmann & Näther (2010, 2011, 2012). For related structures, see: Du *et al.* (2009); Yang *et al.* (2008).



Experimental

Crystal data

$[\text{Co}(\text{NCSe})_2(\text{C}_9\text{H}_7\text{N}_7)_2(\text{H}_2\text{O})_2]$	$V = 2580.4$ (4) Å ³
$M_r = 731.35$	$Z = 4$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 17.5460$ (16) Å	$\mu = 3.54$ mm ⁻¹
$b = 7.2752$ (7) Å	$T = 173$ K
$c = 20.3148$ (19) Å	$0.15 \times 0.14 \times 0.13$ mm
$\beta = 95.691$ (2)°	

Data collection

Bruker APEXII CCD diffractometer	6336 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	2280 independent reflections
$T_{\min} = 0.619$, $T_{\max} = 0.656$	2083 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$	186 parameters
$wR(F^2) = 0.065$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\text{max}} = 0.59$ e Å ⁻³
2280 reflections	$\Delta\rho_{\text{min}} = -0.58$ e Å ⁻³

Table 1

Selected bond lengths (Å).

Co1—N8	2.097 (2)	Co1—O1	2.1434 (18)
Co1—N3	2.122 (2)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A \cdots N2	0.84	2.43	3.017 (3)	127
O1—H1B \cdots N7 ⁱⁱⁱ	0.84	2.00	2.837 (3)	173
C1—H1 \cdots N6 ⁱⁱⁱ	0.95	2.37	3.293 (3)	163
C5—H5 \cdots N8 ^{iv}	0.95	2.56	3.356 (3)	142
C7—H7 \cdots N6 ⁱⁱⁱ	0.95	2.46	3.373 (3)	162
C9—H9 \cdots Se1 ^{iv}	0.95	2.96	3.877 (3)	164

 Symmetry codes: (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$; (iii) $x + \frac{1}{2}, y + \frac{1}{2}, z$; (iv) $-x + 2, -y + 1, -z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL and DIAMOND (Brandenburg, 1999); software used to prepare material for publication: publCIF (Westrip, 2010).

This work was supported financially by Tianjin Educational Committee (20090504, 20110301) and Tianjin Normal University (1E0402B).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZJ2088).

References

- Boeckmann, J. & Näther, C. (2010). *Dalton Trans.* **39**, 11019–11026.
 Boeckmann, J. & Näther, C. (2011). *Chem. Commun.* **47**, 7104–7106.
 Boeckmann, J. & Näther, C. (2012). *Polyhedron*, **31**, 587–595.
 Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Du, Z.-Y., Sun, Y.-H., Liu, Q.-Y., Xie, Y.-R. & Wen, H.-R. (2009). *Inorg. Chem.* **48**, 7015–7017.

Liu, Y. Y., Huang, Y. Q., Shi, W., Cheng, P., Liao, D. Z. & Yan, S. P. (2007). *Cryst. Growth Des.* **7**, 1483–1489.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
Yang, B.-P., Prosvirin, A. V., Guo, Y.-Q. & Mao, J.-G. (2008). *Inorg. Chem.* **47**, 1453–1459.

supplementary materials

Acta Cryst. (2012). E68, m1097–m1098 [doi:10.1107/S1600536812032461]

Diaquabis[2,6-bis(4*H*-1,2,4-triazol-4-yl)pyridine- κ N²]bis(selenocyanato- κ N)cobalt(II)

Yuan-Yuan Liu and Pan Yang

Comment

Previously we have reported on the a series of novel zinc(II) and cadmium(II) compounds based on 2,6-di-(1,2,4-triazole-4-yl)pyridine (Liu *et al.*, 2007). On the other hand, dependent on the nature of the metal cation, anti- or ferromagnetic ordering is observed and for the compounds with Co(II) as cation previous resrarch have found a slow relaxation of the magnetization (Boeckmann & Näther 2010, 2011, 2012). To investigate the influence of the co-ligand on the magnetic properties for the compounds with Co(II), we tried to prepare cobalt(II) compounds based on 2,6-di-(1,2,4-triazole-4-yl)pyridine, which resulted in the formation of the title compound in which the neutral ligands are only terminal N-coordinated. This compound was characterized only by single-crystal X-ray diffraction. In the crystal structure the cobalt(II) cations are coordinated by four nitrogen atoms of two terminal N-bonded seleno-cyanato anions and two terminal bonded 2,6-di-(1,2,4-triazole-4-yl)pyridine co-ligands as well as two water molecules into discrete complexes (Fig. 1). The coordination polyhedron of the Co cations can be described as a slightly distorted octahedron with the Co cation located on a centre of inversion. The discrete cobalt complexes are bridged by intermolecular O—H \cdots N, C—H \cdots N and C—H \cdots Se hydrogen bonds (Yang *et al.*, 2008; Du *et al.*, 2009), wich assemble (I) into a three-dimensional supra-molecular architecture(Fig. 2 and Table 1).

Perspective drawing with the atomic numbering scheme is illustrated in figure 1. Selected geometric parameters (Å, °) for (I) are listed in table 1. Selected hydrogen-bonding geometric parameters (Å, °) for (I) are listed in table 2. The two-dimensional supramolecular framework of (I) is shown in Figure 2.

Experimental

The compound was synthesized under hydrothermal conditions. A mixture of *L* (*L* = 1,4-Bis(2,6-Bis(4*H*-1,2,4-triazol-4-yl)pyridine) (0.3 mmol, 0.0636 g), CoSO₄·7H₂O (0.1 mmol, 0.028 g), KSeCN (0.2 mmol, 0.029 g) and water (10 ml) was placed in a 25 ml acid digestion bomb and heated at 393 K for two days, then equably cooled to room temperature for three days. After washed by 5 ml water for twice, Red block crystals of the compound were obtained..

Refinement

The water H atoms were located in a Fourier difference map and refined subject to an O—H restraint 0.88 (1) Å and an H \cdots H restraint of 1.42 (2) Å. Other H atoms were allowed to ride on their parent atoms with C—H distances of 0.93 Å (*U*_{iso}(H) = 1.2*U*_{eq}(C)). All of the non-hydrogen atoms were refined anisotropically..

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97*

(Sheldrick, 2008); molecular graphics: *SHELXTL* and *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *publCIF* (Westrip, 2010).





Figure 1

The structure of the title complex, showing 50% probability displacement ellipsoids and the atom-numbering schemes. Atoms of the inversion-related half-complex have symmetry code: $(2 - x, y, 1/2 - z)$.


Figure 2

The three-dimensional layer structure of the title complex. Purple Dashed lines indicate O—H···N, C—H···N and C—H···Se hydrogen bonds.

Diaquabis[2,6-bis(4*H*-1,2,4-triazol-4-yl)pyridine- κ N²]bis(selenocyanato- κ N)cobalt(II)
Crystal data

[Co(NCSe)₂(C₉H₇N₇)₂(H₂O)₂]

$M_r = 731.35$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 17.5460$ (16) Å

$b = 7.2752$ (7) Å

$c = 20.3148$ (19) Å

$\beta = 95.691$ (2)°

$V = 2580.4$ (4) Å³

$Z = 4$

$F(000) = 1444$

$D_x = 1.883$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

$\mu = 3.54$ mm⁻¹

$T = 173$ K

Block, red

$0.15 \times 0.14 \times 0.13$ mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.619$, $T_{\max} = 0.656$

6336 measured reflections

2280 independent reflections

2083 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$
 $\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 2.3^\circ$

$h = -20 \rightarrow 9$
 $k = -8 \rightarrow 8$
 $l = -23 \rightarrow 24$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.065$
 $S = 1.05$
 2280 reflections
 186 parameters

0 restraints
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0325P)^2 + 4.1522P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 0.59 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.58 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	1.0000	0.85337 (6)	0.2500	0.01309 (13)
Se1	1.238743 (15)	0.51140 (3)	0.329458 (13)	0.02053 (11)
O1	1.08413 (10)	1.0675 (2)	0.26203 (8)	0.0171 (4)
H1A	1.1136	1.0578	0.2321	0.026*
H1B	1.1116	1.0623	0.2983	0.026*
N1	1.01232 (11)	0.7890 (3)	0.04271 (10)	0.0134 (4)
N2	1.08763 (12)	0.9006 (3)	0.12643 (10)	0.0171 (5)
N3	1.01748 (11)	0.8524 (3)	0.14815 (10)	0.0143 (4)
N4	0.91149 (11)	0.6660 (3)	-0.02481 (10)	0.0125 (4)
N5	0.80409 (12)	0.5469 (3)	-0.08553 (10)	0.0136 (4)
N6	0.69319 (13)	0.4701 (3)	-0.05234 (12)	0.0217 (5)
N7	0.68675 (13)	0.4682 (3)	-0.12169 (11)	0.0182 (5)
N8	1.09050 (12)	0.6689 (3)	0.27341 (10)	0.0180 (5)
C1	1.08281 (14)	0.8611 (3)	0.06371 (12)	0.0156 (5)
H1	1.1228	0.8797	0.0361	0.019*
C2	0.97369 (14)	0.7874 (3)	0.09813 (12)	0.0139 (5)
H2	0.9227	0.7452	0.0998	0.017*
C3	0.98360 (13)	0.7256 (3)	-0.02156 (12)	0.0114 (5)
C4	0.88145 (14)	0.6091 (3)	-0.08405 (12)	0.0135 (5)
C5	0.91929 (15)	0.6072 (3)	-0.14054 (12)	0.0160 (5)
H5	0.8949	0.5656	-0.1817	0.019*
C6	0.99442 (14)	0.6689 (3)	-0.13447 (12)	0.0163 (5)
H6	1.0226	0.6702	-0.1720	0.020*
C7	1.02849 (14)	0.7287 (3)	-0.07389 (12)	0.0150 (5)
H7	1.0801	0.7698	-0.0685	0.018*
C8	0.76306 (16)	0.5177 (3)	-0.03236 (14)	0.0199 (6)

H8	0.7829	0.5305	0.0126	0.024*
C9	0.75337 (15)	0.5140 (3)	-0.13963 (13)	0.0171 (6)
H9	0.7651	0.5233	-0.1842	0.020*
C10	1.14850 (15)	0.6077 (3)	0.29588 (12)	0.0167 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0104 (2)	0.0163 (2)	0.0123 (2)	0.000	-0.00015 (19)	0.000
Se1	0.01776 (16)	0.02176 (16)	0.02165 (17)	0.00585 (11)	-0.00019 (12)	0.00176 (10)
O1	0.0126 (8)	0.0238 (9)	0.0150 (9)	-0.0010 (8)	0.0014 (7)	-0.0007 (7)
N1	0.0111 (10)	0.0154 (10)	0.0138 (11)	-0.0007 (8)	0.0017 (8)	-0.0003 (8)
N2	0.0118 (11)	0.0201 (10)	0.0192 (12)	-0.0021 (9)	0.0015 (9)	-0.0001 (9)
N3	0.0113 (10)	0.0159 (10)	0.0158 (11)	-0.0006 (8)	0.0020 (9)	0.0017 (8)
N4	0.0110 (10)	0.0127 (9)	0.0138 (10)	0.0004 (8)	0.0017 (8)	0.0005 (8)
N5	0.0101 (10)	0.0156 (10)	0.0150 (11)	-0.0016 (8)	0.0012 (9)	-0.0010 (8)
N6	0.0156 (11)	0.0293 (12)	0.0205 (12)	-0.0058 (10)	0.0033 (10)	-0.0008 (10)
N7	0.0151 (11)	0.0208 (11)	0.0185 (12)	-0.0006 (9)	0.0005 (9)	-0.0004 (9)
N8	0.0191 (12)	0.0200 (10)	0.0151 (11)	0.0031 (10)	0.0023 (9)	-0.0022 (9)
C1	0.0121 (12)	0.0177 (12)	0.0171 (13)	-0.0028 (10)	0.0017 (10)	-0.0015 (10)
C2	0.0122 (11)	0.0159 (12)	0.0140 (12)	-0.0007 (10)	0.0032 (10)	0.0016 (10)
C3	0.0105 (11)	0.0089 (11)	0.0145 (12)	0.0003 (9)	-0.0002 (10)	0.0012 (9)
C4	0.0106 (12)	0.0108 (11)	0.0189 (13)	-0.0008 (10)	0.0007 (10)	0.0007 (10)
C5	0.0182 (13)	0.0155 (11)	0.0145 (12)	-0.0021 (10)	0.0021 (10)	-0.0032 (10)
C6	0.0174 (13)	0.0181 (12)	0.0140 (12)	0.0005 (10)	0.0055 (10)	0.0013 (10)
C7	0.0104 (11)	0.0157 (12)	0.0190 (13)	0.0000 (10)	0.0021 (10)	0.0022 (10)
C8	0.0171 (14)	0.0266 (14)	0.0161 (14)	-0.0047 (11)	0.0019 (11)	-0.0026 (10)
C9	0.0157 (13)	0.0193 (13)	0.0160 (14)	-0.0017 (10)	0.0002 (11)	-0.0005 (10)
C10	0.0198 (14)	0.0157 (12)	0.0152 (13)	0.0000 (11)	0.0052 (11)	-0.0036 (10)

Geometric parameters (\AA , $^\circ$)

Co1—N8 ⁱ	2.097 (2)	N5—C8	1.373 (3)
Co1—N8	2.097 (2)	N5—C4	1.428 (3)
Co1—N3 ⁱ	2.122 (2)	N6—C8	1.300 (4)
Co1—N3	2.122 (2)	N6—N7	1.402 (3)
Co1—O1	2.1434 (18)	N7—C9	1.302 (3)
Co1—O1 ⁱ	2.1433 (17)	N8—C10	1.162 (3)
Se1—C10	1.803 (3)	C1—H1	0.9500
O1—H1A	0.8400	C2—H2	0.9500
O1—H1B	0.8400	C3—C7	1.385 (3)
N1—C2	1.371 (3)	C4—C5	1.382 (3)
N1—C1	1.372 (3)	C5—C6	1.387 (4)
N1—C3	1.429 (3)	C5—H5	0.9500
N2—C1	1.301 (3)	C6—C7	1.384 (3)
N2—N3	1.393 (3)	C6—H6	0.9500
N3—C2	1.300 (3)	C7—H7	0.9500
N4—C4	1.331 (3)	C8—H8	0.9500
N4—C3	1.333 (3)	C9—H9	0.9500
N5—C9	1.365 (3)		

N8 ⁱ —Co1—N8	100.44 (12)	C9—N7—N6	107.1 (2)
N8 ⁱ —Co1—N3 ⁱ	92.26 (8)	C10—N8—Co1	161.5 (2)
N8—Co1—N3 ⁱ	87.48 (8)	N2—C1—N1	111.0 (2)
N8 ⁱ —Co1—N3	87.48 (8)	N2—C1—H1	124.5
N8—Co1—N3	92.26 (8)	N1—C1—H1	124.5
N3 ⁱ —Co1—N3	179.60 (11)	N3—C2—N1	109.7 (2)
N8 ⁱ —Co1—O1	171.22 (7)	N3—C2—H2	125.2
N8—Co1—O1	86.66 (7)	N1—C2—H2	125.2
N3 ⁱ —Co1—O1	93.20 (7)	N4—C3—C7	125.2 (2)
N3—Co1—O1	87.09 (7)	N4—C3—N1	113.4 (2)
N8 ⁱ —Co1—O1 ⁱ	86.66 (7)	C7—C3—N1	121.4 (2)
N8—Co1—O1 ⁱ	171.22 (7)	N4—C4—C5	125.0 (2)
N3 ⁱ —Co1—O1 ⁱ	87.09 (7)	N4—C4—N5	114.1 (2)
N3—Co1—O1 ⁱ	93.20 (7)	C5—C4—N5	120.9 (2)
O1—Co1—O1 ⁱ	86.76 (9)	C4—C5—C6	117.0 (2)
Co1—O1—H1A	108.8	C4—C5—H5	121.5
Co1—O1—H1B	113.3	C6—C5—H5	121.5
H1A—O1—H1B	106.9	C7—C6—C5	120.2 (2)
C2—N1—C1	104.6 (2)	C7—C6—H6	119.9
C2—N1—C3	126.1 (2)	C5—C6—H6	119.9
C1—N1—C3	129.3 (2)	C6—C7—C3	116.7 (2)
C1—N2—N3	106.3 (2)	C6—C7—H7	121.6
C2—N3—N2	108.43 (19)	C3—C7—H7	121.6
C2—N3—Co1	129.19 (17)	N6—C8—N5	110.3 (2)
N2—N3—Co1	121.72 (15)	N6—C8—H8	124.8
C4—N4—C3	115.8 (2)	N5—C8—H8	124.8
C9—N5—C8	104.8 (2)	N7—C9—N5	110.6 (2)
C9—N5—C4	127.9 (2)	N7—C9—H9	124.7
C8—N5—C4	127.2 (2)	N5—C9—H9	124.7
C8—N6—N7	107.2 (2)	N8—C10—Se1	179.1 (2)
C1—N2—N3—C2	-0.3 (3)	C4—N4—C3—C7	-1.3 (3)
C1—N2—N3—Co1	171.20 (16)	C4—N4—C3—N1	178.59 (19)
N8 ⁱ —Co1—N3—C2	12.0 (2)	C2—N1—C3—N4	2.0 (3)
N8—Co1—N3—C2	112.4 (2)	C1—N1—C3—N4	-179.1 (2)
N3 ⁱ —Co1—N3—C2	62.2 (3)	C2—N1—C3—C7	-178.1 (2)
O1—Co1—N3—C2	-161.1 (2)	C1—N1—C3—C7	0.8 (4)
O1 ⁱ —Co1—N3—C2	-74.5 (2)	C3—N4—C4—C5	0.3 (3)
N8 ⁱ —Co1—N3—N2	-157.53 (17)	C3—N4—C4—N5	-179.81 (19)
N8—Co1—N3—N2	-57.18 (17)	C9—N5—C4—N4	166.4 (2)
N3 ⁱ —Co1—N3—N2	-107.3 (3)	C8—N5—C4—N4	-9.4 (3)
O1—Co1—N3—N2	29.36 (17)	C9—N5—C4—C5	-13.7 (4)
O1 ⁱ —Co1—N3—N2	115.95 (17)	C8—N5—C4—C5	170.5 (2)
C8—N6—N7—C9	0.3 (3)	N4—C4—C5—C6	0.3 (4)
N8 ⁱ —Co1—N8—C10	-156.3 (7)	N5—C4—C5—C6	-179.6 (2)
N3 ⁱ —Co1—N8—C10	-64.4 (6)	C4—C5—C6—C7	0.1 (4)
N3—Co1—N8—C10	115.9 (6)	C5—C6—C7—C3	-0.9 (3)
O1—Co1—N8—C10	28.9 (6)	N4—C3—C7—C6	1.6 (4)

O1 ⁱ —Co1—N8—C10	-12.6 (10)	N1—C3—C7—C6	-178.2 (2)
N3—N2—C1—N1	0.2 (3)	N7—N6—C8—N5	-0.3 (3)
C2—N1—C1—N2	0.0 (3)	C9—N5—C8—N6	0.3 (3)
C3—N1—C1—N2	-179.1 (2)	C4—N5—C8—N6	176.8 (2)
N2—N3—C2—N1	0.3 (3)	N6—N7—C9—N5	-0.1 (3)
Co1—N3—C2—N1	-170.33 (15)	C8—N5—C9—N7	-0.1 (3)
C1—N1—C2—N3	-0.2 (3)	C4—N5—C9—N7	-176.6 (2)
C3—N1—C2—N3	178.9 (2)	Co1—N8—C10—Se1	-132 (15)

Symmetry code: (i) $-x+2, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1 <i>A</i> ...N2	0.84	2.43	3.017 (3)	127
O1—H1 <i>B</i> ...N7 ⁱⁱ	0.84	2.00	2.837 (3)	173
C1—H1...N6 ⁱⁱⁱ	0.95	2.37	3.293 (3)	163
C5—H5...N8 ^{iv}	0.95	2.56	3.356 (3)	142
C7—H7...N6 ⁱⁱⁱ	0.95	2.46	3.373 (3)	162
C9—H9...Se1 ^{iv}	0.95	2.96	3.877 (3)	164

Symmetry codes: (ii) $x+1/2, -y+3/2, z+1/2$; (iii) $x+1/2, y+1/2, z$; (iv) $-x+2, -y+1, -z$.