

Research article

Open Access

## Cigarette smoking and tooth loss experience among young adults: a national record linkage study

Miki Ojima<sup>1</sup>, Takashi Hanioka<sup>\*2</sup>, Keiko Tanaka<sup>3</sup> and Hitoshi Aoyama<sup>4</sup>

Address: <sup>1</sup>Department of Preventive Dentistry, Graduate School of Dentistry, Osaka University, Osaka, Japan, <sup>2</sup>Department of Preventive and Public Health Dentistry, Fukuoka Dental College, Fukuoka, Japan, <sup>3</sup>Department of Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan and <sup>4</sup>Tochigi Prefectural Medical and Social Welfare College, Utsunomiya, Tochigi, Japan

Email: Miki Ojima - [ojima@dent.osaka-u.ac.jp](mailto:ojima@dent.osaka-u.ac.jp); Takashi Hanioka\* - [haniokat@college.fdcnet.ac.jp](mailto:haniokat@college.fdcnet.ac.jp); Keiko Tanaka - [k-tanaka@fukuoka-u.ac.jp](mailto:k-tanaka@fukuoka-u.ac.jp); Hitoshi Aoyama - [aoyamah01@pref.tochigi.jp](mailto:aoyamah01@pref.tochigi.jp)

\* Corresponding author

Published: 2 November 2007

Received: 13 April 2007

BMC Public Health 2007, 7:313 doi:10.1186/1471-2458-7-313

Accepted: 2 November 2007

This article is available from: <http://www.biomedcentral.com/1471-2458/7/313>

© 2007 Ojima et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Various factors affect tooth loss in older age including cigarette smoking; however, evidence regarding the association between smoking and tooth loss during young adulthood is limited. The present study examined the association between cigarette smoking and tooth loss experience among adults aged 20–39 years using linked data from two national databases in Japan.

**Methods:** Two databases of the National Nutrition Survey (NNS) and the Survey of Dental Diseases (SDD), which were conducted in 1999, were obtained from the Ministry of Health, Labor and Welfare with permission for analytical use. In the NNS, participants received physical examinations and were interviewed regarding dietary intake and health practices including cigarette smoking, whereas in the SDD, participants were asked about their frequency of daily brushing, and received oral examinations by certified dentists. Among 6,805 records electronically linked via household identification code, 1314 records of individuals aged 20 to 39 years were analyzed. The prevalence of I+ tooth loss was compared among non-, former, and current smokers. Multiple logistic regression models were constructed including confounders: frequency of tooth brushing, body mass index, alcohol consumption, and intake of vitamins C and E.

**Results:** Smoking rates differed greatly in men (53.3%) and women (15.5%). The overall prevalence of tooth loss was 31.4% (31.8% men and 31.1% women). Tooth loss occurred more frequently among current smokers (40.6%) than former (23.1%) and non-smokers (27.9%). Current smoking showed a significant association with I+ tooth loss in men (adjusted OR = 2.21 [1.40–3.50], P = 0.0007) and women (1.70 [1.13–2.55], P = 0.0111). A significant positive exposure-related relationship between cigarette smoking status and tooth loss was observed (P for trend < 0.0001 and 0.0004 in men and women, respectively). Current smoking was also associated with the prevalence of decayed teeth (1.67 [1.28–2.20], P = 0.0002).

**Conclusion:** An association between cigarette smoking and tooth loss was evident among young adults throughout Japan. Due to limitations of the available variables in the present databases, further studies including caries experience and its confounders should be conducted to examine whether smoking is a true risk of premature tooth loss in young adults.

## Background

The loss of many teeth often reduces the quality of life; embarrassment and self-consciousness limit social interaction and communication [1]. Because of chewing problems and decreased masticatory function, a limitation in food selection may occur, resulting in nutritionally poor diets. Poor nutrition might contribute to an increased risk of several systemic diseases such as cardiovascular diseases [2] and hypertension [3]. Tooth loss may be a significant problem related to general health and the quality of life.

Many studies demonstrated that cigarette smokers have more missing teeth and experience greater rates of tooth loss than non-smokers. Most of these studies regarding the association between smoking and tooth loss concentrated on middle aged and elderly populations and, among them, several studies observed nationally represented population [4-6] and longitudinal cohort [7]. On the other hand, the effect of smoking in young adults would be more apparent than that in older adults because the effects of confounding factors, which may be age-dependent, would be less apparent.

Smoking is a global problem. In several countries, the association between smoking and tooth loss in young adults has been examined [8-10]; however, the sample population was limited to specific age groups, gender and region. Health consequences of smoking at the national level should be evaluated using nationally representative samples of the country.

Data on cigarette smoking and dental status have been accumulated in two independent national surveys in Japan, and record linkage using a common household number in the two surveys is possible. The aim of the present study was to examine the association between cigarette smoking and tooth loss experience among adults aged 20–39 years using linked data from the two national databases.

## Methods

Databases of the National Nutrition Survey (NNS) and the Survey of Dental Diseases (SDD) from 1999 were obtained from the Ministry of Health, Labor and Welfare with permission for analytical use. The SDD, which involved the same sample population as that of the NNS, is conducted every six years. Data were collected from persons aged 1 year or older from approximately 5000 households in 300 districts. The background and general procedure of the surveys were described previously [11].

Smoking status was defined in the questionnaires of the NNS as: "current smoker", an individual who currently smokes cigarettes daily or occasionally; "former smoker", an individual who has smoked cigarettes at some point in

their life, but who currently does not smoke; "non-smoker", an individual who was an experimental smoker or who has never smoked cigarettes. Current smokers were asked about cigarette consumption in terms of the number of cigarettes per day and duration of smoking.

In the SDD, calibrated dentists examined and recorded the status of each tooth which was present in the oral cavity regardless of the degree of eruption. Lost teeth were defined as a permanent tooth (excluding third molars) lost due to extraction or dropout. Though reasons for extraction were not surveyed, teeth lost through extraction due to orthodontic treatment (e.g., bilateral missing of first premolars) were not included. Decayed and filled teeth were diagnosed with a mirror and a Community Periodontal Index probe in the absence of an air jet. Initial and manifest caries lesions for all teeth were recorded. Secondary caries was categorized into decayed teeth. Coronal and root caries were not differentiated.

We selected several variables *a priori* as possible confounders from the two databases in order to control for the effect of behavioral and lifestyle factors on tooth loss. The frequency of daily brushing was divided into two groups: more than twice and less than twice. The body mass index (BMI) was calculated as an indicator of obesity. BMI was divided into two groups: <25.0 and ≥25.0. Alcohol consumption was classified into three groups: current, former, and never. Based on the Recommended Dietary Allowances for Japanese (6<sup>th</sup> Revision), vitamin C and vitamin E intakes were categorized into two groups: deficiency and sufficiency. The reference value of vitamin C was 100 mg/day, and those of vitamin E for males and females were 10 mg/day and 8 mg/day, respectively.

Records in the SDD database were electronically linked to those in the NNS database using the household identification number as the linking variable. All links were checked via comparison of the information on birth month and sex between the two electronic databases. In the event of a mismatch of these variables, such cases were evaluated by a manual search of records among the same household. Among records of 6903 subjects in the SDD and of 12,763 subjects in the NNS, 6805 records were linked successfully. Analyses were limited to 1314 subjects aged 20 to 39 years without missing values.

The prevalence of 1+ tooth loss was compared among non-, former, and current smokers on the basis of gender and study variables. Bivariate analyses were conducted to determine associations between selected variables as well as smoking and the prevalence of tooth loss using the chi-square test. The multiple logistic regression models were constructed, stratified by sex, including frequency of daily brushing, BMI, current smoking, vitamin C and E intakes,

and alcohol consumption as confounding factors. Adjusted odds ratios (ORs) for tooth loss for each variable were calculated. The dose-response relationship was evaluated using life-time exposure as the Brinkman Index, which was calculated by daily consumption times years of smoking. Former smokers were excluded from this analysis because data on the number of years after cessation of smoking were not available. Trend analysis was performed by entering categories of life-time exposure as a continuous variable. SPSS for Windows (SPSS Inc., Chicago) was utilized.

Since caries is the predominant reason for extraction in the study population [12], the association between smoking status and caries prevalence was further examined to investigate the hypothesis that caries experience may modulate subsequent tooth loss possibly due to smoking.

## Results

### Distribution of smokers by study variables

Table 1 shows the prevalence of current, former, and non-smokers by study variables among adults aged 20–39 years. The overall percentage of current smokers was 29.6% of the study population (55.3% males and 15.5% females). Current smokers were relatively predominant in the following groups in comparison with counterpart groups: daily brushing less than twice (142/286; 49.7% vs 247/1028; 24.0%), BMI  $\geq 25.0$  (101/245; 41.2% vs 288/

1069; 26.9%) and current drinking (169/288; 58.7% vs 203/992; 20.5%).

### Bivariate associations between tooth loss and decayed teeth, and study variables

The prevalence of tooth loss and decayed teeth was compared using the study variables in Table 2. Smoking and each variable, with the exception of gender and vitamin intake, were associated with the prevalence of tooth loss ( $P < 0.05$ ). All variables including smoking were associated with the prevalence of decayed teeth ( $P < 0.05$ ). Subjects who had lost teeth were more frequently those with decayed teeth than those without. Also, subjects with decayed teeth were more likely to be those with lost teeth than those without.

### Prevalence of tooth loss by smoking status

Table 3 shows the prevalence of 1+ tooth loss by smoking status among adults aged 20–39 years. Overall, 31.4% of subjects had lost one or more teeth, and the mean tooth loss was  $0.6 \pm 1.4$  teeth. The distribution of subjects with 1–3 lost teeth, 4–6 and  $\geq 7$  was 27.2%, 3.2% and 1.0%, respectively. The prevalence of tooth loss was similar in males (31.8%) and females (31.1%). Tooth loss occurred more frequently among current smokers (40.6%) than former (23.1%) and non-smokers (27.9%) in the overall population. A significant difference was observed among smoking status groups ( $p < 0.0001$ ). In both genders, significant differences were detected among smoking status groups ( $p = 0.0005$  and  $0.0022$ , in males and females, respectively).

**Table 1: Number of subjects according to smoking status by study variables**

Variables	Total	Smoking status		
		Non-	Former	Current
Gender				
males	465	170	38	257
females	849	677	40	132
Frequency of daily brushing				
more than twice	1028	718	63	247
less than twice	286	129	15	142
BMI				
< 25.0	1069	717	64	288
$\geq 25.0$	245	130	14	101
Alcohol consumption				
never	992	744	45	203
former	34	10	7	17
current	288	93	26	169
Vitamin C intake				
$\geq 100$ mg	596	403	33	160
< 100 mg	718	444	45	229
Vitamin E intake*				
$\geq 10$ or 8 mg	671	451	37	183
< 10 or 8 mg	643	396	41	206
All	1314	847	78	389

\*10 mg for males and 8 mg for females

**Table 2: Results from bivariate analyses of tooth loss and decayed teeth, and study variables**

Study variables	tooth loss	decayed teeth
	p-value <sup>†</sup>	p-value <sup>‡</sup>
Smoking status (non vs. former or current)	0.0002	< 0.0001
Gender (females vs males)	0.7843	< 0.0001
Frequency of brushing ( $\geq 2$ times vs. < 2 times)	0.0390	0.0001
Body mass index (< 25.0 vs. $\geq 25.0$ )	0.0012	< 0.0001
Alcohol consumption (never vs. former or current)	0.0084	< 0.0001
Intake of vitamin C ( $\geq 100$ mg vs. < 100 mg)	0.5403	0.0037
Intake of vitamin E* ( $\geq 10/8$ mg vs. < 10/8 mg)	0.4472	0.0273
Tooth loss (1+ vs. 0)	-	< 0.0001
Decayed teeth (1+ vs. 0)	< 0.0001	-

\*10 mg for males and 8 mg for females

<sup>†</sup> differences in prevalence of 1+ tooth loss

<sup>‡</sup> differences in prevalence of 1+ decayed teeth

**Table 3: Comparison of prevalence (%) of 1+ tooth loss by smoking status**

	Total	Smoking status			P value
		Non-	Former	Current	
Males	31.8 (148/465)	21.8 (37/170)	26.3 (10/38)	39.3 (101/257)	0.0005
Females	31.1 (264/849)	29.4 (199/677)	20.0 (8/40)	43.2 (57/132)	0.0022
Overall	31.4 (412/1314)	27.9 (236/847)	23.1 (18/78)	40.6 (158/389)	<0.0001

**Association between cigarette smoking and prevalence of tooth loss**

Adjusted ORs for the prevalence of tooth loss are presented in Table 4. Current smoking was significantly correlated with tooth loss in males (OR = 2.21, 95% CI; 1.40–3.50) and females (OR = 1.70, 95% CI; 1.13–2.55), following consideration of possible confounding factors. However, the association with tooth loss was not significant with respect to former smoking. BMI and the frequency of daily brushing were significantly associated with the prevalence of 1+ tooth loss in females.

**Exposure-related relationship between cigarette smoking and prevalence of tooth loss**

An exposure-related relationship was demonstrated among non- and current smokers, stratified by gender in Table 5. The prevalence of 1+ tooth loss and adjusted OR increased with lifetime exposure in both genders. Current smoking with a Brinkman Index of  $\geq 200$  was significantly associated with an increased prevalence of 1+ tooth loss. A significant positive exposure-related relationship between cigarette smoking status and tooth loss was

observed (P for trend < 0.0001 and 0.0004 in males and females, respectively).

**Association between cigarette smoking and prevalence of decayed teeth**

Based on the finding that the prevalence of 1+ tooth loss between current and non-smokers clearly differed in adults aged 20–39 years, the following analyses were further performed in terms of caries, because caries is a major reason for tooth extraction in the age group. Only decayed teeth were considered as the dependent variable due to the high prevalence of filled teeth (96.7%). Prevalence and adjusted ORs and 95% CIs for decayed teeth among adults aged 20–39 years are presented in Table 6. The association between smoking and decayed teeth was significant in both genders. Adjusted ORs were 1.87(95% CI; 1.24–2.84), 1.56(95% CI; 1.04–2.33), and 1.67(95% CI; 1.28–2.20), in males, females, and overall, respectively.

**Discussion**

In the present study, a significant association between current smoking and tooth loss prevalence was demonstrated among young adults aged 20–39 years in Japan. Follow-

**Table 4: Adjusted odds ratio (OR) and 95% confidence interval (CI) for prevalence of tooth loss**

Study variable	Criteria	Males		Females	
		Adjusted OR (95% CI)	P value	Adjusted OR (95% CI)	P value
Smoking status	Non-	1.00 (Reference)		1.00 (Reference)	
	Former	1.25 <sup>†</sup> (0.55–2.86)	0.5971	0.52 <sup>†</sup> (0.23–1.18)	0.1197
	Current	2.21 <sup>†</sup> (1.40–3.50)	0.0007	1.70 <sup>†</sup> (1.13–2.55)	0.0111
Frequency of brushing	$\geq 2$ times	1.00 (Reference)		1.00 (Reference)	
	< 2 times	0.97 (0.64–1.48)	0.8945	1.59 (1.03–2.44)	0.0350
Body mass index	< 25.0	1.00 (Reference)		1.00 (Reference)	
	$\geq 25.0$	1.28 (0.83–1.96)	0.2640	2.00 (1.31–3.04)	0.0013
Alcohol consumption	Never	1.00 (Reference)		1.00 (Reference)	
	Former	1.05 (0.34–3.30)	0.9287	1.95 (0.75–5.06)	0.1685
	Current	1.17 (0.77–1.78)	0.4565	1.48 (0.90–2.44)	0.1205
Intake of Vitamin C	$\geq 100$ mg	1.00 (Reference)		1.00 (Reference)	
	< 100 mg	1.01 (0.67–1.53)	0.9559	0.81 (0.59–1.11)	0.1899
Intake of Vitamin E*	$\geq 10/8$ mg	1.00 (Reference)		1.00 (Reference)	
	< 10/8 mg	1.16 (0.77–1.76)	0.4756	1.10 (0.81–1.50)	0.5398

\*10 mg for males and 8 mg for females

<sup>†</sup>Adjusting for frequency of brushing, body mass index, alcohol consumption, intake of Vitamin C and E

**Table 5: Adjusted odds ratio (OR) and 95% confidence interval (CI) tooth loss by lifetime exposure to smoking**

	Lifetime exposure*	Prevalence of 1 + tooth loss (%)	Adjusted OR† (95%CI)	P value
Males	0	21.8 (37/170)	1.00 (Reference)	
	1-199	26.9 (29/108)	1.34 (0.76-2.36)	0.3134
	200-399	42.7 (41/96)	2.75 (1.57-4.83)	0.0004
	400+	58.5 (31/53)	5.17 (2.59-10.3)	<0.0001
	P for trend			
Females	0	29.4 (199/677)	1.00 (Reference)	
	1-199	36.2 (38/105)	1.31 (0.84-2.06)	0.2379
	200+	70.4 (19/27)	5.34 (2.24-12.7)	0.0002
	P for trend			

\*Brinkman Index

†Adjusting for frequency of brushing, body mass index, alcohol consumption, intake of Vitamin C and E

ing adjustment for confounding factors, the odds of 1+ tooth loss was approximately 2 times greater in current smokers than non-smokers. Furthermore, a dose-response relationship with lifetime exposure to cigarette smoking was evident. Smokers are likely to show unhealthy behavior and a negative attitude towards general health [13]. In the present study, after adjusting for behavioral and lifestyle factors, the relation of smoking to tooth loss was evident, consistent with the previous study [8]. Vitamin C and E intake was not independently associated with tooth loss. The effect of vitamin C and E intake on tooth loss, if any, may be observed via periodontitis [14,15]. Caries is the predominant reason for extraction in the study population [12]. The design of the present study was cross-sectional. The time sequence between smoking status and tooth loss could not be assessed because the point in time when the teeth were lost was not available; therefore, a causal association between smoking and tooth loss should not be assumed.

Few studies have suggested an association between smoking and tooth loss in young adults [8-10]; however, the population was not a nationally representative sample

and the analysis was based on self-reported tooth loss. In the present study, 1314 samples throughout Japan were analyzed, and the number of teeth was assessed by clinical examination. The 1314 records consisted of 42% of those subjects aged 20-39 years in the NNS. Participants of the dental examination may be limited and biased. However, this sample was similar to the NNS sample with respect to smoking prevalence by sex and age (data not shown). Several confounders were not controlled in the multivariate models. Socio-economic status (SES) and dental service use were not included in the surveys; consequently, this parameter was not considered in the present investigation. In a cohort study of young New Zealanders, SES inequalities in tooth loss appear to begin early in the life course, and are modified by individuals' SES and dental service use [16]. Few studies have examined the relation between socioeconomic factors and smoking in the Japanese population. Association of socioeconomic status with smoking was reported in the recent article [17]; however, the effect of socioeconomic disadvantage on smoking was controversial and much less than in other industrialized counties in the 1990s [18-20]. No significant difference was found in smoking prevalence by SES

**Table 6: Prevalence and adjusted OR of decayed teeth**

	Smoking status	Prevalence (%) of decayed teeth	Adjusted OR† (95%CI)	P value
Males	Non-	37.1 (63/170)	1.00 (Reference)	
	Former	44.7 (17/38)	1.34 (0.64-2.80)	0.4362
	Current	55.6 (143/257)	1.87 (1.24-2.84)	0.0030
Females	Non-	33.1 (224/677)	1.00 (Reference)	
	Former	40.0 (16/40)	1.10 (0.56-2.19)	0.7780
	Current	46.2 (61/132)	1.56 (1.04-2.33)	0.0308
Overall	Non-	33.9 (287/847)	1.00 (Reference)	
	Former	42.3 (33/78)	1.25 (0.77-2.04)	0.3677
	Current	52.4 (204/389)	1.67 (1.28-2.20)	0.0002

†Adjusting for frequency of brushing, body mass index, alcohol consumption, intake of Vitamin C and E

and the frequency of dental check-ups among Japanese in their twenties in 1997 [21]. The effect of these factors may have been weak at the time when the surveys were conducted, and in the study population.

A national survey in Japan reported that about 85% of permanent tooth extractions were due to dental caries and its sequela, and periodontal disease [12]. Caries-associated tooth loss increased in the late twenties and early thirties in a birth cohort study of New Zealanders [22]. In the study population, current smokers may lose teeth via extraction due to dental caries rather than periodontal disease. Actually, no significant difference was observed in the prevalence of periodontitis between current and non-smokers in the same population [11]. Several hypothesized mechanisms underlie the relationship between active smoking and dental caries [23], e.g., impairment of the immune system, alteration of the bacterial profile, and salivary function. These observations may be associated with an increase of endodontic diseases [24], which is also a reason for extraction in young adulthood.

The association between active smoking and dental caries is controversial [23]. Methodological considerations limit interpretation of the findings of the aforementioned investigations. In particular, many studies did not control for potential confounding factors of these associations, such as less effective dental hygiene and plaque removal [25]. We found that the independent association between cigarette smoking and the prevalence of decayed teeth following adjustment for behavioral factors including oral hygiene practices. Biological and prospective epidemiological studies are further required for assessments of the effect of smoking on carious lesion formation.

## Conclusion

The association between cigarette smoking and tooth loss was evident among young adults throughout Japan. Due to limitations of the available variables in the present databases, further studies should be conducted to examine whether smoking is a true risk of premature tooth loss in young adults by employing variables of caries experience and its confounders.

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

MO performed the statistical analysis and drafted the manuscript. TH conceived of the study, and participated in its design and coordination and helped to draft the manuscript. KT helped to draft the manuscript and analyze the data. HA participated in the design and coordina-

tion of the study. All authors read and approved the final manuscript.

## Acknowledgements

This study was supported by the Health and Labour Sciences Research Grants for Research on Cancer Prevention and Health Services and Clinical Cancer Research by the Ministry of Health, Labour and Welfare, Japan.

## References

1. Steele JG, Sanders AE, Slade GD, Allen PF, Lahti S, Nuttall N, Spencer AJ: **How do age and tooth loss affect oral health impacts and quality of life? A study comparing two national samples.** *Community Dent Oral Epidemiol* 2004, **32**:107-114.
2. Hung HC, Colditz G, Joshipura KJ: **The association between tooth loss and the self-reported intake of selected CVD-related nutrients and foods among US women.** *Community Dent Oral Epidemiol* 2005, **33**:167-173.
3. Taguchi A, Sanada M, Sueti Y, Ohtsuka M, Lee K, Tanimoto K, Tsuda M, Ohama K, Yoshizumi M, Higashi Y: **Tooth loss is associated with an increased risk of hypertension in postmenopausal women.** *Hypertension* 2004, **43**:297-300.
4. Ragnarsson E, Eliasson ST, Olafsson SH: **Tobacco smoking, a factor in tooth loss in Reykjavik, Iceland.** *Scand J Dent Res* 1992, **100**:322-326.
5. Axelsson P, Paulander J, Lindhe J: **Relationship between smoking and dental status in 35-, 50-, 65-, and 75-year-old individuals.** *J Clin Periodontol* 1998, **25**:297-305.
6. Klein BE, Klein R, Knudtson MD: **Life-style correlates of tooth loss in an adult Midwestern population.** *J Public Health Dent* 2004, **64**:145-50.
7. Dietrich T, Maserejian NN, Joshipura KJ, Krall EA, Garcia RI: **Tobacco use and incidence of tooth loss among US male health professionals.** *J Dent Res* 2007, **86**:373-7.
8. Ylöstalo PV, Sakki TK, Laitinen J, Järvelin M-R, Knuutila MLE: **The relation of tobacco smoking to tooth loss among young adults.** *Eur J Oral Sci* 2004, **112**:121-126.
9. Tanaka K, Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, Hirota Y, Oda H, Osaka Maternal and Child Health Study Group: **Active and passive smoking and tooth loss in Japanese women: baseline data from the Osaka maternal and child health study.** *Ann Epidemiol* 2005, **15**:358-364.
10. Susin C, Haas AN, Opermann RV, Albandar JM: **Tooth loss in a young population from south Brazil.** *J Public Health Dent* 2006, **66**:110-115.
11. Ojima M, Hanioka T, Tanaka K, Inoshita E, Aoyama H: **Relationship between smoking status and periodontal conditions: findings from national databases in Japan.** *J Periodontol Res* 2006, **41**:573-579.
12. Aida J, Ando Y, Akhter R, Aoyama H, Masui M, Morita M: **Reasons for permanent tooth extractions in Japan.** *J Epidemiol* 2006, **16**:214-219.
13. Revicki D, Sobal J, DeForge B: **Smoking status and the practice of other unhealthy behaviors.** *Fam Med* 1991, **23**:361-364.
14. Nishida M, Grossi SG, Dunford RG: **Dietary vitamin C and the risk for periodontal disease.** *J Periodontol* 2000, **7**:1215-1223.
15. Cohen ME, Meyer DM: **Effect of dietary vitamin E supplementation and rotational stress on alveolar bone loss in rice rats.** *Arch Oral Biol* 1993, **38**:601-606.
16. Thomson WM, Poulton R, Kruger E, Boyd D: **Socio-economic and behavioural risk factors for tooth loss from age 18 to 26 among participants in the Dunedin Multidisciplinary Health and Development Study.** *Caries Res* 2000, **34**:361-366.
17. Fukuda Y, Nakamura K, Takano T: **Socioeconomic pattern of smoking in Japan: income inequality and gender and age differences.** *Ann Epidemiol* 2005, **15**:365-72.
18. Nakamura Y, Sakata K, Kubo N, Akikawa Y, Nagai M, Yanagawa H: **Smoking habits and socioeconomic factors in Japan.** *J Epidemiol* 1994, **4**:157-161.
19. Anzai Y, Ohkubo T, Nishino Y, Tsuji I, Hisamichi S: **Relationship between health practices and education level in the rural Japanese population.** *J Epidemiol* 2000, **10**:149-156.
20. Martikainen P, Ishizaki M, Marmot MG, Nakagawa H, Kagamimori S: **Socioeconomic differences in behavioral and biological risk**

**factors: A comparison of a Japanese and an England cohort of employed men.** *Int J Epidemiol* 2001, **30**:833-38.

21. Tada A, Hanada N: **Sexual differences in smoking behaviour and dental caries experience in young adults.** *Public Health* 2002, **116**:341-346.
22. Broadbent JM, Thomson WM, Poulton R: **Progression of dental caries and tooth loss between the third and fourth decades of life: a birth cohort study.** *Caries Res* 2006, **40**:459-465.
23. US Department of Health and Human Services: **Dental diseases.** In *The Health Consequences of Smoking. A Report of the Surgeon General* Edited by: Samet JM, Norman LA, Wilbanks C. Washington DC: US Department of Health and Human Services; 2004:732-766.
24. Duncan HF, Pitt Ford TR: **The potential association between smoking and endodontic disease.** *Int Endod J* 2006, **39**:843-854.
25. Andrews JA, Severson HH, Lichtenstein E, Gordon JS: **Relationship between tobacco use and self-reported oral hygiene habits.** *J Am Dent Assoc* 1998, **129**:313-320.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/7/313/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

