

## Invited Mini Review

## Roles of mitochondria in neuronal development

Geurim Son<sup>1</sup> & Jinju Han<sup>1,2,\*</sup><sup>1</sup>Biomedical Science and Engineering Interdisciplinary Program, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Korea, <sup>2</sup>Graduate School of Medical Science and Engineering, KAIST, Daejeon 34141, Korea

**Mitochondria are ubiquitous and multi-functional organelles involved in diverse metabolic processes, namely energy production and biomolecule synthesis. The intracellular mitochondrial morphology and distribution change dynamically, which reflect the metabolic state of a given cell type. A dramatic change of the mitochondrial dynamics has been observed in early development that led to further investigations on the relationship between mitochondria and the process of development. A significant developmental process to focus on, in this review, is a differentiation of neural progenitor cells into neurons. Information on how mitochondria-regulated cellular energetics is linked to neuronal development will be discussed, followed by functions of mitochondria and associated diseases in neuronal development. Lastly, the potential use of mitochondrial features in analyzing various neurodevelopmental diseases will be addressed. [BMB Reports 2018; 51(11): 549-556]**

## INTRODUCTION

All cells undergo cellular respiration, whether it uses oxygen or not, to produce energy for survival. The process using oxygen to respire is called aerobic respiration and all aerobically respiring mammalian cells contain and utilize mitochondria for energy production (1). Dependency on mitochondria for energy production varies on different cell types. High energy-demanding cells rely on mitochondria for adenosine triphosphate (ATP) production because mitochondria; a major powerhouse of cells, generate ATP in the most efficient way via oxidative phosphorylation (OXPHOS) (2). OXPHOS creates a proton gradient that induces mitochondrial membrane potential (MMP), then uses oxygen for the synthesis of ATP (3). On the other hand, certain cell types prefer utilizing cytoplasmic metabolic pathways of

glycolysis and pentose phosphate pathway (PPP) for cellular metabolism even though mitochondria are found in these cells (4, 5). Glycolysis and PPP use glucose then generate pyruvate and NADPH in addition to ATP (Fig. 1A). Pyruvate and NADPH are essential molecules for the synthesis of amino acids and nucleotides required for highly proliferative cells to divide.

Cells in the developmental process change metabolic states to support any newly acquired structural and functional properties (6, 7). Although ATP production is more efficient in OXPHOS, glycolysis is enhanced in actively proliferating cells since diverse metabolic substances (not limited to ATP) are required (8, 9). Cells need a lot of energy to sustain homeostasis and support for specialized functions they have acquired from cellular differentiation. Therefore, energy metabolism shifts from glycolysis to OXPHOS and mitochondrial maturation occurs during cellular differentiation (9).

Mutations on genes necessary for mitochondrial maturation are associated with a failure in metabolic transition that can result in developmental defects (10). Since mitochondria are present everywhere, they can have an impact on all types of tissues with no limits. Studies linking mitochondrial dysfunction and developmental diseases are beginning to re-emerge. Knowing how mitochondria behave in a given condition and which genes regulate mitochondrial dynamics will facilitate an understanding of many developmental disease-etologies. This review will focus on distinct features of mitochondria in neuronal development and diseases. We will address the roles of mitochondria along with the process of neurodevelopment.

## Genes regulating mitochondria

Mitochondria are thought to have been engulfed by an ancestral cell during evolution, via a process named endosymbiosis, for more efficient cell survival (11-13). Mitochondria, therefore, are double-membraned: including a permeable outer membrane that is structurally similar to the plasma membrane and an inner membrane forming cristae that divides the mitochondrial matrix and the intermembrane space. Mitochondrion has its own genome and machinery for its gene expression. Mitochondrial genome encodes 37 genes and only makes 13 polypeptides that belong to the OXPHOS complexes (Fig. 1B) (14).


OXPHOS components excluding all 13 polypeptides synthesized in mitochondria are translated in cytoplasm from the nuclear genome transcripts (10, 15-17). Nuclear genome-

\*Corresponding author. Tel: +82-42-350-4862; Fax: +82-42-350-4240; E-mail: jinjuhan@kaist.ac.kr

<https://doi.org/10.5483/BMBRep.2018.51.11.226>

Received 4 September 2018

**Keywords:** Energy metabolism, Mitochondria, Neural stem cell, Neurodevelopmental diseases, Neuronal development



**Fig. 1.** Mitochondrial proteins, functions and dynamics. (A) Major function of mitochondria is energy production through OXPHOS. Glycolysis occurring in the cytosol produce pyruvate, which is necessary to fuel the tricarboxylic acid (TCA) cycle. The pentose phosphate pathway (PPP) is a shunt for glycolysis. Through the PPP, cells acquire required components for other cellular processes including nucleotide synthesis. In mitochondria, beta-oxidation occurs as the other mechanism of converting lipid to generate energy. (B) Most proteins localized in the mitochondria are produced from the nuclear genome (nDNA) and transported into mitochondria. Mitochondria contain its own genome (mitochondrial DNA, mtDNA) and produce 13 proteins comprising oxidative phosphorylation (OXPHOS) complex. (C) Dynamically changing morphology of mitochondria through continuous cycle of fusion and fission.

derived mitochondrial proteins are transported through TOM (translocase of outer mitochondrial membrane), TIM (translocase of inner mitochondrial membrane), OXA (oxidase assembly machinery), etc. (11, 18-22) The five OXPHOS complexes in the inner mitochondrial membrane (IMM) comprise of NADH dehydrogenase (Complex I), succinate dehydrogenase (Complex II), cytochrome c reductase (Complex III), cytochrome c oxidase (Complex IV), and ATP synthase (Complex V).

Mitochondria mainly facilitate energy production through OXPHOS, consisting of an electron transport chain (ETC) and an ATP synthase (17). The ETC carries electrons step-by-step that triggers proton gradient across the IMM. Keeping a constant MMP and cellular respiration cycle are significant in operating the OXPHOS, since the ATP producing complex, ATP synthase, needs a proton gradient to convert adenosine diphosphates (ADPs) to ATPs. Mitochondria participate in other cellular processes like calcium signaling, trafficking and apoptosis by interacting with many other intracellular organelles such as endoplasmic reticulums, lysosomes, peroxisomes, etc. Proteins necessary for additional mitochondrial functions are originated from the nuclear genome and transported into mitochondria. Exploration of mitochondrial protein composition, localization, and topology are in progress to fully investigate the role of mitochondria (23-28).

Mitochondria change their morphology and localization in cells under given conditions to function properly. Mitochondria are regulated dynamically with a balanced and continuous cycle of fusion and fission (Fig. 1C) (29). Fusion

allows mitochondria to exchange membranous materials including various metabolites and rescue damaged mitochondria. On the other hand, fission can segregate and degrade mitochondria with impaired mitochondrial DNA (mtDNA), only leaving healthy mitochondria inside the cell. Proteins helping the fusion: MFN1/2 and OPA1, and the fission: DRP1 and FIS1, have been identified and studies are ongoing to regulate mitochondrial dynamics (30-32). Mitochondria move to a specific region of a cell where high-energy consumption is required (33). Rho GTPases of mitochondria that affect mitochondrial motility, mitochondrial transport, etc. in mammals have been identified: MIRO-1 and MIRO-2 (34). These are proteins of the outer mitochondrial membrane (OMM) interacting with motor proteins such as KINESIN and transporting mitochondria along microtubules.

Databases demonstrating the localization of mitochondrial proteins in representative tissues have been developed. Databases such as MitoCarta, MitoMiner including Integrated Mitochondrial Protein Index (IMPI), MitoP2, MitoProteome, etc. include protein information mostly obtained by an approach to isolate mitochondria from cells (35-38). Biochemical isolation of mitochondria from cells removes the OMM. This approach identifies proteins of mitochondrial matrix and IMM. Recently, proteins localized in OMM facing the cytosol can be analyzed by an enzymatic method using the engineered ascorbate peroxidase (APEX) that labels proximal and interacting proteins (28). The latest version of MitoCarta, named MitoCarta2.0, includes additional proteomic data reported in literatures and discovered via APEX (35). However,

most of above databases are limited to a number of tissues and only focused on certain complexes. An extensive research that uncovers the entire protein composition of mitochondria, linked to each protein's physiological role, is necessary.

### Developmental process of neurons

Mammalian neurogenesis begins at the prenatal stage and continues to the postnatal stage even for adult brains. During neonatal development, neural stem cells (NSCs) appear by the end of gastrulation and majority of the brain structure is formed by the end of embryogenesis (39). In early fetal development, NSCs called radial glial cells (RGCs) reside in the ventricular zone and produce neurons that assemble the neocortex. Newborn neurons migrate inside-out to the neocortex and form the cortical layer; younger neurons are at the outer layer of the cortex. Following neurogenesis, NSCs produce glial cells as well. In the postnatal brain, certain population of the fetal NSCs are retained in two restricted regions of the brain and maintained as adult NSCs (40,41). While adult NSCs are surrounded by glial cells, adult NSCs keep their multi-potency and produce neurons. One of the neurogenic niches in the adult brain is the dentate gyrus (DG) of the hippocampus. Radial glia-like (RGL) NSCs reside in the subgranular zone and add newborn neurons to the granule cell layer (GCL) of DG with an outside-in pattern. Younger neurons are at the inner GCL. The occurrence of adult hippocampal neurogenesis in human brains is like other mammalian brains except for cetaceans such as whales and dolphins. Although debated recently, adult hippocampal neurogenesis in the human brain has been widely accepted for two decades (42-44).

Fetal and adult NSCs share common features in the process of neuronal development even though the environment of neurogenic niche and the layering patterns of newborn neurons are different from each other. RGCs and RGL NSCs displaying bipolar structures produce intermediate neural progenitor cells (NPCs, also referred to as IPCs) with non-polar structures. NPCs continue to retain stem cell markers such as SOX2 and actively proliferate. Neuronal cell fate becomes more apparent at the NPC stage. Neurons differentiated from NPCs undergo morphological changes via axonal and dendritic arborization, resulting in a change in cell polarity. Newborn neurons migrate to their destination and make connections with pre-existing neurons by forming synapses and integrating into an established neuronal circuit.


Neurogenesis is modulated by diverse molecular mechanisms. A representative mechanism that regulates neurogenesis is the transcriptional gene regulation (45, 46). Transcription factors drive a change in the transcriptome profile of cells during neuronal development. Extrinsic factors such as signaling molecules also affect many steps of neurogenesis. For example, the WNT protein family affects proliferation of NPCs, morphogenesis of newborn neurons, migration of newborn neurons, etc. (47, 48). Recently, interests on studying the

mechanisms of neurogenesis have expanded to lipid metabolism (49, 50). In adult hippocampal neurogenesis, fatty acid oxidation is required for maintenance and proliferation of NPCs and lipogenesis is critical for neuronal differentiation (49). These processes aid the metabolic shift during neuronal development. Lipid, in addition to glucose, can be used as an alternative energy source in (an anaerobic) glucose metabolism. In this regard, significance of the mitochondrial role in neuronal development is now recognized and receiving more attention.

### Mitochondrial dynamics during neuronal development and its potential association with developmental brain diseases

The significance of mitochondrial dynamics in neuronal development has been described in the animal brain. Ablating some genes involved in mitochondrial fission and fusion resulted in defects of brain development, although fusion-and-fission dynamics during neuronal development under physiological conditions is unknown (51-53). Recent studies have reported on morphological changes of mitochondria when NSCs are differentiated in the developing brain and the adult brain (54-56). In the developing brain, mitochondria shape themselves with an elongated morphology in NSCs and a fragmented morphology in NPCs (54). On the other hand, in the adult hippocampus, mitochondria form a mixture of globular and tubular structures in NSCs and a thin and more elongated morphology in NPCs (56). However, divergent observations in fetal and adult brains come to an agreement for neurons. Mitochondria of the differentiated neurons reveal an elongated morphology in the developing brain and a wider and highly elongated morphology in the adult hippocampus (54, 56). Morphological changes of mitochondria during neurogenesis illustrate maturation of mitochondria and reflect metabolic shift of cells from glycolysis to OXPHOS for an increase in bioenergetics (Fig. 2) (57).

Adding on to morphological changes of mitochondria during neurogenesis, RNA expression profiles of single cells during neuronal differentiation demonstrate a metabolic shift from glycolysis to OXPHOS (58, 59). In adult hippocampus, it clearly demonstrates that RGL NSCs highly express glycolytic genes and lose expression of those genes with differentiation (58). Corresponding to this, increased expression of OXPHOS genes are characterized in post-mitotic neurons. Particularly, genes of Complex V in OXPHOS are dramatically increased in their expression level upon neuronal differentiation. Expression levels of ETC genes, referring to other OXPHOS complexes: Complex I-IV, are quite consistent during neuronal development. When cells enter the post-mitotic stage in the developing brain, expression patterns of the metabolic genes also change dramatically (59). For example, expression levels of glycolytic genes such as *ALDOC* (Aldolase C) and *HK2* (Hexokinase 2) decreased once NSCs started to differentiate into neurons. Although changes in the level of some metabolism-related transcripts have been reported, a deeper



**Fig. 2.** Mitochondrial features and bioenergetics during neuronal development. Neural stem cells (NSCs) and intermediate neural progenitor cells (NPCs, also referred to as IPCs) have self-renewing capacities. NSCs are differentiated into NPCs, which are then differentiated into neurons. The changes in mitochondrial morphology during neuronal development should be noted. In corticogenesis in developing brains (A), the mitochondrial morphology change from elongated structure to fragmented, then elongated again, followed by more complex structure due to further elongation. In adult hippocampal neurogenesis (B), the mitochondrial morphology changes from mixed globular and tubular structures to thin and elongated, then elongated more, followed by a wider and more complex structure due to further elongation. Level of glycolysis is decreased in both (A) and (B) when level of OXPHOS is increased along with neuronal differentiation.

analysis on the expression level changes of mitochondria- and metabolism-related genes in neurogenesis will augment mechanism studies of neuronal development. Further investigations to elucidate which mitochondrial genes and proteins contribute to mitochondrial maturation and functions at each stage of neurogenesis will be necessary.

Mitochondrial dynamics and a metabolic shift have also been investigated in human neurogenesis by utilizing NPCs, derived from human pluripotent stem cells (60, 61). Expression patterns of metabolic genes were analyzed at different stages of neuronal differentiation. The expression level of MFN2, a key player of mitochondrial fusion, increase along with the differentiation of NPCs (60). Depletion of MFN2 in NPCs delays neuronal development when the overexpression promotes neuronal development, indicating significance of mitochondrial dynamics in human neurogenesis. Expression of key glycolytic genes, *HK2* and *LDHA*, are decreased while transcript levels of most OXPHOS genes do not change when NPCs are differentiated into neurons that correspond to the results from the study of adult hippocampal neurogenesis (61).

The gene expression pattern indicates that human NPCs undergo a bioenergetic shift from glycolysis to OXPHOS. Increased mass of mitochondria in the process of NPC differentiation also support that NPCs rely on the mitochondrial function with differentiation (62, 63). However, there is a caveat to understanding mitochondrial function based solely on the RNA transcript level. The level of transcripts does not always correlate to the level of proteins

(64, 65). Thus, additional layers of analysis on the translational and post-translational gene regulation should also be considered in order to interpret the functions of mitochondria.

Developing neurons extend neurites and generate axons required for migration of mitochondria from the soma toward axon terminals. Mitochondria supply ATPs to modulate actin filaments at the axon terminal. Regarding the transport system in mammals, two Rho GTPases of mitochondria: MIRO-1 and MIRO-2, have effects on the mitochondrial motility and transport (66). MIROs have calcium binding domains and are proteins of the OMM (67). They interact with calcium and regulate cellular motor proteins, mainly KINESIN-1. MIRO-1 is also known to mediate mitochondrial fusion and fission depending on the level of calcium in human cell lines (68). In fully differentiated neurons, mitochondria are concentrated at the pre-synapse and the post-synapse: location where a lot of energy is necessary (33, 69, 70). Mitochondria at synaptic terminals provide regional ATPs to neurons and modulate cytosolic calcium levels. Thus, if mitochondria are unable to reach the signal exchanging center during development and even after development, neuronal function will be impaired (71).

Mitochondrial dysfunctions due to mutations may affect the proliferation rate of NPCs and also change the efficiency of differentiation into neurons, resulting in delayed or paused neurogenesis (72). Dysfunctional mitochondria are associated with neurological diseases such as Leigh syndrome, Rett syndrome, Angelman syndrome, Autism Spectrum Disorder

(ASD), Schizophrenia, and Bipolar disorder. Leigh syndrome is a representative mitochondrial disease of dysfunctional Complex I or IV and also occurring due to a mutation in *MT-ATP6* gene belonging to Complex V (73-78). Defective Complex I and IV prevents the proton gradient from being maintained, but dysfunctional Complex V will not produce ATP even when sufficient proton gradient is generated. Rett syndrome caused by a mutation in *MECP2* gene on X-chromosome is a neurodevelopmental disease (79-81). A mutation in *MECP2* can alter the epigenetic status of the nuclear genome (82, 83). As most OXPHOS subunit proteins are encoded in the nuclear genome, transcription of these genes can be affected by an *MECP2* mutation. Deficiency in Complex IV activity is observed in animal models of Rett syndrome (84). Angelman syndrome is caused by *UBE3A* loss-of-function (85). Reduced activity of Complex III and change in mitochondrial morphology are observed in the brain of animal models for the Angelman syndrome. ASD is not always associated with mitochondrial dysfunctions (86). However, it is reported to be more severe with mitochondrial dysfunctions and correlated with decreased level of an antioxidant defense mechanism and an elevated level of ROS and lactate (87-91). Schizophrenia can be caused due to defective Complex I, III, and/or IV that result in decreased ATP production, higher anaerobic metabolism of glucose, and increased lactate level (91-93). Bipolar disorder is also affected by oxidative stresses similar to that of schizophrenia: higher lactate level and decreased number of protons in the mitochondrial matrix (89, 94, 95). Hence, studies to reveal the hidden molecular mechanisms of the neurological disease-causing mitochondrial dysfunctions will be necessary.

### Perspectives

Several studies have linked mitochondria to neurological diseases by observing stage-dependent and metabolism-related changes of neurogenesis. This has opened an era of more in-depth investigations on neurometabolic diseases. Here, diverse aspects of metabolism as main factors associated with neurodevelopmental diseases examined by many research groups have been introduced. Although a correlation between mitochondria and neuronal differentiation has been demonstrated the underlying mechanisms in connecting mitochondria and various neurological diseases still remain to be elucidated. Especially, functional implications of mitochondria on neurological diseases are lacking scientific findings that may be applied to clinical settings. However, specific features of mitochondria have been identified and are used as biomarkers or in treatments for some diseases, leaving hope for such application in neurodevelopmental diseases (96-98). Studies using human NSCs and unbiased identifications of functional proteins in mitochondria will bring in novel insights and thought-provoking discoveries to the field. The significance and function of mitochondria in neurodevelopmental diseases should not be underestimated.

### ACKNOWLEDGEMENTS

We sincerely apologize to numerous research groups whose work could not be cited due to space limitations. This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2018R1D1A1B07043238) and Short Term Innovative Research by KAIST (N11180123).

### CONFLICTS OF INTEREST

The authors have no conflicting interests.

### REFERENCES

1. Lunt SY and Vander Heiden MG (2011) Aerobic Glycolysis: Meeting the Metabolic Requirements of Cell Proliferation. *Annu Rev Cell Dev Biol* 27, 441-464
2. Kuznetsov AV, Hermann M, Saks V, Hengster P and Margreiter R (2009) The cell-type specificity of mitochondrial dynamics. *Int J Biochem Cell Biol* 41, 1928-1939
3. Yellen G (2018) Fueling thought: Management of glycolysis and oxidative phosphorylation in neuronal metabolism. *J Cell Biol* 217, 2235-2246
4. Novello F and McLean P (1968) The pentose phosphate pathway of glucose metabolism. Measurement of the non-oxidative reactions of the cycle. *Biochem J* 107, 775-791
5. Marin-Valencia I, Cho SK, Rakheja D et al (2012) Glucose metabolism via the pentose phosphate pathway, glycolysis and Krebs cycle in an orthotopic mouse model of human brain tumors. *NMR Biomed* 25, 1177-1186
6. Zhang H, Menzies KJ and Auwerx J (2018) The role of mitochondria in stem cell fate and aging. *Development* 145, dev143420
7. Teslaa T and Teitell MA (2015) Pluripotent stem cell energy metabolism: an update. *EMBO J* 34, 138-153
8. Lees JG, Gardner DK and Harvey AJ (2017) Pluripotent stem cell metabolism and mitochondria: beyond ATP. *Stem Cells Int* 2017, 2874283
9. Xu X, Duan S, Yi F, Ocampo A, Liu GH and Izpisua Belmonte JC (2013) Mitochondrial regulation in pluripotent stem cells. *Cell Metab* 18, 325-332
10. Suomalainen A and Battersby BJ (2018) Mitochondrial diseases: The contribution of organelle stress responses to pathology. *Nat Rev Mol Cell Biol* 19, 77-92
11. Ott M, Amunts A and Brown A (2016) Organization and regulation of mitochondrial protein synthesis. *Annu Rev Biochem* 85, 77-101
12. Roger AJ, Muñoz-Gómez SA and Kamikawa R (2017) The origin and diversification of mitochondria. *Curr Biol* 27, R1177-R1192
13. Dyall SD, Brown MT and Johnson PJ (2004) Ancient Invasions: From endosymbionts to organelles. *Science* 304, 253-257
14. Anderson S, Bankier AT, Barrell BG et al (1981) Sequence and organization of the human mitochondrial genome.

- Nature 290, 457-465
15. Lightowlers RN, Rozanska A and Chrzanowska-Lightowlers ZM (2014) Mitochondrial protein synthesis: Figuring the fundamentals, complexities and complications, of mammalian mitochondrial translation. *FEBS Lett* 588, 2496-2503
  16. Richter-Dennerlein R, Oeljeklaus S, Lorenzi I et al (2016) Mitochondrial protein synthesis adapts to influx of nuclear-encoded protein. *Cell* 167, 471-483 e10
  17. Rampelt H and Pfanner N (2016) Coordination of two genomes by mitochondrial translational plasticity. *Cell* 167, 308-310
  18. Haynes CM, Yang Y, Blais SP, Neubert TA and Ron D (2010) The Matrix peptide exporter HAF-1 signals a mitochondrial UPR by activating the transcription factor ZC376.7 in *C. elegans*. *Mol Cell* 37, 529-540
  19. Larsson NG (2010) Somatic Mitochondrial DNA mutations in mammalian aging. *Annu Rev Biochem* 79, 683-706
  20. Fox TD (2012) Mitochondrial protein synthesis, import, and assembly. *Genetics* 192, 1203-1234
  21. Dolezal P, Likic V, Tachezy J and Lithgow T (2006) Evolution of the molecular machines for protein import into mitochondria. *Science* 313, 314-318
  22. Wiedemann N and Pfanner N (2017) mitochondrial machineries for protein import and assembly. *Annu Rev Biochem* 86, 685-714
  23. Rhee HW, Zou P, Udeshi ND et al (2013) Proteomic mapping of mitochondria in living cells via spatially restricted enzymatic tagging. *Science* 339, 1328-1331
  24. Lee SY, Kang MG, Park JS, Lee G, Ting AY and Rhee HW (2016) APEX fingerprinting reveals the subcellular localization of proteins of interest. *Cell Rep* 15, 1837-1847
  25. Lee SY, Kang MG, Shin S et al (2017) Architecture mapping of the inner mitochondrial membrane proteome by chemical tools in live cells. *J Am Chem Soc* 139, 3651-3662
  26. Han S, Udeshi ND, Deerinck TJ et al (2017) Proximity biotinylation as a method for mapping proteins associated with mtDNA in living cells. *Cell Chem Biol* 24, 404-414
  27. Hung V, Zou P, Rhee HW et al (2014) Proteomic mapping of the human mitochondrial intermembrane space in live cells via ratiometric APEX tagging. *Mol Cell* 55, 332-341
  28. Hung V, Lam SS, Udeshi ND et al (2017) Proteomic mapping of cytosol-facing outer mitochondrial and ER membranes in living human cells by proximity biotinylation. *Elife* 6, 1-38
  29. Westermann B (2010) Mitochondrial fusion and fission in cell life and death. *Nat Rev Mol Cell Biol* 11, 872-884
  30. Chan DC (2006) Mitochondrial fusion and fission in mammals. *Annu Rev Cell Dev Biol* 22, 79-99
  31. Chen H and Chan DC (2005) Emerging functions of mammalian mitochondrial fusion and fission. *Hum Mol Genet* 14 Spec No. 2, R283-289
  32. Ishihara N, Nomura M, Jofuku A et al (2009) Mitochondrial fission factor Drp1 is essential for embryonic development and synapse formation in mice. *Nat Cell Biol* 11, 958-966
  33. Chang DTW, Honick AS and Reynolds IJ (2006) Mitochondrial trafficking to synapses in cultured primary cortical neurons. *J Neurosci* 26, 7035-7045
  34. Fransson Å, Ruusala A and Aspenström P (2006) The atypical Rho GTPases Miro-1 and Miro-2 have essential roles in mitochondrial trafficking. *Biochem Biophys Res Commun* 344, 500-510
  35. Calvo SE, Clauser KR and Mootha VK (2016) MitoCarta2.0: An updated inventory of mammalian mitochondrial proteins. *Nucleic Acids Res* 44, D1251-1257
  36. Smith AC and Robinson AJ (2016) MitoMiner v3.1, an update on the mitochondrial proteomics database. *Nucleic Acids Res* 44, D1258-1261
  37. Prokisch H and Ahting U (2007) MitoP2, an integrated database for mitochondrial proteins. *Methods Mol Biol* 372, 573-586
  38. Cotter D, Guda P, Fahy E and Subramaniam S (2004) MitoProteome: mitochondrial protein sequence database and annotation system. *Nucleic Acids Res* 32, D463-467
  39. Stiles J and Jernigan TL (2010) The basics of brain development. *Neuropsychol Rev* 20, 327-348
  40. Fuentealba LC, Rompani SB, Parraguez JI et al (2015) Embryonic origin of postnatal neural stem cells. *Cell* 161, 1644-1655
  41. Li G, Fang L, Fernández G and Pleasure SJ (2013) The ventral hippocampus is the embryonic origin for adult neural stem cells in the dentate gyrus. *Neuron* 78, 658-672
  42. Sorrells SF, Paredes MF, Cebrian-Silla A et al (2018) Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature* 555, 377-381
  43. Boldrini M, Fulmore CA, Tartt AN et al (2018) Human hippocampal neurogenesis persists throughout aging. *Cell Stem Cell* 22, 589-599
  44. Kempermann G, Gage FH, Aigner L et al (2018) Human adult neurogenesis: Evidence and remaining questions. *Cell Stem Cell* 23, 25-30
  45. Beckervordersandforth R, Zhang C and Lie DC (2015) Transcription-factor-dependent control of adult hippocampal neurogenesis. *Cold Spring Harb Perspect Biol* 7, a018879
  46. Martynoga B, Drechsel D, Guillemot F et al (2012) Molecular control of neurogenesis : A view from the mammalian cerebral cortex molecular control of neurogenesis : A view from the mammalian cerebral cortex. *Cold Spring Harb Perspect Biol* 4, a008359
  47. Hirabayashi Y (2004) The Wnt/beta-catenin pathway directs neuronal differentiation of cortical neural precursor cells. *Development* 131, 2791-2801
  48. Lie DC, Colamarino SA, Song HJ et al (2005) Wnt signalling regulates adult hippocampal neurogenesis. *Nature* 437, 1370-1375
  49. Knobloch M, Pilz GA, Ghesquière B et al (2017) A fatty acid oxidation-dependent metabolic shift regulates adult neural stem cell activity. *Cell Rep* 20, 2144-2155
  50. Knobloch M, Braun SMG, Zurkirchen L et al (2014) Metabolic control of adult neural stem cell activity by Fasn-dependent lipogenesis. *Nature* 493, 226-230
  51. Smirnova E, Griparic L, Shurland DL and van der Bliek

- AM (2001) Dynamin-related protein Drp1 is required for mitochondrial division in mammalian cells. *Mol Biol Cell* 12, 2245-2256
52. Chen H, Detmer SA, Ewald AJ, Griffin EE, Fraser SE and Chan DC (2003) Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development. *J Cell Biol* 160, 189-200
53. Chen H and Chan DC (2004) Mitochondrial dynamics in mammals. In: Schatten GP, ed. Vol 59. Current topics in developmental biology. Academic Press 59, 119-144
54. Khacho M, Clark A, Svoboda DS et al (2016) Mitochondrial dynamics impacts stem cell identity and fate decisions by regulating a nuclear transcriptional program. *Cell Stem Cell* 19, 232-247
55. Khacho M and Slack RS (2018) Mitochondrial dynamics in the regulation of neurogenesis: From development to the adult brain. *Dev Dyn* 247, 47-53
56. Beckervordersandforth R, Ebert B, Schäffner I et al (2017) Role of mitochondrial metabolism in the control of early lineage progression and aging phenotypes in adult hippocampal neurogenesis. *Neuron* 93, 560-573 e6
57. Alirol E and Martinou JC (2006) Mitochondria and cancer: Is there a morphological connection? *Oncogene* 25, 4706-4716
58. Shin J, Berg DA, Zhu Y et al (2015) Single-cell RNA-seq with waterfall reveals molecular cascades underlying adult neurogenesis. *Cell Stem Cell* 17, 360-372
59. Telley L, Govindan S, Prados J et al (2016) Sequential transcriptional waves direct the differentiation of newborn neurons in the mouse neocortex. *Science* 351, 1443-1446
60. Fang D, Yan S, Yu Q, Chen D and Yan SS (2016) Mfn2 is required for mitochondrial development and synapse formation in human induced pluripotent stem cells/hiPSC derived cortical neurons. *Sci Rep* 6, 1-13
61. Zheng X, Boyer L, Jin M et al (2016) Metabolic reprogramming during neuronal differentiation from aerobic glycolysis to neuronal oxidative phosphorylation. *Elife* 5, 1-25
62. Ebrahimi-Fakhari D, Saffari A, Wahlster L et al (2016) Impaired mitochondrial dynamics and mitophagy in neuronal models of tuberous sclerosis complex. *Cell Rep* 17, 1053-1070
63. Agostini M, Romeo F, Inoue S et al (2016) Metabolic reprogramming during neuronal differentiation. *Cell Death Differ* 23, 1502-1514
64. Fortelny N, Overall CM, Pavlidis P and Freue GVC (2017) Can we predict protein from mRNA levels? *Nature* 547, E19-E20
65. Liu Y, Beyer A and Aebersold R (2016) On the dependency of cellular protein levels on mRNA Abundance. *Cell* 165, 535-550
66. Chen Y and Sheng ZH (2013) Kinesin-1-syntaphilin coupling mediates activity-dependent regulation of axonal mitochondrial transport. *J Cell Biol* 202, 351-364
67. MacAskill AF, Rinholm JE, Twelvetrees AE et al (2009) Miro1 is a calcium sensor for glutamate receptor-dependent localization of mitochondria at synapses. *Neuron* 61, 541-555
68. Nemani N, Carvalho E, Tomar D et al (2018) MIRO-1 determines mitochondrial shape transition upon GPCR activation and Ca<sup>2+</sup> stress. *Cell Rep* 23, 1005-1019
69. Kang JS, Tian JH, Pan PY et al (2008) Docking of axonal Mitochondria by syntaphilin controls their mobility and affects short-term facilitation. *Cell* 132, 137-148
70. Li Z, Okamoto KI, Hayashi Y and Sheng M (2004) The importance of dendritic mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell* 119, 873-887
71. Calkins MJ, Manczak M, Mao P, Shirendeb U and Reddy PH (2011) Impaired mitochondrial biogenesis, defective axonal transport of mitochondria, abnormal mitochondrial dynamics and synaptic degeneration in a mouse model of Alzheimer's disease. *Hum Mol Genet* 20, 4515-4529
72. Lorenz C, Lesimple P, Bukowiecki R et al (2017) Human iPSC-derived neural progenitors are an effective drug discovery model for neurological mtDNA disorders. *Cell Stem Cell* 20, 659-674 e9
73. Zsurka G and Kunz WS (2015) Mitochondrial dysfunction and seizures: The neuronal energy crisis. *Lancet Neurol* 14, 956-966
74. Martin MA, Blazquez A, Gutierrez-Solana LG et al (2005) Leigh syndrome associated with mitochondrial complex I deficiency due to a novel mutation in the NDUFS1 gene. *Arch Neurol* 62, 659-661
75. Marin SE, Mesterman R, Robinson B, Rodenburg RJ, Smeitink J and Tarnopolsky MA (2013) Leigh syndrome associated with mitochondrial complex I deficiency due to novel mutations in NDUFV1 and NDUFS2. *Gene* 516, 162-167
76. Distelmaier F and Koopman WJH, van den Heuvel LP et al (2009) Mitochondrial complex I deficiency: from organelle dysfunction to clinical disease. *Brain* 132, 833-842
77. Saneto R and Ruhoy I (2014) The genetics of Leigh syndrome and its implications for clinical practice and risk management. *Appl Clin Genet* 7, 221-234
78. DiMauro S, Tanji K and Schon EA (2012) The many clinical faces of cytochrome c oxidase deficiency. In: Kadenbach B, ed. *Mitochondrial oxidative phosphorylation: nuclear-encoded genes, enzyme regulation, and pathophysiology*. New York, NY: Springer New York Chapter 14, 341-357
79. Kriaucionis S, Paterson A, Curtis J, Guy J, MacLeod N and Bird A (2006) Gene Expression Analysis Exposes mitochondrial abnormalities in a mouse model of rett syndrome. *Mol Cell Biol* 26, 5033-5042
80. Gibson JH, Slobedman B, KN H et al (2010) Downstream targets of methyl CpG binding protein 2 and their abnormal expression in the frontal cortex of the human Rett syndrome brain. *BMC Neurosci* 11, 53
81. Pecorelli A, Leoni G, Cervellati F et al (2013) Genes related to mitochondrial functions, protein degradation, and chromatin folding are differentially expressed in lymphomonocytes of rett syndrome patients. *Mediators Inflamm* 2013, 137629
82. Su D, Cha YM and West AE (2012) Mutation of Mecp2 alters transcriptional regulation of select immediate-early genes. *Epigenetics* 7, 146-154
83. Chahrouh M, Jung SY, Shaw C et al (2008) MeCP2, a key contributor to neurological disease, activates and

- represses transcription. *Science* 320, 1224-1229
84. Shulyakova N, Andreatza AC, Mills LR and Eubanks JH (2017) Mitochondrial dysfunction in the pathogenesis of rett syndrome: Implications for mitochondria-targeted therapies. *Front Cell Neurosci* 11, 58
  85. Su H, Fan W, Coskun PE et al (2011) Mitochondrial dysfunction in CA1 hippocampal neurons of the UBE3A deficient mouse model for Angelman syndrome. *Neurosci Lett* 487, 129-133
  86. Khemakhem AM, Frye RE, El-Ansary A, Al-Ayadhi L and Bacha A Ben (2017) Novel biomarkers of metabolic dysfunction in autism spectrum disorder: potential for biological diagnostic markers. *Metab Brain Dis* 32, 1983-1997
  87. James SJ, Melnyk S, Fuchs G et al (2009) Efficacy of methylcobalamin and folinic acid treatment on glutathione redox status in children with autism. *Am J Clin Nutr* 89, 425-430
  88. James SJ, Rose S, Melnyk S et al (2009) Cellular and mitochondrial glutathione redox imbalance in lymphoblastoid cells derived from children with autism. *FASEB J* 23, 2374-2383
  89. Perry SW, Norman JP, Litzburg A and Gelbard HA (2004) Antioxidants are required during the early critical period, but not later, for neuronal survival. *J Neurosci Res* 78, 485-492
  90. Chugani DC, Sundram BS, Behen M, Lee ML and Moore GJ (1999) Evidence of altered energy metabolism in autistic children. *Prog Neuro-Psychopharmacology Biol Psychiatry* 23, 635-641
  91. Brennand K, Savas JN, Kim Y et al (2015) Phenotypic differences in hiPSC NPCs derived from patients with schizophrenia. *Mol Psychiatry* 20, 361-368
  92. Maurer I, Zierz S and Möller H (2001) Evidence for a mitochondrial oxidative phosphorylation defect in brains from patients with schizophrenia. *Schizophr Res* 48, 125-136
  93. Prabakaran S, Swatton JE, Ryan MM et al (2004) Mitochondrial dysfunction in schizophrenia: Evidence for compromised brain metabolism and oxidative stress. *Mol Psychiatry* 9, 684-697
  94. Scaini G, Rezin GT, Carvalho AF, Streck EL, Berk M and Quevedo J (2016) Mitochondrial dysfunction in bipolar disorder: Evidence, pathophysiology and translational implications. *Neurosci Biobehav Rev* 68, 694-713
  95. Mertens J, Wang QW, Kim Y et al (2015) Differential responses to lithium in hyperexcitable neurons from patients with bipolar disorder. *Nature* 527, 95-99
  96. Malik AN and Czajka A (2013) Is mitochondrial DNA content a potential biomarker of mitochondrial dysfunction? *Mitochondrion* 13, 481-492
  97. Budnik LT, Kloth S, Baur X, Preisser AM and Schwarzenbach H (2013) Circulating mitochondrial DNA as biomarker linking environmental chemical exposure to early preclinical lesions elevation of mtDNA in human serum after exposure to carcinogenic halo-alkane-based pesticides. *Hoque MO, ed. PLoS One* 8, e64413
  98. Kilbaugh TJ, Lvova M, Karlsson M et al (2015) Peripheral blood mitochondrial DNA as a biomarker of cerebral mitochondrial dysfunction following traumatic brain injury in a porcine model. *Ai J, ed. PLoS One* 10, e0130927