

Data and text mining

# rPanglaoDB: an R package to download and merge labeled single-cell RNA-seq data from the PanglaoDB database

Daniel Osorio<sup>1,\*</sup>, Marieke L. Kuijjer<sup>1,2</sup> and James J. Cai<sup>3,4,5,6</sup>

<sup>1</sup>Centre for Molecular Medicine Norway (NCMM), University of Oslo, 0349 Oslo, Norway, <sup>2</sup>Department of Pathology, Leiden University Medical Center, Leiden, The Netherlands, <sup>3</sup>Department of Veterinary Integrative Biosciences, Texas A&M University, College Station, TX 77843, USA, <sup>4</sup>Department of Electrical & Computer Engineering, Texas A&M University, College Station, TX 77843, USA, <sup>5</sup>Interdisciplinary Program of Genetics, Texas A&M University, College Station, TX 77843, USA and <sup>6</sup>Center for Statistical Bioinformatics, Texas A&M University, College Station, TX 77843, USA

\*To whom correspondence should be addressed.

Associate Editor: Christina Kendzioriski

Received on June 1, 2021; revised on July 14, 2021; editorial decision on July 25, 2021; accepted on July 26, 2021

## Abstract

**Motivation:** Characterizing cells with rare molecular phenotypes is one of the promises of high throughput single-cell RNA sequencing (scRNA-seq) techniques. However, collecting enough cells with the desired molecular phenotype in a single experiment is challenging, requiring several samples preprocessing steps to filter and collect the desired cells experimentally before sequencing. Data integration of multiple public single-cell experiments stands as a solution for this problem, allowing the collection of enough cells exhibiting the desired molecular signatures. By increasing the sample size of the desired cell type, this approach enables a robust cell type transcriptome characterization.

**Results:** Here, we introduce rPanglaoDB, an R package to download and merge the uniformly processed and annotated scRNA-seq data provided by the PanglaoDB database. To show the potential of rPanglaoDB for collecting rare cell types by integrating multiple public datasets, we present a biological application collecting and characterizing a set of 157 fibrocytes. Fibrocytes are a rare monocyte-derived cell type, that exhibits both the inflammatory features of macrophages and the tissue remodeling properties of fibroblasts. This constitutes the first fibrocytes' unbiased transcriptome profile report. We compared the transcriptomic profile of the fibrocytes against the fibroblasts collected from the same tissue samples and confirm their associated relationship with healing processes in tissue damage and infection through the activation of the prostaglandin biosynthesis and regulation pathway.

**Availability and implementation:** rPanglaoDB is implemented as an R package available through the CRAN repository <https://CRAN.R-project.org/package=rPanglaoDB>.

**Contact:** daniecos@uio.no

## 1 Introduction

Merging and integrating the count matrices derived from multiple independent public single-cell RNA sequencing (scRNA-seq) experiments allows for better evaluation of the biological patterns in cell composition of tissues, as well as the identification of patterns of gene expression and gene regulation that are consistent across cells of the same cell type obtained from independent samples (Swamy *et al.*, 2021). A good quality integration of multiple datasets allowing comparison and contrasting of the data across different projects begins with a consistent preprocessing of the samples, using the same reference genome and the same quantification method to generate the count matrices (Lachmann *et al.*, 2018). These steps are

then followed by the batch effect removal, normalization, cell type annotation and characterization of samples and cell types (Butler *et al.*, 2018; Hie *et al.*, 2019; Korsunsky *et al.*, 2019; Luecken *et al.*, 2020; Stuart *et al.*, 2019).

PanglaoDB is a secondary scRNA-seq database that reports the annotated count matrices for thousands of human and mice scRNA-seq experiments deposited in the sequence read archive (SRA) database of the National Center for Biotechnology Information. Samples available in the PanglaoDB database are uniformly processed with the 'alona' package and made available in a web-based unified framework at <https://panglaoDB.se/> (Franzen *et al.*, 2019; Franzen and Bjorkegren, 2020). However, the PanglaoDB database reports each sample on a single web page and does not offer options to



Fibrocytes are associated with fibrosis, autoimmunity, cardiovascular disease and asthma, among other pathologies (Reilkoff et al., 2011). Since, fibrocytes are marrow-derived cells that differentiate into fibroblasts-like phenotypes, they are usually wrongly labeled as fibroblasts. Thus, using the 'getSamples' function in rPanglaoDB, we downloaded all fibroblasts available from dermis samples in the database (SRA accessions: SRS3121028 and SRS3121030) (Lim et al., 2018). We merged a total of 2172 cells and processed the associated scRNA-seq data using the Seurat package recommended pipeline (Stuart et al., 2019). Datasets were integrated using Harmony and further corroboration of the marker genes defining their identity as fibrocytes (*CD34*, *ACTA2*, *COL5A1*, *COL5A2*, *COL5A3*, *FN1*, *FAP*, *SIRPA*, *PTPRC*, *MME* and *SEMA7A*) was assessed using the Nebulosa package (Fig. 1B) (Alquicira-Hernandez and Powell, 2021; Korsunsky et al., 2019; Reilkoff et al., 2011).

Differential expression analysis of the cluster 8, which contained 157 fibrocytes (Fig. 1A), against all the fibroblasts in the samples was performed using the MAST package (Fig. 1C); returning 50 upregulated genes in fibrocytes associated with the *TGF-beta* regulation of extracellular matrix, ECM-receptor interaction, Prostaglandin biosynthesis and regulation, Notch signaling pathway, Interleukin-5 regulation of apoptosis, Integrin beta-5 pathway, Oncostatin M, Hematopoietic cell lineage and Inflammatory response pathway (FDR < 0.05 using the hypergeometric-test through the enrichR package) (Finak et al., 2015; Xie et al., 2021). We cross-validated the enrichment of the genes associated with the Prostaglandin biosynthesis and regulation pathway (*ANXA3*, *S100A10*, *ANXA5*, *ANXA2*, *PTGIS*, *ANXA1*, *S100A6*, *PTGS1* and *HPGD*) using the Gene Set Enrichment Analysis (GSEA) approach included in the fgsea package (FDR = 0.01, Fig. 1D) and the single-sample Gene Set Enrichment Analysis (ssGSEA) included in the GVSAs package (FDR =  $2.18 \times 10^{-18}$ , Fig. 1E) (Hanzelmann et al., 2013; Korotkevich et al., 2021). All other associations did not pass the FDR threshold in all the other approaches applied (GSEA and ssGSEA). Our results show the potential of rPanglaoDB as a tool to collect cells with rare molecular phenotypes from the PanglaoDB database. We anticipate its use in the construction of atlases to characterize the different molecular phenotypes exhibited by different cell types in different tissues and organisms. We also provide the first unbiased, highly specific characterization of the fibrocyte transcriptome in skin wound tissues and confirm their association with healing processes in tissue damage and infection through the activation of the Prostaglandin Biosynthesis and Regulation Pathway (Grieb et al., 2011; Zhang et al., 2018).

## Funding

D.O. was supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement

[801133]. M.L.K. was supported by the Norwegian Research Council, Helse Sør-Øst, and University of Oslo through the Centre for Molecular Medicine Norway (NCMM, 187615).

*Conflict of Interest:* none declared.

## References

- Alquicira-Hernandez, J. and Powell, J.E. (2021) Nebulosa recovers single cell gene expression signals by kernel density estimation. *Bioinformatics*, **37**, 2485–2487.
- Butler, A. et al. (2018) Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.*, **36**, 411–420.
- Finak, G. et al. (2015) Mast: a flexible statistical framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol.*, **16**, 278.
- Franzen, O. and Bjorkegren, J.L.M. (2020) alona: a web server for single-cell RNA-Seq analysis. *Bioinformatics*, **36**, 3910–3912.
- Franzen, O. et al. (2019) PanglaoDB: a web server for exploration of mouse and human single-cell RNA sequencing data. *Database (Oxford)*, **2019**, baz046.
- Grieb, G. et al. (2011) Circulating fibrocytes—biology and mechanisms in wound healing and scar formation. *Int. Rev. Cell Mol. Biol.*, **291**, 1–19.
- Hanzelmann, S. et al. (2013) GSVA: gene set variation analysis for microarray and RNA-Seq data. *BMC Bioinformatics*, **14**, 7.
- Hie, B. et al. (2019) Efficient integration of heterogeneous single-cell transcriptomes using scanorama. *Nat. Biotechnol.*, **37**, 685–691.
- Korotkevich, G. et al. (2021) Fast gene set enrichment analysis. *bioRxiv*.
- Korsunsky, I. et al. (2019) Fast, sensitive and accurate integration of single-cell data with harmony. *Nat. Methods*, **16**, 1289–1296.
- Lachmann, A. et al. (2018) Massive mining of publicly available RNA-Seq data from human and mouse. *Nat. Commun.*, **9**, 1366.
- Lim, C.H. et al. (2018) Hedgehog stimulates hair follicle neogenesis by creating inductive dermis during murine skin wound healing. *Nat. Commun.*, **9**, 4903.
- Luecken, M. et al. (2020) Benchmarking atlas-level data integration in single-cell genomics. *bioRxiv*.
- Pilling, D. et al. (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. *PLoS One*, **4**, e7475.
- Reilkoff, R.A. et al. (2011) Fibrocytes: emerging effector cells in chronic inflammation. *Nat. Rev. Immunol.*, **11**, 427–435.
- Stuart, T. et al. (2019) Comprehensive integration of single-cell data. *Cell*, **177**, 1888–1902.e21.
- Swamy, V.S. et al. (2021) Building the mega single cell transcriptome ocular meta-atlas. *bioRxiv*.
- Xie, Z. et al. (2021) Gene set knowledge discovery with enrichr. *Curr. Protoc.*, **1**, e90.
- Zhang, S. et al. (2018) Prostaglandin e2 hydrogel improves cutaneous wound healing via m2 macrophages polarization. *Theranostics*, **8**, 5348–5361.