



REVIEW

Open Access



Mental health and psychological impacts from the 2011 Great East Japan Earthquake Disaster: a systematic literature review

Nahoko Harada^{1,2*}, Jun Shigemura³, Masaaki Tanichi³, Kyoko Kawaida¹, Satomi Takahashi¹ and Fumiko Yasukata¹

Abstract

Background: On March 11, 2011, Japan experienced an unprecedented combination of earthquake/tsunami/nuclear accidents (the Great East Japan Earthquake; GEJE). We sought to identify mental health and psychosocial consequences of this compound disaster.

Method: A systematic literature review was conducted of quantitative research articles addressing mental health of survivors and the psychological impact of the GEJE. For articles between March 2011 and December 2014, PubMed, PsychINFO, and EMBASE databases were searched with guidance on literature review method.

Results: Forty-nine studies met the inclusion criteria. A substantial proportion of the affected individuals experienced considerable psychological distress. Mental health outcomes included, but were not limited to, posttraumatic stress disorder, depression, and anxiety symptoms. Physical health changes, such as sleeping and eating disturbances, also occurred. In Fukushima, radioactive release induced massive fear and uncertainty in a large number of people, causing massive distress among the affected residents, especially among mothers of young children and nuclear plant workers. Stigma was additional challenge to the Fukushima residents. The review identified several groups with vulnerabilities, such as disaster workers, children, internally displaced people, patients with psychiatric disorders, and the bereaved.

Conclusions: Following the GEJE, a considerable proportion of the population was mentally affected to a significant degree. The affected individuals showed a wide array of mental and physical consequences. In Fukushima, the impact of nuclear disaster was immense and complex, leading to fear of radiation, safety issues, and stigma issues.

Keywords: Disaster, Mental health, Psychological service, Posttraumatic stress disorder, Great East Japan Earthquake, Earthquake, Tsunami, Fukushima Daiichi nuclear accident, Radiation fear

Background

On March 11, 2011, a 9.0-magnitude mega-earthquake hit the islands of Japan at 2:46 pm local time. This earthquake was the strongest recorded earthquake in the country's modern history [1]. The earthquake's epicenter

was located approximately 80 km off the northeastern (Tohoku) region of the island of Honshu, the country's main island. Repeated aftershocks and towering tsunami waves occurred after the mega-earthquake; the tsunami waves were as high as 40 m above sea level and reached 10 km inland (Fig. 1) [2]. A large majority of the damage occurred in three prefectures in the Tohoku area: Iwate, Miyagi, and Fukushima (Fig. 2). As of December 10, 2014, the numbers of dead, missing, or injured were 15,889, 2594, and 6152, respectively [3].

*Correspondence: nahoko-harada@umin.ac.jp

¹ Division of Nursing, School of Medicine, National Defense Medical College, 3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan
Full list of author information is available at the end of the article




Fig. 1 Locations of earthquake epicenter of the 2011 Great East Japan Earthquake and major cities

In Fukushima, the earthquakes and tsunamis triggered a nuclear accident at Tokyo Electric Power Company (TEPCO) Fukushima Daiichi Nuclear Power Plant. Between March 11 and 15, 2011, four of the six reactors experienced explosions, and three reactors escalated to nuclear meltdown and released radioactive materials, requiring a mandatory evacuation of the surrounding region. This crisis became the largest nuclear accident since the 1986 Chernobyl nuclear disaster, and the second accident, after Chernobyl, to measure Level 7 on the International Nuclear Event Scale.

On March 12, 2011, the Japanese government ordered mandatory evacuation of residents in the 20 km radius

of the nuclear plant. As of December 2014, more than 121,000 residents are still on evacuation status [4]. Although there have been no fatalities owing to radiation exposure, safety concerns related to nuclear contamination created enormous fear, burden, and disruption to individuals, groups, communities, and local/national governments. These series of disasters were eventually named as the Great East Japan Earthquake (GEJE).

When evaluating post-disaster outcome studies, it must be noted every disaster is different in the terms of disaster type (e.g., natural, technological, manmade), location (e.g., developed vs. developing country), population (e.g., adults vs. children), intensity (life-threatening vs. non-),




Fig. 2 Three severely affected prefectures of the Tohoku region, Japan; Iwate, Miyagi, and Fukushima

exposure frequency (e.g., single vs. repeated experiences), and many other medical/social/economic conditions. It is also very rare to randomize study samples or to have pre-disaster comparison [5].

Given these limitations, researches have shown a large majority of the people affected by disasters is resilient and will fully recover from their traumatic experiences [6]. A small portion of the affected individuals will result in a wide range of mental, behavioral, and physical health consequences [e.g., depression, posttraumatic stress disorder (PTSD) and other anxiety disorders, suicidal behaviors, alcohol misuse, and sleep disturbances]. Studies also have highlighted at-risk populations for adverse outcomes, such as female gender, preexisting psychiatric

illnesses, presence of children in the home, secondary stressors, and low psychosocial resources [5–7]. Disaster workers, exposed to a variety of traumatic exposure through their work roles, are of significant concern when considering post-disaster mental health [5]. This trend was true for the Chernobyl first responders and cleanup workers as well; their depression and PTSD rates remained high two decades later [8].

A wide majority of these studies revealed the impact on mental health and psychosocial aspects of one particular event (e.g., man-made or natural). However, in the case of large-scale compound disasters, the epidemiological data are scarce. As of writing, GEJE mental health studies are evolving, but scientific review of this disaster's mental

health studies is warranted to increase better understanding of psychosocial outcomes of the people affected by the GEJE.

The aims of this article are to (1) consolidate quantitative and qualitative studies examining mental health and psychosocial impacts in people affected by the GEJE, (2) identify their mental health and psychosocial consequences, (3) ascertain vulnerable populations, and (4) elucidate factors that impact mental health and psychosocial outcomes in populations affected by the earthquake.

Methods

A systematic literature review was conducted of quantitative research articles addressing mental health of survivors and the psychological impact of the 2011 GEJE between March 2011 and December 2014. PubMed, PsychINFO, and EMBASE databases were searched with guidance on literature review method [9]. The searched keywords included Great East Japan Earthquake Disaster, Japan, disaster, health, mental health, psychological, impact, stress, trauma, bereavement, and grief and these items were used either alone or in combination.

All identified articles were examined with the title and abstract whether the article specifically addresses mental health and psychosocial issues related to the GEJE by the investigators (NH and JS). If the abstract unclearly described the study aims, method, or results, NH and JS read the article to determine the relevance of the article. Studies written in a language other than English, situational reports, activity reports, conference reports/abstracts/summaries, letters to the editor (including replies), and bulletins from universities or private organizations were not included in this review. To ensure inclusiveness, the other authors of the current article were encouraged to search for articles manually and, if additional articles met the inclusion criteria, such articles were also included for review.

A total of 382 studies were identified, with 49 articles meeting the inclusion criteria. The identified articles were categorized by four main research topics: (1) mental health outcomes among affected populations (excluding Fukushima), (2) Fukushima resident studies, (3) disaster and support worker researches, and (4) grief studies. We decided to separate the first two topics because of the uniqueness of nuclear disaster and a potentially profound mental health impact among the affected people.

Results

Table 1 summarizes the study results of mental health outcomes among the affected populations (excluding Fukushima). A total of 28 articles met this inclusion criterion [10–37]. A majority of the study populations

were from Miyagi and Iwate [15 (53.6 %) and 3 (10.7 %), respectively]. Other studies included subjects from the Ibaraki, Tochigi, Tokyo, among others. A major portion of the studies were cross-sectional. Regarding outcome measures, nine (32.1 %) studies addressed PTSD, six (21.4 %) assessed general psychological distress, two (7.1 %) examined depressive symptoms. Other outcomes included anxiety, sleep disturbance, social functioning, social isolation, admission rates, suicide rates, and cerebral structure changes.

Table 2 compiles the 12 study results reporting psychosocial consequences of the individuals affected by the Fukushima nuclear disaster [38–49]. Adverse outcome measures were primarily general psychological distress, symptoms of PTSD, depression, as well as anxiety disorders, especially in context with radiation fear. Other outcomes represented the uniqueness of a nuclear disaster, such as concern of radiation and food safety [43, 49], maternal anxiety (including food safety, outdoor safety, radiation effects on embryos, economic issues, distrust towards information disclosure) [47, 49], lowered maternal confidence [47], and stigma owing to their radiation exposures [48].

Table 3 represents eight study outcomes of GEJE disaster and support workers [50–57]. Their outcomes measures were general psychological distress as well as symptoms of PTSD or depression. Of note, a study of Fukushima nuclear plant workers showed discrimination/slurs experience as a key factor for their mental health consequences [56]. This trend was associated with the public criticism to the electric company's post-disaster management. One study focused on potential of fish oil in attenuating PTSD symptoms among DMAT (Disaster Medical Assistant Team) medical workers [53].

Table 4 shows a result of a sole study examining the grief responses affected by the GEJE [58]. This study showed the distinctiveness of complicated grief from symptoms of PTSD or depression. Other articles, not listed in the table, were leaned on narratives and support activity reports. For example, our co-author (ST) launched a support group immediately after the disaster for people in grief and bereavement [59]. This support group aims to (1) provide information about grief, the concept of which is not as popularly known in Japan (much like PTSD), and (2) provide training sessions and workshops in collaboration with the local grief support organizations.

Discussions

Our review compiled a wide array of mental health consequence following the GEJE, an unprecedented compound disaster with a combination of earthquakes, tsunamis, and a series of nuclear accidents. A

Table 1 Mental health outcomes among individuals affected by the GEJE (excluding Fukushima)

Citation (reference no.)	Sample type (location)	N	Data collection ^a	Findings (subject proportions, measurements)	Risk factors of outcome(s)
Yokoyama et al. [13]	Residents (Iwate)	10,025	6 and 11 months	42.6 %; K6 \geq 5	Severe financial problems, displacement, lack of network
Niitsu et al. [12]	Residents (Iwate)	902	11 months	48 %; K6 \geq 5	Female, middle-to-low educational status, unemployment
Koyama et al. [10]	Residents (Miyagi)	281	11 and 12 months	35.9 %; K6 \geq 13	No social support, lower annual income, cohabitating with \geq 6 people
Nagata et al. [11]	Residents of temporary housings (Iwate)	200	10–12 and 19–21 months	No significance in K6 \geq 5 between two time points (37.5 %, 10–12 months vs. 43.5 %, 19–21 months). Sense of isolation higher at 19–21 months	
Sugimoto et al. [14]	National sample	8777	12 months	No significance in K6 \geq 13 between certified vs. non-certified groups for house damage (8.4 vs. 9 %)	Lack of support from family, friends and neighbors
Fujihara et al. [15]	Diabetic patients (Ibaraki)	320	3 months pre- and post-disaster	Worsening of glycemic control was associated with total GHQ scores, interruption of drug regimen, somatic symptoms, and sleep disturbances/anxiety	
Inoue et al. [17]	Household of tsunami-affected houses (Miyagi)	4176	7–12 months	Social isolation	Men <65 years and living alone, low income
Funayama et al. [16]	Psychiatric outpatients (Tochigi)	294	2 months	4.1 % worsened and 1.2 % improved in GAF score	GAF score >50
Saito et al. [18]	Psychiatric in/outpatients (Tokyo)	155	1 week	3.5 % worsening of psychiatric symptoms (increase in epileptic seizure or GAF score)	
Aoki et al. [19]	Psychiatric patients on mandatory admission (Tokyo)	224	6 months pre- and post-disaster	Increased admission cases post-disaster (n = 127) compared to pre-disaster (n = 97)	Schizophrenia
Kato et al. [20]	Psychiatric inpatients with suicide attempt (Kanagawa)	592	6–1 months before and 1–6 months post-disaster	The number of admitted patients on ventilator was higher after the earthquake (Pre, n = 87 vs. post, n = 123)	Jobless, family psychiatric history, precipitating attempt, and alcohol intake
Orui et al. [23]	National government statistics report (Iwate, Miyagi and Fukushima)		24 months pre- and post-disaster	Suicide rates in men decreased during the post-disaster period; rates in women increased in the first 7 months	
Momma et al. [22]	Small and medium enterprise employees (Miyagi)	522	7 months pre- and 5 months post-disaster	14.3 % (men), IES-R \geq 25 (5 months post-disaster) 24.4 % (women), IES-R \geq 25 (5 months post-disaster)	Male: weak bilateral leg extension power, daily drinking habits, and depressive symptoms Female: hypertension and depressive symptoms

Table 1 continued

Citation (reference no.)	Sample type (location)	N	Data collection ^a	Findings (subject proportions, measurements)	Risk factors of outcome(s)
Takeda et al. [24, 25]	Female high school students (Miyagi)	1180	9 months	10 % IES-R ≥ 25 Associated with premenstrual syndrome and premenstrual dysphoric disorder severity	
Iwadare et al. [21]	Junior high school student (Miyagi)	1919	8 and 20 months	Shorter sleep duration and later bedtime at 20 months	Bereavement experience
Usami et al. [36, 37]	Children, 4–15 years (Miyagi)	11,639	8 and 20 months	42.6 % PTSSC-15 ≥ 23 at 8 months	Evacuation, house damage and/or separation from family, female, and not having breakfast
Kuwabara et al. [26]	Children, 6–15 years (Miyagi)	2259	6 months	Students from junior high schools with mortality rate ≥ 4 % had higher PTSSC-15 scores	
Numata et al. [31]	PTSD outpatients (Miyagi)	43	Unspecified	2.5 g of <i>saikokeishikankyo</i> powder 3 times a day for 2 weeks improved IES-R score	
Tuerk et al. [35]	Residents (Ibaraki)	41	40 days post-event	27 % self-reported PTSD symptoms	Subjective health and loss of sense of community
Niitsu et al. [29]	College students (USA)	30	12–14 months	Japanese students (n = 17) reported higher hyper-arousal than did non-Japanese students (n = 13)	Media exposure
Sekiguchi et al. [32]	Non-PTSD residents (Miyagi)	42	Pre-disaster, 3–4 months post-disaster	Regional volume changes in the brain observed after the disaster	
Sekiguchi et al. [33]	Non-PTSD residents (Miyagi)	30	Pre-disaster and 3–4 months post-disaster	Post-disaster anxiety level associated with cerebral structure changes	Smaller GMV in the ACC before the earthquake, decreased GMV in the OFC through the earthquake
Matsubara et al. [27]	Survivors remained at damaged residences (Miyagi)	5454	1–4 months	8.1 % depressive reaction (PHQ-2)	Lower FA in the right anterior cingulum, increased FA in the left anterior cingulum and uncinated fasciculus
Nishigori et al. [30]	Postpartum women (Miyagi)	677	1 month pre- and 7 months post-disaster	20 % Edinburgh Postnatal Depression Scale ≥ 9	House flooding below or above the ground floor, unavailability of gas supply, female, middle aged or elderly, regular intake of psychotropic medicine(s) since before the tsunami, no cohabitant
Matsumoto et al. [28]	Residents (Miyagi)	4176	7–12 months	15 % Sleep disturbance measured by an original scale	Maternal age of under 25 years, child's birth weight under 2.5 kg
Sugjura et al. [34]	Food delivery users (Tokyo and Osaka)	5053	2 months pre- and 1 months post-disaster	Insomnia (original scale), post- vs. pre-disaster odds ratio; Tokyo, 2.0, Osaka, 1.6	Lack of pleasure in life, lack of interaction with neighbors

GEJE Great East Japan Earthquake, K6 Kessler Psychological Distress Scale, GHQ General Health Questionnaires, GAF global assessment of functioning, IES-R impact of event scale-revised, PTSSC-15 posttraumatic stress symptoms for children 15 items, GMV grey matter volume, ACC anterior cingulate cortex, OFC orbitofrontal cortex, FA fractional anisotropy, PHQ-2 Patient Health Questionnaire-2

^a Cross-sectional studies unless otherwise noted

Table 2 Mental health outcomes of Fukushima residents following the GEJE

Citation (reference no.)	Sample type (location)	n	Data collection ^a	Findings	Risk factors of outcome(s)
Matsumoto et al. [39]	Psychiatric outpatient (Fukushima)	1273	1 month	Bipolar I showed worst exacerbation among psychiatric diseases and manic change was prominent	
Wada et al. 2013 [45]	Psychiatric inpatient (Fukushima)	28	7 days	Two-thirds showed no change	
Yabe et al. [46]	Residents (Fukushima)	73,433 (2011), 39495 (2012)	10 and 22 months	14.6 vs. 11.9 %, K6 \geq 13, 21.6 vs. 18.3 %, PCL \geq 44 24.4 vs. 16.6 %, SDQ \geq 16, 4–6 years 22.0 vs. 15.8 %, SDQ \geq 16, 6–12 years (2011 vs. 2012)	
Tsubokura et al. [44]	Residents (Fukushima)	155	1 year pre- and post-disaster	12 %, PHQ-9 \geq 10	
Kukihara et al. [38]	Evacuees (Fukushima)	241	9 months	33.2 %, IES-R \geq 25 14.5 %, Zung Depression Scale \geq 60 Resilience was predicted by employment status, eating/exercise and drinking habits	
Sawa et al. [41, 42]	Internally displaced people from Fukushima (Chiba)	73	5 and 10 months	Compared to a reference group (Chiba residents), the study sample was more likely to have GHQ-12 \geq 3, adverse social dysfunction at both time points and psychological distress at 10 months	
Sugimoto et al. [43]	National sample	5809	1 year	Women were more concerned than men about radiation. Participants from Kanto region (vs. non-Tohoku/Kanto regions) were more concerned about radiation and food safety	
Rubin et al. [40]	British nationals living in Japan	284	9 mo.	16 %, GHQ-12 \geq 4 29.7 %, State-trait anxiety inventory \geq 18 30.4 %, State-trait anger inventory \geq 11	
Goto et al. [47]	Parenting counseling records (Fukushima)	150	1 year pre-, 1 and 13 months post-disaster (qualitative analysis)	Lowered maternal confidence and potential role of public health nurses in risk communication process post-disaster were reported	
Yoshii et al. [49]	Post-partum women (Miyagi)	259	16 months (qualitative analysis)	Seven themes of maternal anxiety for radioactivity from the Fukushima emerged: food safety, outdoor safety, radiation effects on embryos during pregnancy, effects on children, radiation exposures, economic issues and distrust towards disclosing information	
Ben-Ezra et al. [48]	Residents (Hiroshima/Nagasaki, Tokyo and Fukushima)	750	3 years post-disaster	10.6 %, Fukushima, endorsed PTSD symptoms 2.4 %, Hiroshima/Nagasaki and Tokyo, endorsed PTSD symptoms Relations between location, radiation stigma, and number of PTSD symptoms	

GEJE Great East Japan Earthquake, K6 Kessler Psychological Distress Scale, PCL PTSD checklist, SDQ Strengths and Difficulties Questionnaire, PHQ-9 Patient Health Questionnaire-9, IES-R impact of events-revised, GHQ-12 General Health Questionnaires 12

^a Cross-sectional studies unless otherwise noted

Table 3 Studies of the GEJE disaster and support workers

Citation (reference no.)	Sample type (location)	n	Data collection ^a	Findings	Risk factors of outcome(s)
Shigemura et al. [55, 56]	Nuclear plant workers (Fukushima)	1495	2–3 months	42.7 %, K6 \geq 13 25.3 %, IES-R \geq 25	Preexisting illness(es), discrimination/slurs, near-death experience, tsunami evacuation, major property loss, home evacuation
Dobashi et al. [50]	Defense personnel (Miyagi)	606	1 month post deployment	6.2 (\pm 8.1), IES-R 12.8 (\pm 4.4), K10	No identified factors
Matsuoka et al. [52]	Disaster Medical Assistant Team	426	1 month	4.0 %, K6 \geq 13 21.4 %, CES-D \geq 17	Concern over radiation exposure
Nishi et al. [54]	Disaster Medical Assistant Team	173	4 months	6.8 (\pm 8.4), IES-R	PDI score and watching earthquake TV news reports \geq 4 h/day
Nishi et al. [53]	Disaster Medical Assistant Team	172	Baseline and 12-week post-intervention	Fish oil attenuated posttraumatic stress symptoms among female	
Fukasawa et al. [51]	Government workers (Miyagi)	4331	2 months	3.0 %, K6 \geq 13 (group with less property damages) 5.9 %, K6 \geq 13 (group with severe property damages)	Less damaged: working overtime (>100 h/mo. overtime), poor workplace communication Severe damaged: handling residents' complaints, poor workplace communication
Suzuki et al. [57]	Government workers (Miyagi)	3743	7 months	9.6 %, K6 \geq 10 4.4 %, K6 \geq 13	Not taking a non-work day each week

GEJE Great East Japan Earthquake, K6 (K10) Kessler Psychological Distress Scale, IES-R impact of events-revised, CES-D Center for Epidemiologic Studies Depression Scale, PDI peritraumatic distress inventory

^a Cross-sectional studies unless otherwise noted

Table 4 Grief study following the GEJE

Citation (reference no.)	Sample type (location)	N	Data collection ^a	Findings	Risk factors of outcome(s)
Tsutsui et al. [58]	Hospital workers (tsunami-affected area)	82	8 months	9.8 %, ICG \geq 25 29.3 %, IES-R \geq 25 37.8 %, CES-D \geq 16 Prolonged grief disorder in qualitatively distinct from PTSD and major depressive disorder	

GEJE Great East Japan Earthquake, ICG Inventory of Complicated Grief, IES-R impact of events-revised, CES-D Center for Epidemiological Studies Depression Scale

^a Cross-sectional studies unless otherwise noted

considerable proportion of the study population was mentally affected to a substantial degree, and mental health responses ranged from approximately one-tenth to nearly half of the respondents [12, 27]. Mental health outcomes included, but were not limited to, PTSD, depression, and anxiety. Physical health changes, such as sleep and eating disturbances, were also reported.

Although every disaster is different, disasters are large-scale, stressful, and distressing events that affect a significant number of people. Those who experience higher exposure to traumatic events are likely to show higher mental health responses (i.e., dose–response relationship) [60]. For the most people, these acute responses are normal and gradually decrease over time, but a small proportion of the affected individuals will suffer long-term mental health issues. In a review of 160 disaster mental health studies, proportions of subjects with severe impairment were 21.6 % for natural disasters and 18.5 % for technological disaster samples [6]. The articles in this review had relatively higher mental health rates than in previous studies. This trend might be related to the high impact of this disaster as well as the GEJE study timing, because most of the studies were conducted among the direct victims within 2 years after the disaster. Long-term, longitudinal studies are evolving, and they will potentially be useful to understand the trajectories of mental health consequences among these people.

In the region affected by the Fukushima nuclear disaster, invisible and imperceptible nature of radioactive materials has been challenged among the affected people. The residents' responses were diverse and complex; along with high proportions of mental health distress among the Fukushima residents [46], concerns for radiation effect were a prominent concern especially among pregnant women and mothers of young children [43, 49, 61]. Safety issues in food and outdoor activities, along with economic issues and distrust in information disclosure were also reported [43, 49]. Public psychosocial

responses such as discrimination and stigmatization were also reported [48, 56].

These findings are in accordance with a series of Chernobyl studies where a complex relationship between radiation exposure and physical/mental health effect has been an ongoing debate. Physical outcome studies tend to be controversial, although firm evidence can be found only on the deaths of first responders due to acute radiation exposures and high prevalence of thyroid cancer among the exposed children [8]. Still, psychosocial and economic disruptions to the affected people were significant, and the International Atomic Energy Agency regarded mental health as the major public health sequelae of the Chernobyl accident [62]. Mothers of young children and plant clean-up workers were among the two groups of particular concern [63–65]. Psychosocial issues included not only mental health disorders but also stigmatization and discrimination of the affected people [8], suggesting the importance of integrity and accuracy of information as well as risk communication strategies.

Two Fukushima studies reported distress among internally displaced people [38, 41]. Mandatory evacuation measures have been in place for the area surrounding the nuclear plant, and the evacuees potentially have uncertain and ambiguous perspectives on whether or not they will be able to return home [66]. This trend was also compatible with Chernobyl studies reporting challenges in evacuation and resettlement [8]. Future studies will be essential to clarify the effect of evacuation following nuclear disasters.

Fukushima and Chernobyl studies suggest that substantial public health efforts are crucial to establish a system capable of such exposures. Integrity and accuracy of information will be a critical issue for the public to assess their health status. These studies also have implications for other “tangible” disasters, such as emergencies related to bio-chemical weapons and infectious diseases [67]. Long-term studies will be important to increase the psychosocial impact among Fukushima residents, with

special focus on children, mothers, and nuclear plant workers.

A number of studies assessed a considerable degree of mental distress among disaster workers. This is likely to be owing to work-related exposures of these workers. In the case of GEJE, many of the workers were also local disaster victims, and had struggles as survivors along with their work-related exposures. This effect was prominent in several studies [51, 56, 57]. Two worker studies identified experiences of being discriminated against and handling residents' complaints as risk factors for their adverse mental health [56, 57]. In the former study, the Fukushima nuclear plant workers became targets of public criticism because their company was blamed for their post-disaster mismanagement. In the latter, Miyagi Prefecture workers received direct complaints from their residents in a chaotic situation. These results might give hypotheses that mental health of disaster workers is susceptible to their stakeholder's criticisms.

Past literatures identified mortuary work as predictors of PTSD or physical symptoms among disaster workers [68, 69], but in our review, there has yet be an evidence that mortuary work was associated with adverse mental health [50, 51]. Further studies will be needed to elucidate the relationship between dead body exposure and mental health outcomes among this population.

Previous studies highlighted vulnerable populations for post-disaster mental health, such as children, evacuees, the bereaved, and individuals with preexisting physical/mental health conditions [5, 6]. Our compilation overall shows a similar trend, although studies are relatively few, especially in the context of grief.

We recognize several limitations of this paper. The GEJE, especially the Fukushima nuclear accident, is an ongoing disaster, and new studies are emerging. Given the timing and methodology of our literature search, we were not able to include narrative studies, non-English papers, or papers describing long-term disaster impact. Although we made every possible effort to include all related studies, some studies may have been inadvertently omitted.

Given these limitations, this literature review encompasses research on the mental health trajectories of people affected by the GEJE, a complex earthquake/tsunami/nuclear disaster. Along with our review, future studies will be essential for having a better understanding of this disaster, and especially for ascertaining the long-term outcomes and their correlates.

Authors' contributions

NH carried out the initial literature search. NH, JS, and KK performed the manual searches for the additional articles. NH and JS reviewed the potential study papers and made decisions on papers to be included. NH, JS, MT, KK, and ST drafted the manuscript. FY supervised the study process. All authors read and approved the final manuscript.

Author details

¹ Division of Nursing, School of Medicine, National Defense Medical College, 3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan. ² William F. Connell School of Nursing, Boston College, Maloney Hall 140 Commonwealth Avenue, Chestnut Hill, MA 02467, USA. ³ Department of Psychiatry, National Defense Medical College, 3-2 Namiki, Tokorozawa, Saitama 359-8513, Japan.

Acknowledgements

None.

Compliance with ethical guidelines

Competing interests

The authors declare that they have no competing interests.

Received: 8 February 2015 Accepted: 25 August 2015

Published online: 02 September 2015

References

- Lay T, Kanamori H. Insights from the great 2011 Japan earthquake. *Phys Today*. 2011;64(12):33–9. doi:10.1063/PT.3.1361.
- Yokoyama Y, Hirano K, Sato M, Abe A, Uebayashi M, Kishi E, et al. Activities and health status of dispatched public health nurses after the great East Japan earthquake. *Public Health Nurs (Boston, Mass)*. 2014;31(6):537–44. doi:10.1111/phn.12141.
- National Police Agency. Damage Situation and Police Countermeasures associated with 2011 Tohoku district-off the Pacific Ocean Earthquake, December 10, 2014. 2014. http://www.npa.go.jp/archive/keibi/biki/higaijokyo_e.pdf. Accessed Dec 30 2014.
- Fukushima Prefecture. Damage situation report (No. 1344) following the 2011 Tohoku district-off the Pacific Ocean Earthquake. 2015. http://www.pref.fukushima.lg.jp/uploaded/ife/109612_206594_misc.xls. Accessed Jan 5 2015 (in Japanese).
- Galea S, Nandi A, Vlahov D. The epidemiology of post-traumatic stress disorder after disasters. *Epidemiol Rev*. 2005;27:78–91. doi:10.1093/epirev/mxi003.
- Norris FH, Friedman MJ, Watson PJ, Byrne CM, Diaz E, Kaniasty K. 60,000 disaster victims speak: Part I. An empirical review of the empirical literature, 1981–2001. *Psychiatry*. 2002;65(3):207–39.
- Norris FH, Friedman MJ, Watson PJ. 60,000 disaster victims speak: Part II. Summary and implications of the disaster mental health research. *Psychiatry*. 2002;65(3):240–60.
- Bromet EJ, Havenaar JM, Guey LT. A 25 year retrospective review of the psychological consequences of the Chernobyl accident. *Clin Oncol*. 2011;23(4):297–305. doi:10.1016/j.clon.2011.01.501.
- Centre for Reviews and Dissemination. Systematic reviews: CRD's guidance for undertaking reviews in health care. York: Centre for Reviews and Dissemination, University of York; 2009.
- Koyama S, Aida J, Kawachi I, Kondo N, Subramanian SV, Ito K, et al. Social support improves mental health among the victims relocated to temporary housing following the Great East Japan Earthquake and Tsunami. *Tohoku J Exp Med*. 2014;234(3):241–7.
- Nagata S, Matsunaga A, Teramoto C. Follow-up study of the general and mental health of people living in temporary housing at 10 and 20 months after the Great East Japan Earthquake. *Jpn J Nurs Sci Jjns*. 2014;. doi:10.1111/jjns.12051.
- Niitsu T, Takaoka K, Uemura S, Kono A, Saito A, Kawakami N, et al. The psychological impact of a dual-disaster caused by earthquakes and radioactive contamination in Ichinoseki after the Great East Japan Earthquake. *BMC Res Notes*. 2014;7:307. doi:10.1186/1756-0500-7-307.
- Yokoyama Y, Otsuka K, Kawakami N, Kobayashi S, Ogawa A, Tannno K, et al. Mental health and related factors after the Great East Japan earthquake and tsunami. *PLoS One*. 2014;9(7):e102497. doi:10.1371/journal.pone.0102497.
- Sugimoto T, Umeda M, Shinozaki T, Naruse T, Miyamoto Y. Sources of perceived social support associated with reduced psychological distress at 1 year after the Great East Japan Earthquake: nationwide cross-sectional survey in 2012. *Psychiatry Clin Neurosci*. 2014;. doi:10.1111/pcn.12235.

15. Fujihara K, Saito A, Heianza Y, Gibo H, Suzuki H, Shimano H, et al. Impact of psychological stress caused by the Great East Japan Earthquake on glycemic control in patients with diabetes. *Exp Clin Endocrinol Diabetes Off J Ger Soc Endocrinol Ger Diabetes Assoc.* 2012;120(9):560–3. doi:10.1055/s-0032-1314873.
16. Funayama M, Mizushima J. Severity of pre-existing psychiatric illness and response to the Great East Japan Earthquake. *J Psychiatr Res.* 2013;47(10):1479–82.
17. Inoue M, Matsumoto S, Yamaoka K, Muto S. Risk of social isolation among Great East Japan Earthquake survivors living in tsunami-affected Ishinomaki, Japan. *Disaster Med Public Health Prep.* 2014;8(4):333–40. doi:10.1017/dmp.2014.59.
18. Saito M. The short-term effect of the recent great earthquake on the general functioning status of psychiatric patients near Tokyo. *Am J Psychiatry.* 2011;168(6):649. doi:10.1176/appi.ajp.2011.11030450.
19. Aoki A, Aoki Y, Harima H. The impact of the Great East Japan earthquake on mandatory psychiatric emergency hospitalizations in Tokyo: a retrospective observational study. *Transl Psychiatry.* 2012;2:e168. doi:10.1038/tp.2012.98.
20. Kato K, Mikami K, Kimoto K, Kimoto K, Takahashi Y, Sato R, et al. Changes in the frequency and clinical features of suicide attempts in the Midwestern area of Kanagawa after the Great East Japan Earthquake. *J Forensic Sci.* 2014;59(2):417–9.
21. Iwadare Y, Usami M, Ushijima H, Tanaka T, Watanabe K, Kodaira M, et al. Changes in traumatic symptoms and sleep habits among junior high school students after the Great East Japan Earthquake and Tsunami. *Sleep Biol Rhythm.* 2014;12(1):53–61.
22. Momma H, Niu K, Kobayashi Y, Huang C, Otomo A, Chujo M, et al. Leg extension power is a pre-disaster modifiable risk factor for post-traumatic stress disorder among survivors of the Great East Japan Earthquake: a retrospective cohort study. *PLoS One.* 2014;9(4):e96131. doi:10.1371/journal.pone.0096131.
23. Orui M, Harada S, Hayashi M. Changes in suicide rates in disaster-stricken areas following the Great East Japan Earthquake and their effect on economic factors: an ecological study. *Environ Health Prev Med.* 2014;19(6):459–66. doi:10.1007/s12199-014-0418-2.
24. Takeda T, Tadakawa M, Koga S, Nagase S, Yaegashi N. Relationship between dysmenorrhea and posttraumatic stress disorder in Japanese high school students 9 months after the Great East Japan Earthquake. *J Pediatr Adolesc Gynecol.* 2013;26(6):355–7. doi:10.1016/j.jpjag.2013.06.020.
25. Takeda T, Tadakawa M, Koga S, Nagase S, Yaegashi N. Premenstrual symptoms and posttraumatic stress disorder in Japanese high school students 9 months after the great East-Japan earthquake. *Tohoku J Exp Med.* 2013;230(3):151–4.
26. Kuwabara H, Araki T, Yamasaki S, Ando S, Kano Y, Kasai K. Regional differences in post-traumatic stress symptoms among children after the 2011 tsunami in Higashi-Matsushima, Japan. *Brain Dev.* 2014;37(1):130–6. doi:10.1016/j.braindev.2014.02.003.
27. Matsubara C, Murakami H, Imai K, Mizoue T, Akashi H, Miyoshi C, et al. Prevalence and risk factors for depressive reaction among resident survivors after the tsunami following the Great East Japan Earthquake, March 11, 2011. *PLoS One.* 2014;9(10):e109240. doi:10.1371/journal.pone.0109240.
28. Matsumoto S, Yamaoka K, Inoue M, Muto S. Social ties may play a critical role in mitigating sleep difficulties in disaster-affected communities: a cross-sectional study in the Ishinomaki area, Japan. *Sleep.* 2014;37(1):137–45. doi:10.5665/sleep.3324.
29. Niitsu K, Watanabe-Galloway S, Sayles H, Houfek J, Rice M. A Pilot Study of the Psychological Impact of the Great East Japan Earthquake and Tsunami. *J Am Psychiatr Nurs Assoc.* 2014;20(3):194–202. doi:10.1177/1078390314536615.
30. Nishigori H, Sugawara J, Obara T, Nishigori T, Sato K, Sugiyama T, et al. Surveys of postpartum depression in Miyagi, Japan, after the Great East Japan Earthquake. *Arch Women's Mental Health.* 2014;17(6):579–81. doi:10.1007/s00737-014-0459-y.
31. Numata T, Gunfan S, Takayama S, Takahashi S, Monma Y, Kaneko S, et al. Treatment of posttraumatic stress disorder using the traditional Japanese herbal medicine saikokeishikanyoto: a randomized, observer-blinded, controlled trial in survivors of the great East Japan earthquake and tsunami. *Evid Based Complement Alter Med Ecam.* 2014;2014:683293. doi:10.1155/2014/683293.
32. Sekiguchi A, Sugiura M, Taki Y, Kotozaki Y, Nouchi R, Takeuchi H, et al. Brain structural changes as vulnerability factors and acquired signs of post-earthquake stress. *Mol Psychiatry.* 2013;18(5):618–23. doi:10.1038/mp.2012.51.
33. Sekiguchi A, Sugiura M, Taki Y, Kotozaki Y, Nouchi R, Takeuchi H, et al. White matter microstructural changes as vulnerability factors and acquired signs of post-earthquake distress. *PLoS One.* 2014;9(1):e83967. doi:10.1371/journal.pone.0083967.
34. Sugiura H, Akahane M, Ohkusa Y, Okabe N, Sano T, Jojima N, et al. Prevalence of insomnia among residents of Tokyo and Osaka after the great East Japan earthquake: a prospective study. *Interact J Med Res.* 2013;2(1):e2. doi:10.2196/ijmr.2485.
35. Tuerk PW, Hall B, Nagae N, McCauley JL, Yoder M, Rauch SA, et al. Forty days after the Great East Japan Earthquake: field research investigating community engagement and traumatic stress screening in a post-disaster community mental health training. *Int J Psychiatry Med.* 2013;45(2):159–74.
36. Usami M, Iwadare Y, Kodaira M, Watanabe K, Aoki M, Katsumi C, et al. Relationships between traumatic symptoms and environmental damage conditions among children 8 months after the 2011 Japan earthquake and tsunami. *PLoS One.* 2012;7(11):e50721. doi:10.1371/journal.pone.0050721.
37. Usami M, Iwadare Y, Watanabe K, Kodaira M, Ushijima H, Tanaka T, et al. Analysis of changes in traumatic symptoms and daily life activity of children affected by the 2011 Japan earthquake and tsunami over time. *PLoS One.* 2014;9(2):e88885. doi:10.1371/journal.pone.0088885.
38. Kukihara H, Yamawaki N, Uchiyama K, Arai S, Horikawa E. Trauma, depression, and resilience of earthquake/tsunami/nuclear disaster survivors of Hirono, Fukushima, Japan. *Psychiatry Clin Neurosci.* 2014;68(7):524–33. doi:10.1111/pcn.12159.
39. Matsumoto J, Kunii Y, Wada A, Mashiko H, Yabe H, Niwa S. Mental disorders that exacerbated due to the Fukushima disaster, a complex radioactive contamination disaster. *Psychiatry Clin Neurosci.* 2014;68(3):182–7.
40. Rubin GJ, Amlot R, Wessely S, Greenberg N. Anxiety, distress and anger among British nationals in Japan following the Fukushima nuclear accident. *Br J Psychiatry J Mental Sci.* 2012;201(5):400–7. doi:10.1192/bjp.bp.112.111575.
41. Sawa M, Osaki Y, Koishikawa H. Delayed recovery of caregivers from social dysfunction and psychological distress after the Great East Japan Earthquake. *J Affect Disord.* 2013;148(2–3):413–7. doi:10.1016/j.jad.2012.11.011.
42. Sawa M, Takase M, Noju K, Tomiyasu T, Kawakami C, Koishikawa H, et al. Impact of the great East Japan earthquake on caregiver burden: a cross-sectional study. *Psychiatr Ser (Washington, DC).* 2013;64(2):189–91. doi:10.1176/appi.ps.000802012.
43. Sugimoto T, Shinozaki T, Naruse T, Miyamoto Y. Who was concerned about radiation, food safety, and natural disasters after the great East Japan earthquake and Fukushima catastrophe? A nationwide cross-sectional survey in 2012. *PLoS ONE.* 2014;9(9):e106377. doi:10.1371/journal.pone.0106377.
44. Tsubokura M, Hara K, Matsumura T, Sugimoto A, Nomura S, Hinata M, et al. The immediate physical and mental health crisis in residents proximal to the evacuation zone after Japan's nuclear disaster: an observational pilot study. *Disaster Med Public Health Prep.* 2014;8(1):30–6. doi:10.1017/dmp.2014.5.
45. Wada A, Kunii Y, Matsumoto J, Itagaki S, Yabe H, Mashiko H, et al. Changes in the condition of psychiatric inpatients after the complex Fukushima disaster. *Fukushima J Med Sci.* 2013;59(1):39–42.
46. Yabe H, Suzuki Y, Mashiko H, Nakayama Y, Hisata M, Niwa S, et al. Psychological distress after the Great East Japan Earthquake and Fukushima Daiichi Nuclear Power Plant accident: results of a mental health and lifestyle survey through the Fukushima Health Management Survey in FY2011 and FY2012. *Fukushima J Med Sci.* 2014;60(1):57–67.
47. Goto A, Rudd RE, Lai AY, Yoshida K, Suzuki Y, Halstead DD, et al. Leveraging public health nurses for disaster risk communication in Fukushima City: a qualitative analysis of nurses' written records of parenting counseling and peer discussions. *Bmc Health Ser Res.* 2014;14:129. doi:10.1186/1472-6963-14-129.

48. Ben-Ezra M, Shigemura J, Palgi Y, Hamama-Raz Y, Lavenda O, Suzuki M, et al. From Hiroshima to Fukushima: PTSD symptoms and radiation stigma across regions in Japan. *J Psychiatr Res*. 2014;60C:185–6. doi:10.1016/j.jpsychires.2014.10.006.
49. Yoshii H, Saito H, Kikuchi S, Ueno T, Sato K. Report on maternal anxiety 16 months after the great East Japan earthquake disaster: anxiety over radioactivity. *Glob J Health Sci*. 2014;6(6):36862. doi:10.5539/gjhs.v6n6p1.
50. Dobashi K, Nagamine M, Shigemura J, Tsunoda T, Shimizu K, Yoshino A, et al. Psychological effects of disaster relief activities on Japan ground self-defense force personnel following the 2011 great east Japan earthquake. *Psychiatry*. 2014;77(2):190–8. doi:10.1521/psyc.2014.77.2.190.
51. Fukasawa M, Suzuki Y, Obara A, Kim Y. Relationships between mental health distress and work-related factors among prefectural public servants 2 months after the great east Japan earthquake. *Int J Behav Med*. 2014; doi:10.1007/s12529-014-9392-8.
52. Matsuoka Y, Nishi D, Nakaya N, Sone T, Noguchi H, Hamazaki K, et al. Concern over radiation exposure and psychological distress among rescue workers following the Great East Japan Earthquake. *BMC Public Health*. 2012;12:249. doi:10.1186/1471-2458-12-249.
53. Nishi D, Koido Y, Nakaya N, Sone T, Noguchi H, Hamazaki K, et al. Fish oil for attenuating posttraumatic stress symptoms among rescue workers after the great east Japan earthquake: a randomized controlled trial. *Psychother Psychosom*. 2012;81(5):315–7. doi:10.1159/000336811.
54. Nishi D, Koido Y, Nakaya N, Sone T, Noguchi H, Hamazaki K, et al. Peritraumatic distress, watching television, and posttraumatic stress symptoms among rescue workers after the Great East Japan earthquake. *PLoS One*. 2012;7(4):e35248. doi:10.1371/journal.pone.0035248.
55. Shigemura J, Tanigawa T, Nishi D, Matsuoka Y, Nomura S, Yoshino A. Associations between disaster exposures, peritraumatic distress, and post-traumatic stress responses in Fukushima nuclear plant workers following the 2011 nuclear accident: the Fukushima NEWS Project study. *PLoS One*. 2014;9(2):e87516. doi:10.1371/journal.pone.0087516.
56. Shigemura J, Tanigawa T, Saito I, Nomura S. Psychological distress in workers at the Fukushima nuclear power plants. *JAMA*. 2012;308(7):667–9. doi:10.1001/jama.2012.9699.
57. Suzuki Y, Fukasawa M, Obara A, Kim Y. Mental health distress and related factors among prefectural public servants 7 months after the great East Japan Earthquake. *J Epidemiol Jpn Epidemiol Assoc*. 2014;24(4):287–94.
58. Tsutsui T, Hasegawa Y, Hiraga M, Ishiki M, Asukai N. Distinctiveness of prolonged grief disorder symptoms among survivors of the Great East Japan Earthquake and Tsunami. *Psychiatry Res*. 2014;217(1–2):67–71. doi:10.1016/j.psychres.2014.03.001.
59. Japan Disaster Grief Support. Japan Disaster Grief Support (JDGS) Project. 2011. <http://jdgs.jp/>. Accessed Jan 5 2015 (in Japanese).
60. Heir T, Weisaeth L. Acute disaster exposure and mental health complaints of Norwegian tsunami survivors six months post disaster. *Psychiatry*. 2008;71(3):266–76. doi:10.1521/psyc.2008.71.3.266.
61. Goto A, Rudd RE, Bromet EJ, Suzuki Y, Yoshida K, Suzuki Y, et al. Maternal confidence of Fukushima mothers before and after the nuclear power plant disaster in Northeast Japan: analyses of municipal health records. *J Commun Healthcare*. 2014;7(2):106–16. doi:10.1179/1753807614Y.0000000051.
62. 2003–2005 CF. Chernobyl's legacy: health, environmental and socio-economic impacts. Vienna: International Atomic Energy Agency; 2006.
63. Adams RE, Guey LT, Gluzman SF, Bromet EJ. Psychological well-being and risk perceptions of mothers in Kyiv, Ukraine, 19 years after the Chernobyl disaster. *Int J Soc Psychiatry*. 2011;57(6):637–45. doi:10.1177/0020764011415204.
64. Havenaar JM, Rumyantzeva GM, van den Brink W, Poelijoe NW, van den Bout J, van Engeland H, et al. Long-term mental health effects of the Chernobyl disaster: an epidemiologic survey in two former Soviet regions. *Am J Psychiatry*. 1997;154(11):1605–7.
65. Loganovsky K, Havenaar JM, Tintle NL, Guey LT, Kotov R, Bromet EJ. The mental health of clean-up workers 18 years after the Chernobyl accident. *Psychol Med*. 2008;38(4):481–8. doi:10.1017/S0033291707002371.
66. Maeda M, Oe M. Disaster behavioral health: psychological effects of the Fukushima nuclear power plant accident. In: Tanigawa K, Chhem RK, editors. *Radiation disaster medicine*. Heidelberg: Springer; 2014. p. 79–88.
67. Ursano RJ. Post-traumatic stress disorder. *N Engl J Med*. 2002;346(2):130–2. doi:10.1056/NEJM200201103460213.
68. McCarroll JE, Ursano RJ, Fullerton CS, Liu X, Lundy A. Somatic symptoms in Gulf War mortality workers. *Psychosom Med*. 2002;64(1):29–33.
69. Ursano RJ, Fullerton CS, Vance K, Kao TC. Posttraumatic stress disorder and identification in disaster workers. *Am J Psychiatry*. 1999;156(3):353–9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

