



RESEARCH ARTICLE

CaMKII binding to GluN2B is important for massed spatial learning in the Morris water maze [v1; ref status: indexed, <http://f1000r.es/3ud>]

Ivar S. Stein, Michaela S. Donaldson, Johannes W. Hell

Department of Pharmacology, School of Medicine, University of California, Davis, 95616-8636, USA

v1 First published: 12 Aug 2014, 3:193 (doi: [10.12688/f1000research.4660.1](https://doi.org/10.12688/f1000research.4660.1))
 Latest published: 12 Aug 2014, 3:193 (doi: [10.12688/f1000research.4660.1](https://doi.org/10.12688/f1000research.4660.1))

Abstract

Learning and memory as well as long-term potentiation (LTP) depend on Ca²⁺ influx through the NMDA-type glutamate receptor (NMDAR) and the resulting activation of the Ca²⁺ and calmodulin-dependent protein kinase (CaMKII). Ca²⁺ influx via the NMDAR triggers CaMKII binding to the NMDAR for enhanced CaMKII accumulation at post-synaptic sites that experience heightened activity as occurring during LTP. Previously, we generated knock-in (KI) mice in which we replaced two residues in the NMDAR GluN2B subunit to impair CaMKII binding to GluN2B. Various forms of LTP at the Schaffer collateral synapses in CA1 are reduced by 50%. Nevertheless, working memory in the win-shift 8 arm maze and learning of the Morris water maze (MWM) task was normal in the KI mice although recall of the task was impaired in these mice during the period of early memory consolidation. We now show that massed training in the MWM task within a single day resulted in impaired learning. However, learning and recall of the Barnes maze task and contextual fear conditioning over one or multiple days were surprisingly unaffected. The differences observed in the MWM compared to the Barnes maze and contextual fear conditioning suggest a differential involvement of CaMKII and the specific interaction with GluN2B, probably depending on varying degrees of stress, cognitive demand or even potentially different plasticity mechanisms associated with the diverse tasks.

Open Peer Review**Invited Referee Responses**

	1	2
version 1 published 12 Aug 2014	 report	 report

1 Mark L. Dell'Acqua, University of Colorado Denver USA

2 Karl-Ulrich Bayer, University of Colorado Denver USA

Latest Comments

No Comments Yet

Corresponding author: Johannes W. Hell (jwhell@ucdavis.edu)

How to cite this article: Stein IS, Donaldson MS and Hell JW. **CaMKII binding to GluN2B is important for massed spatial learning in the Morris water maze [v1; ref status: indexed, <http://f1000r.es/3ud>]** *F1000Research* 2014, 3:193 (doi: [10.12688/f1000research.4660.1](https://doi.org/10.12688/f1000research.4660.1))

Copyright: © 2014 Stein IS *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](https://creativecommons.org/licenses/by/4.0/) (CC0 1.0 Public domain dedication).

Grant information: This work was supported by NIH grants NS046450 and NS078792 to JWH.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: No competing interests were disclosed.

First published: 12 Aug 2014, 3:193 (doi: [10.12688/f1000research.4660.1](https://doi.org/10.12688/f1000research.4660.1))

First indexed: 26 Aug 2014, 3:193 (doi: [10.12688/f1000research.4660.1](https://doi.org/10.12688/f1000research.4660.1))

Introduction

The acquisition and storage of new tasks as well as the modification of already existing memories depend on the selective strengthening and weakening of synaptic interactions embedded in extensive networks of neurons (Kessels & Malinow, 2009; Martin *et al.*, 2000; Morris, 2013; Neves *et al.*, 2008). Synaptic plasticity and especially long-term potentiation (LTP), a stable increase in synaptic strength, are emerging as the cellular equivalent of learning and memory (Gruart *et al.*, 2006; Whitlock *et al.*, 2006). Two key players involved in LTP are the Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) and the NMDA-type glutamate receptor (NMDAR). Ca²⁺ influx through the NMDAR leads to CaMKII activation and recruitment to the postsynaptic density (PSD), which can be persistent and synapse specific (Otmakhov *et al.*, 2004; Zhang *et al.*, 2008) (but see also (Rose *et al.*, 2009)). CaMKII binding to aa 1290–1309 on the NMDAR GluN2B subunit is required for this activity-dependent translocation (Halt *et al.*, 2012; Leonard *et al.*, 1999; Strack & Colbran, 1998; Strack *et al.*, 2000) and is crucial for LTP (Barria & Malinow, 2005; Halt *et al.*, 2012; Zhou *et al.*, 2007). Stimulation of CaMKII results in its auto-phosphorylation on T286 causing a persistent Ca²⁺-independent activation of CaMKII (Lisman *et al.*, 2002). This auto-phosphorylation is increased after LTP and spatial learning in the Morris water maze (MWM) (Lengyel *et al.*, 2004; Ouyang *et al.*, 1997; Ouyang *et al.*, 1999; Tan & Liang, 1996) and is required for effective binding of CaMKII to GluN2B (Bayer *et al.*, 2001; Strack & Colbran, 1998). T286A CaMKII mutant mice exhibit no hippocampal NMDAR dependent LTP and have impaired MWM learning (Giese *et al.*, 1998).

We recently reported the specific disruption of CaMKII binding to GluN2B by point mutations in the GluN2B gene (Halt *et al.*, 2012). The GluN2BKI mice contain two point mutations in the GluN2B C terminus (L1298A and R1300Q). Each of these mutations abolishes CaMKII binding to this site nearly completely *in vitro* (Strack *et al.*, 2000). *In vivo* the two point mutations entirely abrogate the activity-dependent increase in the interaction with the NMDAR, show reduced hippocampal LTP by 50% and result in a MWM memory deficit, while acquisition of the MWM task and working memory as evaluated in the 8 arm win shift test remain normal (Halt *et al.*, 2012).

Now we report a specific moderate spatial learning deficit in the GluN2B knock in (KI) mice during massed 1-day training in the MWM. At the same time learning and memory recall in the Barnes maze and contextual fear conditioning are unaffected.

Material and methods

Mice

All animal procedures were approved (Protocol #: 15512; PHS/NIH Assurance A3433-01) by the UC Davis Institutional Animal Care and Use Committee (IACUC) and followed NIH guidelines. All experiments were conducted with litter matched wild type (WT) and GluN2B KI mice of mixed sex (evenly distributed gender ratios) and between 2 and 4 months of age; mice within a cohort were not more than 3 weeks apart. For a more detailed characterization and description of the GluN2B KI mice (C57BL/6J background) including genotyping procedures refer to (Halt *et al.*, 2012). The mice were on a 12 h light/dark cycle, housed individually for the behavioral experiments and acclimated for at least one week before all

procedures. After the acclimation period and before the training and testing the mice were extensively handled (7–10 times for 1–2 min on different days). All trainings and testing were performed during the 12 h light cycle.

Morris water maze

The MWM was a circular, enamel coated steel tank, 94 cm in diameter filled with water at 22–24°C. For the visible trial, a black labeled square platform (6 × 6 cm) emerging 2 cm out of the opaque (through addition of non-toxic paint) water surface was used, while the clear Plexiglas square platform (6 × 6 cm) for the training trials was submerged 2 cm below the opaque water surface. On the first day, the mice were trained in one visible trial followed by 12 consecutive training trials, while the platform was kept in a fixed position. The mice were always placed facing the wall of the pool and started randomly from three different starting positions that are equally distributed around the perimeter and are not located within the target quadrant. The mice were allowed to swim freely for 90 s to find the platform. If they failed to locate the platform within time, they were gently guided to it. After the mice climbed on the platform they were allowed to remain 30 s on the platform before they were removed from the pool and placed in their home cage for an inter-trial interval of 6–10 min. The mice were returned to the pool for a 90 s probe test without the platform 1 day and 7 days after the training day. The training session and probe trials were monitored and analyzed using the SMART (Version 2.5.19) real-time video-tracking system.

Barnes maze

The Barnes maze was executed according to (Berta *et al.*, 2007). In short, the maze consisted of a circular platform (92 cm diameter; opaque white) with 20 equally spaced holes (5 cm diameter; 7.5 cm between holes) and was elevated 105 cm above the floor. All holes were 2 cm away from the perimeter of the platform and a dark escape box (10 cm × 12.5 cm × 12.5 cm) was located beneath one of the holes.

The mouse was placed in a cylindrical black start chamber (10 cm × 12.5 cm × 12.5 cm) in the middle of the maze. After 10 s the chamber was lifted and the mouse was exposed to bright light (500 W). For the habituation trial on day 1 the animal was gently guided to the escape box and once it entered, the entry hole was covered and the mouse kept for 2 min in the dark escape box. Between each mouse, before the next trial, the platform was cleaned with a 10% Nolvasan solution to avoid remaining olfactory cues. During the acquisition phase, after the start chamber was lifted, the mouse was allowed to explore the maze for 3 min. The trial ended when the mouse entered the escape box or after 3 min had elapsed. Immediately after the mouse entered the escape box, the entry hole was covered and the animal stayed for 1 min in the dark escape box. If the mouse did not reach the target hole/escape box within 3 min, it was gently guided to it. After 1 min in the escape box, the mouse was placed back in its home cage until the next trial. Each mouse received 4 training trials per day with an inter-trial interval of 10–15 min on 4 consecutive days.

The probe trial was conducted on day 5, i.e., 24 h after the last training day, and on day 12. During the probe test the escape box was removed and the mouse was allowed to explore the maze for a fixed

time of 90 s. The number of pokes (errors) in each hole and the latency and path length to reach the virtual target hole was measured using the SMART (Version 2.5.19) real-time video-tracking system.

Elevated plus maze

The elevated plus maze (EPM) was a four arm maze with each arm measuring 30 × 5 cm and the central platform measured 5 × 5 cm (opaque white). One set of arms, opposing one another, were enclosed completely by grey side walls, 15 cm high, while the other set was open with a ledge of 0.5 cm on either side of the arms. The maze was elevated 100 cm from the floor and illuminated evenly with a 500W light. Mice were placed on the central platform, facing towards a closed arm, and allowed to freely explore the maze for 5 min. The SMART (Version 2.5.19) real-time tracking software was used to record the locomotor activity and the time spent on both the closed and the open arms during the test.

Fear conditioning

Contextual fear conditioning was conducted using the 'Video Tracking of Fear Conditioning System' and the 'Video Freeze Software' from MED Associates Inc. The experimentally naïve mice were placed in the conditioning chamber and received the first foot shock at the end of a 3 min habituation period. The mice remained for an additional minute in the chamber after the last foot shock was delivered. For the 5 shock conditioning protocol, the mice received an electric foot shock (0.75 mA, 1 s) at the end of the 3rd, 4th, 5th, 6th, and 7th minute. During the milder 3 shock (0.5 mA, 1 s) paradigm, the shock was received at the end of the 3rd, 4th and 5th minute. For the 4 day training protocol only one shock (0.75 mA, 1 s) per day was delivered after 2.5 min. In case of the milder 3 shock paradigm and the 4 day training protocol the mice were pre-exposed to the conditioning chamber for 3 min and 10 min respectively. For recall the mice were placed back for 5 min in the same chamber after the indicated periods of time. If the mice were exposed to a different context, the chamber geometry was changed (from square to curved wall), the lights were dimmed, the rod-flooring was covered with white linoleum, and the scent was changed from 0.01% bleach to bubble gum. Using the Video Freeze Program, the freezing time was determined (in %) and was defined as the absence of any movement, except for respiratory motion. The observed freezing response to the foot shock is correlated to the degree of learning, the strength of the aversive stimulus, and the number of presentations (Curzon *et al.*, 2009).

Statistical analysis

All data are represented as mean ± the standard error of the mean (SEM). The data were analyzed by unpaired two-tailed t-test, one-way ANOVA or two-way ANOVA followed by the Bonferroni correction as indicated in the results, using GraphPad Prism 5 software. Statistical significance was considered if $p \leq 0.05$.

Results

Single day spatial learning in the MWM is impaired in GluN2B KI

In order to further dissect the memory deficit observed during the initial MWM experiments (Halt *et al.*, 2012), two independent cohorts of GluN2B KI mice and their WT littermate controls (10 mice of

each genotype per cohort) underwent a massed training protocol on a single day in the MWM. Such a compressed training protocol challenges the learning capacity within a short time period. It also has the potential to reveal deficits in early- versus late-phase consolidation as training is not protracted over 6 days as is the case for the classic MWM. The 1 day protocol thus might expose deficits in early consolidation when testing different cohorts 1 and 7 days after the training day without the complication that in the 6 days training paradigm training and consolidation processes overlap. After habituation on the first day (training day), mice underwent one visible platform trial followed by 12 consecutive training trials, in which the platform was no longer visible but kept in a fixed position. The initial trial with a marked visible platform is hippocampus-independent and was conducted to rule out changes in motivation, coordination, or sensory processing in GluN2B KI mice. The latency in reaching the visible platform was not different between the two genotypes (Figure 1A; WT: 51.48 ± 7.43 s, KI: 46.46 ± 6.70 s). In the following 12 consecutive trials with a submerged platform the KI mice slowly started to show deficits in spatial learning around trial 6, roughly the onset of learning in the WT mice (Figure 1B). The latencies of the 12 training trials are paired into blocks of two for analysis and the GluN2B KI mice display significantly increased average escape latencies during the last two trials (genotype: F1,228=2.63, $p=0.1061$; trial: F5,228=13.30, $p<0.0001$; genotype × trial F5,228=3.08, $p=0.0103$; Bonferroni post hoc test shows that the latency is significantly increased in the KI mice during the last two trials, $M_{diff}=20.30$ s, 95% CI [2.58, 38.01], $p<0.05$; Figure 1B; WT₁₁₋₁₂: 30.31 ± 3.77 s, KI₁₁₋₁₂: 50.61 ± 4.49 s).

The covered distance shows like the escape latency a significant *genotype × trial* interaction (Figure 1C; genotype: F1,228=0.16, $p=0.687$; trial: F5,228=43.34, $p<0.0001$; genotype × trial F5,228=3.13, $p=0.0095$), strengthening the mild learning deficit observed during the last training trials. No difference was observed between the mean velocity of the WT and GluN2B KI mice (Figure 1D).

During the probe test one day after training, WT mice required significantly less time to reach the original location of the platform as reflected by faster primary escape latency (Figure 1E; WT: 38.09 ± 6.15 s, KI: 55.32 ± 6.91 s; $t(38)=2.15$, $p<0.05$). Analysis of the entries into the quadrants and the area of the former platform location also showed a significant difference between the two genotypes in their preference for the target quadrant (Figure 1F). The KI mice showed compared to WT no real preference for the target quadrant D and entered it significantly less often (Figure 1F; WT_D: 9.45 ± 0.98, KI_D: 6.70 ± 1.22; genotype: F1,190=3.89, $p<0.05$; quadrant/platform: F4,190=18.99, $p<0.0001$; genotype × quadrant/platform F4,190=1.46, $p=0.2147$; Bonferroni post hoc test shows that the WT mice search preferentially in the target quadrant compared to the KI mice, $M_{diff}=-2.750$ entries, 95% CI [-5.347, -0.153], $p<0.05$). The covered distance and mean velocity was not different between the two genotypes (Figure 1G), indicating similar activity levels.

During the probe test 7 days after training some of the mice (independent of their genotype) were continuously floating and did not search actively for the escape platform. For the analysis of the 7 day probe trial, these floaters were neglected. The performance in the




Figure 1. Spatial learning in the Morris water maze (MWM) is impaired in GluN2B KI mice. (A) WT and KI mice display no difference in the average escape latency during the visible platform trial. (B) The GluN2B KI mice show a deficit during acquisition of the hidden platform task. The latencies of the 12 training trials are split up into blocks of two and the average escape latency is significantly increased for KI versus WT mice during the last two training trials. (C, D) The covered distance over the blocks of two training trials is also different between KI and WT mice (C), while the mean velocity is not changed (D). (E, F) During the probe test 1 day after training the KI mice exhibit increased primary average escape latency (E) and show significantly fewer crossings into the target quadrant D (F), the former location of the submerged platform, compared to WT mice. (G) Both genotypes cover the same distance during the probe trial and display the same mean velocity. (H, I) The KI mice show a similar tendency during the probe test 7 days after training, with extended primary average escape latency (H) and diminished crossings into target quadrant D (I). (J) The covered distance and mean velocity is again unchanged. (A: adjacent right quadrant to D; B: quadrant opposite to quadrant D; C: adjacent left quadrant to D; platform: area of the former location of the platform). The same two independent cohorts, each consisting of 10 WT and 10 GluN2B KI mice, were tested in the Barnes maze before the MWM. Data represent Mean \pm SEM.

probe test 7 days after training showed a similar tendency compared to the one day test. The GluN2B KI leaned towards a longer primary escape latency (Figure 1H, left; WT: 29.50 ± 8.59 s, KI: 48.50 ± 5.98 s; $t(18)=1.87$, $p=0.078$) and tended to cross less often over into the target quadrant in search for the platform than WT mice (Figure 1I; WT_D: 9.67 ± 1.49 , KI_D: 6.46 ± 0.73 ; genotype: $F_{1,90}=3.12$, $p=0.0805$; quadrant/platform: $F_{4,90}=16.69$, $p<0.0001$; genotype \times quadrant/platform $F_{4,90}=1.85$, $p=0.1259$). Both genotypes again showed no observable difference in the covered distance and mean velocity (Figure 1J).

Individual analysis (one-way-ANOVA) of the search pattern (quadrant crossings) reveals that the WT mice preferentially search in the target quadrant during the 1 day ($F_{3,76}=11.61$, $p<0.0001$) as well as the 7 day probe trial ($F_{3,32}=3.91$, $p=0.0175$). The GluN2B KI mice conversely only show a slight non-significant preference for the target quadrant during the 1 day probe trial and no preference at all during the 7 day probe trial.

Spatial learning and memory is not affected in the Barnes maze

The Barnes maze is a less aversive spatial learning paradigm, which unlike the MWM allows the mice to move freely and explore the environment under conditions that are less stressful. It is solely based on the motivation of the mice to avoid prolonged exposure to an open area under light and to search for a dark and safe hideout, the escape box. Interestingly, the GluN2B KI showed no detectable deficits not only in learning but also in early memory (Figure 2). These findings contrast what had been seen before after spaced MWM learning experiments conducted over a period of 6 days (Halt *et al.*, 2012). Over the 16 training trials, which were executed on 4 consecutive days (4 trials per day) the latency to locate and enter the escape box decreased equally fast for both genotypes and saturated on the fourth day (Figure 2A; WT: 22.99 ± 4.08 s, KI: 20.95 ± 3.95 s). During the probe test 1 day after the last training session there was no difference in the primary escape latency (Figure 2B, left; WT: 20.11 ± 6.02 s, GluN2BKI: 18.68 ± 4.72 s), covered distance (Figure 2B, middle; WT: 355.70 ± 37.25 cm, GluN2BKI: 426.60 ± 43.64 cm) or mean velocity (Figure 2B, right; WT: 3.95 ± 0.41 cm/s, GluN2BKI: 4.74 ± 0.48 cm/s). Both WT and KI mice searched preferentially in the target area and the surrounding holes (areas +1 and -1), indicating that they clearly remembered the former location of the escape box (Figure 2B, lower part). The overall number of zone entries and time in zones for GluN2B KI mice showed a slight increase compared to WT mice, which is probably a reflection of the statistically non-significant increased activity of KI as seen in the covered distance and mean velocity (Figure 2B). The performance in the probe test 7 days after the last training session mirrored the results of the 1 day probe test. There was no significant difference in the primary escape latency (Figure 2C, left; WT: 33.37 ± 7.98 s, KI: 26.80 ± 6.61 s), covered distance (Figure 2C, middle; WT: 265.40 ± 26.47 cm, GluN2BKI: 353.00 ± 49.42 cm) and mean velocity (Figure 2C, right; WT: 2.95 ± 0.29 cm/s, GluN2BKI: 3.92 ± 0.55 cm/s). The slightly increased activity of the GluN2B KI mice is again reflected by an increased number of zone entries and by the time spent in the zones (Figure 2C, bottom).

Basal anxiety levels in the GluN2B KI mice are normal

Basal anxiety levels in the GluN2B KI mice were assessed using the EPM. The EPM relies on the rodents' innate fear of heights and open spaces and their preference for dark and enclosed rooms. Basal anxiety levels in the GluN2B KI mice were not different compared to their WT litter-matched controls. During exposure to the EPM the GluN2B KI mice spent most of the time in the closed arms, like the WT controls, and rarely stayed in the open arms or center (Figure 3A; in % of total time, WT: $92.30 \pm 1.16\%$, KI: $88.68 \pm 3.27\%$). Both genotypes covered comparable minimal distances in the open arms (Figure 3B; WT: 20.63 ± 3.96 cm, GluN2B KI: 31.62 ± 10.73 cm), while the WT mice had the tendency to be more active in the closed arms. This tendency is displayed in an increase in covered distance (Figure 3B; WT: 881.59 ± 86.40 cm, KI: 626.92 ± 59.24 cm; genotype: $F_{1,68}=10.17$, $p=0.0022$; location: $F_{3,68}=120.21$, $p<0.0001$; genotype \times location $F_{3,68}=3.40$, $p=0.0225$; Bonferroni post hoc test shows that the distance is significantly reduced in the KI in the closed arms, $M_{diff}=-254.7$ cm, 95% CI [-459.5, -49.49], $p<0.01$) and a trend towards a higher mean velocity in the closed arms (Figure 3C; WT: 3.19 ± 0.31 cm/s, GluN2B KI: 2.37 ± 0.21 cm/s), also apparent in the overall mean velocity (Figure 3C; WT: 3.34 ± 0.29 cm/s, GluN2B KI: 2.66 ± 0.18 cm/s). The observed normal basal anxiety levels concur with earlier findings in the open field analysis and the same innate fear reaction to trimethyl-thiazoline (TMT) (Halt *et al.*, 2012).

Learning and memory in contextual fear conditioning is not affected in GluN2B KI mice

With basal anxiety levels not being affected in the GluN2B KI mice (Halt *et al.*, 2012) (Figure 3), we investigated the role of the CaMKII/GluN2B interaction in contextual fear conditioning. The experimentally naïve mice were placed in the conditioning chamber and after a 3 min habituation period trained with five consecutive foot shocks (0.75 mA, 1 s duration; aversive unconditioned stimulus). The GluN2B KI mice learned to the same extent and at the same speed as their WT litter-matched controls (Figure 4A, C). A total of four independent cohorts were trained. To be able to distinguish possible differences between memory recall and consolidation, two cohorts were tested after 4 days (Figure 4B) and the other two cohorts after 14 days (Figure 4D). For the probe tests (after 4 or 14 days) the mice were placed back for 5 min into the conditioning chamber and the freezing response to the context was measured. There was no difference in the average time WT and KI mice spent freezing during either the 4 (WT: $51.11 \pm 3.37\%$, KI: $52.23 \pm 3.77\%$) or 14 days (WT: $70.48 \pm 4.75\%$, KI: $67.51 \pm 3.33\%$) probe test.

Contextual fear conditioning performance is independent of the stimulus strength

CaMKII T286A mutant mice have impaired contextual short-term memory (STM) and long-term memory (LTM) formation after a single or three tone-shock pairings, while contextual STM and LTM formation after five pairings are unaffected (Irvine *et al.*, 2011; Irvine *et al.*, 2005). To exclude the possibility that the GluN2B KI mice were over trained with five consecutive foot shocks (0.75 mA, 1 s duration) and potential cognitive deficits masked, we also tested two milder contextual fear conditioning paradigms.




Figure 2. Spatial learning and memory is unaffected in the Barnes maze. (A) The escape latency for both genotypes decreases equally over the 16 training trials and saturates on the fourth day (left panel depicts average latencies over each of the 4 training days whereas right panel shows averages for each individual trial). (B) The primary escape latency during the probe test 1 day after the last training session is not different. KI mice search like the WT mice preferentially in and around the target area for the escape box (bottom panels). The covered distance (middle) and mean velocity (right) is slightly, but not significantly increased in KI mice. (C) The probe test 7 days after the last training session shows, like the 1 day test, no difference in the avg. primary escape latency (left) and the KI mice again exhibited the tendency to increased covered distance (middle) and mean velocity (right). Both genotypes searched in the right location in and around the target area (bottom panels). The same two independent cohorts, each consisting of 10 WT and 10 GluN2B KI mice, were tested in the Barnes maze followed by the MWM. Data are presented as Mean \pm SEM.




Figure 3. Basal anxiety levels in the GluN2B KI mice are normal. (A) The % of time the mice spent in the open and closed arms as well as the center is not different between WT and KI. (B) Both genotypes cover the same minimal distances in the open arms but WT mice cover greater distances than KI mice in the closed arms. (C) The overall mean velocity is not different between both genotypes, while the WT mice show a slightly increased mean velocity in the closed arm coinciding with the longer covered distance. One cohort of 10 WT and 9 GluN2B KI mice was tested. The same cohort was subsequently used for the milder three shock fear conditioning paradigm (Figure 5A-E). Data represent Mean \pm SEM.




Figure 4. Contextual fear conditioning and memory is unaffected in GluN2B KI mice. Fear conditioning was evaluated 4 days after training of two different cohorts of naïve litter-matched WT, heterozygous, and GluN2B KI mice (12–14 WT and KI mice, and 3–4 heterozygous mice per cohort) and 14 days after training with yet another two different cohorts of naïve litter-matched WT, heterozygous, and GluN2B KI mice (10 WT mice, 10–12 KI mice, and 5–7 heterozygous mice per cohort). (A, C) After the 3 min habituation phase, 5 shocks (0.75 mA, 1 s) were delivered at the end of 3rd, 4th, 5th, 6th and 7th minute (asterisks). The increase in the fraction of time spent freezing during conditioning is not different between the GluN2B KI, heterozygous, and WT mice. (B, D) All genotypes also display a comparable fraction of time freezing during the 4 day and the 14 day probe test. Data represent percentage freezing (Mean \pm SEM).

During the first paradigm, after two days of 3 min pre exposure to the conditioning chamber, 3 shocks (0.5 mA, 1 s) were delivered on day 3 at the end of the 3rd, 4th and 5th minute. Conditioning and memory tested after 4 days (WT: 53.82 ± 3.55%, KI: 53.55 ± 6.10%) were unaffected (Figure 5A, B). When the mice were exposed to a different context in the same chamber and room 1 day later, the time spent freezing was reduced, similar to pre-conditioning levels, and did not differ between the two genotypes (Figure 5C; WT: 13.33 ± 3.20%, KI: 18.90 ± 2.55%). Long term memory was tested with the same cohort 11 days after training (Figure 5D) and like the 4 day test showed no difference (WT: 47.63 ± 4.60%, KI: 53.08 ± 6.60%). Figure 5E depicts an overview of the experiment. Figure 5E depicts an overview of the experiment.

In addition, another cohort of GluN2B KI and WT mice were trained over multiple days. The mice were pre exposed to the context on day 1 for 10 min and trained on the following 4 days with one foot shock (each 0.75 mA, 1 s) per day (Figure 5F). Again no learning or memory deficit, which was tested 6 (WT: 75.52 ± 4.64%, GluN2B KI: 62.69 ± 6.21) and 8 days (WT: 64.75 ± 5.68%, KI: 59.06 ± 7.56%) after training, was observed. When the mice were exposed

to a different context in the same chamber and room on day 12, 7 days after the last training, both genotypes exhibited reduced freezing levels (WT: 25.56 ± 5.85%, KI: 22.09 ± 6.93%) similar to those observed before conditioning.

Dataset 1. Data sets for behavioral tasks in wild type and GluN2BKI mice

<http://dx.doi.org/10.5256/f1000research.4660.d34165>

The raw data tables show the performances of the individual wild type and GluN2B KI mice in the various behavioral tasks, which are represented in the text and figures as mean ± the standard error of the mean (SEM). Data sets for 'MWM data', 'Barnes maze data', 'Elevated plus maze data set' and 'Contextual fear conditioning data set' are shown. Specific information can be found in text file.

Discussion

CaMKII and the NMDAR are crucial not only for hippocampal LTP but also for hippocampus dependent spatial learning and memory formation. In a number of studies CaMKII α KO and mutant mice




Figure 5. Contextual fear conditioning with milder conditioning is unaffected in GluN2B KI mice. One cohort of litter-matched GluN2B KI and WT mice (9–10 mice per genotype, previously tested in the EPM) was fear conditioned with three shocks on a single day and a different cohort of naïve litter-matched GluN2B KI and WT mice (10 mice per genotype) was fear conditioned with four shocks over a period of 4 days. (A) KI and WT mice acquired to the same extent fear conditioning as indicated by the time spent freezing after three consecutive mild foot shocks (0.5 mA, 1 s) at the end of the 3rd, 4th and 5th minute (asterisks). (B, D) The time spent freezing during recall of context after 4 d and 11 d is not different between genotypes. (C) Exposure to a different context 5 days after training shows pre-conditioning freezing levels for both genotypes. (E) Summary plot of the experimental data from A-D. (F) Summary plot of the experimental data of the 4 day training (1 shock, 0.75 mA, 1 s per day) experiment. WT and KI mice show comparable learning curves during the 4 d conditioning and similar contextual memory when tested 6 and 8 days later. Exposure to a different context in the same room and chamber, 7 d after the last training day, revealed again pre-conditioning freezing levels. Data represent percentage freezing (Mean ± SEM).

displayed learning deficits, notably mainly in aversively motivated tasks (Elgersma *et al.*, 2002; Giese *et al.*, 1998; Irvine *et al.*, 2005; Silva *et al.*, 1992), while appetitive learning was unaffected (Carvalho *et al.*, 2001). Moreover, the regulation of expression or activity level of CaMKII α seems to have an emotional component and especially impact emotional and anxiety-like behavior and learning. CaMKII α heterozygous KO mice show a decreased freezing in fear conditioning induced by an electric foot shock and are more active in the open field suggesting decreased anxiety-related behaviors (Chen *et al.*, 1994). Consistently, transgenic mice overexpressing CaMKII α exhibit an increase in anxiety-like behaviors in open field, light-dark transition, and elevated zero maze (Hasegawa *et al.*, 2009). CaMKII α T286A mutant mice, which are deficient in autonomous and thereby overall CaMKII activity, spend more time in and entered more often the open arm in the EPM and generally react with hyperactivity to novel stimuli that could be perceived as potentially threatening (Easton *et al.*, 2011). However, the behavior in the non-threatening neutral environment of the home cage is normal (Easton *et al.*, 2011), in accordance with an impaired learning in MWM and fear conditioning (Giese *et al.*, 1998; Irvine *et al.*, 2005) but a normal appetitive instrumental conditioning (Carvalho *et al.*, 2001). Results in our study concur with these findings and argue for the specific importance of CaMKII binding to GluN2B in aversive and more demanding spatial learning task of the MWM, with the less threatening and stressful Barnes maze task being unaffected in the KI mice. Contextual fear conditioning and memory was surprisingly not affected in the GluN2B KI mice. We assume that this discrepancy is due to the higher demands of the MWM task and the differential regulation of spatial and contextual LTM formation (Mizuno & Giese, 2005).

Learning during massed training sessions, i.e., when training trials are given in short order on a single day, generally results in lower levels of learning than spaced training over a period of several days. For instance, in the MWM rats showed faster and better acquisition during spaced (16 trials over 4 days) than massed training (16 trials in 1 day) as well as better memory formation (Commins *et al.*, 2003). During contextual fear conditioning 1 h spaced training also results in better memory formation than massed training, while cued fear conditioning is unaffected (Scharf *et al.*, 2002). Likely because of the more demanding task of massed training, which requires information being stored during a shorter time period than in spaced training, the GluN2B KI mice revealed a mild learning deficit compared to WT mice in the 1 day paradigm (Figure 1) when they did not in the 6 day MWM training protocol (Halt *et al.*, 2012). Figure 1D shows that there is no difference in the swim speed between WT and KI mice, which together with the increased covered distance of the KI mice during the last two trials (Figure 1C) argues against the differences in agility and fatigue in the GluN2B KI. The increased covered distance shows that the KI mice have to swim longer and cover a greater distance in order to find the platform during the later trials, further supporting the extended primary escape latency (Figure 1B) and therefore the KI learning deficit. The learning deficit in the 1 day paradigm suggests that the learning capacity over a more limited time period is reduced in GluN2B KI mice. In addition, the elevated stress level during the massed training in the 1 day protocol could affect learning in the GluN2B KI mice more than in WT contributing to the differential outcome.

The GluN2B KI phenotype is similar to alpha and delta CREB KO mice, which are impaired in massed but not in spaced MWM learning and contextual fear conditioning (Kogan *et al.*, 1997). The improvement in memory after spaced learning is also mimicked in acute hippocampal slices where successive spaced theta burst stimulation resulted in enhanced previously saturated LTP (Kramar *et al.*, 2012). In the study by Kramar *et al.* the induction of LTP in a subset of synapses during the first theta burst train primed the initially unresponsive neighbors (probably in a translation-dependent manner, since the two stimulations have to be spaced 1 h apart; see also (Scharf *et al.*, 2002)), resulting in potentiation after stimulation with the second theta burst train. This mechanism will probably be especially important in cases of impaired LTP induction or expression like in our GluN2B KI mice where various forms of LTP were reduced by ~50% (Halt *et al.*, 2012).

The Barnes maze is a demanding spatial task using visual cues in the room for navigation, which is in that regard similar to the MWM. One major difference between these two spatial learning paradigms is probably the amount of stress to which animals are exposed. The unaffected performance of the GluN2B KI mice in the less stressful Barnes Maze is in line with the known importance of CaMKII selectively in aversive rather than appetitive motivated spatial and contextual learning paradigms (Elgersma *et al.*, 2002; Giese *et al.*, 1998; Irvine *et al.*, 2005; Silva *et al.*, 1992). However, further potential differences include sensory processes and response requirements, which cannot be excluded as a reason for the deficit in massed MWM learning compared to no deficit in the Barnes maze (Figure 2).

Fear conditioning, on the other hand, creates a high level of anxiety. Compared to spatial learning paradigms like the MWM, the contextual conditioning only requires memorization of a far less complex environment, which only has to be recognized and does not require navigation and localization of a certain target. Additionally, at the beginning of the conditioning the mice have time to explore and form a spatial map without being stressed or more anxious than during the normal exposure to a novel environment, before the first shock is delivered at the end of the 3rd minute. During the milder conditioning paradigm (Figure 5) the mice were pre-exposed to the shocking chamber on the previous day, which allowed the formation of an even better spatial representation (Fanselow, 2000; Frankland *et al.*, 2004; Hu *et al.*, 2007). This lack of a demanding spatial learning requirement and possible formation of contextual memory during still low stress levels could explain the normal conditioning and contextual memory (Figure 4 and Figure 5) compared to the moderate learning deficit observed during massed MWM training (Figure 1). In addition, fear conditioning in general constitutes a very strong learning paradigm, which might override and mask any mild learning deficits because of its vigor even in the mildest form tested here. In support of this idea the five tone-shock pairings allow full fear conditioning in CaMKII T286A mutant mice when one and three pairings do not (Irvine *et al.*, 2011; Irvine *et al.*, 2005). Accordingly, the strength of conditioning can mask learning deficits, and for GluN2B KI mice that might even be the case for single shock experiments. Spatial and contextual LTM formation is differently regulated. There are not only divergent regional requirements within the hippocampus,

but also the control of gene transcription varies between spatial and contextual LTM formation (Mizuno & Giese, 2005). Accordingly, MWM and fear conditioning could depend on molecular mechanisms that vary in detail even though both learning paradigms require CaMKII.

Contrasting CaMKII α heterozygous KO, T286A KI mice, or transgenic mice overexpressing CaMKII α , the GluN2B KI mice display no change in basal anxiety levels or anxiety-like behaviors. They show avoidance of the open arms in the EPM (Figure 3), behavior in the open field (Halt *et al.*, 2012), and reaction to TMT (the anxiogenic compound in fox urine) (Halt *et al.*, 2012), a measure of innate fear, that are comparable to WT mice. Thus it appears that CaMKII α affects fear behavior independent of its binding to GluN2B.

Our study suggests a specific requirement for the activity-dependent interaction of CaMKII with the NMDAR during the acquisition of more elaborate spatial learning tasks under modestly aversive, stressful conditions. The results argue not only for a role of CaMKII binding to GluN2B during consolidation (Halt *et al.*, 2012), but also for learning, which becomes obvious under the here applied more demanding MWM conditions. The less stressful Barnes maze neither results in a consolidation deficit nor is the learning itself affected. In addition, the less demanding but more potent contextual fear conditioning and memory was surprisingly normal. Whether reduced recall of the MWM task 1 and 7 days after single day training reflects reduced learning, reduced memory consolidation, or both cannot be answered at this point.

Data availability

F1000Research: Dataset 1. Data sets for behavioral tasks in wild type and GluN2BKI mice, [10.5256/f1000research.4660.d34165](https://doi.org/10.5256/f1000research.4660.d34165) (Stein *et al.*, 2014).

Author contributions

ISS and JWH designed the experiments. ISS with help from MSD carried out the experiments. ISS and JWH prepared the manuscript. All authors approved the final content of the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by NIH grants NS046450 and NS078792

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank the Personnel of the UC Davis Mouse Behavioral Assessment Lab, where all behavioral tests were conducted, for their assistance and help with data analysis, as well as Claire M. Koenig for training in the Barnes maze. We thank Dr. Karl-Peter Giese (Kings College, London, UK) for reading an earlier version of the manuscript.

References

- Barria A, Malinow R: **NMDA receptor subunit composition controls synaptic plasticity by regulating binding to CaMKII.** *Neuron.* 2005; **48**(2): 289–301. [PubMed Abstract](#) | [Publisher Full Text](#)
- Bayer KU, De Koninck P, Leonard AS, *et al.*: **Interaction with the NMDA receptor locks CaMKII in an active conformation.** *Nature.* 2001; **411**(6839): 801–805. [PubMed Abstract](#) | [Publisher Full Text](#)
- Berta S, Gert L, Harald H, *et al.*: **Barnes maze, a useful task to assess spatial reference memory in the mice.** *Protoc exch.* 2007. [Publisher Full Text](#)
- Carvalho MO, Silva JA, Balleine WB: **Evidence of selective learning deficits on tests of pavlovian and instrumental conditioning in α -CaMKII T286 mutant mice.** Human Sciences Press, New York, NY ETATS-UNIS. *Int J Comp Psychol.* 2001; **14**(3): 161–174. [Reference Source](#)
- Chen C, Rainnie DG, Greene RW, *et al.*: **Abnormal fear response and aggressive behavior in mutant mice deficient for alpha-calmodulin kinase II.** *Science.* 1994; **266**(5183): 291–294. [PubMed Abstract](#) | [Publisher Full Text](#)
- Commins S, Cunningham L, Harvey D, *et al.*: **Massed but not spaced training impairs spatial memory.** *Behav Brain Res.* 2003; **139**(1–2): 215–223. [PubMed Abstract](#) | [Publisher Full Text](#)
- Curzon P, Rustay NR, Browman KE: **Cued and Contextual Fear Conditioning for Rodents.** In Buccafusco, J.J. (ed), Boca Raton (FL). *Meth Behav Anal Neurosci.* 2009; 19–37. [PubMed Abstract](#) | [Publisher Full Text](#)
- Easton AC, Lucchesi W, Schumann G, *et al.*: **α CaMKII autophosphorylation controls exploratory activity to threatening novel stimuli.** *Neuropharmacology.* 2011; **61**(8): 1424–1431. [PubMed Abstract](#) | [Publisher Full Text](#)
- Elgersma Y, Fedorov NB, Ikonen S, *et al.*: **Inhibitory autophosphorylation of CaMKII controls PSD association, plasticity, and learning.** *Neuron.* 2002; **36**(3): 493–505. [PubMed Abstract](#) | [Publisher Full Text](#)
- Fanselow MS: **Contextual fear, gestalt memories, and the hippocampus.** *Behav Brain Res.* 2000; **110**(1–2): 73–81. [PubMed Abstract](#) | [Publisher Full Text](#)
- Frankland PW, Josselyn SA, Anagnostaras SG, *et al.*: **Consolidation of CS and US representations in associative fear conditioning.** *Hippocampus.* 2004; **14**(5): 557–569. [PubMed Abstract](#) | [Publisher Full Text](#)
- Giese KP, Fedorov NB, Filipkowski RK, *et al.*: **Autophosphorylation at Thr286 of the alpha calcium-calmodulin kinase II in LTP and learning.** *Science.* 1998; **279**(5352): 870–873. [PubMed Abstract](#) | [Publisher Full Text](#)
- Gruart A, Muñoz MD, Delgado-García JM: **Involvement of the CA3-CA1 synapse in the acquisition of associative learning in behaving mice.** *J Neurosci.* 2006; **26**(4): 1077–1087. [PubMed Abstract](#) | [Publisher Full Text](#)
- Halt AR, Dallapiazza RF, Zhou Y, *et al.*: **CaMKII binding to GluN2B is critical during memory consolidation.** *EMBO J.* 2012; **31**(5): 1203–1216. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hasegawa S, Furuichi T, Yoshida T, *et al.*: **Transgenic up-regulation of alpha-CaMKII in forebrain leads to increased anxiety-like behaviors and aggression.** *Mol Brain.* 2009; **2**: 6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Hu H, Real E, Takamiya K, *et al.*: **Emotion enhances learning via norepinephrine regulation of AMPA-receptor trafficking.** *Cell.* 2007; **131**(1): 160–173. [PubMed Abstract](#) | [Publisher Full Text](#)

- Irvine EE, Danhiez A, Radwanska K, *et al.*: **Properties of contextual memory formed in the absence of α CaMKII autophosphorylation.** *Mol Brain*. 2011; **4**: 8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Irvine EE, Vernon J, Giese KP: **AlphaCaMKII autophosphorylation contributes to rapid learning but is not necessary for memory.** *Nat Neurosci*. 2005; **8**(4): 411–412.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kessels HW, Malinow R: **Synaptic AMPA receptor plasticity and behavior.** *Neuron*. 2009; **61**(3): 340–350.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kogan JH, Frankland PW, Blendy JA, *et al.*: **Spaced training induces normal long-term memory in CREB mutant mice.** *Curr Biol*. 1997; **7**(1): 1–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kramár EA, Babayan AH, Gavin CF, *et al.*: **Synaptic evidence for the efficacy of spaced learning.** *Proc Natl Acad Sci U S A*. 2012; **109**(13): 5121–5126.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lengyel I, Voss K, Cammarota M, *et al.*: **Autonomous activity of CaMKII is only transiently increased following the induction of long-term potentiation in the rat hippocampus.** *Eur J Neurosci*. 2004; **20**(11): 3063–3072.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Leonard AS, Lim IA, Hemsworth DE, *et al.*: **Calcium/calmodulin-dependent protein kinase II is associated with the N-methyl-D-aspartate receptor.** *Proc Natl Acad Sci U S A*. 1999; **96**(6): 3239–3244.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lisman J, Schulman H, Cline H: **The molecular basis of CaMKII function in synaptic and behavioural memory.** *Nat Rev Neurosci*. 2002; **3**(3): 175–190.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Martin SJ, Grimwood PD, Morris RG: **Synaptic plasticity and memory: an evaluation of the hypothesis.** *Annu Rev Neurosci*. 2000; **23**: 649–711.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mizuno K, Giese KP: **Hippocampus-dependent memory formation: do memory type-specific mechanisms exist?** *J Pharmacol Sci*. 2005; **98**(3): 191–197.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Morris RG: **NMDA receptors and memory encoding.** *Neuropharmacology*. 2013; **74**: 32–40.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Neves G, Cooke SF, Bliss TV: **Synaptic plasticity, memory and the hippocampus: a neural network approach to causality.** *Nat Rev Neurosci*. 2008; **9**(1): 65–75.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Otmakhov N, Tao-Cheng JH, Carpenter S, *et al.*: **Persistent accumulation of calcium/calmodulin-dependent protein kinase II in dendritic spines after induction of NMDA receptor-dependent chemical long-term potentiation.** *J Neurosci*. 2004; **24**(42): 9324–9331.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ouyang Y, Kantor D, Harris KM, *et al.*: **Visualization of the distribution of autophosphorylated calcium/calmodulin-dependent protein kinase II after tetanic stimulation in the CA1 area of the hippocampus.** *J Neurosci*. 1997; **17**(14): 5416–5427.
[PubMed Abstract](#)
- Ouyang Y, Rosenstein A, Kreiman G, *et al.*: **Tetanic stimulation leads to increased accumulation of Ca(2+)/calmodulin-dependent protein kinase II via dendritic protein synthesis in hippocampal neurons.** *J Neurosci*. 1999; **19**(18): 7823–7833.
[PubMed Abstract](#)
- Rose J, Jin SX, Craig AM: **Heterosynaptic molecular dynamics: locally induced propagating synaptic accumulation of CaM kinase II.** *Neuron*. 2009; **61**(3): 351–358.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Scharf MT, Woo NH, Lattal KM, *et al.*: **Protein synthesis is required for the enhancement of long-term potentiation and long-term memory by spaced training.** *J Neurophysiol*. 2002; **87**(6): 2770–2777.
[PubMed Abstract](#)
- Silva AJ, Paylor R, Wehner JM, *et al.*: **Impaired spatial learning in alpha-calmodulin kinase II mutant mice.** *Science*. 1992; **257**(5067): 206–211.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Stein IS, Donaldson MS, Hell JW: **Data sets for behavioral tasks in wild type and GlunN2BK1 mice.** *F1000Research*. 2014.
[Data Source](#)
- Strack S, Colbran RJ: **Autophosphorylation-dependent targeting of calcium/calmodulin-dependent protein kinase II by the NR2B subunit of the N-methyl-D-aspartate receptor.** *J Biol Chem*. 1998; **273**(33): 20689–20692.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Strack S, McNeill RB, Colbran RJ: **Mechanism and regulation of calcium/calmodulin-dependent protein kinase II targeting to the NR2B subunit of the N-methyl-D-aspartate receptor.** *J Biol Chem*. 2000; **275**(31): 23798–23806.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Tan SE, Liang KC: **Spatial learning alters hippocampal calcium/calmodulin-dependent protein kinase II activity in rats.** *Brain Res*. 1996; **711**(1–2): 234–240.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Whitlock JR, Heynen AJ, Shuler MG, *et al.*: **Learning induces long-term potentiation in the hippocampus.** *Science*. 2006; **313**(5790): 1093–1097.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Zhang YP, Holbro N, Oertner TG: **Optical induction of plasticity at single synapses reveals input-specific accumulation of alphaCaMKII.** *Proc Natl Acad Sci U S A*. 2008; **105**(33): 12039–12044.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhou Y, Takahashi E, Li W, *et al.*: **Interactions between the NR2B receptor and CaMKII modulate synaptic plasticity and spatial learning.** *J Neurosci*. 2007; **27**(50): 13843–13853.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Referee Status:



Referee Responses for Version 1



Karl-Ulrich Bayer

Department of Pharmacology, University of Colorado Denver, Aurora, USA

Approved: 26 August 2014

Referee Report: 26 August 2014

doi:[10.5256/f1000research.4981.r5919](https://doi.org/10.5256/f1000research.4981.r5919)

Summary: This study substantially extends the previous behavioral analysis of a mouse line with mutant GluN2B that is incapable of CaMKII binding. Individually, both GluN2B and CaMKII are critically involved in regulating synaptic plasticity and learning and memory. Previous studies from the same lab showed that the mutant GluN2B KI mice have impaired hippocampal CA1 LTP (reduced to ~50%) and impaired recall during the consolidation phase of a Morris water maze (MWM) task ¹. Here, learning impairments were found in a modified MWM task that involves massed learning during day 1. All other learning paradigms tested here or previously were normal. Basal behavior was also extensively evaluated, and was found to be normal in all tasks tested here or previously.

Evaluation: The manuscript is very well written. The experiments and the analyses are solid. The interpretation and discussion of the results are both careful and insightful. Overall, it can be very reasonably concluded that learning requires the CaMKII/GluN2B interaction only when the learning task is rather difficult and/or involves some stress (such as in a water maze), consistent also with impaired but not abolished LTP in these mice.

Comments: The conclusions from this study may also inform potential future behavioral studies on these mice. For instance, while the GluN2B KI mice were normal in multiple variations of contextual fear conditioning, the contextual part of the task was made rather easy in all paradigms tested. However, even with shorter exposure to the context, fear conditioning may still be too easy of a learning task to reveal any differences in the GluN2B mice. Thus, while such future studies may be interesting, either outcome would still be consistent with the conclusions reached here.

The behavioral analyses of the GluN2B KI mice were done specifically in order to evaluate the impact of the LTP impairments in these mice. When comparing the GluN2B KI mice with CaMKII T286A or CaMKII knock-out mice in other behavioral tasks in future studies, it may have to be taken into account that the GluN2B KI mice are impaired only in LTP, while the CaMKII mutant mice are additionally impaired in LTD.

One point in which my opinion differs is within the introduction, specifically regarding the statement that “*auto-phosphorylation on T286 is required for effective binding of CaMKII to GluN2B*”. The cited references could indeed be interpreted support this statement^{2 3}. However, newer data indicate that the increased binding observed after ATP addition was largely caused by direct nucleotide binding to the kinase, rather than by auto-phosphorylation^{4 5 6}.

Notably, this point does not affect the validity of any conclusions reached in the manuscript. However, it may significantly mislead the reader about the larger context of CaMKII in synaptic plasticity. Recently, we have shown that CaMKII T286 phosphorylation is required not only for LTP, but also for LTD⁷. Prior to this, other recent evidence had already suggested a more prolonged T286 phosphorylation after LTD than after LTP^{8 9 10} (for review see Coultrap & Bayer, 2012¹¹). If T286 phosphorylation would induce the most effective CaMKII binding to GluN2B (as stated), binding would be expected to be more extensive after LTD compared to LTP. However, this does not appear to be the case⁹, and the CaMKII/GluN2B interaction contributes exclusively to the CaMKII functions in LTP but not in LTD¹.

References

1. Halt AR, Dallpiazza RF, Zhou Y, Stein IS, Qian H, Juntti S, Wojcik S, Brose N, Silva AJ, Hell JW: CaMKII binding to GluN2B is critical during memory consolidation. *EMBO J.* 2012; **31** (5): 1203-1216 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
2. Bayer KU, De Koninck P, Leonard AS, Hell JW, Schulman H: Interaction with the NMDA receptor locks CaMKII in an active conformation. *Nature.* 2001; **411** (6839): 801-805 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Strack S, Colbran RJ: Autophosphorylation-dependent targeting of calcium/ calmodulin-dependent protein kinase II by the NR2B subunit of the N-methyl- D-aspartate receptor. *J Biol Chem.* 1998; **273** (33): 20689-20692 [PubMed Abstract](#) | [Publisher Full Text](#)
4. Barcomb K, Coultrap SJ, Bayer KU: Enzymatic Activity of CaMKII Is Not Required for Its Interaction with the Glutamate Receptor Subunit GluN2B. *Mol Pharmacol.* 2013; **84**: 834-843 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
5. O'Leary H, Liu WH, Rorabaugh JM, Coultrap SJ, Bayer KU: Nucleotides and phosphorylation bi-directionally modulate Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) binding to the N-methyl-D-aspartate (NMDA) receptor subunit GluN2B. *J Biol Chem.* 2011; **286** (36): 31272-31281 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
6. Robison AJ, Bartlett RK, Bass MA, Colbran RJ: Differential modulation of Ca²⁺/calmodulin-dependent protein kinase II activity by regulated interactions with N-methyl-D-aspartate receptor NR2B subunits and alpha-actinin. *J Biol Chem.* 2005; **280** (47): 39316-39323 [PubMed Abstract](#) | [Publisher Full Text](#)
7. Coultrap SJ, Freund RK, O'Leary H, Sanderson JL, Roche KW, Dell'Acqua ML, Bayer KU: Autonomous CaMKII mediates both LTP and LTD using a mechanism for differential substrate site selection. *Cell Reports.* 2014; **6** (3): 431-437 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
8. Lee SJ, Escobedo-Lozoya Y, Szatmari EM, Yasuda R: Activation of CaMKII in single dendritic spines during long-term potentiation. *Nature.* 2009; **458** (7236): 299-304 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
9. Marsden KC, Shemesh A, Bayer KU, Carroll RC: Selective translocation of Ca²⁺/calmodulin protein kinase IIalpha (CaMKIIalpha) to inhibitory synapses. *Proc Natl Acad Sci U S A.* 2010; **107** (47): 20559-20564 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)
10. Mockett BG, Guevremont D, Wutte M, Hulme SR, Williams JM, Abraham WC: Calcium/calmodulin-dependent protein kinase II mediates group I metabotropic glutamate receptor-dependent protein synthesis and long-term depression in rat hippocampus. *J Neurosci.* 2011; **31** (20): 7380-7391 [PubMed Abstract](#) | [Publisher Full Text](#)
11. Coultrap SJ, Bayer KU: CaMKII regulation in information processing and storage. *Trends Neurosci.* 2012; **35** (10): 607-618 [PubMed Abstract](#) | [Free Full Text](#) | [Publisher Full Text](#)

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.



Mark L. Dell'Acqua

Department of Pharmacology, University of Colorado Denver, Denver, USA

Approved: 22 August 2014

Referee Report: 22 August 2014

doi:[10.5256/f1000research.4981.r5845](https://doi.org/10.5256/f1000research.4981.r5845)

This is an interesting article that provides additional, detailed behavioral analysis of hippocampal-dependent spatial and contextual learning and memory performance of knock-in mice that are deficient in CaMKII binding to the GluN2B subunit of the NMDA-subtype ionotropic glutamate receptor. The current findings extend those of the authors in [Halt et al. \(2012\)](#) where spaced training over 6 days in the Morris Water Maze (MWM) revealed no deficits in spatial learning but did find significant deficits in consolidation of long-term memory probed several days after training in these CaMKII binding-deficient GluN2B knock-in mice.

Interestingly, here, using a more intense and demanding one day, massed training version of the MWM, the authors now detect modest deficits in these GluN2B knock-in mice, not only in memory probed one day after training, but also in spatial learning measured on the day of training. However, other single day training spatial/contextual learning and memory tests, that are both more aversive but less spatially challenging (contextual fear conditioning) and less aversive but more spatially challenging (Barnes maze), did not reveal any difference in performance between WT and GluN2B knock-in mice. Thus, an important conclusion of this study is that increased postsynaptic targeting of CaMKII and the increased level of LTP (~50%; see Halt et al.) that are achieved through CaMKII binding to GluN2B are only required for spatial learning when the task is both fairly demanding and at least moderately aversive. These findings should give the field a heightened appreciation that highly-compartmentalized regulation of signaling pathways by postsynaptic protein-protein scaffolding interactions, such as achieved through CaMKII binding to GluN2B during LTP-induction, may not be absolutely essential for spatial learning and memory in general but may have evolved to allow animals to specifically sustain optimal spatial learning and memory capabilities even under demanding and stressful circumstances.

Overall, the manuscript is very well written and the data are rigorously analyzed and presented clearly.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.
