

MEETING ABSTRACT

Open Access

EHMTI-0184. Ictal adiponectin levels are modulated by pain severity and treatment response in episodic migraineurs

NC Chai^{1*}, B Gelaye², GE Tietjen³, PD Dash⁴, BA Gower⁵, LW White¹, TN Ward⁶, AI Scher⁷, BL Peterlin¹

From 4th European Headache and Migraine Trust International Congress: EHMTIC 2014 Copenhagen, Denmark. 18-21 September 2014

Introduction

Adiponectin (ADP) and leptin (LEP) are adipokines with roles in inflammation.

Aim

To assess ADP and LEP levels before and after acute abortive treatment in episodic migraineurs (EM).

Methods

Peripheral blood specimens were collected from EM participants before and after acute abortive treatment with sumatriptan/naproxen sodium versus placebo.

Results

A total of 34 participants (17 responders, 17 non-responders) were included. In all participants, for every 1 point increase in the HMW:T-ADP ratio, pain severity increased by 4.11 (CI: 0.44, 7.77; $p = 0.028$). In responders ($n = 17$), crude T-ADP levels were reduced at 30 min (11.49 ± 3.7 ; $p = 0.001$), 60 min (11.54 ± 3.2 ; $p = 0.001$) and 120 min (11.39 ± 3.7 ; $p = 0.001$) after treatment as compared to onset (12.47 ± 3.6). In non-responders ($n = 17$), crude T-ADP levels were unchanged after treatment. After adjustments, T-ADP levels remained decreased 30-120 min after treatment in responders; additionally, HMW-ADP, and the HMW:T-ADP ratio were decreased and LMW-ADP and the LMW:T-ADP ratio (coeff 0.04; CI: 0.01, 0.07; $p = 0.043$) were increased 120 min after treatment in responders. In non-responders, the adjusted LMW-ADP (coef -0.45; CI: -0.77, -0.14; $p = 0.005$) and the LMW:T-ADP ratio (coef -0.04; CI: -0.07, -0.01; $p = 0.018$) were decreased at 60 min as well as at 120 min after

treatment. Unadjusted and adjusted LEP levels were not modulated by changes in pain severity or treatment response.

Conclusion

Adiponectin, but not leptin, is associated with pain severity and is modulated by treatment response in episodic migraineurs.

Conflict of interest.

Authors' details

¹Neurology, Johns Hopkins University School of Medicine, Baltimore, USA. ²Epidemiology, Harvard School of Public Health, Boston, USA. ³Neurology, University of Toledo, Toledo, USA. ⁴Neurology, Johns Hopkins Community Physicians, Baltimore, USA. ⁵Department of Nutrition Sciences, University of Alabama at Birmingham, Birmingham, USA. ⁶Neurology, Dartmouth Hitchcock Medical Center, Lebanon, USA. ⁷Preventive Medicine and Biometrics, Uniformed Services University, Bethesda, USA.

Published: 18 September 2014

doi:10.1186/1129-2377-15-S1-E25

Cite this article as: Chai et al.: EHMTI-0184. Ictal adiponectin levels are modulated by pain severity and treatment response in episodic migraineurs. *The Journal of Headache and Pain* 2014 **15**(Suppl 1):E25.

¹Neurology, Johns Hopkins University School of Medicine, Baltimore, USA
Full list of author information is available at the end of the article