

# N-Acetylcysteine Compared to Metformin, Improves The Expression Profile of Growth Differentiation Factor-9 and Receptor Tyrosine Kinase c-Kit in The Oocytes of Patients with Polycystic Ovarian Syndrome

Ebrahim Cheraghi, Ph.D.<sup>1</sup>, Malek Soleimani Mehranjani, Ph.D.<sup>2\*</sup>, Seyed Mohammad Ali Shariatzadeh, Ph.D.<sup>2</sup>, Mohammad Hossein Nasr Esfahani, Ph.D.<sup>3</sup>, Behrang Alani, Ph.D.<sup>4</sup>

1. Department of Biology, Faculty of Sciences, University of Qom, Qom, Iran

2. Department of Biology, Faculty of Science, Arak University, Arak, Iran

3. Department of Reproductive Biotechnology, Reproductive Biomedicine Research Center, Royan Institute, ACECR, Isfahan, Iran

4. Department of Applied Cell Science, Faculty of Medicine, Kashan University of Medical Sciences, Kashan, Iran

## Abstract

**Background:** Paracrine disruption of growth factors in women with polycystic ovarian syndrome (PCOS) results in production of low quality oocyte, especially following ovulation induction. The aim of this study was to investigate the effects of metformin (MET), N-acetylcysteine (NAC) and their combination on the hormonal levels and expression profile of GDF-9, BMP-15 and c-kit, as hallmarks of oocyte quality, in PCOS patients.

**Materials and Methods:** This prospective randomized, double-blind, placebo controlled trial aims to study the effects of MET, NAC and their combination (MET+NAC) on expression of GDF-9, BMP-15 and c-kit mRNA in oocytes [10 at the germinal vesicle (GV) stage, 10 at the MI stage, and 10 at the MII stage from per group] derived following ovulation induction in PCOS. Treatment was carried out for six weeks, starting on the third day of previous cycle until oocyte aspiration. The expression of GDF9, BMP15 and c-kit were determined by quantitative real time polymerase chain reaction (RT-qPCR) and western blot analysis. Data were analyzed with one-way ANOVA.

**Results:** The follicular fluid (FF) level of c-kit protein significantly decreased in the NAC group compared to the other groups. Significant correlations were observed between the FF soluble c-kit protein with FF volume, androstenedione and estradiol. The GDF-9 expression in unfertilized mature oocytes were significantly higher in the NAC group compared to the other groups ( $P < 0.001$ ). Similar difference was not observed between the MET, NAC+MET and control groups. The c-kit expression in unfertilized mature oocytes were significantly lower in the NAC group compared to the other groups ( $P < 0.001$ ). Similar difference was not observed between the MET, NAC+MET and control groups (Registration number: IRCT201204159476N1).

**Conclusion:** : We concluded that NAC can improve the quality of oocytes in PCOS.

**Keywords:** Gene Expression, Metformin, N-acetylcysteine, Oocyte, Polycystic Ovarian Syndrome

**Citation:** Cheraghi E, Soleimani Mehranjani M, Shariatzadeh SMA, Nasr Esfahani MH, Alani B. N-acetylcysteine compared to metformin, improves the expression profile of growth differentiation factor-9 and receptor tyrosine kinase c-kit in the oocytes of patients with polycystic ovarian syndrome. *Int J Fertil Steril.* 2018; 11(4): 270-278 . doi: 10.22074/ijfs.2018.5142.

## Introduction

Anovulation associated with polycystic ovary syndrome (PCOS), as a common metabolic disorder, is the major cause of female infertility (1, 2). The principal feature of PCOS is the large number of follicles arresting at early growth stage. The cytoplasmic and nuclear maturity of oocytes is reduced following ovarian stimulation and may account for embryo quality in these couples (3). Exclusive oocyte secreted factors (OSFs), such as growth differentiation factor-9 (GDF-9) and

bone morphogenetic factor-15 (BMP-15), belonging to transforming growth factor- $\beta$  superfamily, are essential for oocyte competence (3-5). The receptor tyrosine kinase c-kit is another OSFs which plays important role in oogenesis and folliculogenesis (6). Recent evidence suggest possible involvement of c-kit and its receptor, kit ligand (KL) in PCOS pathology (7). Indeed, it has been shown that the aberrant or low expression of these exclusive oocyte secreting factors (BMP-15 and GDF-9), lead to over expression of c-kit and its receptor (8-

Received: 11 Dec 2016, Accepted: 18 Jun 2017

\*Corresponding Address: P.O.Box: 381-5688138, Department of Biology, Faculty of Science, Arak University, Sardasht, Arak, Iran  
Email: M-soleimani@araku.ac.ir



Royan Institute  
International Journal of Fertility and Sterility  
Vol 11, No 4, Jan-Mar 2018, Pages: 270-278

10). Therefore, drugs which can modulate the regulation of these intra-ovarian factors may play a role in the clinical management of PCOS.

To improve the quality of oocyte, various protocol have been designed, tested and verified for ovulation induction along with insulin-sensitizing drugs in PCOS patients. But, the risks of poor response, ovarian hyperstimulation, production of low quality oocytes, reduced fertilization rates and poor embryo quality remains among concern to be dealt within PCOS women undergoing *in vitro* fertilization (IVF) or intracytoplasmic sperm injection (ICSI) (11). Metformin, an insulin-lowering agent, has been extensively used for treatment of anovulation and infertility in PCOS patients. However, the efficacy of MET treatment is still disputed (12). In this regard, background studies indicate that MET does not improve the overall outcomes of assisted reproductive procedure in term of the aforementioned parameters (13, 14).

On the contrary, administration of N-acetylcysteine (NAC) has been shown to improve not only the number and also the quality of oocytes in these patients. This phenomenon has been mainly related to the strong antioxidant effect of NAC, which has been shown to reduce follicle atresia and improve the quality of oocyte (15). *In vitro*, NAC plays a key role in cell survival through the production of trophic factor and follicular preservation (16, 17). In line with these reports, Sacchinelli et al. (18) showed that co-administration of inositol and NAC improve ovarian function of PCOS patients. Therefore, considering the fact that oocyte secretory factors are hallmarks of oocyte quality, this study aims to evaluate the effects of NAC, MET and their co-administration on the expression of GDF-9, BMP-15 and c-kit in PCOS individuals undergoing ovarian stimulation in ICSI cycle.

## Materials and Methods

Antibodies directed against c-kit and  $\beta$ -actin was obtained from Cell Signaling Technology (Beverly, MA, USA). BMP-15 antibody was obtained from Abcam Technology (Cambridge, MA, USA) and GDF-9 antibody from Santa Cruz Biotechnology (CA, USA). Other reagents used in this study were obtained from Sigma-Aldrich (St. Louis, MO, USA). Cell culture media and sera were obtained from Gibco BRL (Carlsbad, CA, USA).

### Study design

This study was performed in continuation of our prospective randomized, double-blind, placebo controlled trial, in the IVF Unit of Infertility Research Center of the Academic Center for Education, Culture and Research (ACECR), Qom/Iran. 80 infertile PCOS women at the age of 25-35 years, in the interval between July 2012 and February 2013, who planned to undergo ICSI were included in this study (19). Individuals were

diagnosed as PCOS according to the Rotterdam consensus workshop (20). Based on this consensus, each individual needed to have two out of three criteria: i. Biochemical or clinical hyperandrogenism, ii. Chronic oligo or anovulation and iii. Polycystic ovaries at ultrasound examination. Ethical consideration and further information on this clinical trial are provided in previous studies (19). This study was approved by the Ethics Committee (EC/91/1041) of Royan Institute, Tehran, Iran. The patients provided an informed consent and committed to avoid any changes in their normal physical activity, diet or starting a new medical regimen throughout the study.

### Treatment design and ovulation induction

The female partner of ICSI candidates were examined and randomly divided into 4 groups (n=20): i. Placebo (PLA) receiving oral rehydration solution (ORS, Poursina, Iran), ii. MET receiving MET (Glucophage, Merck, West Drayton, UK, 500 mg), iii. NAC receiving NAC (Holzkirchen, Germany, batch no. 6N5483, 600 mg), and iv. MET+NAC group receiving the combination of MET and NAC at the aforementioned doses. Treatment was carried out three times daily for a period of six weeks. The dose and duration of NAC treatment was chosen according to recent studies (21-23).

Gonadotropin-releasing hormone (GnRH) agonist protocol (18) was used to induce ovulation. The female partner of ICSI candidates randomized to four groups received PLA, MET, NAC or MET+NAC from the third day of last menstrual period (LMP) of previous cycle until the day of oocyte aspiration. Oral contraceptive pills (OCPs) were also included in the regimen for 21 days starting simultaneously with placebo, MET, NAC or MET+NAC on the day 3 of menstrual cycle prior to the treatment cycle. For ovarian down-regulation, daily injections of Bucerelin acetate (1 mg, Suprefact, Aventis, Germany) were administered from the day 19 of the preceding, menstrual cycle until day 2 of the next cycle. On the second day on the next cycle if the endometrial thickness was less than 4 mm, the dose of Bucerelin acetate was reduced to 0.5 mg.

Ovulation induction was induced from the day two of the cycle with average daily injections of 2 ampoules of recombinant follicle stimulating hormone (rFSH, Gonal-f, Merck Serono S.A., Geneva, Switzerland). Vaginal ultrasound (Honda Electronics HS 4000-Japan) was also used to monitor the cycles. 10,000 IU human chorionic gonadotropin (hCG, Pregnyl, Organon, Netherlands) was administered to induce ovulation. 36 hours after the administration of hCG, when at least three follicles had reached the diameters of 16-18 mm, transvaginal oocyte aspiration was performed under ultrasound guidance and general anesthesia. This protocol of induction ovulation was used for all the individuals in the 4 group. During the treatment the par-

Participants were asked to report any probable side effects such as abdominal discomfort, diarrhea and nausea. Due to these side effects, 20 couples (5 per group) were excluded from the study (Fig.1).

**Preparation of oocytes, follicular fluid and blood samples**


Based on our previous study, oocytes and follicular fluid (FF) from multiple follicles, from each subject were pooled as explained (18). Following oocyte retrieval, their cumulus cells were removed by exposure to 20 IU/ml hyaluronidase (ART-4007A, SAGE BioPharma, USA) in HEPES-based medium for 30 seconds followed by mechanical pipetting in HEPES-buffered HTF containing 5 mg/ml human serum albumin (ART-3001, SAGE BioPharma, USA).

The nuclear status of each oocyte was determined under the stereo microscope (Olympus Co., Japan) and classified into three categories: i. Unfertilized mature oocyte [metaphase II (MII)] following ICSI, ii. Germinal vesicle (GV) stage, iii. Without first polar body called metaphase I (MI). For gene expression analysis, in each experimental group, 10 GV, 10 MI, and 10 MII oocytes were separately pooled and washed in phosphate-buffered saline (PBS) and transferred into RNase-

free microcentrifuge tubes. 50 µl of RNAlater, RNA Stabilization Reagent (Qiagen, USA) were added to each tube and all samples were stored in a -80°C freezer until analysis. Only MII oocytes were used for ICSI.

The FF, from the first aspirated with no visible blood contamination was collect and immediately centrifuged at 3000 rpm for 10 minutes, and the supernatants were stored at -70°C for further analysis. Fasting blood sample were also collected from each participant once prior to the start of treatment (day 2 of previous cycle) and once on the day of ovum pick up of ICSI cycle. The samples were immediately centrifuged for 10 minutes at 3000 rpm (Hettich, EBA20, UK) and the resulting serum were stored at -70°C for evaluation.

The levels of luteinizing hormone (LH, mIU/ml), FSH (mIU/ml), total testosterone (TT, ng/ml), Progesterone (ng/ml), estradiol (E2, pg/ml) and androstenedione (ng/ml) in the FF and serum were measured in all samples using the ELISA enzyme immunoassay (Demeditec Diagnostics GmbH, Germany) according to the manufacturer’s protocol. The FF soluble protein level of c-Kit (pg/ml) was measured with the ELISA Kit (Abnova Corporation, Taiwan) by sandwich enzyme immunoassay technique, according to the manufacturer’s protocol.



**Fig.1:** Flowchart of participants in this study. MET; Metformin, NAC; N-acetylcysteine, and PLA; Placebo.

## Gene expression analysis

Total RNA from the oocytes of each group were isolated using the EZ-10 total RNA mini-prep Kit (Bio Basic Inc., Canada), according to the manufacturer's protocol. All samples were stored at  $-80^{\circ}\text{C}$  till analysis. Complementary DNA (cDNA) was synthesized, using random hexamers [using the RevertAid First Strand cDNA synthesis Kit (Thermo scientific, USA)]. To determine the relative expression of target genes, quantitative real time polymerase chain reaction (RT-qPCR) was carried out using SYBR-Green/ROX qPCR master mix assay (Thermo scientific, USA) by gene-specific primers (Table 1). Relative gene expression was calculated as the abundance ratio of each target gene relative to  $\beta$ -actin. The ABI step one plus (ABI, USA) instrument was used for real time PCR experiments and the  $\Delta\Delta\text{Ct}$  method for data calculation.

**Table 1:** Identity and sequence details of polymerase chain reaction (PCR) primers used to analyses mRNA expression in oocytes

Gene	Sequence primer (5'-3')
<i>GDF-9</i>	F: CCAATAGAAGTCACCTC R: GCGATCCAGGTTAAATAGCA
<i>BMP-15</i>	F: CAGTCCTCTATTGCCCTTCT R: AATGGTGCGGTTCTCTCTA
<i>c-Kit</i>	F: ACGAATGAGAATAAGCAGAATGAA R: GAGAGGACAGCGGACCAG
$\beta$ -actin	F: GGACTTCGAGCAAGAGATGG R: AGCACTGTGTTGGCGTACAG

Total proteins from each pool of oocytes were extracted using RIPA lysis and extraction buffer Kit (Cat No: 89900, Thermo Scientific, USA), according to the manufacturer's instruction. Concentration of proteins was determined according to Bradford's method using bovine serum albumin (BSA) as a reference standard (Bradford, 1976). Total proteins were electrophoresed in 12.5% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) gel, transferred to polyvinylidene fluoride membranes, and probed with specific antibodies. Membranes were developed using enhanced chemiluminescence reagents (Amersham Bioscience, USA) and the intensity of immunoreactive polypeptides was analyzed subsequent to visualization of the bands developed on a photographic film. Protein bands on photographic film

were quantified by densitometry scanning after background subtraction. Integrated densities of bands were measured by Image J software.

## Statistics

The normality of continuous variables was confirmed using the Kolmogorov-Smirnov test and data were reported as means  $\pm$  SEM. Data analysis were performed using one-way ANOVA and Tukey's test for post-hoc. Means were considered significantly different at  $P < 0.05$ . Pearson's correlation test defined the relation between variables. All data were analyzed with the statistical software SPSS (version 16.0 for windows, Chicago, IL, USA).

## Results

Patients characteristics including age, body mass index (BMI), the level of LH, FSH, E2 and TT were not significantly different among the groups PCOS prior to treatment.

### Follicular fluid analysis

FF volume and FF level of androstenedione, E2 and progesterone were similar in all groups ( $P > 0.05$ , Table 2), but the level of soluble c-kit protein in the FF significantly decreased in the NAC group compared to other groups ( $P < 0.01$ ). Our results also showed a significant correlation between the soluble c-kit protein in the FF of all the population with the FF volume ( $r = 0.508$ ,  $P = 0.02$ ), androstenedione ( $r = 0.682$ ,  $P = 0.01$ ), and E2 ( $r = 0.638$ ,  $P = 0.01$ ) (Fig.2).


### Evaluation of oocyte and embryo quality

The number of immature oocytes (MI+GV) and abnormal mature oocytes significantly decreased in the NAC group ( $P < 0.01$ ) compared to the other groups. Similar reduction was also observed in MET and MET+NAC groups but the reduction was not significant compared to the placebo group ( $P > 0.05$ ). The fertilization rate of metaphase II oocytes were similar in all groups ( $P > 0.05$ ). The number of good embryo (grade I) on day 3 showed a significant increase in the NAC group ( $P < 0.02$ ) compared to placebo group. This improvement was also observed in the MET and MET+NAC groups when compared to the placebo ( $P > 0.05$ ), but remained insignificant (Fig.3). The percentage of top grad embryos was not different between the three NAC with MET and MET+NAC groups.


**Table 2:** Comparison of the biochemical parameters of follicular fluid in PCOS patients

Parameter	NAC	MET	NAC+MET	PIA
Follicular fluid Volume (ml)	$5.4 \pm 1.13^a$	$5.7 \pm 1.1^a$	$4.96 \pm 1.1^a$	$4.9 \pm 1.1a$
Estradiol (pg/ml)	$466.6 \pm 34.8^a$	$496.6 \pm 44.8^a$	$470.2 \pm 48.4^a$	$436 \pm 40.16^a$
Progesterone (ng/ml)	$3983.7 \pm 353.9^a$	$3916.9 \pm 359.5^a$	$3501.2 \pm 326.9^a$	$3255.3 \pm 414.8^a$
Androstenedione (ng/ml)	$426.2 \pm 30.3^a$	$435.2 \pm 46.7^a$	$487.6 \pm 42.9^a$	$548.3 \pm 42.36^a$
Soluble c-Kit (pg/ml)	$317.8 \pm 27.5^b$	$380.8 \pm 30.3^a$	$429.8 \pm 28.7^a$	$455.2 \pm 28.75^a$

Data are shown as mean  $\pm$  SEM. Analysis was performed by ANOVA and Tukey's test for multiple comparisons. Means without a common letter are significantly different ( $P < 0.05$ ). PCOS; Polycystic ovarian syndrome, MET; Metformin, and NAC; N-acetylcysteine, and PLA; Placebo.



**Fig.2:** Correlation between parameters of follicular fluid (FF) in all the population. **A.** Soluble c-Kit with volume of FF, **B.** Estradiol (E2), and **C.** Androstenedione. R<sup>2</sup>; Determination of coefficient.




**Fig.3:** Distribution of oocytes retrieved, quality of oocytes and embryos in polycystic ovary syndrome patients undergoing treatment of N-acetylcysteine (NAC), metformin (MET), NAC+MET and Placebo (control). Data are the mean  $\pm$  SEM. Statistical analyses were performed by ANOVA followed by Tukey's test for multiple comparisons. Means without a common letter are significantly different ( $P < 0.05$ ).

**Expression profile of GDF-9, BMP-15 and c-Kit in oocytes**


The level of BMP-15 protein in the mature unfertilized oocytes and GV oocytes did not differ significantly among the groups ( $P > 0.05$ , Figs.4, 5A, D, E). The expression level of GDF-9 in the GV oocytes significantly increased in all groups compared to the placebo ( $P < 0.001$ ) (Fig.4B, D, F), while for unfertilized mature oocytes, GDF-9 mRNA and protein levels only significantly increased in the NAC group ( $P < 0.001$ , Fig.5B, D, F). The expression of c-kit in the GV oocytes significantly decreased in the NAC and

MET groups compared to the placebo group ( $P < 0.001$ ), but not in NAC+MET group (Fig.4C, D, G). The c-kit expression in the unfertilized mature oocytes significantly decreased in the NAC group compared to the MET and other treatment groups ( $P < 0.001$ ), but no significant difference was found in the MET and NAC+MET groups when compared to the placebo group ( $P > 0.05$ , Fig.5C, D, G). It is important to note that the results for MI oocytes were similar to GV oocyte; therefore, in this article only the results of GV were presented. This observation is in accordance with pervious literature (24).



**Fig.4:** Effects of NAC and MET on *BMP-15*, *GDF-9* and *c-kit* mRNA and protein expression in immature oocytes (IMO, GV oocytes) of PCOS patients. Results of reverse transcriptase real-time polymerase chain reaction (PCR) for mRNAs of **A.** *BMP-15*, **B.** *GDF-9*, **C.** *c-kit* in GV oocytes, **D.** Immunoblots of BMP-15, GDF-9 and c-kit from oocyte cell lysates. Densities of **E.** BMP-15, **F.** GDF-9, and **G.** c-kit protein bands in the experimental groups are shown. Means without a common letter are significantly different ( $P < 0.05$ ). NAC; N-acetylcysteine, MET; Metformin, GV; Germinal vesicle, and PCOS; Polycystic ovarian syndrome.

## NAC Improves Gene Expression Profile



**Fig.5:** Effects of NAC and MET on *BMP-15*, *GDF-9* and *c-kit* mRNA and protein expression in unfertilized mature oocytes (UMO, MII oocytes) of PCOS patients. Results of reverse transcriptase real-time polymerase chain reaction (PCR) for mRNAs of **A.** *BMP-15*, **B.** *GDF-9*, **C.** *c-kit* in MII oocytes, **D.** Immunoblots of BMP-15, GDF-9 and c-kit from oocyte cell lysates. Densities of **E.** BMP-15, **F.** GDF-9, and **G.** c-kit protein bands in the experimental groups are shown. Means without a common letter are significantly different ( $P < 0.05$ ). NAC; N-acetylcysteine, MET; Metformin, and PCOS; Polycystic ovarian syndrome.

## Discussion

A typical characteristic of PCOS patient commonly observed during induction stimulation for ART cycles is increased number of low quality oocytes which is mainly related to state of endocrine disorder in these individuals (25). Considering the indispensable role of OSFs in oocyte development and maturation, many researches have shown impaired expression of OSFs particularly GDF-9, BMP-15 and c-kit, may account for low quality oocyte in PCOS undergoing ovarian stimulation (26). This may explain, at least a part of the folliculogenesis disorders found in these patients (27-29). Background literature in this filed is very discrepant. Some authors have reported reduce expression of GDF-9 with no significant alteration in the expression of BMP-15 (8), while others have shown no alteration in expression of these two factors both at RNA and protein level (9) in oocyte of PCOS individuals. The exact reason of such discrepancy is not well understood.

In continue to our previous study, we demonstrated that unlike MTE and NAC+MET groups, the administration of NAC compared to placebo group, improves the maturation and quality of oocytes and also embryo development in PCOS patients undergoing ICSI (18). Therefore, in this we aimed to evaluate whether NAC could alter BMP-15, GDF-9 and c-kit levels, as the main OSFs in the oocytes of PCOS patients in comparison to MET and MET+NAC.

Compelling evidence suggest that GDF-9 and BMP-15 members of the TGF $\beta$  superfamily are exclusively expressed in the oocyte and their expression increases as follicle development progresses (30). During postnatal ovarian development, c-kit mRNA and protein are localized in the oocytes (6), and in this regard Brankin et al. (31) has shown a relation between KL/c-kit interaction with antrum formation, steroidogenesis and oocyte quality. Furthermore, genetic and descriptive studies have implicated the involvement of c-kit receptor and its ligand, KL, in oocyte growth (32).

Low GDF-9 levels is associated with abnormally increased KL level in PCOS, which could lead to abnormal ovarian features such as enlarged oocytes and increased follicle numbers (7, 32). In PCOS patients, the GDF-9 mRNA level within the oocytes is lower than in oocyte derived from normal individuals (8), and it is believed that there is a negative association between GDF-9 expression and KL/c-kit expression. Tuck (7) believes that excess androgens may act to further reduce the inhibitory effect of GDF-9, thus resulting in an abnormal increase in the KL/c-kit protein level in PCOS individual. Considering the inverse relationship between c-kit and GDF-9 in PCOS (7, 8, 32), therefore, improving the expression of GDF-9 is expected to cause a reduction in c-kit levels.

Our results displayed a significant increase in the expression of GDF-9 in the unfertilized mature oocytes of PCOS patients after administration of NAC compared to MET, indicating that NAC, as an anti-oxidant/anti-apoptotic agent, could enhance the expression of GDF-9 through inhibiting the activity of NF-kB and AP-1 tran-

scription factors, therefore affecting the activity of MAP-kinase signaling and related genes expression (33), which may be able to alleviate PCOS follicular disorders and prevent follicular developmental detention and atresia.

Our study, in agreement with aforementioned studies, also showed a significant decrease in the expression of c-kit in the oocytes of PCOS patients and also the soluble c-kit protein in the FF following administration of NAC compared to control. In addition, evidence has indicated the relationship between KL/c-kit system with MAPK pathway and/or PI3K/Akt pathway, which are both necessary for follicle development (34). It is likely that NAC decreases the expression of c-kit through interference in MAPK pathway, all of this could be the underlying reason in the role of NAC in preventing follicular developmental detention and atresia and alleviation of follicular disorders in PCOS patients.

Although the FF content may be an invaluable hallmark for PCOS diagnosis, but the NAC ability to modulate these intra-ovarian factors of the oocyte may have interesting pharmacological perspectives for clinical management of PCOS patients. According to literature (7, 8, 32), there is an inverse relationship between c-kit and GDF-9. Therefore, improves expression of GDF-9 by NAC treatment, is expected to follow by a reduction in c-Kit and indeed MET appears to mask this effect of NAC, how this masking effect is performed, remains to be elucidated.

In this regard it has been shown that with increased follicular size and E2 production, the amount of soluble c-kit protein in human FF also increase (35), which is consistent with the correlation observed in this study between soluble c-kit with the FF volume, E2, and androstenedione concentrations.

It has been demonstrated that the excess secretion of anti-mullerian hormone (AMH) in the FF of PCOS patients may directly inhibit the production of OSFs such as GDF-9 and BMP-15, which can explain the low levels of OSFs in PCOS oocytes (36). Our findings showed a reduction in the AMH level in the FF of NAC treated group (18). Although this reduction was not statistically significant but may be considered as an underlying reason for the increased levels of GDF-9 in the NAC treated patients. Moreover, in agreement with other findings (8, 9), our study revealed no significant difference in the level of BMP-15 mRNA among the studies groups.

## Conclusion

Considering the fact that NAC improves oocyte maturation and embryo quality, and decreases the rate of immature oocytes in women with PCOS while being a safe and well-tolerated agent, we suggest the administration of NAC as an alternative to other insulin-sensitizing agents like MET. Therefore, the present study argues that NAC possibly improves the oocyte quality of PCOS patients compared to MET through modulating the c-kit and GDF-9 expression, indicating that NAC supplement may be a therapeutic alternative to the insulin-sensitizing agents in PCOS management.



## Acknowledgements

We thank the members of the IVF unit of the Infertility Research Center of the Academic Center for Education, Culture and Research (ACECR), Qom. This research was financially supported by Arak University. The authors declare that there is no conflict of interest.

## Author's Contributions

E.C., M.S.M., S.M.A.S.; Participated in study design, data collection and evaluation, drafting and statistical analysis. M.H.N.E., E.C.; Performed follicle collection and prepared oocytes for ICSI pertaining to this component of the study. E.C., M.S.M., M.H.N.E.; Contributed extensively in interpretation of the data and the conclusion. B.A., E.C.; Conducted molecular experiments and RT-qPCR analysis. All authors performed editing and approving the final version of this paper for submission, also participated in the finalization of the manuscript and approved the final draft.

## References

- Teede H, Deeks A, Moran L. Polycystic ovary syndrome: a complex condition with psychological, reproductive and metabolic manifestations that impacts on health across the lifespan. *BMC Med.* 2010; 8: 41.
- Balen AH, Rutherford AJ. Managing anovulatory infertility and polycystic ovary syndrome. *BMJ.* 2007; 335(7621): 663-666.
- Qiao J, Feng HL. Extra-and intra-ovarian factors in polycystic ovary syndrome: impact on oocyte maturation and embryo developmental competence. *Hum Reprod Update.* 2011; 17(1): 17-33.
- Gilchrist RB, Lane M, Thompson JG. Oocyte-secreted factors: regulators of cumulus cell function and oocyte quality. *Hum Reprod Update.* 2008; 14(2): 159-177.
- Knight PG, Glister C. TGF- $\beta$  superfamily members and ovarian follicle development. *Reproduction.* 2006; 132(2): 191-206.
- Celestino JJH, Matos MHT, Saraiva MVA, Figueiredo JR. Regulation of ovarian folliculogenesis by Kit Ligand and the c-Kit system in mammals. *Anim Reprod.* 2009; 6(3): 431-439.
- Tuck AR, Robker RL, Norman RJ, Tilley WD, Hickey TE. Expression and localisation of c-kit and KITL in the adult human ovary. *J Ovarian Res.* 2015; 8: 31.
- Teixeira Filho FL, Baracat EC, Lee TH, Suh CS, Matsui M, Chang RJ, et al. Aberrant expression of growth differentiation factor-9 in oocytes of women with polycystic ovary syndrome. *J Clin Endocrinol Metab.* 2002; 87(3): 1337-1344.
- Zhao SY, Qiao J, Chen YJ, Liu P, Li J, Yan J. Expression of growth differentiation factor-9 and bone morphogenetic protein-15 in oocytes and cumulus granulosa cells of patients with polycystic ovary syndrome. *Fertil Steril.* 2010; 94(1): 261-267.
- Persani L, Rossetti R, Di Pasquale E, Cacciatore C, Fabre S. The fundamental role of bone morphogenetic protein 15 in ovarian function and its involvement in female fertility disorders. *Hum Reprod Update.* 2014; 20(6): 869-883.
- Palomba S, Falbo A, La Sala GB. Metformin and gonadotropins for ovulation induction in patients with polycystic ovary syndrome: a systematic review with meta-analysis of randomized controlled trials. *Reprod Biol Endocrinol.* 2014; 12: 3.
- Lord JM, Flight IH, Norman RJ. Metformin in polycystic ovary syndrome: systematic review and meta-analysis. *BMJ.* 2003; 327(7421): 951-953.
- Kjøtrød SB, von Düring V, Carlsen SM. Metformin treatment before IVF/ICSI in women with polycystic ovary syndrome: a prospective, randomized, double blind study. *Hum Reprod.* 2004; 19(6): 1315-1322.
- Tang T, Glanville J, Orsi N, Barth JH, Balen AH. The use of metformin for women with PCOS undergoing IVF treatment. *Hum Reprod.* 2006; 21(6): 1416-1425.
- Liu J, Liu M, Ye X, Liu K, Huang J, Wang L, et al. Delay in oocyte aging in mice by the antioxidant N-acetyl-L-cysteine (NAC). *Hum Reprod.* 2012; 27(5): 1411-1420.
- Fabbri R, Montanaro L, Pasquinelli G, Magnani V, Tamburini F, Takeuchi T, et al. Effects of N-Acetyl-Cystein on GDF-9 expression and BCL-2/BAX ratio in long-term culture of frozen/thawed human ovarian tissue. 25<sup>th</sup> Annual Meeting of ESHRE; 2009 Jun-Jul 28-1; Amsterda; the Netherlands; 2009.
- Mahmoodi M, Soleimani Mehranjani M, Shariatzadeh SM, Eimani H, Shahverdi A. N-acetylcysteine improves function and follicular survival in mice ovarian grafts through inhibition of oxidative stress. *Reprod Biomed Online.* 2015; 30(1): 101-110.
- Sacchinelli A, Venturella R, Lico D, Di Cello A, Lucia A, Rania E, et al. The efficacy of inositol and N-Acetyl cysteine administration (Ovaric HP) in improving the ovarian function in infertile women with PCOS with or without insulin resistance. *Obstet Gynecol Int J.* 2014; 2014: 1-5.
- Cheraghi E, Mehranjani MS, Shariatzadeh MA, Esfahani MH, Ebrahimi Z. N-Acetylcysteine improves oocyte and embryo quality in polycystic ovary syndrome patients undergoing intracytoplasmic sperm injection: an alternative to metformin. *Reprod Fertil Dev.* 2016; 28(6): 723-731.
- Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003. consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril.* 2004; 81(1): 19-25.
- Fulghesu AM, Ciampelli M, Muzj G, Belosi C, Selvaggi L, Ayala GF, et al. N-acetyl-cysteine treatment improves insulin sensitivity in women with polycystic ovary syndrome. *Fertil Steril.* 2002; 77(6): 1128-1135.
- Elnashar A, Fahmy M, Mansour A, Ibrahim K. N-acetyl cysteine vs. metformin in treatment of clomiphene citrate-resistant polycystic ovary syndrome: a prospective randomized controlled study. *Fertil Steril.* 2007; 88(2): 406-409.
- Oner G, Muderris II. Clinical, endocrine and metabolic effects of metformin vs N-acetyl-cysteine in women with polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol.* 2011; 159(1): 127-131.
- Wei LN, Liang XY, Fang C, Zhang MF. Abnormal expression of growth differentiation factor 9 and bone morphogenetic protein 15 in stimulated oocytes during maturation from women with polycystic ovary syndrome. *Fertil Steril.* 2011; 96(2): 464-468.
- Dumesic DA, Abbott DH. Implications of Polycystic ovary syndrome (pcos) on oocyte development. *Semin Reprod Med.* 2008; 26(1): 53-61.
- Surcel M, Stamatian F. Folliculogenesis disturbances within the polycystic ovarian syndrome and possible consequences on oocyte quality. *Acta Endocrinologica (Buc).* 2012; 8(2): 267-287.
- Shimasaki S, Moore RK, Otsuka F, Erickson GF. The bone morphogenetic protein system in mammalian reproduction. *Endocr Rev.* 2004; 25(1): 72-101.
- Wu YT, Tang L, Cai J, Lu XE, Xu J, Zhu XM, et al. High bone morphogenetic protein-15 level in follicular fluid is associated with high quality oocyte and subsequent embryonic development. *Hum Reprod.* 2007; 22(6): 1526-1531.
- Wei LN, Li LL, Fang C, Huang R, Liang XY. Inhibitory effects of controlled ovarian stimulation on the expression of GDF9 and BMP15 in oocytes from women with PCOS. *J Assist Reprod Genet.* 2013; 30(10): 1313-1318.
- Otsuka F, McTavish KJ, Shimasaki S. Integral role of GDF-9 and BMP-15 in ovarian function. *Mol Reprod Dev.* 2011; 78(1): 9-21.
- Brankin V, Hunter MG, Horan TL, Armstrong DG, Webb R. The expression patterns of Mrna-encoding stem cell factor, internal stem cell factor and c-kit in the prepubertal and adult porcine ovary. *J Anat.* 2004; 205(5): 393-403.
- Hutt K, McLaughlin EA, Holland MK. Kit ligand and c-Kit have diverse roles during mammalian oogenesis and folliculogenesis. *Mol Hum Reprod.* 2006; 12(2): 61-69.
- De Flora S, Izzotti A, D'Agostini F, Balansky RM. Mechanisms of N-acetylcysteine in the prevention of DNA damage and cancer, with special reference to smoking-related end-points. *Carcinogenesis.* 2001; 22(7): 999-1013.
- Liu K. Stem cell factor (SCF)-kit mediated phosphatidylinositol 3 (PI3) kinase signaling during mammalian oocyte growth and early follicular development. *Front Biosci.* 2006; 11: 126-135.
- Tanikawa M, Harada T, Mitsunari M, Onohara Y, Iwabe T, Terakawa N. Expression of c-kit messenger ribonucleic acid in human oocyte and presence of soluble c-kit in follicular fluid. *J Clin Endocrinol Metab.* 1998; 83(4): 1239-1242.
- Desforges-Bullet V, Gallo C, Lefebvre C, Pigny P, Dewailly D, Catteau-Jonard S. Increased anti-Müllerian hormone and decreased FSH levels in follicular fluid obtained in women with polycystic ovaries at the time of follicle puncture for in vitro fertilization. *Fertil Steril.* 2010; 94(1): 198-204.