


ORIGINAL RESEARCH

The effect of test modality on dynamic exercise biomarkers in children, adolescents, and young adults

Ronen Bar-Yoseph¹ , Janos Porszasz², Shlomit Radom-Aizik¹, Kim D. Lu¹, Annamarie Stehli¹, Pearl Law¹ & Dan M. Cooper^{1,3}

1 Pediatric Exercise and Genomics Research Center (PERC), Department of Pediatrics, University of California Irvine, Irvine, California

2 Rehabilitation Clinical Trials Center, Division of Respiratory and Critical Care Physiology and Medicine, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, California

3 University of California Irvine Institute for Clinical and Translational Science, Irvine, California

Keywords

Cycle ergometry, peak oxygen uptake (\dot{V}), puberty, submaximal biomarkers of fitness, treadmill.

Correspondence

Ronen Bar-Yoseph, Pediatric Exercise and Genomics Research Center, 101 Academy Drive, Irvine, CA 92617.
Tel: +972-50-206-3131/+972-4777-4360
Fax: +972-4-777-4395
E-mail: r_bar-yoseph@rambam.health.gov.il

Funding Information

This work was supported in part by National Institutes of Health (NIH) P01HD-048721 grant, Pediatric Exercise and Genomic Research Center (PERC) Systems Biology Research Fund, National Center for Advancing Translational Sciences (NCATS) UL1TR001414 grant and NCATS CTSA Collaboration Innovation Award, Project REACH (Revamping Exercise Assessments in Child Health), UL1TR002004.

Received: 2 June 2019; Revised: 23 June 2019; Accepted: 24 June 2019

doi: 10.14814/phy2.14178

Physiol Rep. 7 (14), 2019, e14178.
<https://doi.org/10.14814/phy2.14178>

Correction added on 7 September 2019, after first online publication: New author included

Introduction

The goal of this study was to test hypotheses focused on the effect of the two most common exercise testing modalities, cycle ergometry (CE), and treadmill (TM), on

Abstract

Cardiopulmonary exercise testing (CPET) modalities, treadmill (TM), and cycle ergometer (CE), influence maximal gas exchange and heart rate (HR) responses. Little is known regarding CPET modality effect on submaximal biomarkers during childhood and adolescence. Ninety-four healthy participants (7–34 y.o., 53% female) performed TM and CE CPET to address two major gaps: (1) the effect of modality on submaximal CPET biomarkers, and (2) estimation of work rate in TM CPET. Breath-by-breath gas exchange enabled calculation of linear regression slopes such as $\dot{V}O_2/\Delta HR$ and $\Delta \dot{V}E/\Delta \dot{V}CO_2$. Lean body mass (LBM) was measured with dual X-ray absorptiometry. We tested a novel TM CPET estimate of work rate based on TM velocity², incline, and body mass (VIM). Like the linear relationship between $\dot{V}O_2$ and work rate in CE CPET, $\dot{V}O_2$ increased linearly with TM VIM. TM $\Delta \dot{V}O_2/\Delta HR$ was highly correlated with CE ($r = 0.92$), and each increased substantially with LBM ($P < 0.0001$ for TM and CE). $\Delta \dot{V}O_2/\Delta HR$ was to a small (~8.7%) but significant extent larger in TM (1.6 mL/min/beat, $P = 0.04$). In contrast, TM and CE $\Delta \dot{V}E/\Delta \dot{V}CO_2$ decreased significantly with LBM, supporting earlier observations from CE CPET. For both CE and TM, males had significantly higher $\Delta \dot{V}O_2/\Delta HR$ but lower $\Delta \dot{V}E/\Delta \dot{V}CO_2$ than females. Novel TM CPET biomarkers such as $\Delta VIM/\Delta HR$ and $\Delta \dot{V}O_2/\Delta VIM$ paralleled effects of LBM observed in CE CPET. TM and CE CPET submaximal biomarkers are not interchangeable, but similarly reflect maturation during critical periods. CPET analysis that utilizes data actually measured (rather than estimated) may improve the clinical value of TM and CE CPET.

cardiopulmonary exercise testing (CPET) results in children, adolescents, and young adults. We addressed two key challenges in comparing TM and CE CPET among children, adolescents, and young adults: (1) the difficulty in quantifying the work performed in TM CPET, and (2) useful

approaches to scaling CPET results when body size and physiologic function change so dramatically over the course of growth and development (Cooper et al., 1987; Cooper et al., 2014). CPET biomarkers are used to assess disease severity, progress, and response to therapy (including exercise prescriptions) across an expanding range of childhood diseases and conditions and across the lifespan (Ploeger et al., 2009; Pahkala et al., 2013; Liem et al., 2015; Sule and Fontaine, 2016; Cordingley et al., 2016; Gualano et al., 2017; Li et al., 2017). Despite these factors, CPET has failed to fulfil its promise in child health research and clinical practice (Ashish et al., 2015). A major barrier to more accurate and effective clinical use of CPET in children and adults has been a lack of harmonization of protocol types and exercise modalities (Ashish et al., 2015), factors that influence CPET results (Fredriksen et al., 1998; Beltrami et al., 2012; Bires et al., 2013; May et al., 2014; Cunha et al., 2015). For example, in clinical trials involving CPET in children over the past five years, a PubMed search revealed 40 published studies that used CE and 113 that used TM.

We concentrated on dynamic submaximal physiologic output variables (Fig. 1) that are less effort-dependent than the traditional $\dot{V}O_2\text{max}$ and, arguably, more acceptable in children and adolescents, particularly those with chronic diseases or conditions (Stein et al., 2003; Cooper et al., 2014). Dynamic relationships among CPET variables (such as HR, $\dot{V}E$, $\dot{V}CO_2$, and $\dot{V}O_2$) reveal novel insights into cardiorespiratory function in health and disease (Cooper et al., 1984; Troutman et al., 1998; Moser et al., 2000; Chen et al., 2014; Elbehairy et al., 2015; Hestnes et al., 2017), and can be obtained in both CE and TM modalities without necessarily measuring work rate. The effect of exercise modality on submaximal physiologic output variables has not been adequately studied in children and adolescents.

In CE, the work rate is usually measured directly from the known resistance on the ergometer's flywheel. $\dot{V}O_2$ is linearly related to work rate (Whipp et al., 1981) and, as a consequence, CE protocols can be easily designed to produce a linear relationship between protocol duration (i.e., exercise time) and $\dot{V}O_2$, which simplifies the ultimate analysis of CPET data. In contrast, it is difficult to estimate the relationship between work rate and $\dot{V}O_2$ in TM because of the complexity of both the physics and human mechanical efficiency of treadmill walking and running (Workman and Armstrong, 1963; Kyröläinen et al., 1995; Porszasz et al., 2003; Keir et al., 2012; Azuma, 2014). Treadmill work is determined by kinetic energy [functions of the velocity (V) of the TM and the body mass (M) of the participant] and work against gravity imposed by the TM grade or incline (Ruckstuhl et al., 2010). While body weight is often used to scale CPET values in an effort to compare results among individuals of different size, recent data suggest that LBM is significantly better correlated to size-dependent CPET biomarkers (Cooper et al., 2014). Lean body mass (LBM) is




Figure 1. Overview – Interoperability challenge in CE and TM CPET. Some values are comparable (as $\Delta\dot{V}E/\Delta\dot{V}CO_2$ and $\Delta\dot{V}O_2/\Delta HR$) while others depend on accurate measurement of work rate ($\Delta\dot{V}O_2/\Delta WR$ and $\Delta HR/\Delta WR$) available only for the CE.

a more direct measure of skeletal muscle, the predominant metabolizing tissue in exercise, than body weight. Consequently, we measured LBM using dual X-ray absorptiometry DXA (Bridge et al., 2011), and used LBM in our comparison of the two CPET modalities.

Although the magnitude of peak or maximal $\dot{V}O_2$ is similar in CE and TM CPET, one observation made consistently in both children and adults is that peak $\dot{V}O_2$ tends to be somewhat greater in TM CPET (Turley and Wilmore, 1997). We hypothesized that CPET slope variables would reflect exercise modality differences as well. We further hypothesized that the metabolic response to TM is proportional to the kinetic energy exerted on the center of gravity of the body, therefore an estimate of TM work rate was calculated from body mass, V^2 , and TM incline would be linearly related to $\dot{V}O_2$.

Methods

Participants

The study was approved by the UC Irvine Institutional Review Board. Inclusion criteria included healthy 7–

35 y.o. participants (53% female) without any known respiratory, cardiac, or metabolic disease, and not taking any chronic prescribed medication. BMI of each participant was less than the 95th percentile for children (<http://www.cdc.gov/healthyweight/assessing/bmi/chi>) and BMI less than 30 for adults. Each volunteer visited the laboratory on two occasions. During the first visit, informed consent was obtained (parental consent + child assent for participants <18 y/o), demographic and anthropometric data were recorded, Tanner stage (by questionnaire) was assessed, and DXA was performed. A maximal progressive exercise protocol on a CE was also done on the first visit. At least 48 h later, each participant performed TM CPET with a protocol designed to mimic the rate of power increase found in the CE CPET.

Anthropometric measurement and body composition

Standard calibrated scales and stadiometers were used to determine weight and height. Body composition, including LBM, fat mass, and percent body fat were determined by DXA using a Hologic QDR 4500 densitometer. Participants were scanned while lying supine and wearing light clothing. On the day of each test, the DXA instrument was calibrated using the procedures provided by the manufacturer, and DXA scans were performed and analyzed using pediatric software where appropriate.

CE and TM protocols

The CE protocol consisted of a ramp-type progressive cycle ergometry used previously in this and other laboratories to measure peak $\dot{V}O_2$ in children and adults (Cooper et al., 2016). After a 2-min period of unloaded cycling (0 W), power output was increased by 8–30 W/min. The increase in the ramp W/min was individualized for each participant empirically from the following formula:

$$\text{work rate increment} = (\text{bodyweight} * 3)/10 \quad (1)$$

where work rate increment is in watts per minute and body weight in kg. This formula was derived empirically from the thousands of tests we have performed and is fairly reliable in producing CPET duration of 8–15 min, an interval previously determined to optimize evaluation of breath-by-breath data (Buchfuhrer et al., 1983; Myers and Froelicher, 1993). Participants cycled at a constant pedaling rate between 60 and 70 revolutions per minute (rpm) throughout the test on an electronically braked, servo-controlled cycle ergometer. The increasing work rate was discontinued when the participants indicated that they had reached the limit of their tolerance and/or a drop occurred in

pedaling rate below 60 rpm despite strong verbal encouragement. At this point, the work rate was lowered to 0 W and the participants continued to pedal for at least 5 more minutes while lowering the pedaling rate to below 40 rpm in order to prevent an excessively sudden drop in blood pressure (Kenney and Seals, 1993).

Since our goal was to compare the two modalities using tests in which $\dot{V}O_2$ increased linearly over a duration of about 8–15 min, we used the results of the CE CPET (in which work rate was precisely known) to guide the velocity and incline configuration for the TM CPET performed on a separate data. Our coauthor (Dr. Porszasz) and coworkers (Porszasz et al., 2003) previously developed a TM CPET protocol that linearized the $\dot{V}O_2$ increase over a 10–15 min exercise duration. They found that a protocol combining an initial slow walking speed that progressively increased in concert with a dynamically changing incline met the demands of an initially low exercise metabolic rate and optimum test duration.

In designing the protocol, we first determined the desired work rate for each of the 1-min steps assuming a linear increase both in work rate and speed. The speed-range we used was between 0.5 and 10.5 miles per hour (0.8 km/h and 16.8 km/h, respectively); baseline speed was set to 0.5 mph and every minute was increased by 0.5 mph up to 10.5 mph max. Each step had the same work rate and speed increments; having formulated these, the inclination was determined by the following formula:

$$\text{TM inclination angle} = \arctan[\text{work rate}/(\text{mass} \times g \times v^2)] \quad (2)$$

where inclination angle is in radians; work rate in watts, mass in kg, g as 9.81m/sec^2 , and v in m/sec. The inclination as percentage was then determined and set for each step in the protocol. This resulted in a decreasing incline profile (set individually, with 30% as the highest incline in the cohort), which approached 1% toward the end of the test. The general formula [Equation (2)] was used to calculate the desired change in TM incline to produce a given work rate at a particular velocity and work rate (Porszasz et al., 2003). We assumed that the work rate in TM CPET would be related to the kinetic energy equation, $\text{work} = \frac{1}{2}mv^2$ and the following equation was used to estimate TM work rate:

$$\text{WR(TM)} = k \cdot \text{mass} \cdot v^2 \cdot (I + 1) \quad (3)$$

where mass is body weight (kg), v is treadmill speed (m/sec), I is incline (%), and k is a conversion factor constant. We arbitrarily used the expression $I + 1$ because, at 0% incline, work rate would not be possible to calculate due to multiplication by zero. For convenience, we refer to $k \cdot \text{mass} \cdot v^2 \cdot (I + 1)$ as VIM. Using these equations and

protocol, we achieved a largely linear relationship between $\dot{V}O_2$ and exercise work intensity up to a reasonable level and the test duration was between 8 and 15 min.

Gas exchange measurement

Gas exchange was measured breath-by-breath using the SensorMedics metabolic system (Vmax Encore 229, Yorba Linda, CA). The breath-by-breath gas exchange data were interpolated to 1-sec and 10-sec bin averages were formed and used for all later analyses. Physiologically abnormal data for HR and gas exchange (e.g., HR < 50 beat/min or >230 beat/min, or $\dot{V}O_2 = 0$ L/min or >5 L/min) and outliers, based on each subject, are occasionally observed in breath-by-breath CPET data obtained in children. These data were identified and excluded for slope or peak calculation.

Calculation of submaximal CPET slopes and peak values

Submaximal slopes ($\Delta\dot{V}O_2/\Delta HR$, $\Delta\dot{V}E/\Delta\dot{V}CO_2$, etc.) were calculated using standard linear regression as described previously (Cooper et al., 2014) omitting the first minute and the last 30 sec of the exercise. The peak values were taken as the highest values in 20-sec bin averages over the last 2 min of exercise. There is currently no validated, universally accepted approach for the determination of peak $\dot{V}O_2$ in children. We used a criterion implemented in a large study by Rowland et al. (2008) defined by inability to maintain the pedaling cadence in association with subjective evidence of fatigue (sweating, hyperpnea) and HR >185 bpm (children) or >170 bpm (adults) and/or respiratory exchange ratio (RER, $\dot{V}CO_2/\dot{V}O_2$) >1.00 (children) or >1.10 (adults).

Comparing CE and TM work rate input

We assumed that the linear relationship between work rate and $\dot{V}O_2$ (Whipp et al., 1981) in both exercise modalities have the same slope and intercept, we used the linear regression parameters for the measured $\dot{V}O_2$ on the CE (in which work rate was known) to determine the work rate on the TM (in which the velocity, incline, and participant's weight were known) for each participant. The equivalent work rate on TM exercise was marked as WR'. The above is described adequately by the following two equations:

$$\begin{aligned}\dot{V}O_2(\text{CE}) &= a \cdot \text{WR}(\text{CE}) + b \\ \text{and} \\ \text{WR}' &= [\dot{V}O_2(\text{TM}) - b]/a\end{aligned}$$

We used standard linear regression techniques to estimate the parameters 'a' and 'b' for each participant and calculated the WR' for TM CPET.

Comparison of fitness variables obtained from CE CPET and TM CPET

We compared the CPET variables described above obtained from the two modalities. To eliminate any confounding effect introduced by the estimation of WR' (due to its dependence on the $\dot{V}O_2$ -WR relationship derived from CPET-CE), we also compared CPET values using VIM itself. In addition, we tested the degree to which CE and TM CPET variables scaled to body mass and composition, factors essential for understanding CPET in the growing child. For example, while the numerical value of $\Delta WR/\Delta HR$ and $\Delta VIM/\Delta HR$ will be quite different, we expected that their relationship and correlation to key variables such as age, body weight, lean body mass (LBM), and sex would be quite similar when comparing the two modalities.

Statistical analysis

For each peak $\dot{V}O_2$, $\Delta\dot{V}O_2/\Delta HR$ slope, $\Delta\dot{V}E/\Delta\dot{V}CO_2$ slope, and other slopes, statistical comparisons of CE versus TM were performed using mixed models (via SAS PROC MIXED) to account for subject level intercorrelation between the paired modality measurements (TM and CE). Each model also included puberty group (children tanner 1–2, adolescents 4–5, adults >18 years), sex, puberty \times sex, puberty group \times modality interaction, sex \times modality interaction, and puberty group \times sex \times modality interaction. Post hoc comparisons of model-generated, least-square (LS) means were evaluated where significant fixed effects were found. This was done according to the hierarchy principle such that if an interaction was present, only the appropriate conditional means were compared and interpreted. Significance for the post hoc comparisons was determined by Tukey-adjusted *P*-values of the LS mean differences. We performed a standard Bland-Altman (BA) analysis to compare to peak $\dot{V}O_2$ and submaximal slopes between the two modalities.

Results

Participants characteristics

Representative examples of CE and TM CPET are shown in Figure 2. A total of 111 healthy children and young adults (7–34 y.o.) participated in this study. We excluded 17 of them from the final analysis: five due to technical problems, three due to incompleteness of the study protocol, two due to a submaximal effort on the TM, six due to inability to assess the Tanner score, and one due to exercise-induced bronchoconstriction following CE ramp test. Ninety-four participants were




Figure 2. (A) Relationship between $\dot{V}O_2$ and work rate in early (12 y/o) boy and late (17 y/o) pubertal boy in CE CPET. (B) Relationship between $\dot{V}O_2$ and $V \times I \times M$ (see text) in TM CPET in the two participants. (C) Relationship between $\dot{V}O_2$ and $V^2 \times I \times M$ (see text) in TM CPET in the two participants. By using this approach, we were able to linearize the relationship between oxygen uptake and velocity, incline and mass.

included in the analysis, and demographic and anthropometric data are presented in Table 1. For analysis of peak $\dot{V}O_2$, data from 88 participants were analyzed (six

participants did not meet the criteria for a maximal test as noted above). Submaximal slopes, and peak CPET values are shown in Tables 2, 3, 4. Detailed summary of statistical analyses are shown in Tables 5 and 6.

Linearity of $\dot{V}O_2$ with work rate estimate and exercise duration in the two modalities

Corroborating the previous work by Porszasz et al., (2003, we achieved success in linearizing the relationship between $\dot{V}O_2$ and exercise duration (time). This was evidenced both by visual inspection of the exercise tests (e.g., Fig. 2) as well as by the remarkably high correlation between $\dot{V}O_2$ and time for the TM CPET (mean r was 0.978). We compared the correlation coefficients of two linear regressions: $\dot{V}O_2$ versus $v \times (I + 1) \times \text{mass}$ and $v^2 \times (I + 1) \times \text{mass}$. For the former, the average $R_1^2 = 0.7889$ and for the latter $R_s^2 = 0.9255$. A paired t -test for the mean difference was statistically significant [$\Delta (R_s^2 - R_1^2) = 0.1367$, $P < 0.00001$], suggesting stronger prediction of $\dot{V}O_2$ by V^2IM . The duration of exercise for TM and CE modality in each group of participants is shown in Table 4. For the adolescents and young adults, TM duration was significantly longer than CE. In general, exercise duration for both modalities was longer in males than in females.

Submaximal CPET variables

$\Delta \dot{V}O_2 / \Delta HR$

$\Delta \dot{V}O_2 / \Delta HR$ results are shown in Figure 3 and Table 2. The CE and TM values were highly correlated ($P < 0.0001$, Fig. 3A). A small but significantly higher mean $\Delta \dot{V}O_2 / \Delta HR$ difference (1.7 ± 0.81 mL/beat, about 10%) was found in TM versus CE. For the group as a whole, BA analysis revealed higher $\Delta \dot{V}O_2 / \Delta HR$ for TM CPET (bias of 1.74, 95% CI of 1.06 to 2.42). Statistically significant maturation-dependent differences were observed in both males and females (Table 6). Consistent with our previous study of cycle ergometer exercise in children and adolescents (Cooper et al., 2014), $\Delta \dot{V}O_2 / \Delta HR$ increased with LBM ($r = 0.88$, $P < 0.0001$). For both CE and TM (Fig. 3B), peak $\dot{V}O_2$ was highly correlated to $\Delta \dot{V}O_2 / \Delta HR$. We used the linear regression equations relating $\Delta \dot{V}O_2 / \Delta HR$ and LBM to calculate a predicted value for each participant, then compared the percent predicted from the CE and TM CPET to determine how interchangeable the two modalities were. A moderate correlation was found (correlation coefficient $r = 0.66$, $P < 0.0001$, Fig. 3C).

Table 1. Participant characteristics.

Group	Children	Children	Adolescent	Adolescent	Adult	Adult
Sex	M	F	M	F	M	F
Tanner	1–2	1–2	4–5	4–5	NA	NA
N	18	16	18	23	8	11
Age (year)	10.8 ± 1.7	8.8 ± 1.1	16.9 ± 1.4	15.5 ± 1.8	29.0 ± 2.3	26.3 ± 4.6
Height (cm)	144.0 ± 11.8	132.1 ± 9.4	172.4 ± 7.0	161.4 ± 6.0	178.8 ± 9.0	161.8 ± 4.8
Total body mass (kg)	37.4 ± 12.0	29.7 ± 8.0	62.4 ± 9.4	54.8 ± 8.4	79.5 ± 8.3	58.1 ± 7.6
Lean body mass (kg)	25.2 ± 6.5	19.0 ± 3.8	47.3 ± 6.5	35.4 ± 4.6	59.5 ± 7.0	38.4 ± 5.0
% Body fat	29.0 ± 5.9	32.6 ± 5.6	21.3 ± 5.0	32.6 ± 5.2	22.5 ± 5.2	31.2 ± 4.3
BMI (kg/m ²)	17.5 ± 3.0	16.7 ± 2.4	17.2 ± 3.3	20.9 ± 2.2	24.9 ± 1.9	22.1 ± 2.3
BMI percentile	47.3 ± 30.7	49.4 ± 29.0	49.9 ± 29.4	43.1 ± 23.5	N/A	N/A
Ethnicity (Hispanic or Latino)	0	1	1	2	0	0
Race (White/Asian/African-American)	16/1/1	13/2/1	8/10/0	16/7/0	5/3/0	7/3/1

M, Male, F, Female, BMI, body mass index.
Data are presented as mean ± SD.

Table 2. Submaximal slopes of CPET variables obtainable from both CE and TM modalities.

Group	Children	Children	Adolescent	Adolescent	Adult	Adult
Sex	M	F	M	F	M	F
CE $\Delta\dot{V}E/\Delta\dot{V}CO_2$	31.4 ± 3.5	31.2 ± 2.7	27.9 ± 4.0	30.7 ± 4.0	27.2 ± 3.9	29.4 ± 5.4
TM $\Delta\dot{V}E/\Delta\dot{V}CO_2$	30.6 ± 2.2	29.8 ± 2.7	27.9 ± 2.2	28.7 ± 2.5	25.2 ± 3.7	28.9 ± 3.7
CE $\Delta\dot{V}O_2/\Delta HR$ (mL/beat)	13.7 ± 4.1	9.4 ± 1.9	26.0 ± 6.4	15.8 ± 4.8	29.2 ± 8.0	21.2 ± 5.9
TM $\Delta\dot{V}O_2/\Delta HR$ (mL/beat)	14.9 ± 4.6	10.8 ± 2.6	27.2 ± 7.1	18.8 ± 4.27	31.7 ± 7.6	22.0 ± 5.0

Data are presented as mean ± SD.

Table 3. Similar maturation and sex patterns in heartrate response to work rate in CE CPET (work rate actually measured) and TM CPET (work rate estimated by WR' and VIM).

Group	Children	Children	Adolescent	Adolescent	Adult	Adult
Sex	M	F	M	F	M	F
CE $\Delta WR/\Delta HR$ (watts/beat/min)	1.17 ± 0.41	0.77 ± 0.19	2.33 ± 0.49	1.58 ± 0.34	2.61 ± 0.62	2.09 ± 0.61
TM $\Delta WR'/\Delta HR$ (watts/beat/min)	1.29 ± 0.48	0.90 ± 0.27	2.46 ± 0.55	1.92 ± 0.43	2.85 ± 0.61	2.20 ± 0.64
TM $\Delta VIM/\Delta HR$	123.9 ± 43.4	85.9 ± 22.7	334.2 ± 130.2	198.1 ± 75.5	368.9 ± 110.6	245.1 ± 78.9

Data are presented as mean ± SD.

For CE WR-HR and TM WR'-HR, the sex difference is significant for all three puberty groups. For VIM-HR the sex difference is significant for adolescents and adults, but not children. In males, all three measures, demonstrate a puberty effect with adolescents and adults having larger slopes than children. Adolescents and adult males are not significantly different from each other. In females all three measures also show adolescents and adults having larger slopes than children. In addition, CE WR-HR and TM VIM-HR show adult females having significantly larger slopes than adolescent females, but this was not the case for TM WR'-HR.

$\Delta\dot{V}E/\Delta\dot{V}CO_2$

$\Delta\dot{V}E/\Delta\dot{V}CO_2$ results are shown in Figure 4 and Table 2. The CE and TM values were significantly correlated ($P < 0.0001$, Fig. 4A). A small but significantly higher mean $\Delta\dot{V}E/\Delta\dot{V}CO_2$ difference (1.151 ± 0.527 , about 3.7%) was found in CE vs. TM. For the group as a

whole, BA analysis revealed lower $\Delta\dot{V}E/\Delta\dot{V}CO_2$ in TM CPET (bias of -1.13 , 95%CI of -1.79 to -0.47). In the males only, the values were significantly greater ($P < 0.0001$) in the children compared to the adolescents and the adults. For the participant population as a whole, $\Delta\dot{V}E/\Delta\dot{V}CO_2$ was inversely correlated with LBM ($P < 0.0001$, Fig. 4B).

Table 4. Peak $\dot{V}O_2$ and HR from CE and TM modalities.

Group	Children	Children	Adolescent	Adolescent	Adult	Adult
Sex	M	F	M	F	M	F
CE peak $\dot{V}O_2$ (L/min)	1.89 ± 0.47	1.31 ± 0.27	3.45 ± 0.69	2.20 ± 0.53	3.81 ± 0.87	2.52 ± 0.63
CE peak $\dot{V}O_2$ /weight (mL/kg/min)	51.5 ± 7.2	45.0 ± 7.3	54.3 ± 6.8	40.5 ± 8.0	47.8 ± 9.0	43.4 ± 10.0
CE peak $\dot{V}O_2$ /LBM (mL/kg/min)	74.9 ± 6.1	70.1 ± 6.7	71.0 ± 7.2	62.2 ± 10.6	63.7 ± 10.4	62.2 ± 11.9
TM peak $\dot{V}O_2$ (L/min)	1.93 ± 0.56	1.37 ± 0.27	3.42 ± 0.48	2.41 ± 0.49	4.29 ± 0.64	2.64 ± 0.43
TM peak $\dot{V}O_2$ /weight (mL/kg/min)	52.2 ± 8.1	47.3 ± 7.7	55.5 ± 7.3	44.3 ± 7.6	54.0 ± 5.7	45.9 ± 8.2
TM peak $\dot{V}O_2$ /LBM (mL/kg/min)	76.1 ± 8.9	73.2 ± 9.1	72.6 ± 7.5	68.0 ± 8.9	72.1 ± 5.8	69.0 ± 9.2
CE peak HR (beats/min)	199.8 ± 7.7	194.5 ± 9.9	197.6 ± 11.1	193.3 ± 8.5	189.3 ± 10.9	183.9 ± 8.8
TM peak HR (beats/min)	195.8 ± 7.0	193.3 ± 9.8	198.8 ± 8.6	197.3 ± 7.5	193.3 ± 12.1	187.5 ± 9.2
CE protocol duration	669.4 ± 72.7	577.9 ± 118.1	691.6 ± 97.2	601.6 ± 70.0	689.8 ± 66.8	604.9 ± 62.2
TM protocol duration	700.3 ± 105.1	581.2 ± 92.1	951.0 ± 153.7	785.8 ± 135.4	1013.1 ± 140.8	805.9 ± 169.7

Data are presented as mean ± SD.

Table 5. Fixed effects from mixed model analysis.

CPET variable	$\Delta\dot{V}O_2/\Delta HR$		$\Delta\dot{V}E/\Delta\dot{V}CO_2$		Peak $\dot{V}O_2$ /weight		Peak $\dot{V}O_2$ /LBM	
	F-stat	P	F-stat	P	F-stat	P	F-stat	P
Puberty	107.15	<0.0001	12.03	<0.0001	0.28	0.7549	5.13	0.0078
Sex	85.50	<0.0001	8.16	0.0053	44.09	<0.0001	9.88	0.0023
Modality	4.34	0.0402	4.77	0.0317	5.10	0.0266	7.04	0.0095
Puberty × sex	4.89	0.0098	4.15	0.019	3.90	0.024	2.26	0.1108
Puberty × modality	0.09	0.9098	0.02	0.9816	0.37	0.6907	0.71	0.4968
Sex × modality	0.00	0.9456	0.09	0.7664	0.00	0.9543	0.22	0.6406
Puberty × Sex × Modality	0.33	0.7227	0.78	0.4597	0.53	0.5881	0.88	0.4177

Sex effects on $\Delta\dot{V}O_2/\Delta HR$ and $\Delta\dot{V}E/\Delta\dot{V}CO_2$

Both CE and TM CPET revealed significant sex effects. $\Delta\dot{V}O_2/\Delta HR$ was greater in males, and $\Delta\dot{V}E/\Delta\dot{V}CO_2$ was greater in females. The sex effect was not influenced by CPET modality.

Relationships between HR, WR, WR', and VIM

As shown in Figure 5, we found strong correlations between LBM and either $\Delta WR/\Delta HR$ from CE CPET or $\Delta WR'/\Delta HR$ from TM CPET (Fig. 5A). $\Delta VIM/\Delta HR$ from TM CPET was highly correlated to LBM and to $\Delta WR/\Delta HR$ from CE CPET (Fig. 5B and 5, respectively). As shown in Table 3, the WR, WR', and VIM relationships with HR all reflected comparable patterns within the sub-populations of the participants.

Peak $\dot{V}O_2$ comparison: CE vs. TM and relationship to submaximal CPET variables

Figure 6 and Table 4 shows the correlations between peak $\dot{V}O_2$ for the whole participant population from the two modalities. Peak $\dot{V}O_2$ was highly correlated between CE

and TM (Fig. 6A), and both CE and TM CPET peak $\dot{V}O_2$ demonstrated high correlation with LBM (correlations with weight were high, but not as high as with LBM, Fig. 6B). Overall, a small ($5.9 \pm 1.3\%$) but significantly higher mean peak $\dot{V}O_2$ difference was found in TM. For the group as a whole, BA analysis revealed higher peak $\dot{V}O_2$ for TM CPET (bias of 122 mL/min, 95% CI of 62–184 mL/min). However, within the puberty subgroups, there was no significant difference between CE and TM. Males had higher peak $\dot{V}O_2$ than females at all puberty levels. In males, there were no differences among puberty groups. In females, adolescents had the lowest mean values, statistically significant only younger ages.

We used the linear regression equations relating peak $\dot{V}O_2$ and LBM to calculate a predicted value for peak $\dot{V}O_2$, then compared the percent predicted from the CE and TM CPET to determine how interchangeable the two modalities were. A moderate correlation was found (Fig. 6C).

Discussion

For TM CPET, we were able to design a protocol that linearized the relationship of $\dot{V}O_2$ to both exercise duration

Table 6. Paired comparisons of mixed model least square means.

CPET variable	$\Delta\dot{V}O_2/\Delta HR$ (mL/beat)		$\Delta\dot{V}E/\Delta\dot{V}CO_2$		Peak $\dot{V}O_2$ per weight (mL/min/kg)		Peak $\dot{V}O_2$ per LBM (mL/min/kg)	
	Mean diff	<i>P</i>	Mean diff	<i>P</i>	Mean diff	<i>P</i>	Mean diff	<i>P</i>
Pubertal status (child = C; adolescent = ADO; adult = ADU)								
C vs. ADO	-9.725	<0.0001	2.110	0.0003	0.360	0.7875	3.674	0.0149
C vs. ADU	-13.824	<0.0001	3.105	<0.0001	1.201	0.457	5.334	0.0046
ADO vs. ADU	-4.099	0.0002	0.995	0.1436	0.841	0.5906	1.659	0.3545
Sex (M,F)								
M vs. F	7.448	<0.0001	-1.506	0.0053	8.153	<0.0001	4.380	0.0023
Modality								
CE vs. TM	-1.679	0.0402	1.151	0.0317	-2.773	0.0266	-3.697	0.0095
Pubertal status \times sex								
C: M vs. F	4.219	0.0009	0.539	0.5153	5.673	0.005	*	*
ADO: M vs. F	9.263	<0.0001	-2.130	0.0060	12.523	<0.0001	*	*
ADU: M vs. F	8.861	<0.0001	-2.926	0.0103	6.263	0.0159	*	*
M: C vs. ADO	-12.247	<0.0001	3.444	<0.0001	-3.064	0.1231	*	*
M: C vs. ADU	-16.144	<0.0001	4.837	<0.0001	0.907	0.7061	*	*
M: ADO vs. ADU	-3.898	0.0149	1.393	0.1759	3.971	0.0976	*	*
F: C vs. ADO	-7.203	<0.0001	0.776	0.3240	3.785	0.0384	*	*
F: C vs. ADU	-11.503	<0.0001	1.372	0.1483	1.496	0.4871	*	*
F: ADO vs. ADU	-4.299	0.0023	0.597	0.5000	-2.289	0.2607	*	*

*No significant interactions.

and an estimate of work rate using the participant's body weight and data easily obtained from TM CPET, namely TM speed and incline. The dynamic relationship between the novel VIM estimate of work rate and CPET variables like HR and $\dot{V}O_2$ paralleled the relationships we found using CE CPET in which work rate is measured directly. This can provide investigators with new tools to gauge fitness in children and adolescents using TM CPET. Dynamic submaximal CPET variables (such as $\Delta\dot{V}O_2/\Delta HR$ and $\Delta\dot{V}E/\Delta\dot{V}CO_2$) were highly correlated between the new linear TM and CE CPET protocols: this is the first attempt to analyze these submaximal CPET parameters in a cohort of children, adolescents, and young adults. Furthermore, we found that the relationship of these CPET results to critical exercise-response determinants such as body size were similar in both exercise modalities. Although the HR and gas exchange results of TM and CE exercise were comparable, our data corroborated previous work establishing that CPET TM peak $\dot{V}O_2$ is somewhat and significantly greater than CE CPET. We extended this finding to a submaximal CPET variable, $\Delta\dot{V}O_2/\Delta HR$. The dynamic submaximal relationship between $\dot{V}E$ and $\dot{V}CO_2$, ($\Delta\dot{V}E/\Delta\dot{V}CO_2$) was to a small but significant degree higher in CE CPET.

The mechanisms responsible for the larger $\Delta\dot{V}O_2/\Delta HR$ in TM CPET are not clearly evident. The Fick equation [$\dot{V}O_2 = HR \cdot SV \times (a - v)O_2$, where HR is heart rate, SV is stroke volume, and $(a - v)O_2$ is arteriovenous

oxygen content difference] indicates that a greater increase in $\dot{V}O_2$ per given change in HR can occur only as a result of a greater change in SV or widening of the arteriovenous O_2 concentration difference. A possible mechanism influencing stroke volume could be higher venous return and increased muscle mass involved in exercise during TM versus CE. We reanalyzed the data cited earlier from Turley et al. (1997) who measured $(a - v)O_2$ and SV indirectly and noninvasively in 24 children and 24 adults during both TM and CE progressive exercise. Interestingly, while we could find no systematic differences in SV between cycle and treadmill exercise, we did find that the average $(a - v)O_2$ during exercise was significantly ($P < 0.01$) higher in TM exercise (10.7/100 mL) compared with CE exercise (9.6/100 mL). Further studies will be needed to examine the matching of blood flow distribution in the exercising muscle to determine possible mechanisms leading to greater O_2 extraction during TM exercise, leading to the small but significant differences in the $\Delta\dot{V}O_2/\Delta HR$.

$\Delta\dot{V}E/\Delta\dot{V}CO_2$ values obtained from TM and CE CPET were correlated, but not as strongly as the $\Delta\dot{V}O_2/\Delta HR$ CPET variable (Fig. 3A, 4A). The relationship of $\dot{V}E$ to $\dot{V}CO_2$ during exercise reveals useful clinical information regarding respiratory dead space and the systemic set point of CO_2 concentration that ultimately modulates respiratory control centers in the brainstem and carotid bodies (Armon et al., 1991; Rausch et al., 1991). Clinical




Figure 3. Interoperability of $\Delta\dot{V}O_2/\Delta HR$ derived from CPET-CE and CPET-TM. (A) The slope of the linear regression equation was highly significant, 0.923 ± 0.0419 , $P < 0.0001$; the y -intercept, 3.11 ± 0.82 mL O₂/beat, was significant at $P = 0.0003$, and $r = 0.92$. (B) Relationship of CE and TM $\Delta\dot{V}O_2/\Delta HR$ to peak $\dot{V}O_2$. Both modalities revealed very high correlations. Linear regression parameters for CPET CE (solid line) were: peak $\dot{V}O_2$ (mL/min) = $113.3 \times \Delta\dot{V}O_2/\Delta HR$ (mL/beat) + 361.7, $r = 0.94$; and for CPET TM (dotted line) peak $\dot{V}O_2$ (mL/min) = $115.2 \times \Delta\dot{V}O_2/\Delta HR$ (mL/beat) + 247.0, $r = 0.94$. (C) We calculated the percent predicted peak $\dot{V}O_2$ based on the LBM linear regression (see text) and plotted TM percent predicted vs. CE percent predicted peak $\dot{V}O_2$. The correlation coefficient was $r = 0.66$, $P < 0.0001$.

insights using CPET-derived $\Delta\dot{V}E/\Delta\dot{V}CO_2$ have been gained in children and adults from both TM and CE CPET in diseases ranging from cystic fibrosis to heart




Figure 4. Interoperability of $\Delta\dot{V}E/\Delta\dot{V}CO_2$ derived from CPET-CE and CPET-TM. (A) The slopes were significantly correlated $\Delta\dot{V}E/\Delta\dot{V}CO_2$ (TM) = $0.48 \times \Delta\dot{V}E/\Delta\dot{V}CO_2$ (CE) + 14.4, $r = 0.63$, $P < 0.0001$. (B) The slopes from both modalities were inversely correlated with LBM (CE: $\Delta\dot{V}E/\Delta\dot{V}CO_2 = -0.12 \times LBM$ (kg) + 34.3, $r = -0.40$, $P < 0.0001$; TM: $\Delta\dot{V}E/\Delta\dot{V}CO_2 = -0.11 \times LBM$ (kg) + 32.3, $r = -0.48$, $P < 0.0001$). These data corroborate previous studies showing generally higher $\Delta\dot{V}E/\Delta\dot{V}CO_2$ in younger (smaller) children compared with adolescents and young adults.

failure (Moser et al., 2000; Ingle et al., 2012). We did observe small but statistically significant differences in $\Delta\dot{V}E/\Delta\dot{V}CO_2$, for example, a 3.7% larger value overall for CE exercise. Our study was not configured to determine the mechanism of this difference (e.g., greater ventilatory dead space or a lower CO₂ set point in CE compared with TM CPET). Nonetheless, while the differences were small in this cohort of children and adults with no history of lung disease, one might speculate that variables like $\Delta\dot{V}E/\Delta\dot{V}CO_2$ might become more useful in participants with chronic lung disease.

One reason for the somewhat smaller correlation for the $\Delta\dot{V}E/\Delta\dot{V}CO_2$ variable between the two modalities may be that the range of $\Delta\dot{V}E/\Delta\dot{V}CO_2$ values in our cohort




Figure 5. Interoperability of work rate–heart rate slopes derived from CPET-CE and CPET-TM. (A) The slopes for $\Delta WR/\Delta HR$ and $\Delta WR'/\Delta HR$ as a function of LBM were similar: $\Delta WR/\Delta HR$ (CE) = $0.048 \times LBM - 0.070$, $r = 0.89$, $P < 0.0001$; $\Delta WR'/\Delta HR$ (TM) = $0.050 \times LBM + 0.041$, $r = 0.89$, $P < 0.0001$. (B) Using data derived solely from CPET TM showed a very similar relationship, $\Delta VIM/\Delta HR$ (TM) = $7.24 \times LBM - 50.12$, $r = 0.86$, $P < 0.0001$. (C) Dynamic WR–HR data obtained from the two modalities were highly correlated. $\Delta VIM/\Delta HR$ (TM) = $140.4 \times \Delta WR/\Delta HR$ (CE) – 24.5, $r = 0.90$, $P < 0.0001$.




Figure 6. Interoperability of peak $\dot{V}O_2$ derived from CPET-CE and CPET-TM. (A) The slope of the linear regression equation was highly significant, 0.975 ± 0.0316 , $P < 0.0001$; the y-intercept, 182.81 ± 81.87 mL O_2 /min, was significant at $P = 0.028$, and $r = 0.95$. (B) Relationship of CE and TM peak $\dot{V}O_2$ to lean body mass. Both modalities revealed very high correlations. Linear regression parameters for CPET CE (solid line) were: peak $\dot{V}O_2$ CE (mL/min) = $67.3 \times LBM$ (kg) + 21.8, $r = 0.91$; and for CPET TM (dotted line) peak $\dot{V}O_2$ TM (mL/min) = $71.3 \times LBM$ (kg) + 4.39, $r = 0.94$. The correlation coefficients for LBM were higher than for weight in both modalities: CE, peak $\dot{V}O_2 = 26.4 \times Weight + 46.4$, $r = 0.82$; TM, peak $\dot{V}O_2 = 49.6 \times Weight - 13.0$, $r = 0.86$. (C) We calculated the percent predicted peak $\dot{V}O_2$ based on the LBM linear regression and plotted TM percent predicted versus CE percent predicted peak $\dot{V}O_2$. The correlation coefficient was $r = 0.66$.

was substantially smaller than for other CPET variables. For example, $\Delta\dot{V}O_2/\Delta HR$ ranged from about 7–45 mL O_2 /beat while $\Delta\dot{V}E/\Delta\dot{V}CO_2$ ranged only from about 25–35 (unitless). Within a participant group of healthy individuals with no history of lung or heart disease, a major determinant of many CPET variables is body size, particularly muscle mass (Cooper et al., 2014). Previous studies using CE CPET demonstrated high correlations between body size and $\Delta\dot{V}O_2/\Delta HR$ and weak, but significant, inverse correlations with $\Delta\dot{V}E/\Delta\dot{V}CO_2$.

We found that the relationships between $\Delta\dot{V}O_2/\Delta HR$ and body mass in TM CPET paralleled the relationships we found previously using CE CPET. The correlation between TM CPET-derived $\Delta\dot{V}O_2/\Delta HR$ and body weight was strong, but even stronger when correlated with LBM. These results emphasize the need to scale $\Delta\dot{V}O_2/\Delta HR$ to some metric of body size in order to interpret the results correctly. The TM CPET-derived $\Delta\dot{V}E/\Delta\dot{V}CO_2$ was to a small but significant degree inversely correlated to body size, similar to the earlier studies using CE CPET (Cooper et al., 1984; Nagano et al., 1998). These similar results from the two different exercise modalities bolster the idea that physiologic mechanisms, such as the CO_2 set point or the relationship between dead space and tidal volume, systematically change over childhood and adolescence.

Both modalities revealed similar and significant sex effects in $\Delta\dot{V}O_2/\Delta HR$ and $\Delta\dot{V}E/\Delta\dot{V}CO_2$. The higher oxygen extraction per beat found in male participants reflects, as noted above, the influence of stroke volume and $(a - v)O_2$. In adults, left ventricular size is smaller in females compared to males (Gebhard et al., 2013). Similar observations have been made in children (Vinet et al., 2003). These results might explain the sexual dimorphism of the $\Delta\dot{V}O_2/\Delta HR$. Although not as well studied as heart size, one study in young and middle-aged adults also showed generally higher $\dot{V}E-\dot{V}CO_2$ based parameters in females compared with males (Sun et al., 2002), an observation not seen in one exercise study in younger volunteers (Guerrero et al., 2008). Sexual dimorphism in respiratory control in adults is a known phenomenon, but the impact of sex on respiratory control during exercise in children is not well understood. Whether the generally higher $\Delta\dot{V}E/\Delta\dot{V}CO_2$ that we found in females indicate greater deadspace ventilation or, alternatively, a lower CO_2 set point, has yet to be determined.

CPET typically consists of an ergometer programmed to increase the participant's

work rate coupled with a set of devices capable of measuring physiologic responses such as gas exchange or HR. These physiologic outputs are useful only insofar as they can be scaled. For example, an isolated HR measured during exercise is uninterpretable unless it is dynamically

scaled to a CPET input such as the work rate. We used several approaches to address the challenging problem of estimating work performed during TM CPET. There are very compelling reasons to do this; one of the most potentially impactful would be in reanalyzing fitness data from many studies in children in which TM CPET in some form is used to estimate, rather than measure, peak $\dot{V}O_2$ [e.g., NHANES (Astrand and Ryhming, 1954; Jackson et al., 1990)]. Subsequent calculated estimates of $\dot{V}O_2$ max derived from the submaximal CPET may include variables or constants reflecting levels of habitual physical activity or normative values obtained from studies in adults. Such approaches can contribute to the increasingly recognized problems that confound data interpretation due to misspecification, collinearity, and mathematical coupling (Tu et al., 2004; Aggarwal and Ranganathan, 2016).

An analysis of TM CPET that relies predominantly on actually measured data would advance our ability to accurately gauge fitness from CPET. In the current study, we were able to calculate the WR' TM exercise based on the $\dot{V}O_2$ -WR relationship measured during CE CPET. As shown in Figure 5A, $\Delta WR/\Delta HR$ from CE CPET and $\Delta WR'/\Delta HR$ from TM CPET had virtually identical relationships with LBM (and body weight, data not shown). Additional parallel effects of sex are shown in Table 4.

Using the VIM estimate of TM work rate led to a linear relationship with $\dot{V}O_2$ (Fig. 2), mimicking the well-established relationship between $\dot{V}O_2$ and work rate observed consistently in CE CPET. The potential value of this approach to TM CPET, *which uses only the actually measured data, that is, body mass, HR, and treadmill speed and incline*, is highlighted in Figure 5C, showing the very high correlation between the submaximal $\Delta VIM/\Delta HR$ of TM CPET and $\Delta WR/\Delta HR$ measured in CE CPET. As shown in Table 3, maturation- and sex-related changes in TM CPET-derived $\Delta VIM/\Delta HR$ paralleled, as expected, the changes in CE $\Delta WR/\Delta HR$. Similarly, for exercise biomarkers expected to be relatively size independent, we found, as expected, little or no differences across our subgroups for CE CPET $\Delta\dot{V}O_2/\Delta WR$ and the parallel TM CPET $\Delta\dot{V}O_2/\Delta VIM$.

We found strong correlations between peak $\dot{V}O_2$ obtained from TM and CE CPET (Fig. 6A). Our data are consistent with previous studies demonstrating generally lower values for CE peak $\dot{V}O_2$. For both TM and CE CPET, there was a strong correlation between $\Delta\dot{V}O_2/\Delta HR$ and peak $\dot{V}O_2$ (Fig. 3B). This observation may be particularly useful in instances when a participant or patient does not meet standard criteria for peak $\dot{V}O_2$, not an infrequent occurrence (Paridon et al., 2008). In these cases, investigators or clinicians might consider using $\Delta\dot{V}O_2/\Delta HR$ (a value not dependent on maximal effort) as a surrogate index for fitness.

Myers et al. (2017 and Kaminsky et al. (2017 recently published normative values for CPET in separate populations of adults using TM and CE. Although the investigators found generally lower peak $\dot{V}O_2$ values in CE CPET, they were unable to identify a unique conversion factor that could eliminate differences between the two ergometer types across the age groups of their study. A number of investigators have compared TM and CE CPET in which participants performed both modalities (Jacobs and Sjödén, 1985; Turley and Wilmore, 1997; Basset and Boulay, 2000; Mitchell et al., 2010; Gordon et al., 2012; Itoh et al., 2013) and consistently higher TM CPET peak $\dot{V}O_2$ has been observed. As noted, Turley and Wilmore (1997) studied both children and adults, and found that CE peak $\dot{V}O_2$ in all groups was lower than TM to a small but consistent degree. There are a number of possible explanations for the higher peak $\dot{V}O_2$ values in TM exercise, including the energy cost of maintaining an upright posture (Miles-Chan et al., 2013; Júdeice et al., 2016) and/or factors related to work efficiency, skeletal muscle mass, and activation that occur in TM but not CE CPET. Muscle mass clearly plays a role; for example, peak $\dot{V}O_2$ is, as expected, smaller in upper body ergometry compared with TM or CE CPET (Drescher et al., 2015). It is noteworthy that we could not find significant changes in any of our submaximal slopes in the transition from walking to running on the treadmill, suggesting that the predominant component of energy costs of TM exercise is related to velocity, mass, and incline.

Using the strong relationships of LBM to both TM and CE peak $\dot{V}O_2$ and $\Delta \dot{V}O_2/\Delta HR$, we addressed the question of whether the two modalities reflected similar hierarchies in fitness among the participants. To do this, we used the linear regression relationship between LBM and both peak $\dot{V}O_2$ and $\Delta \dot{V}O_2/\Delta HR$ to determine a predicted value for each participant based on LBM. We then correlated the TM and CE percent predicted value for each participant. As shown in Figures 3C and 6C, we found modest but significant correlations in fitness hierarchy for both submaximal $\Delta \dot{V}O_2/\Delta HR$ and peak $\dot{V}O_2$. In summary, a participant in our study with a relatively high or low peak $\dot{V}O_2$ would likely have respectively high or low $\Delta \dot{V}O_2/\Delta HR$ on both TM and CE modalities. However, the variability in our data also cautions that relative fitness ascertained by CPET biomarkers is not fully interchangeable between the two modalities.

Limitations: Due to the same order of exercise test modalities a sequence effect may influence the second test session. In this study supramaximal test was not performed. In children $\dot{V}O_2$ peak is more commonly used than in adults and supramaximal tests are equivocal. This study focused on submaximal values and novel TM protocol; thus, measuring peak or max values were presented as secondary end point.

In conclusion, our data reveal the effect of the two predominant modalities of laboratory exercise testing in children, adolescents, and young adults on submaximal and peak CPET results. Both modalities similarly reflected effects of body size on $\Delta \dot{V}O_2/\Delta HR$, $\Delta \dot{V}E/\Delta \dot{V}CO_2$, and peak $\dot{V}O_2$. Results from the two modalities, however, are not interchangeable and may reflect the complexities of how external work, particularly on the treadmill, is transduced to physiologic responses such as $\dot{V}O_2$ (Pandolf et al., 1977; Epstein et al., 1987; Hall et al., 2004). The reasons for using TM or CE exercise for assessment of exercise biomarkers in the clinic or in research ultimately depend on a variety of factors, including the skill and experience of the laboratory, available equipment, and perceived capabilities of the targeted participants or patients. We provide a novel approach for analyzing TM CPET data relying on actually measured HR, body mass, and the velocity and incline. This approach might prove useful in reanalyzing existing datasets where such measurements are available and in the future establishment of normative values for CPET testing in children and adolescents, where reliable datasets in large numbers of healthy participants are currently lacking.

Acknowledgments

We thank Dr. Kim Lu for reviewing the manuscript, Hoang Pham for assistance in performing the exercise testing and the children and their families who participated in this study.

Conflict of Interest

The authors declare no conflicts of interest. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

References

- Aggarwal, R., and P. Ranganathan. 2016. Common pitfalls in statistical analysis: The use of correlation techniques. *Perspect Clin Res* 7:187–190.
- Armon, Y., D. M. Cooper, and S. Zanconato. 1991. Maturation of ventilatory responses to 1-minute exercise. *Pediatr. Res.* 29:362–368.
- Ashish, N., M. M. Bamman, F. J. Cerny, D. M. Cooper, P. D'Hemecourt, J. C. Eisenmann, et al. 2015. The clinical translation gap in child health exercise research: a call for disruptive innovation. *Clin. Transl. Sci.* 8:67–76.
- Astrand, P. O., and I. Ryhming. 1954. A nomogram for calculation of aerobic capacity (physical fitness) from pulse rate during sub-maximal work. *J. Appl. Physiol.* 7:218–221.
- Azuma, A. 2014. Effects of a vigorous gait on mechanical work and oxygen uptake during treadmill walking. *Percept. Mot Skills* 119:6–19.

- Basset, F. A., and M. R. Boulay. 2000. Specificity of treadmill and cycle ergometer tests in triathletes, runners and cyclists. *Eur. J. Appl. Physiol.* 81:214–221.
- Beltrami, F. G., C. Froyd, A. R. Mauger, A. J. Metcalfe, F. Marino, and T. D. Noakes. 2012. Conventional testing methods produce submaximal values of maximum oxygen consumption. *Br. J. Sports Med.* 46:23–29.
- Bires, A. M., D. Lawson, T. E. Wasser, and D. Raber-Baer. 2013. Comparison of Bruce treadmill exercise test protocols: is ramped Bruce equal or superior to standard Bruce in producing clinically valid studies for patients presenting for evaluation of cardiac ischemia or Arrhythmia with body mass index equal to or greater than 30?. *J. Nucl. Med. Technol.* 41:274–278. <https://doi.org/10.2967/jnmt.113.124727>.
- Bridge, P., N. A. Pocock, T. Nguyen, C. Munns, C. T. Cowell, N. Forwood, et al. 2011. Validation of longitudinal DXA changes in body composition from pre- to mid-adolescence using MRI as reference. *J. Clin. Densitom.* 14:340–347.
- Buchfuhrer, M. J., J. E. Hansen, T. E. Robinson, D. Y. Sue, K. Wasserman, and B. J. Whipp. 1983. Optimizing the exercise protocol for cardiopulmonary assessment. *J. Appl. Physiol.* 55:1558–1564.
- Chen, C. A., S. Y. Chen, H. H. Chiu, J. K. Wang, C. I. Chang, I. S. Chiu, et al. 2014. Prognostic value of submaximal exercise data for cardiac morbidity in Fontan patients. *Med. Sci. Sports Exerc.* 46:10–15.
- Cooper, D. M., D. Weiler-Ravell, B. J. Whipp, and K. Wasserman. 1984. Aerobic parameters of exercise as a function of body size during growth in children. *J. Appl. Physiol.* 56:628–634.
- Cooper, D. M., M. R. Kaplan, L. Baumgarten, D. Weiler-Ravell, B. J. Whipp, and K. Wasserman. 1987. Coupling of ventilation and CO₂ production during exercise in children. *Pediatr. Res.* 21:568–572.
- Cooper, D. M., S. Y. Leu, P. Galassetti, and S. Radom-Aizik. 2014. Dynamic interactions of gas exchange, body mass, and progressive exercise in children. *Med. Sci. Sports Exerc.* 46:877–886.
- Cooper, D. M., S.-Y. Leu, C. Taylor-Lucas, K. Lu, P. Galassetti, and S. Radom-Aizik. 2016. Cardiopulmonary exercise testing in children and adolescents with high body mass index. *Pediatr. Exerc. Sci.* 28:98–108.
- Cordingley, D., R. Girardin, K. Reimer, L. Ritchie, J. Leiter, K. Russell, et al. 2016. Graded aerobic treadmill testing in pediatric sports-related concussion: safety, clinical use, and patient outcomes. *J. Neurosurg. Pediatr.* 18:693–702.
- Cunha, F. A., A. W. Midgley, T. Gonçalves, P. P. Soares, and P. Farinatti. 2015. Parasympathetic reactivation after maximal CPET depends on exercise modality and resting vagal activity in healthy men. *Springerplus* 4:100. <https://doi.org/10.1186/s40064-015-0882-1>.
- Drescher, U., J. Koschate, and U. Hoffmann. 2015. Oxygen uptake and heart rate kinetics during dynamic upper and lower body exercise: an investigation by time-series analysis. *Eur. J. Appl. Physiol.* 115:1665–1672.
- Elbehairy, A. F., C. E. Ciavaglia, K. A. Webb, J. A. Guenette, D. Jensen, S. M. Mourad, et al. 2015. Canadian respiratory research network. pulmonary gas exchange abnormalities in mild chronic obstructive pulmonary disease. Implications for dyspnea and exercise intolerance. *Am. J. Respir. Crit. Care Med.* 191:1384–1394.
- Epstein, Y., L. A. Stroschein, and K. B. Pandolf. 1987. Predicting metabolic cost of running with and without backpack loads. *Eur. J. Appl. Physiol. Occup. Physiol.* 56:495–500.
- Fredriksen, P. M., F. Ingjer, W. Nystad, and E. Thaulow. 1998. Aerobic endurance testing of children and adolescents—a comparison of two treadmill-protocols. *Scand. J. Med. Sci. Sports* 8:203–207.
- Gebhard, C., B. E. Stähli, C. E. Gebhard, H. Tasnady, D. Zihler, M. B. Wischnowsky, et al. 2013. Age- and gender-dependent left ventricular remodeling. *Echocardiography* 30:1143–1150.
- Gordon, D., M. Mehter, M. Gernigon, O. Caddy, D. Keiller, and R. Barnes. 2012. The effects of exercise modality on the incidence of plateau at VO₂max. *Clin. Physiol. Funct. Imaging* 32:394–399.
- Gualano, B., E. Bonfa, R. M. R. Pereira, and C. A. Silva. 2017. Physical activity for paediatric rheumatic diseases: standing up against old paradigms. *Nat. Rev. Rheumatol.* 13:368–379.
- Guerrero, L., J. Naranjo, and M. D. Carranza. 2008. Influence of gender on ventilatory efficiency during exercise in young children. *J. Sports Sci.* 26:1455–1457.
- Hall, C., A. Figueroa, B. Fernhall, and J. A. Kanaley. 2004. Energy expenditure of walking and running: comparison with prediction equations. *Med. Sci. Sports Exerc.* 36:2128–2134.
- Hestnes, J., H. Hoel, O. J. Risa, H. O. Romstøl, O. Røksund, B. Frisk, et al. 2017. Ventilatory efficiency in children and adolescents born extremely preterm. *Front. Physiol.* 8:499.
- Ingle, L., R. Sloan, S. Carroll, K. Goode, J. G. Cleland, and A. L. Clark. 2012. Abnormalities of the ventilatory equivalent for carbon dioxide in patients with chronic heart failure. *Pulm. Med.* 2012:1–6.
- Itoh, H., R. Ajsaka, A. Koike, S. Makita, K. Omiya, Y. Kato, et al. 2013. Heart rate and blood pressure response to ramp exercise and exercise capacity in relation to age, gender, and mode of exercise in a healthy population. *J. Cardiol.* 61:71–78.
- Jackson, A. S., S. N. Blair, M. T. Mahar, L. T. Wier, R. M. Ross, and J. E. Stuteville. 1990. Prediction of functional aerobic capacity without exercise testing. *Med. Sci. Sport Exerc.* 22:863.
- Jacobs, I., and B. Sjödén. 1985. Relationship of ergometer-specific VO₂ max and muscle enzymes to blood lactate during submaximal exercise. *Br. J. Sports Med.* 19:77–80.
- Júdice, P. B., M. T. Hamilton, L. B. Sardinha, T. W. Zderic, and A. M. Silva. 2016. What is the metabolic and energy

- cost of sitting, standing and sit/stand transitions? *Eur. J. Appl. Physiol.* 116:263–273.
- Kaminsky, L. A., M. T. Imboden, R. Arena, and J. Myers. 2017. Reference standards for cardiorespiratory fitness measured with cardiopulmonary exercise testing using cycle ergometry: data from the fitness registry and the importance of exercise National Database (FRIEND) Registry. *Mayo Clin. Proc.* 92:228–233.
- Keir, D. A., R. Zory, C. Boudreau-Larivière, and O. Serresse. 2012. Mechanical efficiency of treadmill running exercise: Effect of anaerobic-energy contribution at various speeds. *Int. J. Sports Physiol. Perform.* 7:382–389.
- Kenney, M. J., and D. R. Seals. 1993. Postexercise hypotension. Key features, mechanisms, and clinical significance. *Hypertens* 22:653–664.
- Kyröläinen, H., P. V. Komi, and A. Belli. 1995. Mechanical efficiency in athletes during running. *Scand. J. Med. Sci. Sports* 5:200–208.
- Li, C., F. Feng, X. Xiong, R. Li, and N. Chen. 2017. Exercise coupled with dietary restriction reduces oxidative stress in male adolescents with obesity. *J. Sports Sci.* 35:663–668.
- Liem, R. I., M. Reddy, S. A. Pelligra, A. P. Savant, B. Fernhall, M. Rodeghier, et al. 2015. Reduced fitness and abnormal cardiopulmonary responses to maximal exercise testing in children and young adults with sickle cell anemia. *Physiol. Rep.* 3:1–8.
- May, L. J., R. Punn, I. Olson, J. A. Kazmucha, M. Y. Liu, and C. Chin. 2014. Supine cycling in pediatric exercise testing: disparity in performance measures. *Pediatr. Cardiol.* 35:705–710.
- Miles-Chan, J. L., D. Sarafian, J. P. Montani, Y. Schutz, and A. Dulloo. 2013. Heterogeneity in the energy cost of posture maintenance during standing relative to sitting: phenotyping according to magnitude and time-course. *PLoS ONE* 8: e65827.
- Mitchell, J., W. B. Kist, K. Mears, J. Nalls, and K. Ritter. 2010. Does standing on a cycle-ergometer, towards the conclusion of a graded exercise test, yield cardiorespiratory values equivalent to treadmill testing? *Int. J. Exerc. Sci.* 3:117–125.
- Moser, C., P. Tirakitsoontorn, E. Nussbaum, R. Newcomb, and D. M. Cooper. 2000. Muscle size and cardiorespiratory response to exercise in Cystic Fibrosis. *Am. J. Respir. Crit. Care Med.* 162:1823–1827.
- Myers, J., and V. F. Froelicher. 1993. Exercise testing. Procedures and implementation. *Cardiol. Clin.* 11:199–213.
- Myers, J., L. A. Kaminsky, R. Lima, J. W. Christle, E. Ashley, and R. Arena. 2017. A Reference equation for normal standards for VO₂MAX: analysis from the Fitness Registry and the Importance of Exercise National Database (FRIEND Registry). *Prog. Cardiovasc. Dis.* 60:21–29.
- Nagano, Y., R. Baba, K. Kuraishi, T. Yasuda, M. Ikoma, K. Nishibata, et al. 1998. Ventilatory control during exercise in normal children. *Pediatr. Res.* 43:704–707.
- Pahkala, K., T. T. Laitinen, O. J. Heinonen, J. S. A. Viikari, T. Ronnema, H. Niinikoski, et al. 2013. Association of fitness with vascular intima-media thickness and elasticity in adolescence. *Pediatrics* 132:e77–e84.
- Pandolf, K. B., B. Givoni, and R. F. Goldman. 1977. Predicting energy expenditure with loads while standing or walking very slowly. *J. Appl. Physiol.* 43:577–581.
- Paridon, S. M., P. D. Mitchell, S. D. Colan, R. V. Williams, A. Blaufox, J. S. Li, et al. 2008. A cross-sectional study of exercise performance during the first 2 decades of life after the Fontan operation. *J. Am. Coll. Cardiol.* 52:99–107.
- Ploeger, H. E., T. Takken, M. H. G. DeGreef, and B. W. Timmons. 2009. The effects of acute and chronic exercise on inflammatory markers in children and adults with a chronic inflammatory disease: a systematic review. *Exerc. Immunol. Rev.* 15:6–41.
- Porszasz, J., R. Casaburi, A. Somfay, L. J. Woodhouse, and B. J. Whipp. 2003. A treadmill ramp protocol using simultaneous changes in speed and grade. *Med. Sci. Sports Exerc.* 35:1596–1603.
- Rausch, S. M., B. J. Whipp, K. Wasserman, and A. Huszczuk. 1991. Role of the carotid bodies in the respiratory compensation for the metabolic acidosis of exercise in humans. *J. Physiol.* 444:567–578.
- Rowland, T., S. Hagenbuch, D. Poher, and A. Garrison. 2008. Exercise tolerance and thermoregulatory responses during cycling in boys and men. *Med. Sci. Sports Exerc.* 40:282–287.
- Ruckstuhl, H., T. Schlabs, A. Rosales-Velderrain, and A. R. Hargens. 2010. Oxygen consumption during walking and running under fractional weight bearing conditions. *Aviat. Space Environ. Med.* 81:550–554.
- Stein, R., H. Selvadurai, A. Coates, D. L. Wilkes, J. Schneiderman-Walker, and M. Corey. 2003. Determination of maximal voluntary ventilation in children with cystic fibrosis. *Pediatr. Pulmonol.* 35:467–471.
- Sule, S., and K. Fontaine. 2016. Abnormal body composition, cardiovascular endurance, and muscle strength in pediatric SLE. *Pediatr. Rheumatol.* 14:50.
- Sun, X.-G., J. E. Hansen, N. Garatachea, T. W. Storer, and K. Wasserman. 2002. Ventilatory efficiency during exercise in healthy subjects. *Am. J. Respir. Crit. Care Med.* 166:1443–1448.
- Troutman, W. B., T. J. Barstow, A. J. Galindo, and D. M. Cooper. 1998. Abnormal dynamic cardiorespiratory responses to exercise in pediatric patients after fonton procedure. *J. Am. Coll. Cardiol.* 31:668–673.
- Tu, Y.-K., I. H. Maddick, G. S. Griffiths, and M. S. Gilthorpe. 2004. Mathematical coupling can undermine the statistical assessment of clinical research: illustration from the treatment of guided tissue regeneration. *J. Dent.* 32:133–142.
- Turley, K. R., and J. H. Wilmore. 1997. Cardiovascular responses to treadmill and cycle ergometer exercise in children and adults. *J. Appl. Physiol.* 83:948–957.

Vinet, A., S. Mandigout, S. Nottin, L. Nguyen, A.-M. Lecoq, D. Courteix, et al. 2003. Influence of body composition, hemoglobin concentration, and cardiac size and function of gender differences in maximal oxygen uptake in prepubertal children. *Chest* 124:1494–1499.

Whipp, B. J., J. A. Davis, F. Torres, and K. Wasserman. 1981. A test to determine parameters of aerobic function during exercise. *J. Appl. Physiol.* 50:217–221.

Workman, J. M., and B. W. Armstrong. 1963. Oxygen cost of treadmill walking. *J. Appl. Physiol.* 18:798–803.