

Gain-of-Function Polymorphisms in Human Inflammasomes: Implications for Cystic Fibrosis

Cystic fibrosis (CF) is an inherited disease involving chronic infection and inflammation of the lungs (1). CF results from mutations in the *CFTR* (CF transmembrane conductance regulator) gene (2) and is the most common inherited genetic disease in white individuals, affecting ~1 in 3,000 newborns (3). *Pseudomonas aeruginosa* is a common cause of morbidity in patients with CF, triggering excessive inflammation, lung damage, and eventual respiratory failure (4). Several therapeutic approaches have been attempted or are currently in use, including antiinflammatory drugs, antioxidants, antibiotics, CFTR modulators, and gene therapy (1, 5). However, management of infection and lung damage because of chronic inflammation remains a major challenge for patients with CF.

CF lung disease involves elevated expression of proinflammatory cytokines because of dysregulation of the immune response (6). Central regulators of innate immunity are pattern recognition receptors, including TLRs (Toll-like receptors) and NLRs (nod-like receptors) that recognize pathogen-associated molecular patterns and damage-associated molecular patterns. Some NLRs, such as inflammasomes, are self-oligomerizing protein complexes that consist of nucleotide-binding and oligomerization domains, and the ASC protein, which recruits procaspase-1, leading to its activation (7–9). Active caspase-1 is responsible for proteolytic activation of proinflammatory cytokines, IL-1 β and IL-18. The release of these proinflammatory cytokines and subsequent pyroptosis of the cell can lead to inflammatory lung damage (6, 7). In a recent study of a murine CF model using *P. aeruginosa*, the inhibition of the NLRP3 inflammasome was linked to improved bacterial clearance from the lung (10). Unlike animal models, the complexity of inflammasome biology in the human population arises from SNPs of these genes. These SNPs can give rise to inflammasome variants, which alter their activity, leading to changes in proinflammatory cytokine expression. In fact, it has been known for some time that mutations in the gene encoding *NLRP3* can induce a number of autoinflammatory disorders known as cryopyrin-associated periodic syndromes (11). Studies are now beginning to delve into the mechanistic action of specific inflammasome genetic variants. For instance, a recent report on the NLRP3 (Q705K) variant demonstrated an increase in inflammasome activation for the variant over wild-type NLRP3 (12). However, a variant in the *NLRC4* gene was found to correlate with decreased expression of IL-18 and improved lung function in patients with human immunodeficiency virus (HIV) and

tuberculosis coinfection (13). In another study, a missense variant of *NLRC4* was also associated with decreased expression of IL-18 (14).

In this issue of the *Journal*, Graustein and colleagues (pp. 157–166) identify inflammasome SNPs associated with *P. aeruginosa* lung colonization and lung function in CF (15). For this study, the authors genotyped variants from patients in the EPIC (Early *Pseudomonas* Infection Control) observational study. Previous investigations have described SNPs for *NLRP3* and *NLRC4* that alter the activity of these inflammasomes (12–14, 16, 17). However, Graustein and colleagues now show for the first time a correlation between inflammasome SNPs and the clinical outcomes of patients with CF carrying these genetic variants. The authors demonstrate that the NLRP3 (Q705K) variant correlated with an increased rate of *P. aeruginosa* lung infection and decreased lung function over time. However, only children who had previously been infected with *P. aeruginosa* demonstrated this effect. The authors note that a group of older children who were never infected with *P. aeruginosa* could exert a cohort bias against containing the NLRP3 (Q705K) variant, possibly explaining why the effects were only observed in children previously infected.

The authors also identify the novel *NLRC4* (A929S) variant as an SNP associated with protection of lung function for children with CF never infected with *P. aeruginosa*. However, because of the small sample size, the effect of the *NLRC4* (A929S) SNP on *P. aeruginosa* lung colonization could not be directly investigated. Instead, the authors analyzed a group of variants from multiple genes in the *NLRC4* pathway as a single variable and found a protective role against *P. aeruginosa* lung colonization for these alterations. The lung colonization findings include only patients never infected with *P. aeruginosa* before enrollment. As with the *NLRP3* SNP cohorts, this protective *NLRC4* variant could reflect cohort bias against including children previously infected with *P. aeruginosa*. The authors also note that the *NLRC4*-related variants studied were only associated with protection in children with wild-type *CAV2* (caveolin 2), a gene associated with facilitating *P. aeruginosa* infection (18). This finding suggests that the *NLRC4* pathway variants may antagonize the activity of *CAV2*.

To better understand the importance of *NLRP3* and *NLRC4* SNPs in chronic inflammation, the authors used human macrophage-like cell lines expressing wild-type or variant inflammasome constructs for *in vitro* studies. Using these cell lines, the authors demonstrate that the NLRP3 (Q705K) variant-expressing cells are more responsive to

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1165/rcmb.2021-0183ED on May 25, 2021

NLRP3-specific stimulation (nigericin) than wild-type cells. On the other hand, when cells were infected with *P. aeruginosa* strain PAO1, there was no difference in the response between NLRP3 wild-type and variant-expressing cells. The authors were unable to detect any statistical differences in cytokine expression from NLRC4 (A929S) variant-expressing cells. These findings may be due to the properties of the cell line or the *P. aeruginosa* strain used. A study using neutrophils or lung epithelial cells would likely help to further elucidate the mechanism of the NLRP3 (Q705K) and NLRC4 (A929S) variants. Furthermore, it has been reported that NLRP3 is predominantly responsible for IL-1 β processing in CF neutrophils (10). It was also found that patients carrying the NLRP3 (Q705K) and CARD-8 (C10X) SNPs have delayed neutrophil apoptosis, increased IL-1 β levels, and low macrophage phagocytic capacity, which increases the risk of damaging inflammation (17). The release of IL-18 from cells expressing the NLRP3 (Q705K or V198M) variants or the NLRC4 (A929S) variant represents another avenue to investigate inflammasome gain or loss of function. In fact, a previous study with a monocytic cell line expressing wild-type and the Q705K variant of NLRP3 has demonstrated an enhanced expression of IL-18 and IL-1 β in the variant-expressing cells over wild-type in response to stimulation by alum (12).

In conclusion, Graustein and colleagues provide compelling clinical insights into the roles of NLRP3 and NLRC4 SNPs in CF. Importantly, the clinical data presented in this paper are accompanied by mechanistic *in vitro* studies. Together, these analyses shed light on the role of inflammasome variants in altering the balance between beneficial and detrimental excessive inflammation. Specifically, their clinical and *in vitro* findings identify the NLRP3 (Q705K) inflammasome variant as a biomarker for increased inflammation, earlier *P. aeruginosa* colonization, and accelerated disease progression in CF. Finally, the authors provide evidence in support of the concept that NLRP3 inflammasome inhibition is a viable approach to slow the progression of CF and potentially other pulmonary inflammatory diseases. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Duane Jeansonne, Ph.D.
Center for Lung Biology and Disease
and

Department of Pathobiological Sciences Louisiana State University School of Veterinary Medicine
Baton Rouge, Louisiana

Samithamby Jeyaseelan, D.V.M., Ph.D.
Center for Lung Biology and Disease

Department of Pathobiological Sciences Louisiana State University School of Veterinary Medicine
Baton Rouge, Louisiana
and

Department of Medicine Louisiana State University Health Sciences Center
New Orleans, Louisiana

References

- Ralhan A, Laval J, Lelis F, Ballbach M, Grund C, Hector A, *et al*. Current concepts and controversies in innate immunity of cystic fibrosis lung disease. *J Innate Immun* 2016;8:531–540.
- Choi JY, Muallem D, Kiselyov K, Lee MG, Thomas PJ, Muallem S. Aberrant CFTR-dependent HCO₃⁻ transport in mutations associated with cystic fibrosis. *Nature* 2001;410:94–97.
- O'Sullivan BP, Freedman SD. Cystic fibrosis. *Lancet* 2009;373:1891–1904.
- Davies JC, Alton EW, Bush A. Cystic fibrosis. *BMJ* 2007;335:1255–1259.
- Almughem FA, Aldossary AM, Tawfik EA, Alomary MN, Alharbi WS, Alshahrani MY, *et al*. Cystic fibrosis: overview of the current development trends and innovative therapeutic strategies. *Pharmaceutics* 2020;12:616.
- Iannitti RG, Napolioni V, Oikonomou V, De Luca A, Galosi C, Pariano M, *et al*. IL-1 receptor antagonist ameliorates inflammasome-dependent inflammation in murine and human cystic fibrosis. *Nat Commun* 2016;7:10791.
- Ravi Kumar S, Paudel S, Ghimire L, Bergeron S, Cai S, Zemans RL, *et al*. Emerging roles of inflammasomes in acute pneumonia. *Am J Respir Crit Care Med* 2018;197:160–171.
- Zheng D, Liwinski T, Elinav E. Inflammasome activation and regulation: toward a better understanding of complex mechanisms. *Cell Discov* 2020;6:36.
- Leissinger M, Kulkarni R, Zemans RL, Downey GP, Jeyaseelan S. Investigating the role of nucleotide-binding oligomerization domain-like receptors in bacterial lung infection. *Am J Respir Crit Care Med* 2014;189:1461–1468.
- McElvaney OJ, Zaslon Z, Becker-Flegler K, Palsson-McDermott EM, Boland F, Gunaratnam C, *et al*. Specific inhibition of the NLRP3 inflammasome as an antiinflammatory strategy in cystic fibrosis. *Am J Respir Crit Care Med* 2019;200:1381–1391.
- Kastner DL. Hereditary periodic fever syndromes. *Hematology Am Soc Hematol Educ Program* 2005;2005(1):74–81.
- Verma D, Särndahl E, Andersson H, Eriksson P, Fredrikson M, Jönsson JI, *et al*. The Q705K polymorphism in NLRP3 is a gain-of-function alteration leading to excessive interleukin-1 β and IL-18 production. *PLoS One* 2012;7:e34977.
- Ravimohan S, Maenetje P, Auld SC, Ncube I, Mlotshwa M, Chase W, *et al*. A common NLRC4 gene variant associates with inflammation and pulmonary function in human immunodeficiency virus and tuberculosis. *Clin Infect Dis* 2020;71:924–932.
- Johansson Å, Eriksson N, Becker RC, Storey RF, Himmelmann A, Hagström E, *et al*; PLATO Investigators. NLRC4 inflammasome is an important regulator of interleukin-18 levels in patients with acute coronary syndromes: genome-wide association study in the PLATOlet inhibition and patient outcomes trial (PLATO). *Circ Cardiovasc Genet* 2015;8:498–506.
- Graustein AD, Berrington WR, Buckingham KJ, Nguyen FK, Joudeh LL, Rosenfeld M, *et al*. Inflammasome genetic variants, macrophage function, and clinical outcomes in cystic fibrosis. *Am J Respir Cell Mol Biol* 2021;65:157–166.
- Verma D, Lerm M, Blomgran Julinder R, Eriksson P, Söderkvist P, Särndahl E. Gene polymorphisms in the NALP3 inflammasome are associated with interleukin-1 production and severe inflammation: relation to common inflammatory diseases? *Arthritis Rheum* 2008;58:888–894.
- Blomgran R, Brodin VP, Verma D, Bergström I, Söderkvist P, Sjöwall C, *et al*. Common genetic variations in the NALP3 inflammasome are associated with delayed apoptosis of human neutrophils. *PLoS One* 2012;7:e31326.
- Zaas DW, Swan ZD, Brown BJ, Li G, Randell SH, Degan S, *et al*. Counteracting signaling activities in lipid rafts associated with the invasion of lung epithelial cells by *Pseudomonas aeruginosa*. *J Biol Chem* 2009;284:9955–9964.