

## Research Article

# The Critical Gene Screening to Prevent Chromophobe Cell Renal Carcinoma Metastasis through TCGA and WGCNA

Cheng Yan,<sup>1</sup> Yan Yang,<sup>1</sup> Yunxiang Tang,<sup>2</sup> Xiaojing Zheng,<sup>1</sup> and Bin Xu<sup>1</sup> <sup>2</sup>

<sup>1</sup>Department of Oncology and Hematology, Chongqing Emergency Medical Center, Chongqing University Central Hospital, Chongqing 400014, China

<sup>2</sup>Department of Nuclear Medicine, Chongqing Emergency Medical Center, Chongqing University Central Hospital, Chongqing 400014, China

Correspondence should be addressed to Bin Xu; [doctorxubinnuclear@163.com](mailto:doctorxubinnuclear@163.com)

Received 22 August 2022; Revised 16 September 2022; Accepted 26 September 2022; Published 15 October 2022

Academic Editor: Muhammad Muddassir Ali

Copyright © 2022 Cheng Yan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Common chromophobe renal cell carcinoma (chRCC) has a good prognosis when cured by surgery. However, clinical practice shows that a small number of patients with chRCC will produce metastasis, and the prognosis after metastasis is poor. In this regard, we try to find potential biological targets to prevent CRCC metastasis. In this experiment, we analyzed the clinical traits and gene expression data of chRCC samples which were provided by the TCGA database by the WGCNA method. On this basis, we selected METan, a module with a significant positive correlation with the M phase of chRCC, for subsequent analysis. The METan module genes in the biological process of chRCC were mainly related to steroid metabolic process, cholesterol metabolic process and STEM cell differentiation. KEGG analysis showed that these genes were mainly enriched in cancer-related signaling pathways, such as Neuroactive Ligand–receptor interaction, cAMP signaling pathway, and Wnt signaling pathway. Subsequently, we mapped the PPI interaction network and screened the key gene beta-arrestin 2 (ARRB2). Expression analysis showed that there was a significantly increased expression of ARRB2 in chRCC patients in comparison to the normal group. Expression survival analysis indicated that ARRB2 was inversely associated with overall survival. We firmly believe that the key genes identified in this study would be able to provide new clues and research basis for the treatment of chRCC.

## 1. Introduction

Over 400,000 cases of renal cell carcinoma (RCC) are diagnosed each year in the world, making it one of the most common renal malignancies [1]. Pathologically, RCC is divided into three types: clear cell renal cell carcinoma (ccRCC), papillary carcinoma (pRCC), and chromophobe carcinoma (chRCC). ChRCC is the third subtype of RCC recognized by the World Health Organization (WHO) in 2016 [2]. An estimated 5-10% of all kidney cancers are chRCC, which are equally common in men and women, with a higher incidence in those aged 50-60 [3–5]. ChRCC behaves differently than other types of renal cell carcinomas. Recent statistics indicate an increase in chRCC incidence [6, 7]. Patients with chRCC may present with hematuria or tumor compression symptoms, and a few show diffuse growth and invasion of the perirenal region [8, 9].

A large number of clinical practices have shown that chRCC is usually cured by surgery, and the prognosis of patients is good, with 5-year survival rates of 78-100% and 10-year survival rates of 80-90%; however, there are still 5-10% of patients with chRCC who will develop metastases [10, 11]. Approximately 14 percent of patients with metastatic RCC will survive five years, similar to those with definite metastatic chRCC [12]. Therefore, an in-depth study of the genes related to the pathogenesis of chRCC will comprehensively explain the pathogenesis and disease progression of chRCC, which is of great significance for its treatment and prevention.

There is currently no more comprehensive tumor gene expression profile database than the Cancer Genome Atlas (TCGA), which is distinguished by its large sample size and rich clinical information [13]. An analysis of gene-phenotype relationships called Weighted Gene Coexpression Network Analysis (WGCNA) has gained popularity for its

ability to investigate complex relationships between genes and phenotypes. With the WGCNA method, researchers are able to transform gene expression data into coexpression modules and provide insights into signaling networks that may be responsible for the phenotypic characteristics of the object of interest [14].

Data from gene chips related to chRCC disease were integrated and analyzed using bioinformatics technology: GO and KEGG pathway enrichment analyses were performed first to filter out the differential genes; then, we commenced WGCNA to analyze the clinical characteristics and gene expression data of chRCC samples provided by TCGA database, and made PPI interaction network to find the key genes in the pathogenesis and development of renal chromophobe cell carcinoma; the final step of our study was to investigate the survival of the genes mined to reconstruct the mechanism of renal chromophobe cell carcinoma.

## 2. Materials and Methods

**2.1. Data Capturing.** The TCGA Datasets (<https://www.cancer.gov/about-nci/organization/ccg/research/structural-genomics/tcga>) in the database were accessed with the keyword of chromophobe takes cell carcinoma to search, and the genome data of renal color cell cancer was downloaded. The data included 65 chromophobe cell carcinoma tumor samples and 25 normal tissue samples.

**2.2. DEG Capturing.** Standardizing and analyzing renal chromophobe cell carcinoma datasets were done by using the DESeq2 algorithm in R software. A difference factor ( $\log_2$ ) absolute value higher than 1 was used to screen upregulated genes. Ggplot2 software package was used for data visualization.

**2.3. WGCNA Analysis.** WGCNA provides R functions that help analyze gene expression data using weighted correlation networks.

The source code and other materials for this R package are available for free at <http://genetics.ucla.edu/labs/horvath/CoexpressionNetwork/Rpackages/WGCNA>. Our coexpression network was built using the WGCNA R package. In the first step, clustering the samples was performed to identify any outliers. Next, the coexpression network was constructed using the automatic network construction function. Coexpression similarity is proposed to reckon the adjacency with the R function `pickSoftThreshold`.

**2.4. Module-Trait Relationship Analysis.** The corresponding gene modules were sorted according to the WGCNA modules; then, the ME for each module was calculated and correlated with clinical parameters, with statistical significance defined as  $P < 0.05$ .

**2.5. Differentially Expressed Genes Enriched in GO and KEGG.** DAVID database (DAVID; <https://david.ncifcrf.gov>) was used to analyze GO enrichment and KEGG pathway enrichment of significant different genes screened. The R software and clusterProfiler package were used for annotation and visualization, and a  $P$  value less than 0.05 was considered statistically significant.


**2.6. Screening for Hub Genes in the PPI Network and Construction of a Protein-Protein Interaction Network.** Interactions between proteins were identified and predicted using the STRING database (<https://string-db.org/>). Protein-protein interaction (PPI) networks were constructed using STRING for analysis of differentially expressed genes, and screening for hub genes in the STRING PPI network was performed using the Cytoscape plug-in in Cytoscape software.

**2.7. Key Gene Survival Analysis.** R software was used to analyze the survival of the selected key genes, and an analysis to Kaplan-Meier survival curves was carried out to determine the relationship between the key genes and renal chromophobe cell carcinoma recurrence. An evaluation of the survival difference between key genes was conducted via a log-rank test and the overall survival rate for renal chromophobe cell carcinoma patients was  $P < 0.05$ , deemed significant.


## 3. Results

**3.1. Differentially Expressed Genes Analysis.** An analysis of the transcriptome data from TCGA database was conducted on 65 chromophobe cell carcinoma tumor samples and 25 normal tissue samples. The DESeq2 tool identified 13472 DEGs, of which 6066 were upregulated and 7406 were down-regulated (Figures 1(a) and 1(b)). We ran KEGG enrichment analyses on the top 30 DEGs with a  $P < 0.05$  standard, and results showed that they mainly concentrated on pathways of cAMP, Cytokine–cytokine receptor interaction, Calcium, and Neuroactive Ligand–receptor interaction, etc. (Figures 2(a) and 2(b)). A GO enrichment analysis identified three biological processes associated with DEGs: ion transmembrane transport, membrane potential regulation, and organic anion transport; cell composition included an apical area, extracellular matrix containing collagen, and synaptic membrane; there were several molecular functions that were examined, such as passive transmembrane transporter activity, channel activity, receptor ligand activity, and signaling receptor activator activity (Figures 2(c) and 2(d)).

**3.2. A Weighted Coexpression Network Analysis.** Our first step in constructing the WGCNA network was to calculate the soft threshold power  $\beta$ . It was determined that the soft threshold power was 3-; the scale independence was 0.9, and the average connectivity was relatively high (Figure 3(a)). Our gene network construction and module identification was done via the WGCNA R package's one-step network construction function. Figure 3(b) displayed the color-coded coexpressed gene modules identified via WGCNA method, where the grey by default was those genes that could not be classified into any module. It was found that these modules could be classified into two categories and 23 subclasses, and that there was correlation amid these modules (Figures 3(c) and 3(d)). A second purpose of WGCNA is to analyze the correlation between modules and clinical parameters ( $R$  value). Analysis to the correlation amid the module genes and chRCC showed that the modules




(a)




(b)

FIGURE 1: DEG analysis. (a) heat maps; (b) volcanic map.



(a)




Gene\_number


- 50
- 100
- 150

(b)

FIGURE 2: Continued.




(c)




(d)

FIGURE 2: DEG pathway enrichment analysis. (a and b) KEGG enrichment analysis; (c and d) GO enrichment analysis.



(a)



(b)

FIGURE 3: Continued.




FIGURE 3: Continued.




FIGURE 3: DEG analysis was carried out by weighted gene coexpression network analysis (WGCNA) method and gene cluster tree analysis of modular feature genes. (a) Scale-free exponential analysis to various soft threshold powers ( $\beta$ ). (b) The color of the module represented by each dendrogram of the cluster module of the DEG (top) and the color band (bottom). (c) Clustering dendrograms of different genes based on topological overlap, and the colors assigned to the corresponding modules. (d) Correlation analysis onto different modules. (e) Correlation analysis onto modules and traits. ME: module characteristic gene.

MEblack, MEgreen, and MEtan were significantly positively correlated with the M phase of chRCC, and the correlation coefficients  $r$  were 0.28, 0.26, and 0.32, respectively ( $P < 0.05$ , Figure 3(e)). According to Figure 3(e), MEdarkred was positively correlated with T phase, and the correlation coefficient  $r$  was 0.21 ( $P < 0.05$ ). Figure 3(e) illustrated a positive correlation between MEdarkred and T phase ( $r = 0.21$ ;  $P < 0.05$ ).

**3.3. Module MEtan Gene Functional Enrichment Analysis.** The above analysis led us to select MEtan for further analysis, because it has a significant positive correlation with the M phase of chRCC. GO analysis revealed that steroid metabolism, cholesterol metabolism, and stem cell differentiation were the top chRCC biological processes of MEtan module genes (Figures 4(a) and 4(b)). Genes enriched in cancer-related pathways, such as Neuroactive–Ligand receptor interaction, cAMP signaling pathway, and Wnt signaling pathway, were identified in KEGG analysis. (Figures 4(c) and 4(d)).

**3.4. Screening to Hub Genes.** With the help of the STRING online database and Cytoscape software, DEGs from MEtan modules were analyzed, and PPI networks were constructed in order to identify key genes. Cytoscape’s CytoHubba plugin was used to screen the PPI network for key genes. MAG, CHRM1, and ARR2 were in the center of the 36 nodes in the PPI network for module MEtan (Figure 5(a)).


Finally, ARR2 and MAG were the main genes we screened out (Figure 5(b)).

**3.5. Survival Analysis.** In contrast to the normal group, chRCC patients expressed significantly more ARR2 than do normal individuals (Figure 6(a),  $P < 0.05$ ). Kaplan–Meier survival curves were constructed to analyze chRCC’s overall survival rate. All chRCC samples were divided into high expression group and low expression group of key genes, and compared with the median value of key genes; according to expression survival analysis, ARR2 was negatively correlated with overall survival (Figure 6(b),  $P > 0.05$ ).


## 4. Discussion

ChRCC develops from dark cells in the collecting duct epithelium of the kidney [15]. There was 89.3% recurrence-free survival (RFS) and 93% cancer-specific survival (CSS) rates for chRCC after 5 years [16]. Metastatic disease accounts for only 6% of chRCC patients [17]. However, patients with metastatic chRCC illness have a poor prognosis, who more frequently show nodular characteristics and have a low incidence of treatment response [18, 19]. On postoperative follow up, Geramizadeh et al. found that only 20 (16%) of 123 CRCC patients progressed (local recurrence, metastasis, or death) [20]. Therefore, an in-depth study to related genes coexpressed in various stages and links of






(a)




(b)

FIGURE 4: Continued.



(c)



(d)

FIGURE 4: GO and KEGG analysis for module genes. (a and b) GO-term analysis and (c and d) KEGG enrichment analysis of METan module.






FIGURE 6: Overall survival analysis. (a) expression analysis to ARRB2 the key gene in MEtan module; (b) correlation analysis to ARRB2 expression and chRCC patients' overall survival; KICH: kidney chromophobe.

chRCC and discovery of genes that play a crucial regulatory role in its occurrence; furthermore, the development of the disease is indispensable for understanding its mechanism and improving treatment measures.

An in-depth analysis of the key genes involved in renal chromophobe cell carcinoma development and progression was undertaken in this study. A total of 13466 differentially expressed genes of renal chromophobe cell carcinoma were screened and mined by searching TCGA database, among which 6066 genes were upregulated and 7406 genes were downregulated. Several of these DEGs converged on the signaling pathways involving cAMP, cytokine-cytokine receptor interaction, calcium signaling pathway, and Neuroactive Ligand-receptor interaction.

An advantage of the WGCNA method is that it explores the association between clinical traits and coexpression modules, with higher reliability and biological significance [21]. TCGA database samples were analyzed through the WGCNA method to analyze clinical traits and gene expression data. According to the results, the modules MEblack, MEgreen, and MEtan were positively correlated with the M phase of chRCC; the module MEdarkred was positively correlated with the T phase of chRCC; moreover, MEsalmon is also positively correlated with the stage of chRCC. MEtan, which has a significant positive correlation with the M phase of chRCC, was selected for further analysis. Main chRCC

biological processes of MEtan module genes include steroid metabolic process, cholesterol metabolic process, and STEM cell differentiation; besides, KEGG analysis revealed that these genes were primarily enriched in cancer-related signaling pathways such as Neuroactive Ligand-receptor interaction, cAMP signaling pathway, and Wnt signaling pathway.

Studies have shown that cyclic adenosine monophosphate (cAMP) plays an important role in controlling cell proliferation [22]. A total of 19 secreted glycoproteins make up the Wnt family, which regulates cell proliferation, differentiation, survival, migration, and stem cell self-renewal [23, 24]. There is an association between high Wnt1 expression in ccRCCs, increased tumor diameter, and more advanced stages [25]. A significant increase in WNT10A expression was also observed in RCC cells and tissues, and it plays an oncogenic role [26].

With the help of the STRING online database and Cytoscape software, DEGs from MEtan modules were analyzed, and PPI networks were constructed in order to identify key genes, and the key gene was ARRB2. In comparison with the normal group, ARRB2 expression was significantly higher in chRCC patients. ARRB2 expression was negatively correlated with overall survival, according to an expression survival analysis. There is a widespread expression of Arrb2, a multifunctional protein that regulates the desensitization and intracellular transport of G protein-

coupled receptors (GPCRs) [27, 28]. Furthermore, Arrb2 is involved in a variety of signaling pathways, including those that involve extracellular signal-regulated kinases (ERK) and protein kinase B (Akt) [29, 30].ARRB2 has been shown to be involved in the metastasis of a variety of cancer cells. Defective SUMOylation of ARRB2 inhibits the migration of breast cancer cells and has been shown to be involved in ARRB2-dependent metabolic regulation of breast cancer cells [31]. ARRB2 plays a negative regulatory role in glioma growth, invasion, and metastasis by reducing HIF-1 $\alpha$  expression and inhibiting angiogenesis [32]. It was found that inhibition of ARRB2 expression reduced local and metastatic RCC tumor growth [33]. In summary, ARRB2 may consider as a target for therapeutic intervention against tumour development and metastasis in the studies of future. This study provides a reference for the clinical application of ARRB2 as a prognostic biomarker and potential therapeutic target, and we will enrich its mechanism of action in chRCC through more experiments in the future.

## 5. Conclusion

This study screened TCGA databases for genes associated with chRCC occurrence and development and discussed key genes related to chRCC. A possible therapeutic target and prognostic marker for renal chromophobe cell carcinoma may be ARRB2. However, since there have been no studies on the gene level related to chRCC, there is an urgent need for more research into the biological role of chRCC in renal chromophobe cell carcinoma pathogenesis, so that new clues and directions will be offered for the treatment of renal chromophobe cell carcinoma.

## Data Availability

The data that support this study are in the article.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Authors' Contributions

Yan Cheng and Yang Yan contributed equally to this work.

## Acknowledgments

This study was supported by Chongqing Science and Health Joint Medical Research Project "FMRI's early monitoring of the efficacy of EGFR-TKI in the treatment of brain metastasis in non-small cell lung cancer" (Project no. 2019MSXM011).

## References

- [1] F. Bray, J. Ferlay, I. Soerjomataram, R. L. Siegel, L. A. Torre, and A. Jemal, "Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries," *CA: a Cancer Journal For Clinicians*, vol. 68, no. 6, pp. 394–424, 2018.
- [2] P. A. Humphrey, H. Moch, A. L. Cubilla, T. M. Ulbright, and V. E. Reuter, "The 2016 WHO classification of tumours of the urinary system and male genital organs—part B: prostate and bladder tumours," *European Urology*, vol. 70, no. 1, pp. 106–119, 2016.
- [3] J. Casuscelli, M. F. Becerra, K. Seier et al., "Chromophobe renal cell carcinoma: results from a large single-institution series," *Clinical Genitourinary Cancer*, vol. 17, no. 5, pp. 373–379.e4, 2019.
- [4] M. B. Amin, G. P. Paner, I. Alvarado-Cabrero et al., "Chromophobe renal cell carcinoma: histomorphologic characteristics and evaluation of conventional pathologic prognostic parameters in 145 cases," *The American Journal of Surgical Pathology*, vol. 32, no. 12, pp. 1822–1834, 2008.
- [5] A. M. Badowska-Kozakiewicz, M. P. Budzik, P. Koczkodaj, and J. Przybylski, "Selected tumor markers in the routine diagnosis of chromophobe renal cell carcinoma," *Archives of Medical Science: AMS*, vol. 12, no. 4, pp. 856–863, 2016.
- [6] W. M. Linehan and C. J. Ricketts, "The cancer genome atlas of renal cell carcinoma: findings and clinical implications," *Nature Reviews Urology*, vol. 16, no. 9, pp. 539–552, 2019.
- [7] L. Li, X. Chen, L. Hao, Q. Chen, H. Liu, and Q. Zhou, "Exploration of immune-related genes in high and low tumor mutation burden groups of chromophobe renal cell carcinoma," *Bioscience Reports*, vol. 40, no. 7, 2020.
- [8] G. Martignoni, M. Pea, M. Chilosi et al., "Parvalbumin is constantly expressed in chromophobe renal carcinoma," *Modern Pathology*, vol. 14, no. 8, pp. 760–767, 2001.
- [9] L. Zini, X. Leroy, L. Lemaitre et al., "Tumour necrosis in chromophobe renal cell carcinoma: clinical data to distinguish aggressive variants," *European Journal of Surgical Oncology: the Journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*, vol. 34, no. 6, pp. 687–691, 2008.
- [10] J. Zhou, X. Yang, L. Zhou, P. Zhang, and C. Wang, "Combined immunohistochemistry for the "three 7" markers (CK7, CD117, and Claudin-7) is useful in the diagnosis of chromophobe renal cell carcinoma and for the exclusion of mimics: diagnostic experience from a single institution," *Disease Markers*, vol. 2019, Article ID 4708154, 9 pages, 2019.
- [11] J. C. Cheville, C. M. Lohse, H. Zincke, A. L. Weaver, and M. L. Blute, "Comparisons of outcome and prognostic features among histologic subtypes of renal cell carcinoma," *The American Journal of Surgical Pathology*, vol. 27, no. 5, pp. 612–624, 2003.
- [12] J. Bellmunt, J. Puente, J. Garcia de Muro et al., "SEOM clinical guidelines for the treatment of renal cell carcinoma," *Clinical & Translational Oncology*, vol. 16, no. 12, pp. 1043–1050, 2014.
- [13] J.-G. Zhou, J. Yang, S.-H. Jin et al., "Development and validation of a gene signature for prediction of relapse in stage I testicular germ cell tumors," *Frontiers in Oncology*, vol. 10, p. 1147, 2020.
- [14] J. Zhong, S. Shi, W. Peng et al., "Weighted gene co-expression network analysis (WGCNA) reveals the functions of syndecan-1 to regulate immune infiltration by influenced T cells in glioma," *Frontiers in Genetics*, vol. 13, article 792443, 2022.
- [15] Z. B. Sari, M. F. Açikalin, D. Arik, A. Özen, C. Can, and E. Çolak, "The role of CK7, S100A1, and CD82 (KAI1) expression in the differential diagnosis of chromophobe renal cell carcinoma and renal oncocytoma," *Applied Immunohistochemistry*

- & *Molecular Morphology: AIMM*, vol. 29, no. 7, pp. 534–540, 2021.
- [16] Y. Ayari, S. Ben Rhouma, H. Boussaffa et al., “Metachronous isolated locally advanced pancreatic metastasis from chromophobe renal cell carcinoma,” *International Journal of Surgery Case Reports*, vol. 60, pp. 196–199, 2019.
- [17] Y. Ged, Y.-B. Chen, A. Knezevic et al., “Metastatic chromophobe renal cell carcinoma: presence or absence of sarcomatoid differentiation determines clinical course and treatment outcomes,” *Clinical Genitourinary Cancer*, vol. 17, no. 3, pp. e678–e688, 2019.
- [18] V. Michalaki and C. Gennatas, “Chromophobe renal cell carcinoma with prolonged response to targeted therapy: a case report,” *Journal of Medical Case Reports*, vol. 6, no. 1, p. 115, 2012.
- [19] S. R. Granter and A. A. Renshaw, “Fine-needle aspiration of chromophobe renal cell carcinoma. Analysis of six cases,” *Cancer*, vol. 81, no. 2, pp. 122–128, 1997.
- [20] B. Geramizadeh, M. Ravanshad, and M. Rahsaz, “Useful markers for differential diagnosis of oncocytoma, chromophobe renal cell carcinoma and conventional renal cell carcinoma,” *Indian Journal of Pathology & Microbiology*, vol. 51, no. 2, pp. 167–171, 2008.
- [21] W.-C. Chou, A.-L. Cheng, M. Brotto, and C.-Y. Chuang, “Visual gene-network analysis reveals the cancer gene co-expression in human endometrial cancer,” *BMC Genomics*, vol. 15, no. 1, p. 300, 2014.
- [22] M. Massimi, F. Ragusa, S. Cardarelli, and M. Giorgi, “Targeting cyclic AMP signalling in hepatocellular carcinoma,” *Cells*, vol. 8, no. 12, 2019.
- [23] J. N. Anastas and R. T. Moon, “WNT signalling pathways as therapeutic targets in cancer,” *Nature Reviews Cancer*, vol. 13, no. 1, pp. 11–26, 2013.
- [24] K. Willert and K. A. Jones, “Wnt signaling: is the party in the nucleus?,” *Genes & Development*, vol. 20, no. 11, pp. 1394–1404, 2006.
- [25] S. Kruck, C. Eyrich, M. Scharpf et al., “Impact of an altered Wnt1/ $\beta$ -catenin expression on clinicopathology and prognosis in clear cell renal cell carcinoma,” *International Journal of Molecular Sciences*, vol. 14, no. 6, pp. 10944–10957, 2013.
- [26] R.-J. Hsu, J.-Y. Ho, T.-L. Cha et al., “WNT10A plays an oncogenic role in renal cell carcinoma by activating WNT/ $\beta$ -catenin pathway,” *PLoS One*, vol. 7, no. 10, article e47649, 2012.
- [27] F. K. Kuhr, Y. Zhang, V. Brovkovich, and R. A. Skidgel, “ $\beta$ -arrestin 2 is required for B1 receptor-dependent post-translational activation of inducible nitric oxide synthase,” *FASEB Journal: Official Publication of the Federation of American Societies For Experimental Biology*, vol. 24, no. 7, pp. 2475–2483, 2010.
- [28] L. Wang, Y. Zhu, L. Wang et al., “Effects of chronic alcohol exposure on ischemia-reperfusion-induced acute kidney injury in mice: the role of  $\beta$ -arrestin 2 and glycogen synthase kinase 3,” *Experimental & Molecular Medicine*, vol. 49, no. 6, article e347, 2017.
- [29] J. Kim, L. Zhang, K. Peppel et al., “Beta-arrestins regulate atherosclerosis and neointimal hyperplasia by controlling smooth muscle cell proliferation and migration,” *Circulation Research*, vol. 103, no. 1, pp. 70–79, 2008.
- [30] K. Taguchi, T. Matsumoto, K. Kamata, and T. Kobayashi, “G protein-coupled receptor kinase 2, with  $\beta$ -arrestin 2, impairs insulin-induced Akt/endothelial nitric oxide synthase signaling in Ob/Ob mouse aorta,” *Diabetes*, vol. 61, no. 8, pp. 1978–1985, 2012.
- [31] C. Dong, Y. Li, Q. Niu et al., “Sumoylation involves in  $\beta$ -arrestin-2-dependent metabolic regulation in breast cancer cell,” *Biochemical and Biophysical Research Communications*, vol. 529, no. 4, pp. 950–956, 2020.
- [32] W.-Y. Bae, J.-S. Choi, S. Nam, and J.-W. Jeong, “ $\beta$ -arrestin 2 stimulates degradation of HIF-1 $\alpha$  and modulates tumor progression of glioblastoma,” *Cell Death and Differentiation*, vol. 28, no. 11, pp. 3092–3104, 2021.
- [33] J. Masannat, H. T. Purayil, Y. Zhang et al., “ $\beta$ arrestin2 mediates renal cell carcinoma tumor growth,” *Scientific Reports*, vol. 8, no. 1, p. 4879, 2018.