

# Targeting Mitochondrial Homeostasis: The Role of Acupuncture in Depression Treatment

Haiyang Chen<sup>1,\*</sup>, Chenlin Wu<sup>2,\*</sup>, Qin Lv<sup>2</sup>, Mingjie Li<sup>1</sup>, Lu Ren<sup>2,3</sup>

<sup>1</sup>Department of Acupuncture and Moxibustion, Liaoning University of Traditional Chinese Medicine, Shenyang, 110847, People's Republic of China; <sup>2</sup>Graduate School, Liaoning University of Traditional Chinese Medicine, Shenyang, 110847, People's Republic of China; <sup>3</sup>Mental Disorders Research Laboratory, Liaoning University of Traditional Chinese Medicine, Shenyang, 110847, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Lu Ren, Email [lnzxkc@sina.com](mailto:lnzxkc@sina.com)

**Background:** Depression is a common mental health disorder characterized by persistent feelings of sadness, loss of interest or pleasure, and a range of physical and cognitive symptoms. It affects people of all ages and can significantly impact their daily functioning and quality of life. Mitochondrial homeostasis plays an important role in the pathogenesis of depression. Mitochondrial homeostasis includes mitophagy, mitochondrial oxidative stress, mitoptosis, mitochondrial biogenesis, and mitochondrial dynamics. The regulation of mitochondrial homeostasis is the key link in the prevention and treatment of depression.

**Methods:** In this article, we focus on the core link of depression-mitochondrial homeostasis and summarize the research progress of acupuncture targeting mitochondrial homeostasis in the treatment of depression in recent years, so as to provide ideas and experimental basis for the research and formulation of more appropriate depression treatment strategies.

**Results:** Acupuncture has been found to regulate mitochondrial homeostasis (by modulating mitochondrial autophagy, reducing mitochondrial oxidative stress, inhibiting mitochondrial fission, inducing mitochondrial biogenesis, and maintaining mitochondrial dynamics), alleviate depression-like behavior, and regulate signal pathways and key proteins.

**Conclusion:** Here, we highlight the role of acupuncture in the treatment of depression. A comprehensive exploration of the impact of acupuncture on mitochondrial homeostasis could potentially present a novel mechanism for treating depression and offer fresh perspectives for the treatment of patients with clinical depression.

**Keywords:** depression, mitochondria, homeostasis, acupuncture, mitophagy, review

## Introduction

Depression has become one of the most common public health diseases in the world and the incidence is increasing year by year. People suffering from the disease often show self-neglect, loss of interest, eating disorders, insomnia, despair, lack of concentration, and even suicidal tendencies, and the recurrence rate and disability rate account for a considerable proportion. The disease affects more than 350 million people around the world.<sup>1</sup> With the increase of population and the sharp increase of modern social pressure, the social problems caused by depression are becoming more and more serious. A study predicts that depression will become the second largest disease burden in the world in 2030.<sup>2</sup>

The pathogenesis of depression is very complex, including hypothalamus-pituitary-adrenal axis regulation disorder, insufficient serotonin content, inflammation and cytokines, brain-gut axis imbalance, hypothalamus-pituitary-thyroid axis inhibition, etc. The neural substrate of depression refers to the abnormalities of related neuron activities and neurotransmitters in the brains of patients with depression. Studies have shown that patients with depression often have abnormalities related to neurotransmitters, including dysregulation of neurotransmitters such as serotonin, norepinephrine and dopamine.<sup>3,4</sup> These neurotransmitters play an important role in regulating mood and mental state, and abnormal levels may contribute to the development of depression. The study also found that synaptic plasticity in patients with depression may be affected by abnormal release and reuptake of neurotransmitters, changes in synaptic structure,

inflammation and stress response. These factors may lead to the imbalance of synaptic plasticity, which in turn affects emotional regulation and cognitive function.<sup>5</sup>

Drug intervention is still one of the main methods for the treatment of depression. Monoamine drugs, such as selective serotonin reuptake inhibitors (SSRIs) and selective norepinephrine reuptake inhibitors (SNRI), were first introduced at the turn of the century and have since become standard treatments for depression.<sup>6</sup> Currently, SSRI antidepressants have replaced tricyclic antidepressants (TCAs) as the first choice of antidepressants. There are 6 kinds of SSRIs commonly used in clinic: fluoxetine, paroxetine, sertraline, fluvoxamine, citalopram and escitalopram. SSRI and SNRI drugs are thought to work by increasing the number of neurotransmitters in the brain, such as 5-hydroxytryptamine (5-HT) and dopamine (DA) in synaptic spaces.<sup>7–10</sup> Since the application of molecular biology in medicine, non-monoamine drugs, such as ketamine and angomelatine, have become one of the most popular drugs, which are effective in relieving symptoms and improving prognosis of patients with depression.<sup>11,12</sup> Despite this, the therapeutic effect of existing mainstream antidepressants still has considerable limitations, such as SSRIs taking several weeks to take effect, the drug cycle is long, and it is easy to relapse, and some patients do not respond to various drugs. Therefore, it is particularly important to find new treatment strategies.

As a non-drug therapy, acupuncture has great potential in the treatment of depression and can replace or assist drug therapy to improve the curative effect.<sup>13</sup> Compared with drug therapy, acupuncture has the advantages of low cost and less side effects.<sup>14,15</sup> As early as 1984, a study confirmed that acupuncture has a unique therapeutic effect on depression.<sup>16</sup> The report shows that the overall efficacy of various acupuncture methods in the treatment of depression after 6 weeks is about 70% to 83.7%, which is equivalent to or even higher than that of antidepressant western drugs.<sup>17</sup> The results of Meta-analysis showed that acupuncture could significantly reduce the degree of depression compared to the routine treatment group, the sham acupuncture group, and the antidepressant treatment group, and there was a significant correlation between the increase in acupuncture treatment times and the decrease in severity of depression.<sup>18</sup> The latest research has proved that acupuncture can treat depression in many ways (Table 1).

New research has shown that acupuncture can treat depression in a variety of ways, including targeting mitochondrial homeostasis. Mitochondria are one of the most important organelles in the human body, it was once regarded as a simple “living energy metabolism factory”, which synthesizes adenosine triphosphate to supply energy for the body.<sup>39</sup> After nearly a century of research, it has been found that it has many functions: regulating gene expression in the nucleus, regulating synaptic transmission in the brain, releasing inflammatory factors and triggering a systemic inflammatory response, affecting the regulation of complex physiological systems.<sup>40</sup> People with depression often have abnormal energy metabolism, including mitochondrial dysfunction. The study found that there are morphological and functional changes in the mitochondria in patients with depression compared to normal people. These mitochondrial abnormalities may lead to disorders of energy metabolism, leading to symptoms of depression. The abnormal function of mitochondria may also lead to the increase of oxidative stress, inflammation and apoptosis. These changes may be closely related to the occurrence and development of depression.<sup>41,42</sup> In addition, mitochondria are involved in the synthesis and regulation of neurotransmitters. Neurotransmitters are chemicals in the brain that transmit neural signals. The abnormal function of mitochondria can lead to abnormal synthesis and release of neurotransmitters, affect the transmission of neural signals and communication between brain regions, and then affect emotional and cognitive function. The study also found that some antidepressants may play a therapeutic role by regulating mitochondrial function. This suggests that repairing mitochondrial function may be a new direction in the treatment of depression.

Mitochondria are highly dynamic organelles that undergo continuous fission and fusion through mitochondrial biogenesis and mitochondrial autophagy to maintain mitochondrial homeostasis.<sup>43</sup> Mitochondrial homeostasis consists of many aspects: mitochondrial autophagy, mitochondrial oxidative stress, mitoptosis, mitochondrial biogenesis, mitochondrial dynamics (Figure 1). Studies have shown that the destruction of mitochondrial homeostasis is an important pathological mechanism for the occurrence and development of depression. In the process of depression, chronic stress will lead to an imbalance of homeostasis of human physiological function and damage the morphology and function of mitochondria through a variety of pathological pathways.

**Table 1** Summary of Researches Regarding the Effect of Acupuncture in Depression


Ref.	Model	Number of Animals	Sex	Acupoint	Type/Frequency/Time	Days	Treatment Country/Region	Year	Major Efficacy
[19]	CUMS rat model	36	Male	Fengfu (GV16)	MA/none/20min	28	China	2023	Inhibit activation of HMGB1/TLR4 in amygdala of CUMS rats
[20]	FS mouse model	200	Male	Shangxing (GV23) Baihui (GV20)	MA/none/20min	10	Japan	2023	Inhibit inflammation Increased the expression of neurotrophic factors (NGF, BDNF, NT-3, NT-4/5)
[21]	LPS-induced depression mouse model	18	Male	Yintang (GV29) Taichong (LR3)	EA/2Hz/50Hz/20min	14	China	2022	Inhibit collagen type IV trimer, extra cellular matrix organization, and collagen formation
[22]	CUMS rat model	36	Male	Yintang (GV29) Baihui (GV20) Hegu (LI4) Fengfu (GV16)	MA/none/20min	14	China	2022	Enhance MMP1 and MMP9 Promote synaptic plasticity Suppress NLRP3, ASC, caspase-1, GSDMD, IL-1 $\beta$ , IL-18, HMGB1, IFN- $\gamma$ , IL-6 and TNF- $\alpha$
[23]	CUMS rat model	40	Male	Shangxing (GV23) Fengfu (GV16)	MA/none/20min	14	China	2022	Reduce pyroptosis and inflammation Reduce oxidative stress
[24]	CMS mouse model	40	Male	Shangxing (GV23) Baihui (GV20) Shenshu (BL23) Taixi (KI3)	EA/2Hz/15min	21	China	2022	Prevent neuronal apoptosis Inhibit NF- $\kappa$ B/NLRP3 inflammatory pathway
[25]	CUMS rat model	32	Male	Baihui (GV20) Yintang (GV29)	MA/none/10min	36	China	2021	Inhibit NLRP3 inflammasome activation and apoptosis
[26]	SI-induced depression rat model	40	Male	Baihui (DU20)	AC/none/none	-	Mexico	2021	Improve sex hormones release  Increase BDNF hippocampal levels
[27]	CUMS rat model	32	Male	Yintang (EX-HN3) Shenshu (BL23) Pishu (BL20) Ganshu (BL18) Xinshu (BL15) Guanyuan (Ren4) Shangxing Daling	MA/none/20min	21	China	2021	Increase the level of DA and 5-HT Increase the expression of brain-derived neurotrophic factor signaling and the astrocytes in the hippocampus

(Continued)

Table 1 (Continued).

Ref.	Model	Number of Animals	Sex	Acupoint	Type/Frequency/Time	Days	Treatment Country/Region	Year	Major Efficacy
[28]	CRS mouse model	27	Male	Yingu (KI10) Ququan (LR8) Jingqu (LU8) Zhongfeng (LR4)	MA/none/30sec	7	Korea	2021	Decrease AST, IL-1 $\beta$ , TNF- $\alpha$ Attenuate leptin insensitivity
[29]	CUMS rat model	40	Male	Baihui (GV20) Yintang (GV29)	EA/2Hz/30min	14	China	2021	Up-regulate the expression level of FGF2 Increase GFAP and the mean optical density of GFAP-immunoreactive astrocyte (GFAP-ir astrocyte)
[30]	CUMS rat model	48	Male	Hegu (LI4)	EA/15Hz/30min	21	China	2021	Decrease hippocampal glutamate, Bax, caspase 3, CaMKII, NR2B Increase Bcl-2, NR2A
[31]	CUMS rat model	120	Male	Taichong (LR3) Baihui (GV20) Shenting (GV24)	EA/2Hz/20min	21	China	2020	Increase CaMKII and CaMKIV
[32]	HIS rat model	75	Male	Zusanli (ST36)	EA/100Hz/2Hz/20min	9	China	2020	Regulate glucose metabolism
[33]	CUMS rat model	40	Male	Baihui (GV20) Yintang (GV29)	EA/2Hz/30min	14	China	2020	Upregulate 5-HT1A receptor Improve synaptic plasticity in the hippocampus
[34]	OVX rat model	32	Female	Sanyinjiao (SP6)	MAI/none/1min	8	Korea	2019	Mitigate ER stress and oxidative stress in amygdala
[35]	CRS mouse model	-	Male	Yingu (KI10)  Ququan (LR8) Jingqu (LU8) Zhongfeng (LR4) Shaofu (HT8) Xingjian (LR2)	MA/none/30sec	7/14	Korea	2019	Increase expression of 5-HT1A receptor in the cortex, hippocampus, thalamus, and the hypothalamus  Increase 5-HT1B in the cortex and thalamus
[36]	WKY rat model	24	Male	Baihui (GV20)  Yintang (EX-HN3)	EA/2Hz/15min	21	China	2018	Decrease 5-HTT protein expression in the hippocampus CA1 region Downregulate the expression of the 5-HT1A protein in the hippocampus CA1 region Restore hippocampus CA1 synaptic plasticity
[37]	CUS rat model	33	Male	Baihui (Du20)  Yanglingquan (GB34)	EA/2/100Hz/30min	14	China	2018	Decrease NLRP3, IL-1 $\beta$ , P2X7 receptor, Iba-1, IL-18, TNF $\alpha$ and IL-6 Increase GFAP Alleviate the hippocampal neuroinflammation
[38]	CUMS rat model	40	Male	Baihui (GV20)  Yintang (GV29)	MA/none/20min	28	China	2018	Upregulate BDNF in serum and hippocampus Upregulate hippocampal acH3K9 Downregulate hippocampal HDAC2

**Abbreviations:** CUMS, chronic unpredictable mild stress; FS, forced swimming; LPS, lipopolysaccharide; CMS, chronic mild stress; SI, social isolation; CRS, chronic restraint stress; HIS, heterogeneous intermittent stress; OVX, ovariectomize; WKY, Wistar Kyoto; EA, electroacupuncture; MA, manual acupuncture; AC, acupoint catgut; MAI, mechanical acupuncture instrument; NGF, nerve growth factor; BDNF, brain-derived neurotrophic factor; NT, neurotrophin; CaMKII, calcium/calmodulin-dependent protein kinase II; GFAP, glial fibrillary acidic protein; acH3K9, acetylation levels in histone H3 lysine 9.



**Figure 1** The schematic diagram of mitochondrial homeostasis. (I) The damaged mitochondria are specifically wrapped in autophagosomes and fused with lysosomes to complete the degradation of mitochondria and maintain mitochondrial homeostasis. (II) The imbalance between oxidation and antioxidation in mitochondria produces a large amount of mROS, which leads to mitochondrial oxidative stress. (III) Mitochondrial biogenesis is defined as the process by which cells increase the mass of their individual mitochondria. (IV) With the continuous increase of ROS in mitochondria, the increase of Ca<sup>2+</sup> released to mediate the imbalance of mitochondrial energy metabolism homeostasis, which led to the release of pro-apoptotic protein cytochrome c (Cyto c) and the activation of apoptotic caspase pathway, the mitoptosis was elicited. (V) Mitochondrial dynamics refers to the dynamic balance between fusion and fission of mitochondria.

Therefore, this paper reviews the related research in order to clarify the new mechanism of acupuncture in the treatment of depression: targeting mitochondrial homeostasis and providing a new strategy for the treatment of patients with depression.

## Regulate Mitophagy

Mitophagy is a regulatory mechanism that controls the quality and quantity of mitochondria and maintains the general homeostasis of the structure and function of mitochondria in the cellular environment. It specifically recognizes dysfunction or excess mitochondria, recruits the autophagy structure in the cytoplasm to wrap the mitochondria to be degraded to form autophagosomes, and finally binds to lysosomes to complete the degradation of autophagosomes.<sup>44</sup> Abnormal mitophagy is one of the pathogenesises of depression.<sup>45</sup> When the disturbance of mitophagy occurs, the undegradable damaged mitochondria accumulate in the cell, which leads to the decrease of membrane potential and triggers downstream events such as apoptosis and oxidative stress that destroy the homeostasis of the internal environment.<sup>46</sup>

An animal-model study involving 60 SD rats demonstrated that electroacupuncture partially inhibits mitophagy by partially inhibiting the level of mitophagy in the hippocampus of CUMS rats, reducing the number of autolysosomes and the level of Lc3-II/Lc3-I, thus exerting the antidepressant effect.<sup>47</sup> Mitophagy and structural changes of mitochondria are the main causes of neuronal dysfunction.<sup>48</sup> Ge et al<sup>49</sup> established a diabetic model of adult SD rats by using

streptozotocin (STZ) have shown that electroacupuncture promotes mitophagy and improves the learning and memory function of rats by increasing the expression of Disrupted-in-Schizophrenia 1 (DISC1). The down-regulation of DISC1 leads to mitochondrial dysfunction and synaptic plasticity, resulting in cognitive impairment.<sup>50</sup> Studies have shown that electroacupuncture activates mitophagy by regulating the steps in the process of mitophagy, including autophagy induction, phage elongation, vacuole formation and substrate degradation.<sup>51</sup> Zhong et al<sup>52</sup> made the rat model of brain injury induced by middle cerebral artery occlusion (MCAO) demonstrated that electroacupuncture inhibits the activation of NLRP3 inflammatory bodies by regulating melatonin-mediated mitophagy, thus improving the cognitive impairment in stroke rats. It has been found that EA ameliorates nitro/oxidative stress-induced mitochondrial functional damage and decreases the accumulation of damaged mitochondria via Pink1/Parkin-mediated mitophagy clearance to protect cells against neuronal injury in cerebral I/R.<sup>53</sup>

In summary, electroacupuncture can effectively regulate mitophagy, improve mitochondrial dysfunction, and play an antidepressant role in a variety of ways. However, although the mitophagy of depression has been well studied and a certain number of acupuncture and mitophagy studies have been published, the comprehensive study on the effect of acupuncture on mitophagy of depression is still limited.

## Reduce Mitochondrial Oxidative Stress

Mitochondria are the main source of reactive oxygen species (ROS), and more than 95% of the ROS in the body come from mitochondria.<sup>54</sup> Extensive studies have shown that mitochondrial reactive oxygen species (mROS) are essential for healthy cell function.<sup>55</sup> ROS is a component of the cellular signal transduction pathway, which can activate various transcription factors and lead to the expression of proteins that regulate inflammation, cell transformation, tumor cell survival, tumor cell proliferation and invasion, angiogenesis, and metastasis. ROS also controls the expression of various tumor suppressor genes (p53, Rb, and PTEN genes) and inhibits tumor progression. Most chemotherapy and radiotherapy drugs act by increasing ROS stress in tumor cells. Sies<sup>56</sup> introduced the concept of oxidative stress, that is, the destruction of oxidative-antioxidant balance. Oxidative stress represents an imbalance between the production and expression of ROS and the ability of biological system to detoxify intermediates or repair damage. Oxidative stress refers to the imbalance caused by excessive ROS or oxidants exceeding the effective antioxidant response of cells. Oxidative stress is harmful because it can lead to cell biomolecule damage and mutation and growth inhibition, and is involved in the occurrence of various diseases such as atherosclerosis, diabetes, cancer, neurodegenerative diseases, and aging.<sup>57</sup> Over the past decade, oxidative stress has emerged as a leading cause of depression.<sup>58,59</sup> Increased ROS and amplified expression of genes controlled by oxidative stress are closely associated with the development of depression.<sup>60</sup>

Previous studies have shown that acupuncture improves depression-like behavior in CUMS rats, which is associated with oxidative stress in the hippocampus. Acupuncture reduces oxidative stress products by regulating the Nrf2/HO-1 signaling pathway, thereby preventing neuronal apoptosis and playing an antidepressant role.<sup>23</sup> Acupuncture has become a common complementary and alternative treatment of depression, an animal experiment in the OVX-induced depression rat model, showed that acupuncture can relieve oxidative stress in the amygdala, change in oxidative stress-related protein (8-OHdG, BNP, pJNK), play a significant antidepressant effect.<sup>34</sup> Fang et al<sup>61</sup> established a model of ovarian dysfunction by gavage of Tripterygium Glycosides suspension (50 mg·kg<sup>-1</sup>·d<sup>-1</sup>) for 14 successive days, demonstrated that acupuncture can increase SOD content, decrease MDA content, and improve the antioxidant stress ability of rats.

In summary, the incidence of depression is closely related to oxidative stress, which can lead to oxidative damage of proteins, lipids and DNA. Studies have confirmed that oxidative stress may be the main influencing factor of cognitive impairment in depression.<sup>62</sup> Both electroacupuncture and manual acupuncture can inhibit oxidative stress.<sup>63</sup> Therefore, an in-depth study of the exact mechanism of acupuncture stimulation of the antidepressant effect by inhibiting oxidation is particularly important for the treatment of depression.

## Inhibit Mitoptosis

Apoptosis, also known as programmed cell death, is an active death process in order to better adapt to the living environment. Mitochondria are the center of energy metabolism in eukaryotic cells, and the loss of their normal function will lead to apoptosis, and the caspase feedback cycle in mitochondria further amplifies the effect of apoptosis. Therefore,



mitochondria are called the “life-and-death switch” of apoptosis. Mitochondrial apoptosis pathways include exogenous and endogenous pathways. The stress signals of these two apoptosis pathways are concentrated in mitochondria and regulate apoptosis through proteins of the Bcl-2 family. Bcl-2 in mitochondria induces the opening of the transition pore of permeability of the mitochondrial membrane or precisely changes the permeability of the mitochondrial membrane through the release of  $\text{Ca}^{2+}$ , thus initiating cell apoptosis procedures. Exogenous stress signal induces caspase-8-mediated Bid activation by binding to death receptors on the cell membrane. In the endogenous mitochondrial apoptosis pathway, BH3 activation is different due to different BH3 structures in different cells, but all can up-regulate the activity of the pro-apoptotic protein BH3. Therefore, whether the activation of Bid in the exogenous pathway or the up-regulation of BH3 activity in the endogenous pathway, can activate the pro-apoptotic molecule Bak, thus forming pores on the mitochondrial outer membrane to release apoptotic factors, that is, the permeability of the mitochondrial outer membrane. In this process, apoptosis-related factors released from mitochondria, such as cytochrome C (CytC), apoptosis-inducing factor (AIF), and mitochondrial-derived second caspase activator (Smac), can trigger apoptosis once these factors are released from mitochondria.

Mitoptosis was first introduced into the scientific literature by Vladimir P. Skulachev in 1999 to represent the programmed death (elimination) of mitochondria in living cells. He proposed that mitoptosis is the driving force behind apoptosis.<sup>64,65</sup> Mitoptosis is a kind of programmed destruction of mitochondria independent of the cystatin pathway, which often leads to the opening of mitochondrial membrane permeability transition pores mediated by reactive oxygen species, structural changes, rupture and fragmentation of mitochondrial membrane system, and its serious consequences can lead to cell death.

It has been found that acupuncture can reduce oxidative stress products, down-regulate caspase 3 expression, up-regulate Bcl-2 expression, inhibit neuronal apoptosis and exert antidepressant effect by regulating Nrf2/HO-1 signal pathway.<sup>23</sup> Guo et al<sup>30</sup> confirmed the antidepressant effect of electroacupuncture in CUMS rats. Electroacupuncture can reduce Bax and caspase 3 levels in the hippocampus, increase Bcl-2 expression, and regulate apoptosis. Dai et al<sup>66</sup> detected the apoptosis rate of hippocampal cells by Annexin V fluorescein isothiocyanate (FITC) / Propidium iodide (PI) (Annexin V-FITC/PI) double-staining, which confirmed that electroacupuncture can inhibit hippocampal cell apoptosis. A scalp acupuncture study also demonstrated the effect of acupuncture on inhibiting cell apoptosis, which found that scalp acupuncture can increase the expression of Beclin1, Parkin, PINK1, NIX in brain tissue, reduce caspase 3, and thus inhibit nerve cell apoptosis.<sup>67</sup>

To sum up, acupuncture can inhibit apoptosis by regulating signal pathways and the expression of key proteins, thus playing an antidepressant role. At present, there are few studies focusing on the regulation of mitoptosis by acupuncture. Combined with the above contents, the mitoptosis is crucial in the pathogenesis of depression. If further studies on the antidepressant effect of the regulation of the mitochondrial apoptosis pathway by acupuncture will certainly promote people's understanding of depression. If we make an in-depth study on the antidepressant effect of acupuncture regulating mitoptosis, it will certainly promote people's understanding of depression.

## Induce Mitochondrial Biogenesis

Mitochondrial biogenesis refers to the increase in the number and/or quality of mitochondria and the individual and systematic production of new mitochondria in cells.<sup>68</sup> The increase of the number of mitochondria depends on the biogenesis of mitochondria to produce new mitochondria. The biogenesis of mitochondria is largely regulated at the transcriptional level. A series of transcription factors and transcriptional coactivators are involved in the regulation of mitochondrial biogenesis. It is currently acknowledged that the coordination of mitochondrial biogenesis is achieved by the peroxisome proliferator-activated receptor- $\gamma$  coactivator-1 $\alpha$  (PGC-1 $\alpha$ ), which plays an integral role in inducing the transcription of genes encoded by both the nuclear and mitochondria.<sup>69</sup> Puigserver et al<sup>70</sup> first found that the ectopic expression of PGC-1 $\alpha$  in white adipocytes activated the expression of uncoupling protein-1 (UCP-1) and key mitochondrial enzymes in the respiratory chain, which increased the content of mitochondrial DNA (mtDNA) in adipocytes. Upstream AMP-activated protein kinase (AMPK) binds and activates PGC-1 $\alpha$  in muscle by directly phosphorylating two key residues threonine-177 and serine-538.<sup>71</sup> P38 mitogen activated protein kinase (p38MAPK) phosphorylates amino acid residues 262,265,298 directly, inhibits the degradation of PGC-1 $\alpha$  protein and increases the half life of PGC-1 $\alpha$ .<sup>72</sup>

PGC-1 $\alpha$  then binds to the amino acid 180–403 of downstream nuclear respiratory factor 1 (NRF-1) and strongly induces the expression of the NRF-1/2 gene.<sup>73</sup> NRF-1 and NRF-2 directly stimulate the mitochondrial transcription factor A (TFAM) promoter through the binding sites of TFAM-76/58 and TFAM 34/13, respectively, and participate in the activation of the TFAM promoter, which promotes the transcription and replication of mtDNA, and enhances the level of mitochondrial biogenesis.<sup>74,75</sup> In addition to TFAM, PGC-1 $\alpha$ -NRF-1/2 pathway also upregulates mitochondrial transcription factors B1 and B2 (TFB1M / TFB2M) and significantly enhances mtDNA transcription.<sup>76,77</sup> PGC-1 $\alpha$  promotes mtDNA replication and mitochondrial biogenesis by interacting with its upstream and downstream factors.

A study shows that mitochondrial biogenesis is reduced in patients with major depression. In this study, the expression of genes related to mitochondrial biogenesis in patients' blood monocytes was analysed. Researchers found that the expression of genes related to mitochondrial biogenesis (PGC-1 $\alpha$ , TFAM, NRF1) decreased, the expression of antioxidant genes (CuZnSOD and MnSOD) was downregulated, and the level of ATP decreased significantly, indicating that the mitochondrial ability of patients with severe depression is reduced.<sup>78</sup> Tang et al<sup>79</sup> found that electroacupuncture can increase the protein expression of PGC-1 $\alpha$ , TFAM, UCP-1, SIRT-1 and PPAR $\gamma$ , and induce mitochondrial biogenesis through PGC-1 $\alpha$ -TFAM-UCP1 pathway. Electroacupuncture can up-regulate NRF1 and TFAM expression, increase mtDNA level and mitochondrial volume and number, improve mitochondrial function, and induce mitochondrial biogenesis through CB1R/PGC- $\alpha$ .<sup>80</sup>

To sum up, the decrease of mitochondrial biogenesis is an important cause of depression. Acupuncture can effectively improve the function of mitochondria, increase the level of ATP, up-regulate the expression of PGC-1 $\alpha$ , TFAM and NRF-1, and induce mitochondrial biogenesis. Further study on the exact mechanism of mitochondrial biogenesis induced by acupuncture is helpful to solve the difficult problem of depression treatment.

## Maintain Mitochondrial Dynamics

Mitochondria adapt to a variety of stress conditions to meet cell energy metabolism and other biological needs through continuous fusion and fission. This biological process is called mitochondrial dynamics. Mitochondria regulate their morphology, quantity, distribution, and function through fusion and fission, and regulate the morphology of mitochondria in two diametrically opposite ways.<sup>81,82</sup> When fusion increases or fission decreases, it promotes the formation of an extended tubular structure of mitochondria, while when fusion decreases and fission increases, mitochondria show a granular structure.<sup>83</sup> The fusion and fission of mitochondria can be regulated by the highly conserved GTPase protein family. The GTPase protein family completes the remodeling of the inner and outer membrane of mitochondria by self-assembly and hydrolysis of GTP. The fusion of mitochondria can be divided into two parts: the fusion of outer membrane and the fusion of inner membrane. The fusion of the outer membrane of mitochondria is accomplished by GTPase proteins mitofusin 1 (Mfn1) and mitofusin 2 (Mfn2) located in the outer membrane of mitochondria, while the fusion of the inner membrane of mitochondria is accomplished by optic atrophy 1 (OPA1). Different from mitochondrial fusion, mitochondrial fission is mediated by a dynamin-1-like protein 1 (Drp1) with a molecular weight of 80 kDa located in the cytoplasmic matrix. The receptor proteins that can recruit Drp1 are distributed on the outer membrane of mitochondria: mitochondrial fission factor (Mff), mitochondrial dynamics protein of 49 and 51 kDa (Mid49/51) and mitochondrial fission protein 1 (Fis1). These proteins can recruit Drp1 to the outer membrane of mitochondria and further form a spiral ring structure, and then drive mitochondrial constriction and transection by hydrolyzing GTP.<sup>84</sup> Abnormal changes in mitochondrial dynamics can mediate the occurrence of depression; especially the overactivation of Drp1 can lead to increased mitochondrial division, resulting in functional abnormalities, damage to the synaptic microenvironment, and inducing depression-like behaviors.<sup>85</sup>

It has been found that electroacupuncture can regulate the mitochondrial dynamics mediated by the HO-1/PINK1 pathway, increase the content of ATP and the level of mitochondrial fusion protein, and maintain mitochondrial homeostasis.<sup>86</sup> Yong et al<sup>87</sup> found that electroacupuncture at “Zusanli” can improve the level of fission and fusion of mitochondria, and have a certain effect on the dynamic balance of mitochondria. By increasing the expression of fusion protein Opa1 and mitotic protein Drp1, a new balance between fusion and fission of mitochondria is established. This balance is beneficial to the synthesis of mitochondrial ATP and the improvement of mitochondrial function.



In summary, when the balance between mitochondrial fusion and fission is disrupted, mitochondrial dynamics will be abnormal and depression will be induced. Electroacupuncture intervention can help mitochondrial fusion and fission to restore balance, increase ATP content, and maintain mitochondrial homeostasis. The research on acupuncture-mitochondrial dynamics-depression is still shallow, and in-depth research on the exact mechanism of acupuncture regulating mitochondrial dynamics may provide new ideas for the treatment of depression.

## Regulatory Neuroplasticity

Changes in neuroplasticity play an important role in depression. Depression is an emotional and psychological disorder, which is related to the abnormal function of the nervous system.<sup>88</sup> Acupuncture can interfere with neuroplasticity in many ways so as to assist the treatment of depression. Acupuncture can regulate the levels of a variety of neurotransmitters, such as 5-HT, dopamine and glutamic acid. These neurotransmitters play an important role in the development and occurrence of depression.<sup>89</sup> By regulating the balance of neurotransmitters, acupuncture can affect the neural circuits in the brain associated with mood regulation. In addition, acupuncture can promote the growth and regeneration of nerve cells, which is very important in restoring neurological function and improving symptoms of depression. Acupuncture stimulation can stimulate the brain to release growth factors, such as Neurotrophic factors (NTFs), through signal transduction of the nervous system, thus stimulating the growth and differentiation of nerve cells.<sup>20</sup> Studies have found that patients with depression are often accompanied by chronic inflammation. Acupuncture can improve the symptoms of depression by regulating the function of the immune system and reducing the inflammatory response.<sup>90</sup> Acupuncture can also change the electrical activity pattern of the brain and regulate the interaction between brain regions.<sup>91</sup> This is beneficial for the treatment of depression, because the brain activity of patients with depression is often abnormal. In summary, acupuncture can affect neuroplasticity through many mechanisms and improve symptoms of depression.

## Limitations

Currently, there are no human studies to support our proposed mechanism of acupuncture in the treatment of depression by regulating mitochondrial homeostasis. This paper only analyzes the animal experiments, which may have a certain deviation on the results.

## Conclusions

Depression seriously harms human physical and mental health, with the symptoms of depression, lack of interest or pleasure, and decreased will and behavior. With the acceleration of the pace of modern life and the increase of social pressure, the incidence rate has increased year by year in recent years. Depression has a high suicide rate, which brings a heavy burden to patients and families. At present, the commonly used drugs for the treatment of depression and other mental diseases have a variety of adverse reactions such as gastrointestinal tract, cardiovascular system and central nervous system, as well as the safety of drug overdose, drug withdrawal syndrome and many other deficiencies. Acupuncture, as a convenient and quick treatment method, can prevent and cure diseases by stimulating the biological function of the human body, significantly improving symptoms, and having no side effects. Studies have demonstrated the efficacy and safety of acupuncture in antidepressant treatment.<sup>15,92</sup> The research on the antidepressant mechanism of acupuncture has become the focus of many studies. At present, studies in various countries have focused on the effects of acupuncture on behavior, neuroendocrine, neurotransmitters, cytokines, neurotrophic regeneration, cellular signal transduction pathways, and related gene proteins. This paper reviews the research on the antidepressant effect of acupuncture by regulating mitochondrial homeostasis. It is found that acupuncture can regulate mitochondrial homeostasis (regulate mitochondrial autophagy, reduce mitochondrial oxidative stress, inhibit mitoptosis, induce mitochondrial biogenesis, and maintain mitochondrial dynamics), reduce depression-like behavior, regulate signal pathways and key proteins. However, the exact mechanism and relationship of acupuncture in the treatment of depression by regulating mitochondrial homeostasis is not clear. Therefore, in-depth study of acupuncture targeting mitochondrial homeostasis may be a new mechanism for the treatment of depression and provide new ideas for the treatment of patients with clinical depression.

## Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This study was financially supported by the National Natural Science Foundation of China (No. 82274665), the National High Technology Research and Development Program of China (No. 2018YFC1704300), and the Special Professor Project of Liaoning Province (Liao Jiao fa [2015] No.153).

## Disclosure

All authors declare that there are no conflicts of interest regarding the publication of this review.

## References

- Borhannejad F, Shariati B, Naderi S, et al. Comparison of vortioxetine and sertraline for treatment of major depressive disorder in elderly patients: a double-blind randomized trial. *J Clin Pharm Ther.* 2020;45(4):804–811. doi:10.1111/jcpt.13177
- Mathers CD, Loncar D, Samet J. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med.* 2006;3(11):e442. doi:10.1371/journal.pmed.0030442
- Battaglia S, Di Fazio C, Vicario CM, Avenanti A. Neuropharmacological modulation of N-methyl-D-aspartate, noradrenaline and endocannabinoid receptors in fear extinction learning: synaptic transmission and plasticity. *Int J Mol Sci.* 2023;24(6):5926. doi:10.3390/ijms24065926
- Battaglia S, Nazzi C, Thayer JF. Fear-induced bradycardia in mental disorders: foundations, current advances, future perspectives. *Neurosci Biobehav Rev.* 2023;149:105163. doi:10.1016/j.neubiorev.2023.105163
- Battaglia S, Cardellicchio P, Di Fazio C, Nazzi C, Fracasso A, Borgomaneri S. Stopping in (e)motion: reactive action inhibition when facing valence-independent emotional stimuli. *Front Behav Neurosci.* 2022;16:998714. doi:10.3389/fnbeh.2022.998714
- Suchting R, Tirumalajuru V, Gareeb R, et al. Revisiting monoamine oxidase inhibitors for the treatment of depressive disorders: a systematic review and network meta-analysis. *J Affect Disord.* 2021;282:1153–1160. doi:10.1016/j.jad.2021.01.021
- Idova GV, Alperina EL, Chaido MA. Contribution of brain dopamine, serotonin and opioid receptors in the mechanisms of neuroimmunomodulation: evidence from pharmacological analysis. *Int Immunopharmacol.* 2012;12(4):618–625. doi:10.1016/j.intimp.2012.02.010
- Mawe GM, Hoffman JM. Serotonin signalling in the gut—functions, dysfunctions and therapeutic targets. *Nat Rev Gastroenterol Hepatol.* 2013;10(8):473–486. doi:10.1038/nrgastro.2013.105
- He F, Zhang P, Zhang Q, et al. Dopaminergic projection from ventral tegmental area to substantia nigra pars reticulata mediates chronic social defeat stress-induced hypolocomotion. *Mol Neurobiol.* 2021;58(11):5635–5648. doi:10.1007/s12035-021-02522-7
- Li YF. A hypothesis of monoamine (5-HT) - Glutamate/GABA long neural circuit: aiming for fast-onset antidepressant discovery. *Pharmacol Ther.* 2020;208:107494. doi:10.1016/j.pharmthera.2020.107494
- Perrine SA, Ghodoussi F, Michaels MS, Sheikh IS, McKelvey G, Galloway MP. Ketamine reverses stress-induced depression-like behavior and increased GABA levels in the anterior cingulate: an 11.7 T 1H-MRS study in rats. *Prog Neuropsychopharmacol Biol Psychiatry.* 2014;51:9–15. doi:10.1016/j.pnpbp.2013.11.003
- Lorman WJ. Pharmacology corner: esketamine (Spravato)—a new novel medication to treat depression—but with a strong warning. *J Addict Nurs.* 2019;30(4):282–283. doi:10.1097/JAN.0000000000000315
- Smith CA, Armour M, Lee MS, Wang LQ, Hay PJ. Acupuncture for depression. *Cochrane Database Syst Rev.* 2018;3(3):CD004046. doi:10.1002/14651858.CD004046.pub4
- Wang Z, Wang X, Liu J, et al. Corrigendum to “Acupuncture treatment modulates the corticostriatal reward circuitry in major depressive disorder” [J. Psychiatr. Res. 84 (2017) 18–26]. *J Psychiatr Res.* 2017;87(87):105. doi:10.1016/j.jpsychires.2016.12.017
- Chan YY, Lo WY, Yang SN, Chen YH, Lin JG. The benefit of combined acupuncture and antidepressant medication for depression: a systematic review and meta-analysis. *J Affect Disord.* 2015;176:106–117. doi:10.1016/j.jad.2015.01.048
- Luo HC, Zhan L. The observation of therapeutic effects on the treatment of depressive. *Chin Acupunct Moxibustion.* 1984;01:1–4.
- Luo HC, Halbrich U, Shen YC. Comparative study of electroacupuncture and fluoxetine for treatment of depression. *Chin J Psychiatry.* 2003;04:26–30.
- Armour M, Smith CA, Wang LQ, et al. Acupuncture for depression: a systematic review and meta-analysis. *J Clin Med.* 2019;8(8):1140. doi:10.3390/jcm8081140
- Chen W, Chen Y, Cheng W, et al. Acupuncture exerts preventive effects in rats of chronic unpredictable mild stress: the involvement of inflammation in amygdala and brain-spleen axis. *Biochem Biophys Res Commun.* 2023;646:86–95. doi:10.1016/j.bbrc.2023.01.046
- Yamamoto T, Kawanokuchi J, Nagaoka N, et al. Antidepressant effects of acupuncture in a murine model: regulation of neurotrophic factors. *Acupunct Med.* 2023;41(1):38–47. doi:10.1177/09645284221085279
- Feng Y, Zhu G, Chen R, et al. Electroacupuncture remodels the extracellular matrix and promotes synaptic plasticity in a mouse model of depression. *Biochem Biophys Res Commun.* 2022;626:44–50. doi:10.1016/j.bbrc.2022.07.077
- Chen Y, Hao C, Chen W, et al. Anti-depressant effects of acupuncture: the insights from NLRP3 mediated pyroptosis and inflammation. *Neurosci Lett.* 2022;785:136787. doi:10.1016/j.neulet.2022.136787

23. Cheng WJ, Li P, Huang WY, et al. Acupuncture relieves stress-induced depressive behavior by reducing oxidative stress and neuroapoptosis in rats. *Front Behav Neurosci.* 2021;15:783056. doi:10.3389/fnbeh.2021.783056
24. Wang Q, Bi H, Huang H, et al. Electroacupuncture prevents the depression-like behavior by inhibiting the NF- $\kappa$ B/NLRP3 inflammatory pathway in hippocampus of mice subjected to chronic mild stress. *Neuropsychobiology.* 2022;81(3):237–245. doi:10.1159/000521185
25. Li X, Wang H, Li C, et al. Acupuncture inhibits NLRP3 inflammasome activation in the prefrontal cortex of a chronic stress rat model of depression. *Anat Rec.* 2021;304(11):2470–2479. doi:10.1002/ar.24778
26. Dávila-Hernández A, González-González R, Guzmán-Velázquez S, Hernández Hernández OT, Zamudio SR, Martínez-Mota L. Antidepressant-like effects of acupuncture via modulation of corticosterone, sex hormones, and hippocampal BDNF expression in male rats. *Brain Res Bull.* 2021;173:53–65. doi:10.1016/j.brainresbull.2021.05.007
27. Li P, Huang W, Yan YN, et al. Acupuncture can play an antidepressant role by regulating the intestinal microbes and neurotransmitters in a rat model of depression. *Med Sci Monit.* 2021;27:e929027. doi:10.12659/MSM.929027
28. Jung J, Lee SM, Lee BH, et al. Lipidomics reveals that acupuncture modulates the lipid metabolism and inflammatory interaction in a mouse model of depression. *Brain Behav Immun.* 2021;94:424–436. doi:10.1016/j.bbi.2021.02.003
29. Yao Z, Zhang Z, Zhang J, et al. Electroacupuncture alleviated the depression-like behavior by regulating FGF2 and astrocytes in the hippocampus of rats with chronic unpredictable mild stress. *Brain Res Bull.* 2021;169:43–50. doi:10.1016/j.brainresbull.2021.01.005
30. Guo Q, Lin XM, Di Z, Zhang QA, Jiang S. Electroacupuncture ameliorates CUMS-induced depression-like behavior: involvement of the glutamatergic system and apoptosis in rats. *Comb Chem High Throughput Screen.* 2021;24(7):996–1004. doi:10.2174/1386207323666201027121423
31. Bai L, Zhang D, Cui TT, et al. Mechanisms underlying the antidepressant effect of acupuncture via the CaMK signaling pathway. *Front Behav Neurosci.* 2020;14:563698. doi:10.3389/fnbeh.2020.563698
32. Ma FQ, Sun CJ, Wei JJ, Wang YD, Shen JC, Chang JJ. Electro-acupuncture regulates glucose metabolism in chronic stress model rats. *Sci Rep.* 2020;10(1):11281. doi:10.1038/s41598-020-68132-w
33. Chen L, Yao Z, Qu S, et al. Electroacupuncture improves synaptic plasticity by regulating the 5-HT1A receptor in hippocampus of rats with chronic unpredictable mild stress. *J Int Med Res.* 2020;48(5):300060520918419. doi:10.1177/0300060520918419
34. Seo SY, Kang SY, Kwon OS, et al. A mechanical acupuncture instrument mitigates the endoplasmic reticulum stress and oxidative stress of ovariectomized rats. *Integr Med Res.* 2019;8(3):187–194. doi:10.1016/j.imr.2019.07.001
35. Lee MJ, Ryu JS, Won SK, et al. Effects of acupuncture on chronic stress-induced depression-like behavior and its central neural mechanism. *Front Psychol.* 2019;10:1353. doi:10.3389/fpsyg.2019.01353
36. Han X, Wu H, Yin P, et al. Electroacupuncture restores hippocampal synaptic plasticity via modulation of 5-HT receptors in a rat model of depression. *Brain Res Bull.* 2018;139:256–262. doi:10.1016/j.brainresbull.2018.03.004
37. Yue N, Li B, Yang L, et al. Electro-acupuncture alleviates chronic unpredictable stress-induced depressive- and anxiety-like behavior and hippocampal neuroinflammation in rat model of depression. *Front Mol Neurosci.* 2018;11:149. doi:10.3389/fnmol.2018.00149
38. Jiang H, Zhang X, Lu J, et al. Antidepressant-like effects of acupuncture-insights from DNA methylation and histone modifications of brain-derived neurotrophic factor. *Front Psychiatry.* 2018;9:102. doi:10.3389/fpsyg.2018.00102
39. Friedman JR, Nunnari J. Mitochondrial form and function. *Nature.* 2014;505(7483):335–343. doi:10.1038/nature12985
40. Picard M, Wallace DC, Burrelle Y. The rise of mitochondria in medicine. *Mitochondrion.* 2016;30:105–116. doi:10.1016/j.mito.2016.07.003
41. Tanaka M, Szabó Á, Spekker E, Polyák H, Tóth F, Vécsei L. mitochondrial impairment: a common motif in neuropsychiatric presentation? The link to the tryptophan-kynurenine metabolic system. *Cells.* 2022;11(16):2607. doi:10.3390/cells11162607
42. Tanaka M, Á S, Vécsei L. Preclinical modeling in depression and anxiety: current challenges and future research directions. *Adv Clin Exp Med.* 2023;32(5):505–509. doi:10.17219/acem/165944
43. Ma K, Chen G, Li W, Kepp O, Zhu Y, Chen Q. Mitophagy, mitochondrial homeostasis, and cell fate. *Front Cell Dev Biol.* 2020;8:467. doi:10.3389/fcell.2020.00467
44. Gustafsson ÅB, Dorn GW. Evolving and expanding the roles of mitophagy as a homeostatic and pathogenic process. *Physiol Rev.* 2019;99(1):853–892. doi:10.1152/physrev.00005.2018
45. Palikaras K, Lionaki E, Tavernarakis N. Mechanisms of mitophagy in cellular homeostasis, physiology and pathology. *Nat Cell Biol.* 2018;20(9):1013–1022. doi:10.1038/s41556-018-0176-2
46. Youle RJ, Narendra DP. Mechanisms of mitophagy. *Nat Rev Mol Cell Biol.* 2011;12(1):9–14. doi:10.1038/nrm3028
47. Zhang Z, Cai X, Yao Z, et al. EA ameliorated depressive behaviors in CUMS rats and was related to its suppressing autophagy in the hippocampus. *Neural Plast.* 2020;2020:8860968. doi:10.1155/2020/8860968
48. Reddy PH, Yin X, Manczak M, et al. Mutant APP and amyloid beta-induced defective autophagy, mitophagy, mitochondrial structural and functional changes and synaptic damage in hippocampal neurons from Alzheimer's disease. *Hum Mol Genet.* 2018;27(14):2502–2516. doi:10.1093/hmg/ddy154
49. Ge X, Wang L, Cui Q, et al. Electroacupuncture improves cognitive impairment in diabetic cognitive dysfunction rats by regulating the mitochondrial autophagy pathway. *J Physiol Sci.* 2022;72(1):29. doi:10.1186/s12576-022-00854-0
50. Wang ZT, Lu MH, Zhang Y, et al. Disrupted-in-schizophrenia-1 protects synaptic plasticity in a transgenic mouse model of Alzheimer's disease as a mitophagy receptor. *Aging Cell.* 2019;18(1):e12860. doi:10.1111/aclel.12860
51. Hsu WT, Chen YH, Yang HB, Lin JG, Hung SY. Electroacupuncture Improves motor symptoms of parkinson's disease and promotes neuronal autophagy activity in mouse brain. *Am J Chin Med.* 2020;48(7):1651–1669. doi:10.1142/S0192415X20500822
52. Zhong X, Chen B, Li Z, et al. Correction to: electroacupuncture ameliorates cognitive impairment through the inhibition of NLRP3 inflammasome activation by regulating melatonin-mediated mitophagy in stroke rats. *Neurochem Res.* 2022;47(7):1931–1933. doi:10.1007/s11064-022-03590-4
53. Wang H, Chen S, Zhang Y, Xu H, Sun H. Electroacupuncture ameliorates neuronal injury by Pink1/Parkin-mediated mitophagy clearance in cerebral ischemia-reperfusion. *Nitric Oxide.* 2019;91:23–34. doi:10.1016/j.niox.2019.07.004
54. Krivoruchko A, Storey KB. Forever young: mechanisms of natural anoxia tolerance and potential links to longevity. *Oxid Med Cell Longev.* 2010;3(3):186–198. doi:10.4161/oxim.3.3.12356
55. Sena LA, Chandel NS. Physiological roles of mitochondrial reactive oxygen species. *Mol Cell.* 2012;48(2):158–167. doi:10.1016/j.molcel.2012.09.025
56. Sies H. Oxidative stress: oxidants and antioxidants. *Exp Physiol.* 1997;82(2):291–295. doi:10.1113/expphysiol.1997.sp004024
57. Liochev SI. Reactive oxygen species and the free radical theory of aging. *Free Radic Biol Med.* 2013;60:1–4. doi:10.1016/j.freeradbiomed.2013.02.011

58. Lindqvist D, Dhabhar FS, James SJ, et al. Oxidative stress, inflammation and treatment response in major depression. *Psychoneuroendocrinology*. 2017;76:197–205. doi:10.1016/j.psyneuen.2016.11.031
59. Maes M, Galecki P, Chang YS, Berk M. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011;35(3):676–692. doi:10.1016/j.pnpbp.2010.05.004
60. Mellon SH, Wolkowitz OM, Schonemann MD, et al. Alterations in leukocyte transcriptional control pathway activity associated with major depressive disorder and antidepressant treatment. *Transl Psychiatry*. 2016;6(5):e821. doi:10.1038/tp.2016.79
61. Fang CC, Xu QQ, Shen J, Li Q, Shen MH. 针刺对雷公藤多苷片所致卵巢功能减退大鼠氧化应激及凋亡的影响 [Effect of acupuncture on oxidative stress and apoptosis in ovarian hypofunction induced by tripterygium glycosides]. *Zhen Ci Yan Jiu*. 2019;44(11):810–816. Chinese. doi:10.13702/j.1000-0607.180576
62. Bhatt S, Nagappa AN, Patil CR. Role of oxidative stress in depression. *Drug Discov Today*. 2020;25(7):1270–1276. doi:10.1016/j.drudis.2020.05.001
63. Wu YY, Liu MJ, Yin SJ, Wang AY, Li JG. 针刺对溃疡性结肠炎大鼠氧化应激和内质网应激的影响 [Acupuncture reduce colonic inflammation by suppressing oxidative stress and endoplasmic reticulum stress in rats with ulcerative colitis]. *Zhen Ci Yan Jiu*. 2020;45(1):8–13. Chinese. doi:10.13702/j.1000-0607.1806066
64. Skulachev VP. Mitochondrial physiology and pathology; concepts of programmed death of organelles, cells and organisms. *Mol Aspects Med*. 1999;20(3):139–184. doi:10.1016/s0098-2997(99)00008-4
65. Skulachev VP. Phenoptosis: programmed death of an organism. *Biochemistry*. 1999;64(12):1418–1426.
66. Dai W, Li WD, Lu J. 电针对慢性应激抑郁大鼠海马神经元凋亡及JNK信号转导通路的影响 [Effect of electroacupuncture on hippocampal apoptosis and JNK signal pathway in chronic stress depression rats]. *Zhen Ci Yan Jiu*. 2010;35(5):330–334. Chinese.
67. Liu P, Yu X, Dai X, et al. Scalp acupuncture attenuates brain damage after intracerebral hemorrhage through enhanced mitophagy and reduced apoptosis in rats. *Front Aging Neurosci*. 2021;13:718631. doi:10.3389/fnagi.2021.718631
68. Attardi G, Schatz G. Biogenesis of mitochondria. *Annu Rev Cell Biol*. 1988;4(1):289–333. doi:10.1146/annurev.cb.04.110188.001445
69. Fernandez-Marcos PJ, Auwerx J. Regulation of PGC-1 $\alpha$ , a nodal regulator of mitochondrial biogenesis. *Am J Clin Nutr*. 2011;93(4):884S–890S. doi:10.3945/ajcn.110.001917
70. Puigserver P, Wu Z, Park CW, Graves R, Wright M, Spiegelman BM. A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell*. 1998;92(6):829–839. doi:10.1016/s0092-8674(00)81410-5
71. Jäger S, Handschin C, St-Pierre J, Spiegelman BM. AMP-activated protein kinase (AMPK) action in skeletal muscle via direct phosphorylation of PGC-1 $\alpha$ . *Proc Natl Acad Sci U S A*. 2007;104(29):12017–12022. doi:10.1073/pnas.0705070104
72. Puigserver P, Rhee J, Lin J, et al. Cytokine stimulation of energy expenditure through p38 MAP kinase activation of PPAR $\gamma$  coactivator-1. *Mol Cell*. 2001;8(5):971–982. doi:10.1016/s1097-2765(01)00390-2
73. Wu Z, Puigserver P, Andersson U, et al. Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell*. 1999;98(1):115–124. doi:10.1016/S0092-8674(00)80611-X
74. Virbasius JV, Scarpulla RC. Activation of the human mitochondrial transcription factor A gene by nuclear respiratory factors: a potential regulatory link between nuclear and mitochondrial gene expression in organelle biogenesis. *Proc Natl Acad Sci U S A*. 1994;91(4):1309–1313. doi:10.1073/pnas.91.4.1309
75. Ekstrand MI, Falkenberg M, Rantanen A, et al. Mitochondrial transcription factor A regulates mtDNA copy number in mammals. *Hum Mol Genet*. 2004;13(9):935–944. doi:10.1093/hmg/ddh109
76. Gleyzer N, Vercauteren K, Scarpulla RC. Control of mitochondrial transcription specificity factors (TFB1M and TFB2M) by nuclear respiratory factors (NRF-1 and NRF-2) and PGC-1 family coactivators. *Mol Cell Biol*. 2005;25(4):1354–1366. doi:10.1128/MCB.25.4.1354-1366.2005
77. Falkenberg M, Gaspari M, Rantanen A, Trifunovic A, Larsson NG, Gustafsson CM. Mitochondrial transcription factors B1 and B2 activate transcription of human mtDNA. *Nat Genet*. 2002;31(3):289–294. doi:10.1038/ng909
78. Alcocer-Gómez E, Núñez-Vasco J, Casas-Barquero N, et al. Gene expression profile in major depressive disorder shows reduced mitochondrial biogenesis. *CNS Neurosci Ther*. 2016;22(7):636–638. doi:10.1111/cns.12568
79. Tang Q, Lu M, Xu B, et al. Electroacupuncture regulates inguinal white adipose tissue browning by promoting sirtuin-1-dependent PPAR $\gamma$  deacetylation and mitochondrial biogenesis. *Front Endocrinol*. 2020;11:607113. doi:10.3389/fendo.2020.607113
80. Sun S, Jiang T, Duan N, et al. Activation of CB1R-dependent PGC-1 $\alpha$  is involved in the improved mitochondrial biogenesis induced by electroacupuncture pretreatment. *Rejuvenation Res*. 2021;24(2):104–119. doi:10.1089/rej.2020.2315
81. Chan DC. Mitochondrial fusion and fission in mammals. *Annu Rev Cell Dev Biol*. 2006;22(1):79–99. doi:10.1146/annurev.cellbio.22.010305.104638
82. Chan DC. Mitochondria: dynamic organelles in disease, aging, and development. *Cell*. 2006;125(7):1241–1252. doi:10.1016/j.cell.2006.06.010
83. Shutt TE, McBride HM. Staying cool in difficult times: mitochondrial dynamics, quality control and the stress response. *Biochim Biophys Acta*. 2013;1833(2):417–424. doi:10.1016/j.bbamcr.2012.05.024
84. Hoppins S, Lackner L, Nunnari J. The machines that divide and fuse mitochondria. *Annu Rev Biochem*. 2007;76:751–780. doi:10.1146/annurev.biochem.76.071905.090048
85. Deng D, Cui Y, Gan S, et al. Sinisan alleviates depression-like behaviors by regulating mitochondrial function and synaptic plasticity in maternal separation rats. *Phytomedicine*. 2022;106:154395. doi:10.1016/j.phymed.2022.154395
86. Zhang Y, Meng Z, Wu L, et al. Protective effect of electroacupuncture on the barrier function of intestinal injury in endotoxemia through HO-1/PINK1 pathway-mediated mitochondrial dynamics regulation. *Oxid Med Cell Longev*. 2023;2023:1464853. doi:10.1155/2023/1464853
87. Yong RL, Dong JZ, Zhang L, Wu L. 电针“足三里”对脾气虚大鼠骨骼肌组织超微结构及线粒体动力学的影响 [Effects of electroacupuncture at “Zusanli” (ST36) on ultrastructure and mitochondrial dynamics of skeletal muscle in rats with spleen qi deficiency syndrome]. *Zhen Ci Yan Jiu*. 2020;45(1):15–20. Chinese. doi:10.13702/j.1000-0607.1903376
88. Sánchez-Rodríguez I, Temprano-Carazo S, Jeremic D, et al. Recognition memory induces natural LTP-like hippocampal synaptic excitation and inhibition. *Int J Mol Sci*. 2022;23(18):10806. doi:10.3390/ijms231810806
89. Li P, Huang W, Chen Y, et al. Acupuncture alleviates CUMS-induced depression-like behaviors by restoring prefrontal cortex neuroplasticity. *Neural Plast*. 2023;2023:1474841. doi:10.1155/2023/1474841
90. Liao HY, Satyanarayanan SK, Lin YW, Su KP. Clinical efficacy and immune effects of acupuncture in patients with comorbid chronic pain and major depression disorder: a double-blinded, randomized controlled crossover study. *Brain Behav Immun*. 2023;110:339–347. doi:10.1016/j.bbi.2023.03.016

91. Huang X, Zhuo Y, Wang X, et al. Structural and functional improvement of amygdala sub-regions in postpartum depression after acupuncture. *Front Hum Neurosci.* 2023;17:1163746. doi:10.3389/fnhum.2023.1163746
92. Li M, Niu J, Yan P, et al. The effectiveness and safety of acupuncture for depression: an overview of meta-analyses. *Complement Ther Med.* 2020;50:102202. doi:10.1016/j.ctim.2019.102202

### Neuropsychiatric Disease and Treatment

Dovepress

### Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>