



POSTER PRESENTATION

Open Access

The impact of age on right ventricular morphology and function late after repair of Tetralogy of Fallot: a cardiac magnetic resonance study

Benedetta Leonardi^{1*}, Marcello Chinali¹, Aurelio Secinaro¹, Emanuela Valsangiacomo Buechel², Valentina Silvestri¹, Sonia B Albanese¹, Adriano Carotti¹, Giacomo Pongiglione¹

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Cardiac magnetic resonance (CMR) is the gold standard to evaluate right ventricular (RV) hemodynamics after repair of Tetralogy of Fallot (TOF). CMR markers, such as severe RV dilation and biventricular dysfunction, are independent predictors of death, sustained VT and heart failure. However, the relationship between patient's age, type and time of corrective surgery, severity of PR and RV/LV dimensions and function has not been fully explained.

Methods

We enrolled patients who underwent hemodynamic evaluation by CMR late after TOF repair (by either transanular or infundibular RV outflow tract reconstruction) in two pediatric cardiology centers, between March 2008 and March 2013. Surgical data (time and types of repair) and CMR parameters were collected.

Results

The study included 208 patients (62% males) aged 17.5 ± 6.6 (range 5-38) years, with age at repair 16.3 ± 21.7 months (range 0.2-146 months). RV end-diastolic volume indexed (RVEDVi) was mildly negatively correlated with patient's age (figure; $p = 0.019$, $r = -0.163$). In details the negative correlation between RV dilation was stronger when limiting the analysis to patients older than 20 years (figure). On the contrary, LVEDVi was

mildly positively related to patient's age ($p = 0.020$, $r = 0.161$). RVEDVi was also correlated with time interval from corrective surgery, PRF and RVOT obstruction (all $p < 0.05$). Eventually, older age at repair and RVEDVi significantly affected RVEF (all $p < 0.05$). LVEF was significantly correlated with RVEF ($p < 0.001$) and the time elapsed from the repair ($p = 0.016$, $r = 0.171$).

Conclusions

RV dilation in repaired TOF patients might have an unexpected behavior over time, with older patients showing a progressive reduction in RV volume. Longitudinal follow-up studies are needed to verify our findings and define the role of a hypothetical RV "restrictive physiology" in predicting the adverse outcomes of patients with repaired TOF.

Funding

No disclosure to make.

Authors' details

¹Department of Cardiology and Cardiac Surgery, Bambino Gesu' Children's Hospital IRCCS, Rome, Rome, Italy. ²Department of Diagnostic Imaging, University Children's Hospital, Zurich, Switzerland.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P125

Cite this article as: Leonardi et al.: The impact of age on right ventricular morphology and function late after repair of Tetralogy of Fallot: a cardiac magnetic resonance study. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P125.

¹Department of Cardiology and Cardiac Surgery, Bambino Gesu' Children's Hospital IRCCS, Rome, Rome, Italy

Full list of author information is available at the end of the article