

Secondary Metabolites from Leaves of *Manilkara subsericea* (Mart.) Dubard

Fernanda Borges de Almeida¹, Caio Pinho Fernandes^{1,2}, Wanderson Romao^{3,4}, Gabriela Vanini³, Helber Barcelos Costa³, Hildegardo Seibert França⁴, Marcelo Guerra Santos⁵, José Carlos Tavares Carvalho⁶, Deborah Quintanilha Falcão⁷, Leandro Rocha^{2,8}


¹Laboratório de Nanobiotecnologia Fitofarmacêutica, Colegiado de Farmácia, Universidade Federal do Amapá, Campus Universitário Marco Zero do Equador, Rodovia Juscelino Kubitschek de Oliveira, KM, 02 Bairro Zerão, CEP: 68902-280, Macapá, AP, ²Programa de Pós, Graduação em Biotecnologia Vegetal, Centro de Ciências da Saúde, Bloco K, 2º Andar, Sala 032, Universidade Federal do Rio de Janeiro, UFRJ, Av. Brigadeiro Trompowski s/n, CEP: 21941-590, Ilha do Fundão, RJ, ³Laboratório de Petrolômica e Forense, Departamento de Química, Universidade Federal do Espírito Santo, CEP: 29075-910, Vitória, ES, ⁴Instituto Federal de Educação, Ciência e Tecnologia do Espírito Santo, CEP: 29106-010, Vila Velha, ES, ⁵Faculdade de Formação de Professores, UERJ, Rua: Dr. Francisco Portela, 1470, Patronato, CEP: 24435-005, São Gonçalo, Rio de Janeiro, ⁶Laboratório de Pesquisa em Fármacos, Colegiado de Farmácia, Universidade Federal do Amapá, Campus Universitário Marco Zero do Equador, Rod. Juscelino Kubitschek de Oliveira, KM-02, Bairro Zerão, CEP 68902-280 Macapá, AP, ⁷Laboratório de Tecnologia Farmacêutica, Departamento e Tecnologia Farmacêutica, Faculdade de Farmácia, Universidade Federal Fluminense, UFF Rua: Mario Viana, 523, CEP: 24241-000, Santa Rosa, Niterói, ⁸Laboratório de Tecnologia de Produtos Naturais, LTPN, Departamento e Tecnologia Farmacêutica, Faculdade de Farmácia, Universidade Federal Fluminense, UFF Rua: Mario Viana, 523, CEP: 24241-000, Santa Rosa, Niterói, RJ, Brazil

ABSTRACT

Background: *Manilkara subsericea* (Sapotaceae) is a species widely spread in the sandbanks of Restinga de Jurubatiba National Park (Rio de Janeiro, Brazil). It is commonly known as “maçaranduba,” “maçarandubinha” and “guracica,” being used in this locality as food, and timber. However, *M. subsericea* remains almost unexplored regarding its chemical constituents, including secondary metabolites from the leaves. **Objective:** Identify the chemical constituents from the leaves of *M. subsericea*. **Materials and Methods:** Leaves were macerated with ethanol (96% v/v), and dried crude ethanolic extract was sequentially washed with the organic solvents in order to obtain an ethyl acetate fraction. Substances from this fraction were identified by different techniques, such as negative-ion electrospray ionization Fourier and ¹H and ¹³C nuclear magnetic resonance (NMR). Fresh leaves from *M. subsericea* were also submitted to hydrodistillation in order to obtain volatile substances, which were identified by gas chromatograph coupled to mass spectrometer. **Results:** NMR¹H and ¹³C spectra allowed for the identification of the compounds *myricetin*, *quercetin*, and *kaempferol* from the ethyl acetate fraction. The negative-ion electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry mass spectrum also revealed the presence in this fraction of a polyhydroxytriterpene acid (pomolic acid), and some flavonoids, such as *quercitrin*, and *myricitrin*. In all 34 volatile compounds were identified by gas chromatography-mass spectrometry, including monoterpenes, sesquiterpenes, and long chain hydrocarbons. **Conclusion:** This study describes the first reports concerning the phytochemical information about leaves from *M. subsericea*. **Key words:** Essential oil, flavonoids, *Manilkara subsericea*, mass spectrometry, pomolic acid, sapotaceae

SUMMARY

- Manilkara subsericea fruits proved to be a rich source of triterpenes. However, no phytochemical studies were carried out with leaves. Thus, we described identification of volatile substances from its essential oils, in addition to non-reported triterpene and flavonoids from this species.



Correspondence:

Dr. Caio Pinho Fernandes, Universidade Federal do Amapá, Campus Universitário Marco Zero do Equador, Rodovia Juscelino Kubitschek de Oliveira, KM, 02 Bairro Zerão, CEP: 68902-280, Macapá, AP, Brazil.

E-mail: caiofernandes@unifap.br

DOI : 10.4103/0973-1296.172957

Access this article online

Website: www.phcog.com

Quick Response Code:



INTRODUCTION

Sapotaceae family contains 58 genus, and approximately 1250 species with morphological variation, ranging from shrubs to medium, and giant trees.^[1] Brazil comprises of 11 genera, and 231 species, including 1 endemic genus, and 104 endemic species.^[2] This family has the following synapomorphies, well-developed, elongate laticifers with white latex; 2-branched hairs, brownish, T-shaped; berry fruits, seeds usually with a hard shiny testa, and large hilum.^[3]

The genus *Manilkara* Adans. is constituted by 30 species in the neotropics, been approximately 20 species found in Africa, and 12 species found in Asia, and Pacific.^[4] Brazil has 18 species, being 15 endemic to this country.^[5] The genus *Manilkara* is characterized by calyx of 2 whorls of 3 sepals, presence of staminodes and hilum shape seed.^[5-7] Due to this

genus circumscription, some species of the genera *Achras*, and *Mimusops* were included in *Manilkara*.^[6]

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: de Almeida FB, Fernandes CP, Romao W, Vanini G, Costa HB, França HS, et al. Secondary metabolites from leaves of *Manilkara subsericea* (Mart.) Dubard. Phcog Mag 2015;11:533-7.

Manilkara subsericea (Mart.) Dubard is an endemic species from Brazilian Atlantic Rain Forest,^[5] commonly known as “maçaranduba”, “maçarandubinha”, and “guracica”. It is widely distributed at Restinga de Jurubatiba National Park (Rio de Janeiro State, Brazil), being used in this locality as food, and timber.^[8] This species also develops a main role in the ecology of Restinga de Jurubatiba, being an important host plant for some *Lepidoptera* species,^[9] and for porcupine (*Chaetomys subspinosus*) feeding.^[10] Despite some biological and chemical investigations were carried out with *M. subsericea*, to our knowledge, there is no information concerning secondary metabolites from leaves of this species. Thus, the aim of the present study was to perform phytochemical characterization of leaves from *M. subsericea* using spectroscopy methods such as nuclear magnetic resonance (NMR) ¹H, and ¹³C, negative-ion electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry (ESI(-) FT-ICR-MS) MS, and gas chromatography (GC/MS) mass spectrum.

MATERIALS AND METHODS

Plant material

Leaves of *M. subsericea* were collected at Restinga de Jurubatiba National Park, Rio de Janeiro State, Brazil (22°14'46"S-41°34'56"W), October 2010 by Dr. Caio Pinho Fernandes. Identification was performed by the botanist Dr. Marcelo Guerra Santos, and voucher specimen of *M. subsericea* was deposited at the herbarium of the Faculdade de Formação de Professores (Universidade do Estado do Rio de Janeiro, Brazil) under the register number RFFP 15316. The nomenclatural update was realized in Lista de Espécies da Flora do Brasil (<http://floradobrasil.jbrj.gov.br/jabot/listaBrasil/PrincipalUC/PrincipalUC.do>), and the plant list: A working list of all plant species (<http://www.theplantlist.org/>).

Preparation of extracts

Leaves (0.84 kg) were dried at 40°C for 2 days, and then extensively crushed, and macerated (15 L) in ethanol (EtOH) 96% (v/v) at room temperature. After filtration, the ethanolic extract was concentrated under vacuum using a rotary evaporator equipped with the water bath (35°C) (Fisatom, SP) in order to obtain 109.5 g of the ethanolic crude extract from leaves. This dried extract was sequentially washed with n-hexane (5 × 2000 mL), and dichloromethane (5 × 2000) in order to remove the less polar constituents from the extract. The insoluble fraction was washed with ethyl acetate (5 × 2000 mL). The ethyl acetate-soluble fraction was filtered, and concentrated under the vacuum using the rotary evaporator, affording 17.3 g of the ethyl acetate fraction from leaves (EAL).

Isolation of substances

The ethyl acetate-soluble fraction from leaves was fractionated through column chromatography using the Amberlite XAD-2 resin (Sigma-Aldrich, St. Louis). Elution was performed with water, methanol/water mixtures (5:95 → 9:1), methanol, and acetone. Fractions 28–40 were pooled together according to the thin layer chromatography (TLC) profile, and purified on Sephadex LH-20 using methanol as mobile phase, affording 1 (40.2 mg), 2 (35.7 mg), and 3 (11.9 mg). Fractions 8–19 were pooled together according to the TLC profile, and chromatographed in C-18 reversed phase silica gel (Sigma-Aldrich, St. Louis) using gradient of mobile phase constituted by methanol solutions (v/v) in water (60%→63%). Final purification on Sephadex LH-20 using methanol as mobile phase afforded a fraction (7.4 mg) containing 4 and 5. Fractions 41–52 was submitted to silica gel chromatography column using an isocratic system of mobile phase (n-hexane: ethyl acetate: methanol, 5:5:1), affording a fraction (9.7 mg) constituted by 6.

Essential oil extraction

Fresh leaves (340 g) from *M. subsericea* were ground with distilled water using an automatic blender (Ética Equipamentos Científicos SA, Brazil). Hydrodistillation method was employed for 3 h using Clevenger apparatus, and plant material was placed in a 5 L flask. At the end of the extraction, the essential oil was collected with n-hexane, dried over a hydrous sodium sulfate, and stored at 4°C for further analyzes.

Chemical analysis

¹H and ¹³C NMR spectra of 1, 2, 3, and 6 were recorded at 500, and 125 MHz, respectively, on a Varian VNMRs 500 MHz spectrometer. Deuterated methanol was used for solubilization of flavonoids while deuterated DMSO was used for solubilization of triterpene fraction. The solvents were obtained from Cambridge Isotope Laboratories (USA), and TMS peak was used as an internal standard.

Substances 4, 5, and 6 were analyzed by negative-ion electrospray ionization fourier transform ion cyclotron resonance MS, ESI(-)-FT-ICR MS.^[11-13] Briefly, each sample was diluted to ≅ 1.0 mg/mL in acetonitrile (containing 0.1% w/v of NH₄OH). The resulting solution was directly infused at a flow rate of 5 µL/min into the ESI source. The mass spectra were acquired over a mass range of m/z 200–2000. The ESI source conditions were as follows: nebulizer gas pressure of 0.5–1.0 bar, the capillary voltage of 2.5–3.5 kV, and transfer capillary temperature of 250°C. All mass spectra were externally calibrated using a NaTFA (m/z from 200 to 1200). A resolving power ($m/\Delta m_{50\%}$) ≈ 530,000, in which $\Delta m_{50\%}$ is the full peak width at half-maximum peak height) of m/z 400 and a mass accuracy (mass error) of <1 ppm provided unambiguous molecular formula assignments for singly charged molecular ions. Mass spectra were acquired and processed using the software package, Compass Data Analysis (Bruker Daltonics, Bremen, Germany). ESI(-)-MS/MS experiments were collected after 4–40 eV collision-induced dissociations (CID) with argon. Selection of ion was performed by quadrupole, using a unitary m/z window, and collisions were performed in the Rf-only hexapole collision cell, followed by mass analysis of product ions by the ultra-high resolution ICR analyzer.

The essential oil was analyzed by a GC/MS-QP2010 (SHIMADZU) gas chromatograph equipped with a mass spectrometer using electron ionization. The GC conditions were as follows: Essential oil: Injector temperature, 260°C; detector temperature, 290°C; carrier gas (Helium), flow rate 1 mL/min, and split injection with split ratio 1:40. Oven temperature was initially 60°C and then raised to 290°C at a rate of 3°C/min. The sample was diluted with n-hexane (1:100, v/v) and injected at RTX-5 column (i.d. = 0.25 mm, length 30 m, film thickness = 0.25 µm). The MS conditions were voltage, 70 eV, and scan rate; 1 scan/s. The retention indices were calculated by the interpolation of retention times of the substances to the retention times of a mixture of aliphatic hydrocarbons (C7-C40) (Sigma) analyzed in the same conditions.^[14] The identification of substances was performed by comparison of their retention indices and mass spectra with those reported in the literature.^[15] MS fragmentation pattern of compounds was also checked with NIST mass spectra libraries. Quantitative analysis of the chemical constituents was performed by GC-flame ionization detector (GC/FID), under same conditions of GC/MS analysis and percentages obtained by FID peak area normalization method.

RESULTS AND DISCUSSION

Chemical structures from the substances identified after fractionation of ethyl acetate-soluble fraction from leaves of *M. subsericea* are presented in Figure 1. ¹H NMR spectrum of 1 (CD₃OD, 500 MHz) showed the two

doublets at δ_{H} 6.18 ($J = 1.45$ Hz) and δ_{H} 6.38 ($J = 1.45$ Hz) attributable, respectively, to the protons H-8 and H-6, and a two-proton singlet observed at δ_{H} 7.34, corresponding to the B-ring aromatic protons H-2' and H-6' of a flavonol. The signals observed in the ^1H and ^{13}C NMR spectrum are in accordance with the literature data for *myricetin*.^[16] The substances 2 and 3 had their ^1H and ^{13}C NMR spectra compared with the literature data^[17,18] and, therefore, were respectively, identified as *quercetin*, and *kaempferol*.

The remaining flavonoids, 4 and 5, were analyzed by ESI(-)-FT-ICR MS [Figure 2a-d]. For FL4 [Figure 2a], ESI(-)-FT-ICR mass spectrum, detected the presence of the *O*-glycosideo-flavonol-*myricitrin* ($M = \text{C}_{21}\text{H}_{20}\text{O}_{12}$), ions of m/z




Figure 1: Chemical structures of flavonoids and triterpene from leaves of *Manilkara subsericea* (1: Myricetin R1 = R2 = R3 = R4 = OH, 2: Quercetin R1 = R2 = R4 = OH, R3 = H, 3: Kaempferol R1 = R3 = H, R2 = R4 = OH, 4: Myricitrin R1 = R2 = R3 = OH, R4 = Orha, 5: Quercitrin R1 = R2 = OH, R3 = H, R4 = Orha, 6: Pomolic acid)

463.0882 ($[\text{M} - \text{H}]^-$), 499.0651 ($[\text{M} + \text{Cl}]^-$) and 927.1844 ($[\text{2M} - \text{H}]^-$) as deprotonated molecule, chlorine adduct and dimer, respectively. A double bound equivalents (DBE) of 12 found for ions $[\text{M} - \text{H}]^-$ and $[\text{M} + \text{Cl}]^-$ agrees with the chemical structure of *myricitrin*, that presents two aromatic rings (DBE = 8), one rhamnose (DBE = 1), and a ring containing one double bond, and a ketone group (DBE = 3). To confirm the *myricitrin* detection, ESI(-)-MS/MS was performed for the ion of m/z 463 [Figure 2c]. The CID of $[\text{M} - \text{H}]^-$ of m/z 463 agrees well with its structure and connectivity, producing fragments with m/z 316, corresponding to the neutral loss of $\text{C}_6\text{H}_{11}\text{O}_4$. For FL5 [Figure 2b], ESI(-)-FT-ICR mass spectrum identified, simultaneously, ion $[\text{M} - \text{H}]^-$ of m/z 463.0882, and ions $[\text{N} - \text{H}]^-$ and $[\text{N} + \text{Cl}]^-$ of m/z 447.0933 and 483.0703, respectively, where $\text{N} = \text{C}_{21}\text{H}_{20}\text{O}_{11}$, corresponding to the *O*-glycoside flavonol *quercitrin*. CID experiment of the ion of m/z 447 [Figure 2d], produces fragments of m/z 301, and 284 corresponding to the neutral losses of $\text{C}_6\text{H}_{10}\text{O}_4$ and one rhamnose molecule.

Some flavonoids are isolated and identified on the present study, have also been found in this genus. *Quercetin* was identified on *Mimusops manilkara* G. Don (*Manilkara kauki* (L.) Dubard), and *Mimusops littoralis* Kurz (*M. littoralis* (Kurz) Dubard),^[19,20] while *myricetin* was identified on *Achras zapota* L. (*Manilkara zapota* (L.) P. Royen).^[21] The glycosylated flavonoids *quercitrin* and *myricitrin* were isolated from *M. zapota*.^[22]

Chromatographic fractionation of the EAL also afforded a white powder. ESI(-) FT-ICR mass spectrum of the fraction revealed the presence of precursor ions $[\text{C}_{30}\text{H}_{48}\text{O}_4 - \text{H}]^-$, $[\text{C}_{30}\text{H}_{48}\text{O}_5 - \text{H}]^-$ and $[\text{C}_{30}\text{H}_{48}\text{O}_6 - \text{H}]^-$ of m/z 471, 487 and 503, being attributed to dihydroxy, trihydroxy, and tetrahydroxytriterpene acids [Figure 3], which is in accordance with the derivatives from ursolic, and oleanolic acids. ^{13}C NMR spectra of




Figure 2: Negative-ion electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry mass spectra for (a) FL4 and (b) FL5 samples and collision-induced dissociations experiments for ion of m/z 463 and 447 corresponding to (c) myricitrin and (d) quercitrin

Table 1: Relative abundance of essential oil constituents from leaves of *Manilkara subsericea*

Substance	RI	Relative abundance (%)	Substance	RI	Relative abundance (%)
3-hexen-1-ol	850	0.2	Trans-2-decenal	1262	0.2
Hexanol	862	0.2	Beta-damascenone	1388	0.2
2-heptanone	889	0.1	Beta-damascene	1418	0.3
Heptanal	902	0.2	Beta-caryophyllene	1423	0.5
Heptanol	966	0.3	Farnesene	1510	0.3
Octanal	1003	0.2	Caryophyllene oxide	1588	0.7
(3E)-3-hexenyl acetate	1006	0.6	Hexadecanoic acid	1969	8.6
Beta-ocimene	1047	7.3	Eicosene	1994	0.2
Octanol	1069	1.3	Heneicosane	2100	0.3
Linalool oxide	1089	0.5	Phytol	2116	15.6
2-nonanone	1092	0.2	Docosene	2194	0.4
Linalool	1100	27.6	Tricosane	2300	0.6
Terpineol	1193	0.9	Pentacosane	2500	0.4
Methyl salicylate	1197	1.9	Heptacosane	2700	0.9
Safranal	1202	0.3	Squalene	2832	2.6
Beta-cyclocitral	1223	0.3	Nonacosane	2900	8.9
Geraniol	1256	0.3	Untriacontane	3100	3.4
Total identified	86.5				

RI: Retention indices calculated by interpolation of retention times of the substances to the retention times of a mixture of aliphatic hydrocarbons. GC conditions: Injector temperature, 260°C; detector temperature, 290°C; carrier gas, Helium; flow rate, 1 mL/min; split ratio, 1:40. Initial temperature, 60°C; final temperature, 290°C; rate of 3°C/min. RTX-5 column (i.d.=0.25 mm, length 30 m, film thickness=0.25 µm). MS conditions: Voltage, 70 eV; scan rate; 1 scan/s. GC: Gas chromatographic; MS: Mass spectrometry




Figure 3: Negative-ion electrospray ionization Fourier transforms ion cyclotron resonance mass spectrometry of triterpene acids present in the leaves *Manilkara subsericea*

this fraction showed the typical set of signals for triterpene mixtures. It was observed that a typical signal at δ_c 181.0, due to the C-28 carboxyl group of triterpene acid. The presence of C-12/C-13 olefinic carbons at δ_c 128.4/140.3, as well as the signals of carbons bonded to hydroxyl groups at δ_c 79.8 (C-3), and 73.1 (C-19) suggested that the pomolic acid (6) may be the main constituent of this fraction. Assignments for pomolic acid are in accordance with the literature data and allowed establishment of relative configuration.^[23] This triterpene is structurally related to ursolic acid and was already found in the family *Sapotaceae*.^[24] Considering the predominance of pentacyclic triterpenes in *M. subsericea*^[25,26] and wide distribution of these substances through genus *Manilkara*,^[19,27-30] identification of pomolic acid contributes to the concept of chemotaxonomic significance of pentacyclitriterpenes in the genus *Manilkara*.

Essential oil obtained from the fresh leaves of *M. subsericea* was analyzed by GC-MS in order to determine its chemical composition. In all, 34

substances were identified, mainly comprising of monoterpenes, sesquiterpenes, and long chain hydrocarbons, corresponding to 86.5% of the total relative composition of the oil [Table 1]. The monoterpene linalool was the major substance found, corresponding to 27.6% of the total relative composition of the essential oil. The relative amount of each substance found is presented in Table 1. It was also observed a substance with molecular ion peak (M^+) at m/z 296, and characteristic fragments at m/z 278, due to the loss of 18 units of mass ($M^+ - H_2O$), m/z 123 and 71 (base peak), being in accordance with the literature data for phytol.^[31] Squalene was detected, presenting main fragments at m/z 341, 95, 81, and 69 (base peak), which is in accordance with the literature data.^[32]

CONCLUSION

Despite the ecological significance, great abundance, used as food, and literature data concerning its biological activities, *M. subsericea* remains almost chemically unexplored. To our knowledge, only secondary metabolites from fruits of this species were previously reported. As part of our ongoing studies, the present study allowed the identification of substances from the leaves of this species, being all substances identified for the first time on *M. subsericea*.

Acknowledgment

The authors would like to thank CNPQ for the financial support.

Financial support and sponsorship

The authors would like to thank CNPQ (407768/2013-0) and FAPEAP (PRODETEC Araguari – process no. 250.203.035/2013) by the financial support.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Bartish IV, Antonelli A, Richardson JE, Swenson U. Vicariance or long-distance dispersal: Historical biogeography of the pantropical subfamily Chrysophylloideae (*Sapotaceae*). *J Biogeogr* 2011;38:177-90.

2. Carneiro CE, Alves-Araujo A, Almeida EB Jr, Terra-Araujo MH. *Sapotaceae*. In: Lista de Espécies da Flora do Brasil. , Brazil: Jardim Botânico do Rio de Janeiro; 2008. Available from: <http://www.floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB217>. [Last accessed on 2015 Nov 05].
3. Judd WS, Campell CS, Kellogg EA, Stevens PF, Donoghue MJ. *Sistemática Vegetal: Um Enfoque Filogenético*. 3rd ed. PortoAlegre, Brazil: Artmed Editora; 2009.
4. Pennington TD. *Sapotaceae*. Flora da Reserva Ducke, Amazonas, Brasil: Sapotaceae. Rodriguésia; 2006. p. 251-366.
5. Almeida EB Jr. *Manilkara*. In: Lista de Espécies da Flora do Brasil. Brazil: Jardim Botânico do Rio de Janeiro; 2008. Available from: <http://www.floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB14473>. [Last accessed on 2015 Nov 05].
6. Almeida EB Jr. *Diversidade de Manilkara Adans. (Sapotaceae) Para o Nordeste do Brasil*. PhD Thesis, Universidade Federal Rural de Pernambuco. Pernambuco, Brasil; 2010.
7. Pennington TD. *Flora Neotropica, Monographyes no 52 (Sapotaceae)*. New York, Botanical Gardens: Published for Organization for Flora Neotropica; 1990.
8. Santos MG, Fevereiro PC, Reis GL, Barcelos JI. Recursos vegetais da Restinga de Carapebus. *Rev Biol Neotrop* 2009;6:35-54.
9. Monteiro RF, Macedo MV, Nascimento MS, Cury RS. Composição, abundância e not as so brea ecologia de espécies de larvas de lepidópteros associadas a cinco espécies de plant ashospedeiras no Parque Nacional da Restinga de Jurubatiba, RJ. *Rev Bras Entomol* 2007;15:476-83.
10. Oliveira PA, Lima RB, Chiarello AG. Home range, movements and diurnal roosts of the endangered thin-spined porcupine, *Chaetomys subspinosus (Rodentia: Erethizontidae)*, in the Brazilian Atlantic Forest. *Mamm Biol* 2012;77:97-107.
11. Colati KA, Dalmaschio GP, Castro EV, Gomes AO, Vaz BG, Romão W. Monitoring the liquid/liquid extraction of naphthenic acids in Brazilian crude oil using electrospray ionization FT-ICR mass spectrometry (ESI FT-ICR MS). *Fuel* 2013;108:647-55.
12. Dalmaschio GP, Malacarne MM, Almeida VM, Pereira TM, Gomes AO, Castro EV, *et al.* Characterization of polar compounds in atree boiling point distillation system using electrospray Ionization FT-ICR mass spectrometry. *Fuel* 2014;115:190-202.
13. Freitas S, Malacarne MM, Romão W, Dalmaschio GP, Castro EV, Celante VG, *et al.* Analysis of the heavy oil distillation cuts corrosion by electrospray ionization FTICR mass spectrometry, electrochemical impedance spectroscopy, and scanning electron microscopy. *Fuel* 2013;104:656-63.
14. Van den Dool H, Kratz PD. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *J Chromatogr* 1963;11:463-71.
15. Adams RP. *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*. 4th ed. Carol Stream, Illinois: Allured Publishing; 2007.
16. Ceruks M, Romoff P, Fávero OA, Lago JH. Constituintes fenólicos polares de *Schinus terebinthifolius* Raddi (*Anacardiaceae*). *Quim Nova* 2007;30:597-9.
17. Koolen HH, Soares ER, Silva FM, Souza AQ, Filho ER, Souza AD. Triterpene and flavonoids from the roots of *Mauritia flexuosa*. *Braz J Pharmacogn* 2012;22:189-92.
18. Olennikov DN, Tankhaeva LM, Partilkhav VV, Rokhin AV. Chemical constituents of *Caragana bungei* shoots. *Braz J Pharmacogn* 2012;22:490-6.
19. Jahan N, Ahmad W, Malik A. Phytochemistry of the genus *Mimusops*. *J Chem Soc Pak* 1996;18:51-5.
20. Misra G, Mitra CR. *Mimusops manilkara*, constituents of fruit and seed. *Phytochemistry* 1969;8:249-52.
21. Subramanian SS, Nair AG. Myricetin and myricetin-3-O-L-rhamnoside from the leaves of *Madhucaindica* and *Achras zapota*. *Phytochem Rep* 1972;11:3090-1.
22. Ma J, Luo XD, Protiva P, Yang H, Ma C, Basile MJ, *et al.* Bioactive novel polyphenols from the fruit of *Manilkara zapota* (Sapodilla). *J Nat Prod* 2003;66:983-6.
23. Mahato SB, Kundu AP. ¹³C spectra of pentacyclitriterpenoids – A compilation and some salient features. *Phytochemistry* 1994;37:1517-75.
24. Lee TZ, Juang SH, Hsu FL, Wu CY. Triterpene acids from the leaves of *Planchonella duclitan* (Blanco) Bakhuzan. *J Chin Chem Soc* 2005;52:1275-80.
25. Fernandes CP, Corrêa AL, Cruz RA, Botas GS, Silva-Filho MV, Santos MG, *et al.* Anticholinesterasic activity of *Manilkara subsericea* (Mart.) Dubard triterpenes. *Lat Am J Pharm* 2011;30:1631-4.
26. Fernandes CP, Corrêa AL, Lobo JF, Caramel OP, de Almeida FB, Castro ES, *et al.* Triterpene esters and biological activities from edible fruits of *Manilkara subsericea* (Mart.) Dubard, *Sapotaceae*. *Biomed Res Int* 2013;2013:280810.
27. Ahmed R, Ifzal SM, Usmanghani K. Studies on *Achras sapota* L: part IV – Chemistry and pharmacology of wood. *Pak J Pharm Sci* 2001;14:39-40.
28. Fayek NM, Monem AR, Mossa MY, Meselhy MR. New triterpenoid acyl derivatives and biological study of *Manilkara zapota* (L.) Van Royen fruits. *Pharmacognosy Res* 2013;5:55-9.
29. Misra G, Nigam SK, Mitra CR. Constituents of *Mimusops manilkara* leaves and saponins of *Mimusops* seed kernels. *Phytochemistry* 1969;8:2255-6.
30. Rhourri-Frih B, Renimel I, Chaimbault P, André P, Herbet G, Lafosse M. Pentacyclic triterpenes from *Manilkara bidentata* resin. Isolation, identification and biological properties. *Fitoterapia* 2013;88:101-8.
31. Siqueira DS, Pereira AS, Neto FR, Cabral JA, Ferreira CA, Simoneit BR, *et al.* Determination of high molecular mass compounds from Amazonian plant's leaves. *Quim Nova* 2003;26:633-40.
32. Bhatia A, Bharti SK, Tewari SK, Sidhu OP, Roy R. Metabolic profiling for studying chemotype variations in *Withania somnifera* (L.) Dunal fruits using GC-MS and NMR spectroscopy. *Phytochemistry* 2013;93:105-15.



Leandro Rocha



Caio Pinho Fernandes

ABOUT AUTHORS

Leandro Rocha earned his doctorate in Pharmacy at Université de Lausanne and is permanente lecturer and researcher at Fluminense Federal University (Brazil). He has great experience in phytochemistry and published several original papers in scientific journals regarding natural products.

Caio Pinho Fernandes earned his doctorate in Plant Biotechnology at Rio de Janeiro Federal University (Brazil) and is permanente lecturer and researcher at Amapá Federal University (Brazil). He has great experience in phytopharmaceutical nanobiotechnology and published several original papers in scientific journals regarding natural products