

## EDITORIAL



## Hydroxychloroquine for the Prevention of Covid-19 — Searching for Evidence

Myron S. Cohen, M.D.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus that causes coronavirus disease 2019 (Covid-19), has generated a worldwide pandemic. The interruption of its spread depends on a combination of pharmacologic and non-pharmacologic interventions. Initial SARS-CoV-2 prevention includes social distancing, the use of face masks, environmental hygiene, and hand washing.<sup>1</sup> Although the most important pharmacologic interventions to prevent SARS-CoV-2 infection are likely to be vaccines, the repurposing of established drugs for short-term prophylaxis is another, more immediate option.

Some researchers have promoted chloroquine and hydroxychloroquine for the treatment and prevention of illness from a variety of microorganisms, including SARS-CoV.<sup>2</sup> Hydroxychloroquine can inhibit replication of SARS-CoV-2 in vitro.<sup>3</sup> Some observational studies have suggested benefits of hydroxychloroquine for the treatment of Covid-19, whereas other treatment reports have described mixed results.<sup>4</sup>

Boulware et al. now report in the *Journal* the results of a randomized trial testing hydroxychloroquine as postexposure prophylaxis for Covid-19.<sup>5</sup> This is described by the investigators as a “pragmatic” trial in which participants were recruited through social media and almost all data were reported by the participants. Adults who described a high-risk or moderate-risk exposure to someone with Covid-19 in their household or an occupational setting were provided hydroxychloroquine or placebo (by mail) within 4 days after the reported exposure, and before symptoms would be expected to develop. The authors enrolled 821 participants; an illness that was considered to be

consistent with Covid-19 developed in 107 participants (13.0%) but was confirmed by polymerase-chain-reaction assay in less than 3% of the participants. The incidence of a new illness compatible with Covid-19 did not differ significantly between participants receiving hydroxychloroquine (49 of 414 [11.8%]) and those receiving placebo (58 of 407 [14.3%]). Although participant-reported side effects were significantly more common in those receiving hydroxychloroquine (40.1%) than in those receiving placebo (16.8%), no serious adverse reactions were reported.

This trial has many limitations, acknowledged by the investigators. The trial methods did not allow consistent proof of exposure to SARS-CoV-2 or consistent laboratory confirmation that the symptom complex that was reported represented a SARS-CoV-2 infection. Indeed, the specificity of participant-reported Covid-19 symptoms is low,<sup>6</sup> so it is hard to be certain how many participants in the trial actually had Covid-19. Adherence to the interventions could not be monitored, and participants reported less-than-perfect adherence, more notably in the group receiving hydroxychloroquine. In addition, those enrolled in the trial were younger (median age, 40 years) and had fewer coexisting conditions than persons in whom severe Covid-19 is most likely to develop,<sup>7</sup> so enrollment of higher-risk participants might have yielded a different result.

The trial design raises questions about the expected prevention benefits of hydroxychloroquine. Studies of postexposure prophylaxis are intended to provide an intervention in the shortest possible time to prevent infection. In a small-animal model of SARS-CoV-2 infection,<sup>8</sup> prevention of infection

or more severe disease was observed only when the experimental antiviral agent was given before or shortly after exposure. In the current trial, the long delay between perceived exposure to SARS-CoV-2 and the initiation of hydroxychloroquine ( $\geq 3$  days in most participants) suggests that what was being assessed was prevention of symptoms or progression of Covid-19, rather than prevention of SARS-CoV-2 infection.

Drugs for the prevention of infections must have an excellent safety profile. When hydroxychloroquine was initially promoted as a possible solution to SARS-CoV-2 infection, the safety of the drug was emphasized.<sup>2</sup> Under closer scrutiny, however, the potential for cardiac toxic effects and overall adverse outcomes have been emphasized, especially in persons with underlying coexisting conditions that increase the risk of severe Covid-19.<sup>9</sup> Boulware et al. report frequent mild side effects of hydroxychloroquine, but cardiac toxic effects could not be assessed.

So, what are we to do with the results of this trial? The advocacy and widespread use of hydroxychloroquine seem to reflect a reasonable fear of SARS-CoV-2 infection. However, it would appear that to some extent the media and social forces — rather than medical evidence — are driving clinical decisions and the global Covid-19 research agenda.<sup>10</sup> On June 1, 2020, ClinicalTrials.gov listed a remarkable 203 Covid-19 trials with hydroxychloroquine, 60 of which were focused on prophylaxis. An important question is to what extent the article by Boulware et al. should affect planned or ongoing hydroxychloroquine trials. If postexposure prophylaxis with hydroxychloroquine does not prevent symptomatic SARS-CoV-2 infection (with recognition of the limitations of the trial under discussion), should other trials of postexposure prophylaxis with hydroxychloroquine continue unchanged? Do the participants in these trials need to be informed of these results? Do these trial results with respect to postexposure prophylaxis affect trials of preexposure prophylaxis with hydroxychloroquine, some of which are very large (e.g., the Healthcare Worker

Exposure Response and Outcomes of Hydroxychloroquine [HERO-HCQ] trial, involving 15,000 health care workers; ClinicalTrials.gov number, NCT04334148)? The results reported by Boulware et al. are more provocative than definitive, suggesting that the potential prevention benefits of hydroxychloroquine remain to be determined.

Disclosure forms provided by the author are available with the full text of this editorial at NEJM.org.

From the Institute for Global Health and Infectious Diseases, University of North Carolina at Chapel Hill, Chapel Hill.

This editorial was published on June 3, 2020, at NEJM.org.

1. Cohen MS, Corey L. Combination prevention for COVID-19. *Science* 2020;368:551.
2. Rolain J-M, Colson P, Raoult D. Recycling of chloroquine and its hydroxyl analogue to face bacterial, fungal and viral infections in the 21st century. *Int J Antimicrob Agents* 2007;30:297-308.
3. Yao T-T, Qian J-D, Zhu W-Y, Wang Y, Wang G-Q. A systematic review of lopinavir therapy for SARS coronavirus and MERS coronavirus — a possible reference for coronavirus disease-19 treatment option. *J Med Virol* 2020;92:556-63.
4. Hernandez AV, Roman YM, Pasupuleti V, Barboza JJ, White CM. Hydroxychloroquine or chloroquine for treatment or prophylaxis of COVID-19: a living systematic review. *Ann Intern Med* 2020 May 27 (Epub ahead of print).
5. Boulware DR, Pullen MF, Bangdiwala AS, et al. A randomized trial of hydroxychloroquine as postexposure prophylaxis for Covid-19. *N Engl J Med*. DOI: 10.1056/NEJMoa2016638.
6. Tostmann A, Bradley J, Bousema T, et al. Strong associations and moderate predictive value of early symptoms for SARS-CoV-2 test positivity among healthcare workers, the Netherlands, March 2020. *Euro Surveill* 2020;25(16):2000508.
7. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020 February 24 (Epub ahead of print).
8. Sheahan TP, Sims AC, Zhou S, et al. An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* 2020;12:12(541):eabb5883.
9. Magagnoli J, Narendran S, Pereira F, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. April 16, 2020 (<https://www.medrxiv.org/content/10.1101/2020.04.16.20065920v2>). preprint.
10. Sayare S. He was a science star. Then he promoted a questionable cure for COVID-19. *New York Times Magazine*. May 12, 2020 (<https://www.nytimes.com/2020/05/12/magazine/didier-raoult-hydroxychloroquine.html>).

DOI: 10.1056/NEJMe2020388

Copyright © 2020 Massachusetts Medical Society.