

Pain after laparoscopic surgery: Focus on shoulder-tip pain after gynecological laparoscopic surgery

Chih-Hsuan Sao^a, Maresol Chan-Tiopianco^b, Kai-Cheng Chung^a, Yi-Jen Chen^{a,c,d},
Huann-Cheng Horng^{a,c}, Wen-Ling Lee^{e,*}, Peng-Hui Wang^{a,c,d,f,g,*}

^aDepartment of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan, ROC; ^bDepartment of Obstetrics and Gynecology, San Lazaro Hospital, Manila, Philippines; ^cDepartment of Obstetrics and Gynecology, National Yang-Ming University, Taipei, Taiwan, ROC; ^dInstitute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan, ROC; ^eDepartment of Medicine, Cheng-Hsin General Hospital, Taipei, Taiwan, ROC; ^fDepartment of Medical Research, China Medical University Hospital, Taichung, Taiwan, ROC; ^gFemale Cancer Foundation, Taipei, Taiwan, ROC

Abstract: Laparoscopy, one of minimally invasive procedures, is a commonly used procedure in diagnosis and management of various kinds of clinical problems, including gynecologic organ-related diseases. Compared with conventional exploratory laparotomy, the benefits of laparoscopic surgery include reduction of surgical wound, decreasing in postoperative pain, shortening hospital stay, rapid recovery, and a better cosmetic result. However, there are still up to 80% of patients after laparoscopic surgery complaining of high levels of pain and needing pain relief. Postlaparoscopic pain can be separated into distinct causes, such as surgical trauma- or incision wound-associated inflammatory change, and pneumoperitoneum (carbon dioxide [CO₂])-related morphological and biochemical changes of peritoneum and diaphragm. The latter is secondary to irritation, stretching, and foreign body stimulation, leading to phrenic neuropraxia and subsequent shoulder-tip pain (STP). STP is the most typical unpleasant experience of patients after laparoscopic surgery. There are at least 11 strategies available to attempt to decrease postlaparoscopic STP, including (1) the use of an alternative insufflating gas in place of CO₂, (2) the use of low-pressure pneumoperitoneum in place of standard-pressure pneumoperitoneum, (3) the use of warmed or warmed and humidified CO₂, (4) gasless laparoscopy, (5) subdiaphragmatic intraperitoneal anesthesia, (6) local intraperitoneal anesthesia, (7) actively expelling out of gas, (8) intraperitoneal drainage, (9) fluid instillation, (10) pulmonary recruitment maneuvers, and (11) others and combination. The present article is limited in discussing postlaparoscopic STP. We extensively review published articles to provide a better strategy to reduce postlaparoscopic STP.

Keywords: Pain; Postlaparoscopic surgery; Prevention; Reduction; Shoulder-tip pain

1. INTRODUCTION

The evolution of minimally invasive techniques for surgical intervention, including the purpose of diagnosis and therapy has resulted in the dramatic and significant change of the use of surgery in management of various kinds of diseases.¹⁻¹¹ Minimally invasive surgery (MIS), including laparoscopic surgery, has been widely accepted in place of conventional exploratory laparotomy in management of various kinds of benign gynecologic diseases.¹²⁻¹⁵ Among these, MIS is reported the best choice in

the specific-type diseases, such as endometriosis.¹⁶⁻²⁰ MIS is as effective as conventional laparotomy for the management of certain-type gynecological malignancies, because of sharing similar advantages of MIS for benign gynecologic diseases but absence in deteriorated oncologic outcome.²¹⁻²⁶ Some recent reports have much concerned about the issue of safety in the use of MIS for cervical cancers.²⁷⁻³⁰ We still believe that much effort has been developed to overcome the potential limitations of MIS with enthusiasm. MIS has many advantages, such as significant reduction in surgical incision wound, less wound-related pain, less analgesia use, minimally trauma and injury, better cosmetic results, a shorter hospital stay, rapid recovery time, and earlier return to daily activities and work compared with conventional exploratory laparotomy.^{12-15,21-26}

Despite these advantages of MIS, there are still up to 80% of patients (ranging from 35% to 80%) who experience severe pain and require pain relief for their unpleasant feeling or suffering.^{1,31-69} The characteristics of pain are different between MIS, especially gas laparoscopic surgery (keyhole surgery), and exploratory laparotomy. Shoulder-tip pain (STP) and upper abdomen pain are the best examples.⁵⁷⁻⁶³ Early discharge and shorter hospital stay is popular in patients undergoing MIS; however, contributing to the high possibility of unfamiliarity to the post-MIS pain by both physicians and patients, it results in

*Address Correspondence: Dr. Wen-Ling Lee, Department of Medicine, Cheng-Hsin General Hospital, 45, Cheng-Hsin Street, Taipei 112, Taiwan, ROC.; Dr. Peng-Hui Wang, Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, 201, Section 2, Shi-Pai Road, Taipei 112, Taiwan, ROC. E-mail address: phwang@vghtpe.gov.tw; pongpongwang@gmail.com (P.-H. Wang).

Conflicts of interest: The authors declare that they have no conflicts of interest related to the subject matter or materials discussed in this article.

Journal of Chinese Medical Association. (2019) 82: 819-826.

Received August 27, 2019; accepted August 27, 2019.

doi: 10.1097/JCMA.000000000000190.

Copyright © 2019, the Chinese Medical Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

missing the diagnosis, and subsequently inadequate evaluation and improper management. Some are intolerable post-MIS pain, leading to a significant unnecessary increase of analgesia use, slower recovery, longer hospital stays, and rarely, readmission.³²

A better understanding of the mechanism and/or pathophysiology of post-MIS pain can be presumed and developed from the increased number of clinical trials and a better strategy algorithm to maintain a regulated and orchestrated prevention and management.³¹⁻⁶⁹ The present article is a discussion of post-MIS pain, which limited to postoperative STP. We extensively review published articles to provide a better strategy to reduce postoperative STP.

2. PAIN

As shown above,^{24-26,53,59,60} MIS is generally considered to be less painful than exploratory laparotomy. However, post-MIS pain still affects the quality of life and is one of the important reasons for delayed discharge or interference of coming back to normal activities.⁵⁵ Post-MIS pain can be separated into incisional pain, STP, and/or upper abdominal pain.

Pain is a sensory dimension (intensity) and an emotional dimension experience (unpleasantness) associated with actual or potential tissue damage.⁷⁰⁻⁷² Acute pain is the normal predicted physiological response to an adverse chemical, thermal, or mechanical stimulus, resulting from activation of the pain receptors (nociceptors) at the site of damage, which plays a critical and vital role in providing warning signals to avoid further damage and possibly be rescued from the damage.⁷⁰ Acute pain is also accompanied with the activation of sympathetic system of the autonomic nervous system which is reflective by tachycardia, diaphoresis, shallow and rapid respiratory pattern, restlessness, irritability, facial grimacing, anxiety, pallor, pupil dilation, and/or hypertension.⁷⁰ Since MIS-related (laparoscopic gynecological surgery-related) pain involves skin, peritoneal and visceral organs, the pain can be originated from nociceptive pain or non-nociceptive pain. The former includes somatic pain (well localized topographically) and visceral pain (diffuse and poorly localized, referable, accompanied with motor and autonomic reflexes, such as nausea and vomiting), and the later includes neuropathic or idiopathic pain.⁷⁰

Nociceptive pain is an initiation of releasing and producing numerous factors, such as globulin, protein kinases, arachidonic acid, histamine, nerve growth factor, substance P, calcitonin gene-related peptide, among others, and stimulates transducer channels by damaged tissue, and then activates or sensitizes nociceptors in the periphery, followed by transducing noxious stimulus into electrochemical impulse and transmitting to the dorsal horn of the spinal cord and crossing over to the contralateral side to the higher rostral centers in the central nervous system (thalamus or other brain areas like dorsolateral pons).⁷⁰⁻⁷² Acute and well-localized fast pain is mediated through medium diameter myelinated afferents, including A-delta, also calling type I with characteristics of higher temperature thresholds but low threshold to mechanical and chemical stimuli and type II characterized with a much greater sensitivity to heat but a very high mechanical threshold.⁷² By contrast, a poorly localized and slow pain is mediated by small diameter unmyelinated fibers (polymodal fibers responding to both mechanical and thermal noxious stimuli).⁷²

3. INCISIONAL WOUND PAIN

Incisional wounds, ranging from one incision (single port wound) to multiple and separate incisions (multiple port wounds) need a cut in the skin, subcutaneous tissues, and peritoneal tissues.¹¹ It causes damage of the tissues (producing wounds), subsequently

resulting in the stimulation of peripheral nociceptors to produce pain sensation.⁷⁰⁻⁷² In addition, inflammation of the wound can stimulate the nociceptors with resultant and exacerbating pain.⁷³⁻⁷⁷ Therefore, aspirin, and selective and nonselective non-steroidal anti-inflammatory drugs, having been known as analgesic, antipyretic and anti-inflammatory effects, are reported to inhibit prostaglandin E2 production and act as effective pain-killers.⁷⁸⁻⁸³ In addition, some opioid types analgesia or central types of pain-relieving agents, such as paracetamol, or steroid drugs are also frequently used in patients after surgery.^{77,84-87} The above-mentioned medication can be provided by oral, intravenous, or intramuscular routes; however, only oral form can be taken after discharge. Although oral medication is convenient,⁸⁸ gastrointestinal tract irritation and possible side effects, such as allergy may occur.⁸⁹⁻⁹⁵ Therefore, immediate and adequate pain control to minimize the pain scores (visual analog scale [VAS] ranging from 0 to 10 cm) and no need of further care after discharge is welcomed.⁹⁶⁻¹⁰⁰ Therefore, wound infiltration with local anesthetic agents done at the end of the surgery is tested. One meta-analysis study showed the effectiveness of the use of local anesthesia infiltration to the incisional wound in the reduction of wound pain after laparoscopic surgery.⁵⁶ The pain scores were statistically significantly lower in the incisional wound by local anesthetic infiltration than no local anesthesia group at 4 to 8 hours (13 trials containing 806 subjects with mean difference [MD] -1.33 cm on the VAS; 95% CI, -1.54 to -1.12) and 9 to 24 hours (12 trials containing 756 subjects; MD -0.36 cm on the VAS; 95% CI, -0.53 to -0.20), respectively, contributing to higher proportion of patients who were discharged as day surgery (66% vs 42.6%).⁵⁶ However, the length of hospital stay was not different between the local anesthetic agent treatment group and the control group,⁵⁶ supporting the concept that significantly less wound pain occurs in patients undergoing MIS and wound pain after MIS is not a remarkable issue for both physicians and patients. Therefore, the clinical importance of this reduction in pain (incisional wound pain after laparoscopic surgery) is likely to be small.⁵⁶

4. CAUSES OF STP

The cause of post-MIS STP is not fully elucidated, and supposed multifactorial and possibly referred pain.¹ Probable explained theories are least three. The first theory is carbonic acid production inducing a reduction in the peritoneal pH to damage and irritate the peritoneal and diaphragmatic nerves, leading to STP.^{32,60} This irritative effect of carbonic acid on the peritoneum and diaphragm is due to the conversion of carbon dioxide (CO₂) gas to carbonic acid by carbonic anhydrase,¹⁰¹⁻¹⁰³ which occurs in the moist surface of peritoneum and diaphragm.⁶⁰ The further evidence is supported by the use of carbonic anhydrase inhibitor-acetazolamide in the reduction of STP dramatically,¹⁰⁴ although the similar results cannot always be reproduced by other studies.^{32,66,105}

The second theory is the residual pockets of gas in the abdominal cavity (also calling visceral ligament traction), which is supported by the followings: (1) presence of CO₂ gas pockets between the liver and diaphragm leading to loss of negative pressure in the peritoneal cavity, and thereby the loss of suction support of the liver and diaphragm, allowing traction on the triangular and coronary ligaments of the liver, leading to sub-diaphragmatic pain and STP; (2) a close correlation between the amount of residual gas or CO₂ bubble volume under the right hemidiaphragm and STP; (3) the positive correlation between delayed absorption of CO₂ and longer STP; (4) the positional nature of STP occurring when women are sitting up and mobilizing; and (5) the STP occurring generally more than four hours after procedure.⁶⁰

The last is the tissue trauma theory (also calling neuropraxia theory). The stretching and/or injury of the peritoneum and diaphragm by pneumoperitoneum results in tearing of blood vessels, traction of nerves (eg, phrenic nerve), and release of inflammatory mediators, which elicit the referred pain to the shoulder.⁶⁰ The correlation of the degree of stretching and the severity of STP has been reported before.¹⁰⁶

After discussing the possible causes of postlaparoscopic STP, the next section focuses on the strategy in the prevention and reduction of severity of STP. The strategies include the use of an alternative insufflating gas in establishing pneumoperitoneum; the use of warmed or humidified insufflating gas, low-pressure pneumoperitoneum; intraperitoneal fluid instillation; the use of intraperitoneal anesthetic agents; the use of intraperitoneal drains; and specific methods for expelling out of gas, such as active suction of gas or manually forcing gas out of the abdominal cavity at the end of surgery and pulmonary recruitment maneuver (PRM).³¹⁻⁶⁹

5. THE USE OF AN ALTERNATIVE INSUFFLATING GAS IN THE REDUCTION OF STP

To perform the laparoscopic surgery, the establishment of pneumoperitoneum is a crucial element. It is an initial step to provide sufficient working and viewing space and ensure adequate visualization of camera and manipulation of instruments when laparoscopic surgery is performed.⁴⁰ To establish pneumoperitoneum, two steps are needed, including the initial set up of port wound (entry into the abdominal cavity via trocar) and the following insufflation of gas and of course, reverse of them (insufflation of gas initially via Veress needle and the following trocar insertion) is also used in routinely clinical practice.⁴⁰

The ideal gas to establish pneumoperitoneum should fulfill the following criteria, including cheap, easy to obtain, colorless, nonflammable, nonexplosive, easily excreted, and completely nontoxic to patients.⁴⁰ There are many gases available to act the resource for establishing pneumoperitoneum. They are CO₂, helium, argon, nitrogen, nitrous oxide, room air, and others.⁴⁰ Carbon dioxide is the most popular and well-known gas for this purpose, although CO₂ is still not an ideal form. There are many concerns about the use of CO₂ for laparoscopic surgery because CO₂ is soluble and can be absorbed by the peritoneum and delivered directly to the lung by circulation, which may induce metabolic and respiratory changes.⁴⁰ This acid-based balance might be destroyed during laparoscopic surgery with CO₂. Therefore, the risk of hypercapnia, acidosis, and cardiopulmonary complications, such as tachycardia, cardiac arrhythmia, and lung edema may be increased.⁴⁰ In addition, this metabolic and respiratory imbalance might destroy the normal immune response of the host. Finally, as shown above, CO₂ is a potential cause of postlaparoscopic STP. All of these might be more apparent in the elderly population.⁴⁰ During laparoscopy, monitoring of end-tidal CO₂ concentration is mandatory.

However, the risk of venous or arterial air embolization might be lower in laparoscopic surgery with CO₂, based on the characteristics of CO₂ with easy solubility and quick absorption. In contrast, other gases, such as helium, argon, nitrogen, nitrous oxide, and room air, are not so soluble as CO₂ and need much more time to be absorbed. At the end of surgery, special attention should be done when pneumoperitoneum is established. In addition, it is necessary to maintain low intra-abdominal gas pressure during operation and try much effort to expel the gas out of the abdominal cavity at the end of surgery when these relatively insoluble gases are used. All will increase the risk of air embolization during operation or after operation. That is why there are many studies trying to evaluate the feasibility and

safety of the using alternative gases in place of CO₂ for laparoscopic surgery. Unfortunately, results are not consistent.^{40,107-112} Of most importance, safety issue of these alternative gases is always concerned.^{40,107-112}

6. THE USE OF LOW-PRESSURE PNEUMOPERITONEUM IN THE REDUCTION OF STP

As shown before, to permit organs and structures within the abdominal cavity to be viewed, inflation of the abdominal cavity with CO₂ (pneumoperitoneum) should be given during laparoscopic surgery.⁶⁰ Pneumoperitoneum (with increasing intra-abdominal pressure) significantly decreases venous return, preload, and cardiac output, as well as increases heart rate, systemic and pulmonary vascular resistance, resulting in the stimulation of neurohumoral vasoactive system, regardless of which type of gas is given.¹¹³

The gas pressure for gynecologic laparoscopic surgery usually ranges from 12 mmHg to 14 mmHg (no exceeding 15 mmHg) because the study showed that in ASA (the American Society of Anesthesiologists classification) I and II patients, the hemodynamic and circulatory effects of these pressures are generally not clinically relevant.¹¹³ However, this pressure is one of causes contributing to the unwanted side effects, such as STP, and sometimes, this standard pressure might increase the risk of intraoperative hemodynamic instability in ASA III and IV patients.¹¹³ Therefore, if technically feasible, gasless or low-pressure pneumoperitoneum should be considered for patients with limited cardiac and pulmonary function, contributing to the consideration of using low-pressure in place of the standard-pressure to establish pneumoperitoneum during laparoscopic surgery.^{60,113}

As expected, low pressure results in inadequate pneumoperitoneum, and inadequate operative field, contributing to impairment of identification of normal organs and target lesions, and of most importance, low pressure may be correlated with technique difficulty and subsequently increasing risk of intraoperative complication and possible life-threatening situation. Previous trial has shown that pain can be successful reduced by low-pressure pneumoperitoneum but operative time is longer, and hemorrhage is also increased compared with standard-pressure and high-pressure pneumoperitoneum,⁵⁷ supporting uncertainty of safety of low pressure pneumoperitoneum when we perform laparoscopic surgery.

In 2014, a Cochrane review found that approximately 90% of people could be successfully managed by low-pressure laparoscopic cholecystectomy, but the authors concluded no evidence to support the use of low-pressure pneumoperitoneum in routine and uncertainty of safety of low-pressure pneumoperitoneum.⁶⁰ In 2015 and 2016, review also found that low pressures were associated with worse visualization of the surgical field (risk ratio, 10.31; 95% CI, 1.29-82.38), although a statistically significant but modest diminution in postoperative pain of MD, -0.38 cm on the VAS (95% CI, -0.67 to -0.08) during the immediate postoperative period when using low pressure of 8 mmHg compared with \geq 12 mmHg and of MD, 0.50 cm on the VAS (95% CI, -0.80 to -0.21) 24 hours after the laparoscopic surgery.^{114,115} Therefore, the authors did not suggest the use of low pressure during gynecologic laparoscopy because of minimal improvement of pain scores but significantly compromising visualization of the surgical field.^{114,115}

Two recent prospective randomized trials tried to evaluate the feasibility of the use of low-pressure laparoscopic surgery in the management of benign gynecologic pathology.^{1,46} Low-pressure was set up as 7-8 mmHg, and standard-pressure was 15 mmHg CO₂. The results of both studies showed patients undergoing low-pressure laparoscopic surgery had significantly

lower postlaparoscopic abdominal pain and STP scores.^{1,46} In addition, fewer vegetative alterations, lower pain medication requirements, a shorter postoperative hospital stay, and lower intra- and postoperative arterial partial pressure of carbon dioxide values are found in the women undergoing low pressure laparoscopic surgery compared to those undergoing the standard-pressure laparoscopic surgery,¹ suggesting low-pressure laparoscopic surgery for benign lesions is a feasible and safe technique.^{1,46}

7. GASLESS LAPAROSCOPY AND ACTIVELY EXPELLING OUT OF GAS TO REDUCE STP

It is interesting to find studies to use different strategies to decrease residual gas within the abdominal cavity at the end of laparoscopic surgery.^{49,116,117} One is an earlier prospective randomized trial in 1998 to show there is no statistical difference in scores for STP between gasless (a Laprolift system) and standard-pressure laparoscopic surgery for tubal ligation.¹¹⁶ The other study found that the active gas aspiration has lowered pain intensity of STP scores than the simple gas evacuation does, and these statistically significant results are found at 6, 12, and 24 hours after surgery, suggesting active gas aspiration is recommended in routine for decreasing postoperative STP.⁴⁹ To remove residual air within the abdominal cavity as much as possible, the air can be first vacuumed from the pelvic cavity in Trendelenburg position and then the patients are put in anti-Trendelenburg position, where the remaining gas can be shifting toward subdiaphragmatic area, and the suction tube is shifted to a position next to the camera canal and the remaining air is suctioned.¹¹⁷ Although the results are not consistent, we still favor the effort to remove the gas in the abdominal cavity as much as possible to minimize the residual volume of gas at the end of laparoscopic surgery.

8. THE USE OF WARMED OR WARMED AND HUMIDIFIED CO₂ IN THE REDUCTION OF STP

Carbon dioxide used in laparoscopic surgery is typical at 21°C with 0 percent relative humidity.¹¹⁷ This cold and dry gas may cause hypothermia, and postoperative pain or fatigue.¹¹⁷ Therefore, it is rationale to suppose that the use of warmed and humidified gas in place of cold and dry gas might decrease the postlaparoscopic STP. In fact, this hypothesis was supported by the earlier meta-analysis.¹¹⁸ A statistically significant reduction of pain (MD, -0.39 cm on the VAS, 95% CI, -0.67 to -0.18 within 6 hours, MD, -0.34 cm on the VAS, 95% CI, -0.61 to -0.07 on day 1, and MD, -0.88 cm on the VAS, 95% CI, -1.30 to -0.45 on day 3, respectively) was found in the humidification and warming of the insufflated CO₂ group compared to conventional insufflation of cold and dry CO₂ group.¹¹⁸ The study further supported the benefits of the use of heated humidified CO₂ in place of dry and cold CO₂ for laparoscopic surgery, because the former is associated with lesser postoperative pain, lower risk of postoperative hypothermia, and lower analgesic requirements.¹¹⁹ It is reasonable to use warmed and humidified CO₂ for laparoscopic surgery, but this suggestion is still not popular in routinely clinical practice. In fact, a recent meta-analysis focusing on gynecologic laparoscopy, including three studies^{120–122} to assess severity of STP and postoperative analgesia usage found there was no evidence of a difference in the incidence, severity or analgesia requirements between women treated with warming, or warmed and humidified insufflation gas and gas in routine use.³² We favor the use of warmed and humidified insufflation

gas for laparoscopic surgery if applicable, but the benefit/cost ratio is still uncertain.

9. THE USE OF SUBDIAPHRAGMATIC INTRAPERITONEAL ANESTHESIA OR LOCAL INTRAPERITONEAL ANESTHESIA IN THE REDUCTION OF STP

As shown before, pain secondary to laparoscopic surgery has been attributed to stretching of the intra-abdominal cavity, peritoneal inflammation, and establishment of pneumoperitoneum and dissection of the abdominal and pelvic viscera.^{32,123} Local administration of anesthesia might have lowered risk of sedation, nausea, gastrointestinal irritation or injury, as well as paralysis, respiratory depression, and allergic effects than systematic use of analgesia has.^{123,124}

Intraperitoneal analgesia after laparoscopic surgery might directly acts at the site by causing a reversible interruption of nervous conduction, and subsequent inhibiting the visceral afferent signaling. Meta-analysis showed that patients treated with intraperitoneal local anesthesia may have a statistically significant reduction of pain scores within the first 6 hours after laparoscopic surgery (MD, -1.82 cm on the VAS, 95% CI, -2.55 to -1.08 at 1 to 2 hours, and MD, -2.00 cm on the VAS, 95% CI, -3.64 to -0.35 at 4 to 6 hours postoperatively, respectively).¹²³ However, this effect was not statistically significant difference at 24 hours postoperatively.¹²³ A meta-analysis published in 2019, including many trials,^{125–127} also supported the evidence that intraperitoneal local anesthesia (not spreading to sub-diaphragm) can be associated with a statistically significant reduction of incidence of STP (odds ratio [OR], 0.23; 95% CI, 0.06–0.92).³² Similar to the reduction of STP in the patients treated with intraperitoneal local anesthesia, patients treated with subdiaphragmatic intraperitoneal local anesthesia also had a statistically significant reduction of severity of STP at 8 hours postoperatively (MD, -0.95 cm on the VAS, 95% CI, -1.70 to -0.19).³² However, it is interesting to find that pain reduction at the different postoperative time points to evaluate incidence or severity of STP seems to be varied greatly. The effectiveness seems to occur immediately after laparoscopic surgery and the effect of pain reduction disappears in the later. That is to say, the reduced pain scores occur transiently and cannot be maintained.

10. THE USE OF INTRAPERITONEAL DRAINAGE IN THE REDUCTION OF STP

Drains are commonly used after surgeries and can be classified as either active or passive by mechanisms and either informative (prophylactic) or therapeutic by purpose.^{128–137} In addition, vacuum drains can further be classified as high negative pressure (typical bottled vacuum drains, such as Redi-vac™ drain (Redivac, Inc, Daventry, Northamptonshire, UK), with advantages of being sealed, closed-circuit system allowing for easy monitoring and safe disposal of the drainage) and low negative pressure, such as bulk-shaped suction devices (eg, Jackson-Pratt, compression of bulb to force air out to create negative pressure in the system) and collapsible four-channel vacuum drains J Vac (Blake drain, Ethicon, Inc, Somerville, NJ).¹³² After laparoscopic surgery, a dead space and pneumoperitoneum is created, and body has a natural tendency to fill this space with fluid or air. Therefore, prophylactic low-negative-pressure drains can be used to work gently to evacuate excess fluid and air, although it is not routinely recommended. An earlier meta-analysis did not support the routine use of a peritoneal gas drain following gynecological laparoscopy because of very little evidence of an overall benefit from this approach, and in addition, no association with a reduction in the requirement of analgesia and

anti-emetics for STP and total pain when compared to no use of peritoneal gas drain group.¹³⁵ A recent meta-analysis showed there is associated between an intraperitoneal drain and a reduction in the incidence of STP when compared with no intraperitoneal drain at all time points assessed postoperatively with OR, 0.47 (95% CI, 0.25-0.86) at 3-4 hours; OR, 0.08 (95% CI, 0.02-0.36) at 12 hours; OR, 0.3 (95% CI, 0.2-0.46) at 24 hours; and OR, 0.4 (95% CI, 0.21-0.74) at 48 hours postoperatively, respectively.³² Not only is incidence of STP decreased when drain is applied after laparoscopic surgery, but also severity of STP decreased with MD, -1.69 cm on the VAS, 95% CI, -2.2 to -1.19 at 12 hours postoperatively; MD, -1.85 cm on the VAS, 95% CI, -2.15 to -1.55 at 24 hours after operation; MD, -0.7 cm on the VAS, 95% CI, -0.95 to -0.44 at 48 hours postoperatively; and MD, -0.8 cm on the VAS, 95% CI, -1.15 to -0.05 at 72 hours postoperatively, respectively.³²

11. THE USE OF PRM IN THE REDUCTION OF STP

Since CO₂ or gas pneumoperitoneum is one of most common causes inducing postoperative STP, specific technique for releasing the pneumoperitoneum (expelling out of gas as much as possible to minimize the residual gas within the abdominal cavity after laparoscopic surgery) may be effective in reducing postoperative STP. PRM involve positive pressure ventilation (40-60 cm H₂O for five breaths with the final inflation breath being held for a maximum of 5 seconds) with combination of gentle abdominal pressure at the completion of the laparoscopic surgery whilst the patients are still in a Trendelenburg position to expel CO₂ out via open trocars in the abdominal wall.^{32,61,64} Since PRM seems to be an easily performed and potentially preventive strategy of postoperative STP, many physicians highly recommended its routine use. In fact, an earlier meta-analysis also supported the benefits of the use of PRM in the reduction of postoperative STP at the end of laparoscopic surgery, because PRM significantly decreased postoperative STP 12 hours (MD, -1.55 cm on the VAS, 95% CI, -2.01 to -1.10), 24 hours (MD, -1.59 cm on the VAS, 95% CI, -2.00 to -1.18), and 48 hours postoperatively (MD, -0.93 cm on the VAS, 95% CI, -1.37 to -0.50), respectively.⁵⁰ A recent meta-analysis failed to show the reduction of incidence of postoperative STP in patients treated with PRM (OR, 0.77; 95% CI, 0.57-1.05 and OR, 0.79; 95% CI, 0.56-1.11) compared to those treated with standard control.³² However, in agreement with an earlier meta-analysis to have a reduction of severity of STP in PRM group,⁵⁰ a statistically significant reduction of STP is found at all times investigated, including MD, -0.29 cm on the VAS, 95% CI, -0.48 to -0.09 at 3-6 hours; MD, -0.58 cm on the VAS, 95% CI, -0.78 to -0.37 at 12 hours; MD, -0.66 cm on the VAS, 95% CI, -0.82 to -0.50 at 24 hours; MD, -1.26 cm on the VAS, 95% CI, -2.23 to -0.29 at 36 hours; and MD, -0.72 cm on the VAS, 95% CI, -0.99 to -0.45 at 48 hours after laparoscopic surgery, respectively.³²

12. THE USE OF FLUID INSTILLATION IN THE REDUCTION OF STP

Intraperitoneal fluid instillation with 1000-1500 mL of warm saline (or 15-30 mL/kg body weight) into the abdominal cavity was performed in patients at the end of the gynecological laparoscopic procedures whilst patients are still in Trendelenburg positioning until it "spilled out of the remaining open trocars".^{32,61,64} Meta-analysis seems to support the value of fluid instillation in the reduction of STP, because of lower incidence of STP in the fluid instillation group with OR, 0.67; 95% CI, 0.39-1.14 at 12 hours; OR, 0.38; 95% CI, 0.22-0.66; and OR, 0.38; 95% CI, 0.21-0.67, respectively; and less severity of STP in the fluid instillation group with MD, -1.69 cm on the VAS, 95% CI, -2.55 to -0.83 at 12 hours; MD, -2.27 cm on the VAS, 95% CI,

-3.06 to -1.48 at 24 hours; and MD, -1.44 cm on the VAS, 95% CI, -2.07 to -0.81 at 48 hours postoperatively, respectively compared to no fluid instillation group.^{32,61,64}

13. THE USE OF COMBINATION METHODS IN THE REDUCTION OF STP

As shown comments by Dr. Sharp HT⁶² for our previous publication,⁶¹ most of us have tried some methods to reduce one clinical problem, but there are some patients still suffering from other clinical problems. STP and surgical pain might be different, and many clinical problems might be complicated and multifactorial. Therefore, combination of many methods to become a brand new strategy might be a better choice, which may cover more clinical situations. In fact, recent prospective randomized studies seem to be conducted as the use of different combination of all effective tools in the management of troublesome clinical problems. Our previous study using the combination of PRM and NS instillation can successfully and effectively decrease upper abdominal pain and STP,⁶¹ although the recent study by van Dijk et al³⁴ cannot reproduce our findings. However, in the study by van Dijk et al³⁴, the mean VAS score for abdominal pain at 8 hours after surgery was indeed significantly lower in the intervention group compared with the control group (3.2 vs 4.2; $p = 0.02$),³⁴ suggesting feasibility and acceptability of this approach for the patients who undergo laparoscopic gynecologic surgery.

In conclusion, MIS is popular and well-come procedures, which can be used as better alternatives in the management of various kinds of gynecological diseases or other surgical interventions. To achieve the better road to take care of the patients who need surgical intervention, any improvement of quality of life after operation should be taken into consideration. Pain is the fifth vital sign, emphasizing the need to reduce patient suffering.⁶² Safety is a priority for all managements, therapies, and surgeries. Any inexpensive low-technology method, eg, instillation of normal saline at the conclusion of laparoscopic gynecologic procedures to reduce postoperative STP is welcome. We should always be cautious about the fluid overload for patients who are risky in cardiovascular diseases. In fact, instillation of normal saline within the abdominal cavity, this approach has been used very commonly for adhesion prevention at the end of surgery.⁶² Based on the current review, a specific technique for releasing the pneumoperitoneum; intraperitoneal fluid instillation; an intraperitoneal drain; and local anesthetic applied to the peritoneal cavity or to the subdiaphragmatic area, and warmed and humidified insufflating gas may play a role on the reduction of postoperative STP, although evidence needs more studies to verify.

ACKNOWLEDGMENTS

This study was supported by grants from the Ministry of Science and Technology (MOST 106-2314-B-075-061-MY3) and the Taipei Veterans General Hospital (Grant VGH108C-085), Taipei, Taiwan. The authors also appreciate very much financial support from the Female Cancer Foundation, Taipei, Taiwan.

REFERENCES

1. Radosa JC, Radosa MP, Schweitzer PA, Radosa CG, Stotz L, Hamza A, et al. Impact of different intraoperative CO₂ pressure levels (8 and 15 mmHg) during laparoscopic hysterectomy performed due to benign uterine pathologies on postoperative pain and arterial pCO₂: a prospective randomised controlled clinical trial. *BJOG* 2019;126:1276-85.
2. Tu YA, Chang WC, Wu CJ, Sheu BC. Improved hemostasis with plasma kinetic bipolar sealing device in the vaginal steps of laparoscopic-assisted vaginal hysterectomy. *Taiwan J Obstet Gynecol* 2019;58:64-7.
3. Lee CL, Wu KY, Huang CY, Yen CF. Comparison of ligasure™ tissue fusion system and a conventional bipolar device in hysterectomy via

- natural orifice transluminal endoscopic surgery (NOTES): a randomized controlled trial. *Taiwan J Obstet Gynecol* 2019;58:128–32.
4. Chen CW, Chang HC, Huang TF, Liao CC, Huang RL, Lai HC. Transition from multiport to single-site surgery: a single institution experience in robotic supracervical hysterectomy for benign gynecological diseases. *Taiwan J Obstet Gynecol* 2019;58:514–9.
 5. Horng HC, Tsui KH, Wang PH. The powerful hemostatic devices are one of the milestones for successful laparoscopic surgery. *J Chin Med Assoc* 2018;81:92–3.
 6. Huang HY, Liu YC, Li YC, Kuo HH, Wang CJ. Comparison of three different hemostatic devices in laparoscopic myomectomy. *J Chin Med Assoc* 2018;81:178–82.
 7. Chen YC, Lin HH, Hsiao SM. Comparison of robotic assisted laparoscopic myomectomy with barbed sutures and traditional laparoscopic myomectomy with barbed sutures. *Taiwan J Obstet Gynecol* 2018;57:709–12.
 8. Moon HS. Tips on robotic single-site surgery suture technique: screwing and clockwise direction suture technique for robotic single-site surgery. *Taiwan J Obstet Gynecol* 2018;57:432–4.
 9. Wang SC, Chen CC, Yang CK, Hung SW, Jan YJ, Ou YC. Pathological outcomes in men with prostate cancer who are eligible for active surveillance. *J Chin Med Assoc* 2018;81:348–51.
 10. Jiang H, Jiang O, Xia X, Su S, Li B. Laparoscopic versus open hepatectomy approach for regional hepatolithiasis: a meta-analysis. *J Chin Med Assoc* 2018;81:429–36.
 11. Sun HD, Horng HC, Liu CH, Hsiao SM, Chen YJ, Chang WH, et al. Comparison of single-port and three-port laparoscopic salpingectomy in the management for tubal pregnancy. *J Chin Med Assoc* 2018;81:469–74.
 12. Wen KC, Sung PL, Chang WH, Horng HC, Chen YJ, Lee WL, et al. A case-control study to compare the outcome of women treated by two minimally invasive procedures-ultraminilaparotomy myomectomy and laparoscopic myomectomy. *Taiwan J Obstet Gynecol* 2018;57:264–9.
 13. Wen KC, Chen YJ, Sung PL, Wang PH. Comparing uterine fibroids treated by myomectomy through traditional laparotomy and 2 modified approaches: ultraminilaparotomy and laparoscopically assisted ultraminilaparotomy. *Am J Obstet Gynecol* 2010;202:144.e1–8.
 14. Lee WL, Liu WM, Cheng MH, Chao HT, Fuh JL, Wang PH. Uterine vascular occlusion in management of leiomyomas: laparoscopy vs laparotomy. *J Minim Invasive Gynecol* 2009;16:562–8.
 15. Wang PH, Liu WM, Fuh JL, Chao HT, Yuan CC, Chao KC. Symptomatic myoma treated with laparoscopic uterine vessel occlusion and subsequent immediate myomectomy: which is the optimal surgical approach? *Fertil Steril* 2009;92:762–9.
 16. Chen PL, Lin HH, Hsiao SM. Predictors of subsequent pregnancy in women who underwent laparoscopic cornuostomy or laparoscopic wedge resection for interstitial pregnancy. *J Chin Med Assoc* 2019;82:138–42.
 17. Nezhat C, Li A, Falik R, Copeland D, Razavi G, Shakib A, et al. Bowel endometriosis: diagnosis and management. *Am J Obstet Gynecol* 2018;218:549–62.
 18. Singh SS, Suen MW. Surgery for endometriosis: beyond medical therapies. *Fertil Steril* 2017;107:549–54.
 19. Tan SJ, Chen CH, Yeh SD, Lin YH, Tzeng CR. Pregnancy following robot-assisted laparoscopic partial cystectomy and gonadotropin-releasing hormone agonist treatment within three months in an infertile woman with bladder endometriosis. *Taiwan J Obstet Gynecol* 2018;57:153–6.
 20. Fong YF, Hon SK, Low LL, Lim Mei Xian K. The clinical profile of young and adolescent women with laparoscopically diagnosed endometriosis in a Singapore tertiary hospital. *Taiwan J Obstet Gynecol* 2017;56:181–3.
 21. Deura I, Shimada M, Azuma Y, Komatsu H, Nagira K, Sawada M, et al. Comparison of laparoscopic surgery and conventional laparotomy for surgical staging of patients with presumed low-risk endometrial cancer: the current state of Japan. *Taiwan J Obstet Gynecol* 2019;58:99–104.
 22. Lee CL, Kusunoki S, Huang CY, Wu KY, Lee PS, Huang KG. Surgical and survival outcomes of laparoscopic staging surgery for patients with stage I ovarian cancer. *Taiwan J Obstet Gynecol* 2018;57:7–12.
 23. Li YT, Horng HC, Wang PH. The role of complete staging surgery for pure endometrioid-type endometrial cancer. *J Chin Med Assoc* 2018;81:663–4.
 24. Galaal K, Donkers H, Bryant A, Lopes AD. Laparoscopy versus laparotomy for the management of early stage endometrial cancer. *Cochrane Database Syst Rev* 2018;10:CD006655.
 25. Asher R, Obermair A, Janda M, Gebski V. Disease-free and survival outcomes for total laparoscopic hysterectomy compared with total abdominal hysterectomy in early-stage endometrial carcinoma: a meta-analysis. *Int J Gynecol Cancer* 2018;28:529–38.
 26. Wang YR, Lu HF, Huo HC, Qu CP, Sun GX, Shao SQ. A network meta-analysis of comparison of operative time and complications of laparoscopy, laparotomy, and laparoscopic-assisted vaginal hysterectomy for endometrial carcinoma. *Medicine (Baltimore)* 2018;97:e0474.
 27. Cusimano MC, Baxter NN, Gien LT, Moineddin R, Liu N, Dossa F, et al. Impact of surgical approach on oncologic outcomes in women undergoing radical hysterectomy for cervical cancer. *Am J Obstet Gynecol* 2019. doi: 10.1016/j.ajog.2019.07.009. [Epub ahead of print]
 28. Doo DW, Kirkland CT, Griswold LH, McGwin G, Huh WK, Leath CA 3rd, et al. Comparative outcomes between robotic and abdominal radical hysterectomy for IB1 cervical cancer: results from a single high volume institution. *Gynecol Oncol* 2019;153:242–7.
 29. Melamed A, Margul DJ, Chen L, Keating NL, Del Carmen MG, Yang J, et al. Survival after minimally invasive radical hysterectomy for early-stage cervical cancer. *N Engl J Med* 2018;379:1905–14.
 30. Ramirez PT, Frumovitz M, Pareja R, Lopez A, Vieira M, Ribeiro R, et al. Minimally invasive versus abdominal radical hysterectomy for cervical cancer. *N Engl J Med* 2018;379:1895–904.
 31. Lee J, Park C, Kim J, Ki Y, Cha SH, Kim JY. The effect of low-pressure pulmonary recruitment maneuver on post-laparoscopic shoulder pain: a randomized controlled trial. *J Minim Invasive Gynecol* 2019. doi: 10.1016/j.jmig.2019.03.020. [Epub ahead of print]
 32. Kaloo P, Armstrong S, Kaloo C, Jordan V. Interventions to reduce shoulder pain following gynaecological laparoscopic procedures. *Cochrane Database Syst Rev* 2019;1:CD011101.
 33. Kwack JY, Kwon YS. Immediate postoperative pain control with ropivacaine following laparoscopic-assisted vaginal hysterectomy: a randomized double-blind pilot study. *Taiwan J Obstet Gynecol* 2018;57:654–8.
 34. van Dijk J, Dedden SJ, Geomini P, van Kuijk S, van Hanegem N, Meijer P, et al. Randomised controlled trial to estimate reduction in pain after laparoscopic surgery when using a combination therapy of intraperitoneal normal saline and the pulmonary recruitment manoeuvre. *Bjog* 2018;125:1469–76.
 35. Tu FF. Focus on optimising laparoscopic surgery. *Bjog* 2018;125:1477.
 36. Daniels J. Reducing post-laparoscopy shoulder pain: are simple strategies enough? *Bjog* 2018;125:1478.
 37. Kendall MC, Castro-Alves LJ. Re: randomised controlled trial to estimate reduction in pain after laparoscopic surgery when using a combination therapy of intraperitoneal normal saline and the pulmonary recruitment manoeuvre. *Bjog* 2018;125:1493.
 38. Güngördük K, Aşcıoğlu O, Özdemir İA. Effect of the pulmonary recruitment maneuver on pain after laparoscopic gynecological oncologic surgery: a prospective randomized trial. *J Gynecol Oncol* 2018;29:e92.
 39. Yong L, Guang B. Intraperitoneal ropivacaine instillation versus no intraperitoneal ropivacaine instillation for laparoscopic cholecystectomy: a systematic review and meta-analysis. *Int J Surg* 2017;44:229–43.
 40. Yu T, Cheng Y, Wang X, Tu B, Cheng N, Gong J, et al. Gases for establishing pneumoperitoneum during laparoscopic abdominal surgery. *Cochrane Database Syst Rev* 2017;6:CD009569.
 41. van Dijk JEW, Dedden SJ, Geomini PMAJ, Meijer P, van Hanegem N, Bongers MY. Postlaparoscopic reduction of pain by combining intraperitoneal normal saline and the pulmonary recruitment maneuver (POLAR BEAR trial). RCT to estimate reduction in pain after laparoscopic surgery when using a combination therapy of intraperitoneal normal saline and the pulmonary recruitment maneuver. *BMC Womens Health* 2017;17:42.
 42. Mirhosseini H, Avazbakhsh MH, Hosseini Amiri M, Entezari A, Bidaki R. Effect of oral clonidine on shoulder tip pain and hemodynamic response after laparoscopic cholecystectomy: a randomized double blind study. *Anesth Pain Med* 2017;7:e61669.
 43. Ryu K, Choi W, Shim J, Song T. The impact of a pulmonary recruitment maneuver to reduce post-laparoscopic shoulder pain: a randomized controlled trial. *Eur J Obstet Gynecol Reprod Biol* 2017;208:55–60.
 44. Warlé MC, Dahan A. Does deep neuromuscular block affect pain after laparoscopic surgery? *Eur J Anaesthesiol* 2017;34:23–4.
 45. Unterbuchner C, Werkmann M. Postoperative shoulder pain after laparoscopic hysterectomy with deep neuromuscular blockade and low-pressure pneumoperitoneum. *Eur J Anaesthesiol* 2017;34:25–6.
 46. Sroussi J, Elies A, Rigouzzo A, Louvet N, Mezzadri M, Fazel A, et al. Low pressure gynecological laparoscopy (7 mmHg) with Airseal® system versus a standard insufflation (15 mmHg): a pilot study in 60 patients. *J Gynecol Obstet Hum Reprod* 2017;46:155–8.

47. Madsen MV, Istre O, Staehr-Rye AK, Springborg HH, Rosenberg J, Lund J, et al. Postoperative shoulder pain after laparoscopic hysterectomy with deep neuromuscular blockade and low-pressure pneumoperitoneum: a randomized controlled trial. *Eur J Anaesthesiol* 2016;33:341–7.
48. Zhang H, Liu X, Jiang H, Liu Z, Zhang XY, Xie HZ. Parecoxib increases muscle pain threshold and relieves shoulder pain after gynecologic laparoscopy: a randomized controlled trial. *J Pain Res* 2016;9:653–60.
49. Leelasuwattanakul N, Bunyavehchevin S, Sriprachittichai P. Active gas aspiration versus simple gas evacuation to reduce shoulder pain after diagnostic laparoscopy: a randomized controlled trial. *J Obstet Gynaecol Res* 2016;42:190–4.
50. Pergialiotis V, Vlachos DE, Kontzoglou K, Perrea D, Vlachos GD. Pulmonary recruitment maneuver to reduce pain after laparoscopy: a meta-analysis of randomized controlled trials. *Surg Endosc* 2015;29:2101–8.
51. Choi GJ, Kang H, Baek CW, Jung YH, Kim DR. Effect of intraperitoneal local anesthetic on pain characteristics after laparoscopic cholecystectomy. *World J Gastroenterol* 2015;21:13386–95.
52. Valadan M, Banifatemi S, Yousefshahi F. Preoperative gabapentin to prevent postoperative shoulder pain after laparoscopic ovarian cystectomy: a randomized clinical trial. *Anesth Pain Med* 2015;5:e31524.
53. Bogani G, Martinelli F, Ditto A, Chiappa V, Lorusso D, Ghezzi F, et al. Pneumoperitoneum pressures during pelvic laparoscopic surgery: a systematic review and meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2015;195:1–6.
54. Bogani G, Uccella S, Cromi A, Serati M, Casarin J, Pinelli C, et al. Low vs standard pneumoperitoneum pressure during laparoscopic hysterectomy: prospective randomized trial. *J Minim Invasive Gynecol* 2014;21:466–71.
55. Gurusamy KS, Vaughan J, Davidson BR. Low pressure versus standard pressure pneumoperitoneum in laparoscopic cholecystectomy. *Cochrane Database Syst Rev* 2014;3:CD006930.
56. Loizides S, Gurusamy KS, Nagendran M, Rossi M, Guerrini GP, Davidson BR. Wound infiltration with local anaesthetic agents for laparoscopic cholecystectomy. *Cochrane Database Syst Rev* 2014;3:CD007049.
57. Topcu HO, Cavkaytar S, Kokanalı K, Guzel AI, Islımye M, Doganay M. A prospective randomized trial of postoperative pain following different insufflation pressures during gynecologic laparoscopy. *Eur J Obstet Gynecol Reprod Biol* 2014;182:81–5.
58. Ingelmo PM, Bucciero M, Somaini M, Sahillioglu E, Garbagnati A, Charton A, et al. Intraperitoneal nebulization of ropivacaine for pain control after laparoscopic cholecystectomy: a double-blind, randomized, placebo-controlled trial. *Br J Anaesth* 2013;110:800–6.
59. Donatsky AM, Bjerrum F, Gögenur I. Surgical techniques to minimize shoulder pain after laparoscopic cholecystectomy. A systematic review. *Surg Endosc* 2013;27:2275–82.
60. Donatsky AM, Bjerrum F, Gögenur I. Intraperitoneal instillation of saline and local anesthesia for prevention of shoulder pain after laparoscopic cholecystectomy: a systematic review. *Surg Endosc* 2013;27:2283–92.
61. Tsai HW, Wang PH, Yen MS, Chao KC, Hsu TF, Chen YJ. Prevention of postlaparoscopic shoulder and upper abdominal pain: a randomized controlled trial. *Obstet Gynecol* 2013;121:526–31.
62. Sharp HT. Intraperitoneal carbon dioxide after laparoscopy: are two interventions better than one? *Obstet Gynecol* 2013;121:505–6.
63. Rattner DM. The 0.9% solution? Comment on “Maneuvers to decrease laparoscopy-induced shoulder and upper abdominal pain.” *Arch Surg* 2011;146:1366–7.
64. Tsai HW, Chen YJ, Ho CM, Hseu SS, Chao KC, Tsai SK, et al. Maneuvers to decrease laparoscopy-induced shoulder and upper abdominal pain: a randomized controlled study. *Arch Surg* 2011;146:1360–6.
65. Pappas-Gogos G, Tsimogiannis KE, Zikos N, Nikas K, Manataki A, Tsimoyiannis EC. Preincisional and intraperitoneal ropivacaine plus normal saline infusion for postoperative pain relief after laparoscopic cholecystectomy: a randomized double-blind controlled trial. *Surg Endosc* 2008;22:2036–45.
66. Phelps P, Cakmakaya OS, Apfel CC, Radke OC. A simple clinical maneuver to reduce laparoscopy-induced shoulder pain: a randomized controlled trial. *Obstet Gynecol* 2008;111:1155–60.
67. Esin S, Culha ZK, Kara OF, Yildirim H. A simple clinical maneuver to reduce laparoscopy-induced shoulder pain: a randomized controlled trial. *Obstet Gynecol* 2008;112(2 Pt 1):380; author reply 380.
68. Sarli L, Costi R, Sansebastiano G, Trivelli M, Roncoroni L. Prospective randomized trial of low-pressure pneumoperitoneum for reduction of shoulder-tip pain following laparoscopy. *Br J Surg* 2000;87:1161–5.
69. Pier A, Benedic M, Mann B, Buck V. [Postlaparoscopic pain syndrome. Results of a prospective, randomized study]. *Chirurg* 1994;65:200–8.
70. Talbot K, Madden VJ, Jones SL, Moseley GL. The sensory and affective components of pain: are they differentially modifiable dimensions or inseparable aspects of a unitary experience? A systematic review. *Br J Anaesth* 2019;123:e263–72.
71. Anwar K. Pathophysiology of pain. *Dis Mon* 2016;62:324–9.
72. Chen JS, Sehdev JS. Physiology, Pain. StatPearls [Internet]. Treasure Island, FL: StatPearls Publishing; 2019.
73. Wang PH, Huang BS, Horng HC, Yeh CC, Chen YJ. Wound healing. *J Chin Med Assoc* 2018;81:94–101.
74. Horng HC, Chang WH, Yeh CC, Huang BS, Chang CP, Chen YJ, et al. Estrogen effects on wound healing. *Int J Mol Sci* 2017;18:E2325.
75. Feng CJ, Lin CH, Tsai CH, Yang IC, Ma H. Adipose-derived stem cells-induced burn wound healing and regeneration of skin appendages in a novel skin island rat model. *J Chin Med Assoc* 2019;82:635–42.
76. Tsai HW, Wang PH, Tsui KH. Mesenchymal stem cell in wound healing and regeneration. *J Chin Med Assoc* 2018;81:223–4.
77. Chen GY, Chang CP, Wang PH. Burn wound and therapeutic challenge. *J Chin Med Assoc* 2019;82:748–9.
78. Chang WK, Tai YH, Lin SP, Wu HL, Tsou MY, Chang KY. An investigation of the relationships between postoperative pain trajectories and outcomes after surgery for colorectal cancer. *J Chin Med Assoc* 2019;82:865–71.
79. Wei J, Zhu HB, Wang F, Fan Y, Zhou HJ. Clinical utility of flexion-extension ratio measured by surface electromyography for patients with nonspecific chronic low-back pain. *J Chin Med Assoc* 2019;82:35–9.
80. Lai HC, Lu CH, Wong CS, Lin BF, Chan SM, Kuo CY, et al. Baicalein attenuates neuropathic pain and improves sciatic nerve function recovery in rats with partial sciatic nerve transection. *J Chin Med Assoc* 2018;81:955–63.
81. Lin PH. Sonographic findings of painful hemiplegic shoulder after stroke. *J Chin Med Assoc* 2018;81:657–61.
82. Yang CW, Fuh JL. Screening tools for neuropathic pain. *J Chin Med Assoc* 2018;81:1–3.
83. Wang PH, Horng HC, Chen YJ, Hsieh SL, Chao HT, Yuan CC. Effect of a selective nonsteroidal anti-inflammatory drug, celecoxib, on the reproductive function of female mice. *J Chin Med Assoc* 2007;70:245–8.
84. Lin WY, Cheng YT, Huang YH, Lin FS, Sun WZ, Yen CT. Synergistic symptom-specific effects of ketorolac-tramadol and ketorolac-pregabalin in a rat model of peripheral neuropathy. *J Chin Med Assoc* 2019;82:457–63.
85. Liou JY, Wang HY, Tsou MY, Chang WK, Kuo IT, Ting CK. Opioid and propofol pharmacodynamics modeling during brain mapping in awake craniotomy. *J Chin Med Assoc* 2019;82:390–5.
86. Tsou MY, Liou JY, Ting CK, Lin SP. Patient response prediction with logistic regression in gastrointestinal endoscopy under midazolam-alfentanil sedation performed as well as response surface model. *J Chin Med Assoc* 2018;81:1071–6.
87. Maged AM, Deeb WS, Elbaradie S, Elzayat AR, Metwally AA, Hamed M, et al. Comparison of local and intra venous dexamethasone on post operative pain and recovery after caesarean section. A randomized controlled trial. *Taiwan J Obstet Gynecol* 2018;57:346–50.
88. Lee WL, Huang BS, Chen YJ, Wang PH. Overcoming the barriers of osteoporosis treatment—a better route and a longer use. *J Chin Med Assoc* 2015;78:567–8.
89. Badawi MS. Histological study of the protective role of ginger on piroxicam-induced liver toxicity in mice. *J Chin Med Assoc* 2019;82:11–8.
90. Lin XH, Lin CC, Wang YJ, Luo JC, Young SH, Chen PH, et al. Risk factors of the peptic ulcer bleeding in aging uremia patients under regular hemodialysis. *J Chin Med Assoc* 2018;81:1027–32.
91. Kung CC, Chen SS, Yang HJ, Lai CJ, Chen LK. Pharmacogenetic study of pruritus induced by epidural morphine for post cesarean section analgesia. *Taiwan J Obstet Gynecol* 2018;57:89–94.
92. Yokota J, Kyotani S. Influence of nanoparticle size on the skin penetration, skin retention and anti-inflammatory activity of non-steroidal anti-inflammatory drugs. *J Chin Med Assoc* 2018;81:511–9.
93. Hu L, Pan J, Zhang S, Yu J, He K, Shu S, et al. Propofol in combination with remifentanyl for cesarean section: placental transfer and effect on mothers and newborns at different induction to delivery intervals. *Taiwan J Obstet Gynecol* 2017;56:521–6.
94. Mallappallil M, Sabu J, Friedman EA, Salifu M. What do we know about opioids and the kidney? *Int J Mol Sci* 2017;18:E223.

95. McDauid C, Maund E, Rice S, Wright K, Jenkins B, Woolcott N. Paracetamol and selective and non-selective non-steroidal anti-inflammatory drugs (NSAIDs) for the reduction of morphine-related side effects after major surgery: a systematic review. *Health Technol Assess* 2010;14:1–153, iii–iv.
96. Cheng MF, Hung SH, Su YP, Chiang CC, Chang MC, Chiu FY. Displaced isolated greater tuberosity fractures of elder adults treated with plate osteosynthesis. *J Chin Med Assoc* 2019;82:318–21.
97. Hung MJ, Tsai CP, Lin YH, Huang WC, Chen GD, Shen PS. Hyaluronic acid improves pain symptoms more than bladder storage symptoms in women with interstitial cystitis. *Taiwan J Obstet Gynecol* 2019;58:417–22.
98. Behmanesh E, Delavar MA, Kamalinejad M, Khafri S, Shirafkan H, Mozaffarpur SA. Effect of eryngo (eryngium caucasicum trautv) on primary dysmenorrhea: a randomized, double-blind, placebo-controlled study. *Taiwan J Obstet Gynecol* 2019;58:227–33.
99. Juho YC, Wu ST, Kao CC, Meng E, Cha TL, Yu DS. Anatomic mapping of the internal spermatic vein via subinguinal varicocelectomy with intraoperative vascular doppler ultrasound. *J Chin Med Assoc* 2019;82:115–9.
100. Karacin O, Mutlu I, Kose M, Celik F, Kanat-Pektas M, Yilmazer M. Serum vitamin D concentrations in young Turkish women with primary dysmenorrhea: a randomized controlled study. *Taiwan J Obstet Gynecol* 2018;57:58–63.
101. Taskin MI, Bilen C, Ergun A, Gencer N, Inceboz U. In vitro effects of estrogen and progesterone containing drugs on human erythrocyte carbonic anhydrase I and II isozymes in women smokers and non-smokers. *J Chin Med Assoc* 2015;78:513–9.
102. Lee WL, Huang BS, Wang PH. The role of red blood cells in cardiovascular disease. *J Chin Med Assoc* 2015;78:499–500.
103. Rieder E, Gay CV, Schraer H. Autoradiographic localization of carbonic anhydrase in the developing chorioallantoic membrane. *Anat Embryol (Berl)* 1980;159:17–31.
104. Riedel HH, Semm K. [The post-laparoscopic pain syndrome (author's syndrome)]. *Geburtshilfe Frauenheilkd* 1980;40:635–43.
105. Lipscomb GH, Summit RL Jr, McCord ML, Ling FW. The effect of nitrous oxide and carbon dioxide pneumoperitoneum on operative and postoperative pain during laparoscopic sterilization under local anesthesia. *J Am Assoc Gynecol Laparosc* 1994;2:57–60.
106. Wallace DH, Serpell MG, Baxter JN, O'Dwyer PJ. Randomized trial of different insufflation pressures for laparoscopic cholecystectomy. *Br J Surg* 1997;84:455–8.
107. Rammohan A, Manimaran AB, Manohar RR, Naidu RM. Nitrous oxide for pneumoperitoneum: no laughing matter this! A prospective single blind case controlled study. *Int J Surg* 2011;9:173–6.
108. Karapolat S, Gezer S, Yildirim U, Dumlu T, Karapolat B, Ozaydin I, et al. Prevention of pulmonary complications of pneumoperitoneum in rats. *J Cardiothorac Surg* 2011;6:14.
109. Tsereteli Z, Terry ML, Bowers SP, Spivak H, Archer SB, Galloway KD, et al. Prospective randomized clinical trial comparing nitrous oxide and carbon dioxide pneumoperitoneum for laparoscopic surgery. *J Am Coll Surg* 2002;195:173–9; discussion 179–80.
110. Sietses C, von Blomberg ME, Eijssbouts QA, Beelen RH, Berends FJ, Cuesta MA. The influence of CO₂ versus helium insufflation or the abdominal wall lifting technique on the systemic immune response. *Surg Endosc* 2002;16:525–8.
111. Neuhaus SJ, Watson DI, Ellis T, Lafullarde T, Jamieson GG, Russell WJ. Metabolic and immunologic consequences of laparoscopy with helium or carbon dioxide insufflation: a randomized clinical study. *ANZ J Surg* 2001;71:447–52.
112. Bongard FS, Pianim NA, Leighton TA, Dubecz S, Davis IP, Lippmann M, et al. Helium insufflation for laparoscopic operation. *Surg Gynecol Obstet* 1993;177:140–6.
113. Neudecker J, Sauerland S, Neugebauer E, Bergamaschi R, Bonjer HJ, Cuschieri A, et al. The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surg Endosc* 2002;16:1121–43.
114. Kyle EB, Maheux-Lacroix S, Boutin A, Lemyre M. Low vs standard pressures in gynecologic laparoscopy: a systematic review. *JLS* 2016;20:e2015.00113.
115. Kyle EB, Maheux-Lacroix S, Boutin A, Lemyre M. Complications of low compared to standard pneumoperitoneum pressures in laparoscopic surgery for benign gynecologic pathology: a systematic review protocol. *Syst Rev* 2015;4:96.
116. Guido RS, Brooks K, McKenzie R, Gruss J, Krohn MA. A randomized, prospective comparison of pain after gasless laparoscopy and traditional laparoscopy. *J Am Assoc Gynecol Laparosc* 1998;5:149–53.
117. Chaichian S, Moazzami B, Haghgoo A, Sheibani K. A new approach to an old concept for reducing shoulder pain caused by gynecological laparoscopy. *J Reprod Infertil* 2018;19:56–60.
118. Sasmour T, Kahokehr A, Hill AG. Meta-analysis of the effect of warm humidified insufflation on pain after laparoscopy. *Br J Surg* 2008;95:950–6.
119. Sajid MS, Mallick AS, Rimpel J, Bokari SA, Cheek E, Baig MK. Effect of heated and humidified carbon dioxide on patients after laparoscopic procedures: a meta-analysis. *Surg Laparosc Endosc Percutan Tech* 2008;18:539–46.
120. Herrmann A, De Wilde RL. Insufflation with humidified and heated carbon dioxide in short-term laparoscopy: a double-blinded randomized controlled trial. *Biomed Res Int* 2015;2015:412618.
121. Manwaring JM, Readman E, Maher PJ. The effect of heated humidified carbon dioxide on postoperative pain, core temperature, and recovery times in patients having laparoscopic surgery: a randomized controlled trial. *J Minim Invasive Gynecol* 2008;15:161–5.
122. Kissler S, Haas M, Strohmeier R, Schmitt H, Rody A, Kaufmann M, et al. Effect of humidified and heated CO₂ during gynecologic laparoscopic surgery on analgesic requirements and postoperative pain. *J Am Assoc Gynecol Laparosc* 2004;11:473–7.
123. Marks JL, Ata B, Tulandi T. Systematic review and metaanalysis of intraperitoneal instillation of local anesthetics for reduction of pain after gynecologic laparoscopy. *J Minim Invasive Gynecol* 2012;19:545–53.
124. Argo M, Favela J, Phung T, Huerta S. Local VS. other forms of anesthesia for open inguinal hernia repair: a meta-analysis of randomized controlled trials. *Am J Surg* 2019. doi: 10.1016/j.amjsurg.2019.06.024. [Epub ahead of print]
125. Roy KK, Subbaiah M, Naha M, Kumar S, Sharma JB, Jahagirdar N. Intraperitoneal bupivacaine for pain relief after minilaparoscopy in patients with infertility. *Arch Gynecol Obstet* 2014;289:337–40.
126. Malhotra N, Chanana C, Roy KK, Kumar S, Rewari V, Riwari V, et al. To compare the efficacy of two doses of intraperitoneal bupivacaine for pain relief after operative laparoscopy in gynecology. *Arch Gynecol Obstet* 2007;276:323–6.
127. Loughney AD, Sarma V, Ryall EA. Intraperitoneal bupivacaine for the relief of pain following day case laparoscopy. *Br J Obstet Gynaecol* 1994;101:449–51.
128. Chen CF, Tsai SW, Wu PK, Chen CM, Chen WM. Does continued aspirin mono-therapy lead to a higher bleeding risk after total knee arthroplasty? *J Chin Med Assoc* 2019;82:60–5.
129. Tsuei YS, Liao CH, Lee CH, Liang YJ, Chen WH, Yang SF. Intraoperative arterial perforation during neuroendovascular therapy: preliminary result of a dual-trained endovascular neurosurgeon in the neurosurgical hybrid operating room. *J Chin Med Assoc* 2018;81:31–6.
130. Cheng-Yen Lai J, Lai KJ, Yi-Yung Yu E, Hung ST, Chu CY, Wang KL. Sentinel lymphatic mapping among women with early-stage cervical cancer: a systematic review. *Taiwan J Obstet Gynecol* 2018;57:636–43.
131. Kim Y, Jeon GS, Choi SY, Kim MD, Lee SJ. Evaluation of sclerotherapy for the treatment of infected postoperative lymphocele. *Taiwan J Obstet Gynecol* 2017;56:477–81.
132. Durai R, Ng PC. Surgical vacuum drains: types, uses, and complications. *Aorn J* 2010;91:266–71; quiz 272–4.
133. Asgari Z, Hosseini R, Rastad H, Hosseini L. Does peritoneal suction drainage reduce pain after gynecologic laparoscopy? *Surg Laparosc Endosc Percutan Tech* 2018;28:73–6.
134. Haghgoo A, Chaichian S, Ghahremani M, Nooriardabili S, Akbaian A, Moazzami B. The use of peritoneal suction drainage to reduce shoulder pain caused by gynecological laparoscopy. *Arch Iran Med* 2016;19:173–8.
135. Craciunas L, Stirbu L, Tsampras N. The use of a peritoneal gas drain following gynecological laparoscopy: a systematic review. *Eur J Obstet Gynecol Reprod Biol* 2014;179:224–8.
136. Shen CC, Wu MP, Lu CH, Kung FT, Huang FJ, Huang EY, et al. Effects of closed suction drainage in reducing pain after laparoscopic-assisted vaginal hysterectomy. *J Am Assoc Gynecol Laparosc* 2003;10:210–4.
137. Abbott J, Hawe J, Srivastava P, Hunter D, Garry R. Intraperitoneal gas drain to reduce pain after laparoscopy: randomized masked trial. *Obstet Gynecol* 2001;98:97–100.