

POSTER PRESENTATION

Open Access

Extracting activity of individual cell populations from multielectrode recordings


Jan Potworowski, Helena Głąbska, Szymon Łęski, Daniel K Wójcik*

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

The low-frequency part of the extracellular electric signals, the local field potentials (LFP), carries information about dendritic processing in neuronal populations. However, the long-range nature of electric field makes the analysis of LFP difficult, as typically an electrode records activity of many sources. Modern multielectrodes allow for increased spatial resolution, hence also the need for effective data analysis methods which would allow to get more insight. These methods include Current Source Density (CSD) analysis and source separation methods.

In [1] we have combined inverse CSD method [2] with Independent Component Analysis (ICA) [3] to decompose activity recorded in the rat forebrain on a grid of 140 positions obtaining physiologically plausible components across a group of seven animals. The question

remains how the obtained components are connected to the activity of neuronal populations. To study this problem we enriched the thalamocortical model [4,5] by adding the spatial information (Figure 1A) and used it to simulate the LFP generated by a single cortical column. We used the kernel CSD method and spatio-temporal ICA to decompose the LFP measured on a regular grid. We compared the resulting components to the activity of the twelve cortical populations included in the model. We found that the recorded evoked activity was dominated by two populations of pyramidal neurons, which were well separated by ICA (Figure 1B-1G). The activity from other populations was hardly visible on top of the main two dipoles and we were also not able to obtain them through ICA.



* Correspondence: d.wojcik@nencki.gov.pl
Laboratory of Neuroinformatics, Dept. of Neurophysiology, Nencki Institute,
02-093, Poland
Full list of author information is available at the end of the article

Acknowledgements

Supported by grants PBZ/MNiSW/07/2006/11, 5428/B/P01/2010/39 from the Polish Ministry of Science and Higher Education and by a grant from the Polish Ministry of Regional Development, POIG.02.03.00-00-003/09.

Published: 18 July 2011

References

1. Łęski S, Kublik E, Świejkowski DA, Wróbel A, Wójcik DK: **Extracting functional components of neural dynamics with ICA and iCSD.** *J Comput Neurosci* 2010, **29**:459-473.
2. Łęski S, Wójcik DK, Tereszczuk J, Świejkowski DA, Kublik E, Wróbel A: **Inverse Current-Source Density in three dimensions.** *Neuroinformatics* 2007, **5**:207-222.
3. Stone JV, Porrill J, Porter NR, Wilkinson IW: **Spatiotemporal Independent Component Analysis of Event-Related fMRI Data Using Skewed Probability Density Functions.** *NeuroImage* 2002, **15**:407-421.
4. Traub RD, Contreras D, Cunningham MO, Murray H, LeBeau FEN, Roopun A, Bibbig A, Wilent WB, Higley MJ, Whittington MA: **Single-Column Thalamocortical Network Model Exhibiting Gamma Oscillations, Sleep Spindles, and Epileptogenic Bursts.** *J Neurophysiol* 2005, **93**:2194-2232.
5. [<http://senselab.med.yale.edu/ModelDb/ShowModel.asp?model=82894>].

doi:10.1186/1471-2202-12-S1-P374

Cite this article as: Potworowski *et al.*: Extracting activity of individual cell populations from multielectrode recordings. *BMC Neuroscience* 2011 **12**(Suppl 1):P374.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

