



POSTER PRESENTATION

Open Access

# Cartilage-protective effects of C-type natriuretic peptide over expression in K/BxN TCR arthritis model

Hulya Bukulmez<sup>1\*</sup>, Cynthia F Bartels<sup>1</sup>, Kabita Nanda<sup>3</sup>, Tariq M Haqqi<sup>2</sup>, Jean F Welter<sup>1</sup>

From 2011 Pediatric Rheumatology Symposium sponsored by the American College of Rheumatology  
Miami, FL, USA. 2-5 June 2011

## Purpose

The c-type natriuretic peptide (CNP) signaling pathway is known as a major contributor to skeletal growth in children. CNP is produced and secreted by both growth plate and joint chondrocytes, and has a paracrine regulatory effect on cartilage tissue. CNP increases matrix production by chondrocytes and promotes their proliferation. In this study, we investigated whether over-expression of CNP in chondrocytes would be protective of joint cartilage degradation during chronic inflammatory arthritis *in vivo*.

## Methods

We first developed transgenic mice that over-express CNP ( $CNP^{col2a1TG}$ ) in chondrocytes under the control of the collagen 2a1 promoter and enhancer. Then, we obtained K/BxN TCR transgenic mice from a collaborator and analyzed both transgenic mice for their joint cartilage clinical and histologic findings over time. We crossed  $CNP^{col2a1TG}$  mouse with K/BxN TCR transgenic mouse to produce mice with both K/BxN TCR and  $CNP^{col2a1TG}$  backgrounds. The degree of arthritis and cartilage damage in the offspring was analyzed using a clinical scoring system and two histological scoring systems. Differences between the scores were analyzed using the Student's t-test.

## Results

Mice that carried the transgene for both  $CNP^{col2a1TG}$  and K/BxN TCR showed less severe clinical and histologic arthritis findings in the joint cartilage compared to wild type littermates. Between the ages of 6-14 weeks, the average arthritis score of K/BxN TCR transgenic

mice that over-expressed CNP was  $4.37 \pm 1.38$  (n=8), while the average arthritis score of K/BxN TCR arthritic mice of the same age was  $8.66 \pm 3.26$  (n=14), (p<0.05). Histological staining and morphometry did not show any evidence of cartilage degradation in the joint cartilage of  $CNP^{col2a1TG}$  mice. The knee and ankle cartilage of  $CNP^{col2a1TG}$  mice was thick and showed increased proteoglycan content by Safranin-O staining. However, the double-transgenic offspring mice (K/BxN/ $CNP^{col2a1TG}$ ) developed less cartilage damage and less chondrocyte disorganization while still developing inflammatory changes (pannus) in the synovium, similar to the K/BxN TCR mice. We adapted the ICRS histological scoring system and gave scores to the knee joint cartilage of the 8-week-old male mice. K/BxN mice (n=7) scored significantly lower for both chondrocytic cell distribution (III) and chondrocyte matrix content (II) (p<0.001 and p<0.05, respectively) than the (K/BxN/ $CNP^{col2a1TG}$  mice (n=12).

## Conclusion

K/BxN TCR arthritic mice over-expressing CNP did not have joint cartilage damage due to chronic inflammation. We conclude that excess paracrine production of CNP in the joint cartilage of double-transgenic K/BxN/ $CNP^{col2a1TG}$  arthritic mice was able to overcome the effects of pro-inflammatory cytokines on joint cartilage *in vivo*. CNP and the effector molecules of CNP signaling pathway may have therapeutic potential in protecting cartilage homeostasis during chronic inflammatory arthritis.

## Disclosure

Hulya Bukulmez: None; Cynthia F. Bartels: None; Kabita Nanda: None; Tariq M. Haqqi: None; Jean F. Welter: None.

<sup>1</sup>Skeletal Research Center and MetroHealth Medical Center, Case Western Reserve University, Cleveland, OH, USA

Full list of author information is available at the end of the article

**Author details**

<sup>1</sup>Case Western Reserve University, Cleveland, OH, USA. <sup>2</sup>MetroHealth Medical Center, Case Western Reserve University, Cleveland, OH, USA. <sup>3</sup>Rainbow Babies and Children, Cleveland, OH, USA. <sup>4</sup>Skeletal Research Center and MetroHealth Medical Center, Case Western Reserve University, Cleveland, OH, USA.

Published: 13 July 2012

doi:10.1186/1546-0096-10-S1-A109

**Cite this article as:** Bukulmez et al.: Cartilage-protective effects of C-type natriuretic peptide over expression in K/BxN TCR arthritis model. *Pediatric Rheumatology* 2012 **10**(Suppl 1):A109.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

