

Imidazole as a Promising Medicinal Scaffold: Current Status and Future Direction

Sahar S Alghamdi ^{1,2}

Rasha S Suliman ^{1,2}

Khlood Almutairi¹

Khawla Kahtani¹

Dimah Aljatli¹

¹College of Pharmacy, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Kingdom of Saudi Arabia;

²Medical Research Core Facility and Platforms, King Abdullah International Medical Research Center (KAIMRC), Ministry of National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia

Abstract: Various imidazole-containing compounds have been tested for their medical usefulness in clinical trials for several disease conditions. The rapid expansion of imidazole-based medicinal chemistry suggests the promising and potential therapeutic values of imidazole-derived compounds for treating incurable diseases. Imidazole core scaffold contains three carbon atoms, and two nitrogen with electronic-rich characteristics that are responsible for readily binding with a variety of enzymes, proteins, and receptors compared to the other heterocyclic rings. Herein, we provide a thorough overview of the current research status of imidazole-based compounds with a wide variety of biological activities including anti-cancer, anti-microbial, anti-inflammatory and their potential mechanisms including topoisomerase IIR catalytic inhibition, focal adhesion kinase (FAK) inhibition, c-MYC G-quadruplex DNA stabilization, and aurora kinase inhibition. Additionally, a great interest was reported in the discovery of novel imidazole compounds with anti-microbial properties that break DNA double-strand helix and inhibit protein kinase. Moreover, anti-inflammatory mechanisms of imidazole derivatives include inhibition of COX-2 enzyme, inhibit neutrophils degranulation, and generation of reactive oxygen species. This systemic review helps to design and discover more potent and efficacious imidazole compounds based on the reported derivatives, their ADME profiles, and bioavailability scores that together aid to advance this class of compounds.

Keywords: imidazole, apoptosis, kinase inhibitors, tubulin polymerization, topoisomerase II

Introduction

In the last century, nitrogen-based heterocycles, particularly the imidazole ring which was discovered in the 1840s, attracted many researchers as this core scaffold demonstrated promising anti-cancer, anti-microbial, and anti-inflammatory activities.¹ The structural features of the imidazole ring enhance their ability to form multiple drug–ligand interactions via hydrogen bonds, van der Waals, and hydrophobic forces.² Moreover, the imidazole core scaffold is a part of several naturally derived compounds, such as histamine, histidine, biotin, alkaloids, and nucleic acid, and a part of multiple classes of FDA-approved drugs. Because of their important properties as therapeutics, fused imidazole derivatives have held a prominent role in the medical field. Methotrexate, Metronidazole, and Omeprazole are the current imidazole derivatives available on market as anti-cancer, anti-microbial, and anti-inflammatory agents, respectively. Nevertheless, these drugs have shown painful and unacceptable side effects that caused treatment failure.³ To discover potent compounds with better safety profiles, Imidazole-based compounds are under intensive scientific exploration as these derivatives have

Correspondence: Rasha S Suliman
Tel +966 114299999 Ext 99570
Email sulimanr@ksau-hs.edu.sa

Received: 18 February 2021
Accepted: 22 April 2021
Published: 29 July 2021



achieved amazing progression in medicinal chemistry research. The literature summarized in this review aids in the design and development of novel and highly potent compounds with less harmful side effects for the treatment of lethal human diseases.

Anti-Cancer Properties of Imidazole Derivatives

Cancer is the second major cause of death after cardiac disease nationally and internationally. The annual number of new cases is estimated to be 439.2 per 100,000 people.⁴ At the current global rate, the estimated number of cancer is rapidly increasing and predicted to reach approximately 27.5 million new cancer cases per year by 2040.⁵ Until today, the detailed mechanism of cancer is not fully understood and with a simple definition, it is considered to be a genetic disease that starts with abnormal and continuous cell growth, with the chance of metastasis.⁶ The current chemotherapeutic agents available such as Methotrexate, Etoposide, and Paclitaxel demonstrated potent activity and to some extent, good survival rates, however, several challenges need to be overcome, such as drug resistance, toxicity, and intolerable side effects.⁷ Thus, a new rationale for anticancer agents should be investigated to discover new targeted therapy with lower systemic side effects.⁸ According to some reported studies, imidazole has the potential to overcome the unresolved disadvantages of currently available clinical drugs and could be utilized as a chemical scaffold for novel anticancer agents with several potential mechanisms of action. Therefore, in the following sections, we will be discussing the multiple mechanisms of several imidazole derivatives that possess anti-cancer activity.

In this regard, Baviskar et al have investigated five imidazole derivatives that demonstrated potent catalytic inhibition of DNA and authors believed that these compounds were highly specific to the Topoisomerase IIR with improved potency compared to Etoposide and 5-fluorouracil in kidney cancer cells.⁹ An MTT assay was performed to examine the cytotoxic activity in both kidney cancer cells (HEK 293) and Vero cells. After 48 hours of treatment, the cell viability of HEK 293 decreased significantly compared to Vero cells. As a result, compound C1 demonstrated the highest potency with LC₅₀ of 25 μ M in HEK 293 cells and 62 μ M in Vero cells.

Additionally, to seek more powerful and lower toxic anticancer agents, six imidazole compounds were synthesized by Zhao et al and tested in vitro. Among them, compound C2

showed higher activity against breast cancer cells (MCF-7) with IC₅₀ of 0.75 μ M compared to Doxorubicin. Compound C2 had potent anticancer activity, with IC₅₀ of 4.37 μ M, against lung cancer cells (A549) and authors concluded that these derivatives mediated anti-cancer activity via inducing apoptosis and suppressing cancer cell proliferation.¹⁰ Moreover, using in vitro experiments, Dao et al evaluated twenty-six synthesized imidazole compounds, and study results revealed that compound C3 showed the greatest inhibition with IC₅₀ values of 50 μ M against cancer cells expressing high levels of focal adhesion kinase (FAK), including Brain (U87-MG), Colon (HCT-116), Breast (MDA-MB)-231, and Prostate (PC-3) cancer cell lines.¹²

Roopashree et al synthesized new series of imidazole derivatives to investigate the antiproliferative activity on Hela cells and with Sorafenib as a reference drug. The results showed that compound C4 was a very potent inhibitor with an IC₅₀ of 25.3 μ M. Using an in-vivo model, C4 was significantly able to decrease angiogenic properties against Ehrlich ascites tumor (EAT) bearing mice.¹⁴ In 2011, Alkahtani et al synthesized several imidazole derivatives to identify the anti-proliferative activity on the colon (HCT-116) and breast (MCF-7) cancer cells. It was concluded that compound C5 has anti-proliferative activity by inhibiting cyclin-dependent kinase 6 (CDK6) and inducing apoptosis on myeloid cell leukemia 1 (Mcl-1) protein.¹⁵

In another study, Chung et al found that only eleven compounds among the twenty-six imidazole derivatives showed good activity against human umbilical vein endothelial cells (HUVECs) and smooth muscle cells (SMCs), using mycophenolic acid as a standard control. The results suggested that only three compounds, including compound C6, activated the p38 signaling pathway, and were selective inhibitors of endothelial cell proliferation specifically for human umbilical vein endothelial cells (HUVECs).¹⁶ Using in vitro experiments, Xue et al investigated twenty compounds for their cytotoxicity, compound C7 exhibited equal or more potent cytotoxic activity compared to docetaxel in a dose-dependent manner. Additionally, C7 arrested the cell cycle on the S phase at 0.01 μ M, and induced cell apoptosis at 0.1–1.0 μ M. Moreover, an additional amino group at the C-3 position on the B ring was beneficial for potency against human myeloid leukemia (HL-60) and human myeloid leukemia (K562) cells. Using in vivo model, compound C7 showed significant anticancer activity on murine S-180 which was similar to cyclophosphamide (CTX), and higher inhibition on murine H-22 tumor-bearing models was observed.¹⁷

Another study reported that compound C8 bearing imidazole ring demonstrated the highest anti-proliferative activity with an average IC_{50} value of 7.219 μM against four cancer cells, including MCF-7, H1299, HeLa, and B16-F10. The result suggested that C8 has a dose-dependent effect in inhibiting tubulin polymerization and suppressing the cell cycle in the G2/M phase.²¹

On investigating the cytotoxicity of imidazole compounds against cervical cancer, it was reported that C9 possesses a potent anticancer activity with IC_{50} of 0.08 μM , which was comparable to 5-fluorouracil with IC_{50} 0.09 μM . The potential mechanism of C9 was via suppressing the cell cycle at the G1 phase and inducing apoptosis. Furthermore, compound C9 exhibited a remarkable inhibition effect on the carbonic anhydrase IX enzyme with an IC_{50} of 0.12 μM comparable to Dorzolamide HCL with IC_{50} of 0.09 μM , as standard drug.²² Meenakshisundaram et al draw our attention to the role of imidazole derivatives on estrogen receptors by screening 30 compounds, remarkably compound C10 was the most potent derivative with higher anticancer activity against breast cancer cells relative to Adriamycin, with GI_{50} 0.3 and GI_{50} 0.51 μM , respectively. The addition of the nitrogen group in the imidazole ring was beneficial for improving cytotoxic activity. Using in vivo model, C10 demonstrated similar anticancer activity comparable to Tamoxifen. The authors concluded that C10 could bind to the estrogen receptors (ERs) as consequence the receptors dimerized in the breast cancer cells that resulted in cytotoxicity and apoptosis.¹⁸

Recently, Wu et al synthesized and evaluated six derivatives that demonstrated superior anticancer activity against a variety of tumor cells, specifically nasopharyngeal carcinoma (CNE-1) cells. Compound C11 yielded an IC_{50} of 1.1 μM with more potent antitumor activity through stabilizing DNA c-MYC G-quadruplex relative to Doxorubicin and with less harmful side effects against the human normal cell line.¹⁹ In another attempt, Fan et al synthesized and experimentally screened twenty imidazole derivatives. Among them, compound C12 had the highest anti-proliferative activity against breast cancer (MDA-MB-231), prostate cancer (PC3), and neuroblastoma (SH-SY5Y) cells, with an IC_{50} value of 0.38, 1.09, and 0.77 μM , respectively, through inhibition of aurora kinase that resulted in cell apoptosis.²⁰

According to a recent report conducted by Ali et al, the synthesized imidazole derivatives were investigated for potential anti-cancer activity on human cancer cells using Dabrafenib as a reference agent. The results showed that

between nineteen compounds, compound C13 has the greatest activity against NCI60 human cancer cell line and melanoma with mean IC_{50} of 2.4 and 1.8 μM , respectively. Compound C13 possesses a potential anti-cancer activity against both NCI 60 and melanoma human cancer cell lines via BRAF^{V600E} kinase inhibition.²³







In a work reported in 2020, anti-cancer activity of a newly synthesized novel imidazole derivatives was investigated against multiple cancer cell lines including human colon carcinoma (Caco-2 and HCT-116), human cervical carcinoma (HeLa), and human breast adenocarcinoma (MCF-7). Compound C14 demonstrated the highest cytotoxic activity against all cancer cell lines specifically against MCF-7 with IC_{50} 0.38 μM . Moreover, most of the synthesized compounds were non-carcinogenic. Overall, compound C14 showed the best cytotoxic activity through the inhibitory effect of glycogen synthase kinase 3 beta (GSK-3 β) and could be a potential anti-cancer agent with less toxicity.²⁴

On the other hand, several studies had promising results of imidazole compounds as anticancer agents but unfortunately, their mechanisms of action were not determined. As seen in Chen et al study which investigated seventeen imidazole derivatives with potential anti-proliferative activity in two cell lines. Among them, compound C15 demonstrated IC_{50} of 16.1 and 31.6 μM in human (A375) and mouse (B16) melanoma cells, respectively.¹¹ Moreover, Sarkarzadeh et al have shown that among forty compounds, C16 demonstrated promising anti-proliferative activities compared to Cisplatin which emphasizes the importance of imidazole derivatives particularly on the Jurkat cell line.¹³ Besides, a study carried out by Zhang et al evaluated the anticancer activity of synthesized pyrrole-imidazole series against human pancreatic cancer cell lines (PANC and ASPC-1) and prostate cancer cells. The results showed that compound C17 has the highest anti-cancer activity with more selectivity towards PANC and ASPC-1 relative to prostate cancer cells with IC_{50} of 0.063 and 0.062 μM , respectively.²⁵ All the discussed Imidazole derivatives with anti-cancer activities are summarized in Table 1.

Anti-Microbial Properties of Imidazole Derivatives




For several years, humans have been continuously exposed to different pathogens on daily basis and such invasive microbial infections are considered major problems particularly in

Table 1 The Anti-Cancer Activities of Imidazole Derivatives Using Several Cancer Cell Lines

Chemical Structure	Cell Line	Activity	Mechanism	Reference
 <p>C1</p>	Kidney cancer cells (HEK 293) Breast cancer cells (MCF7)	LC₅₀ (μM) 15 at both cell lines	Topoisomerase II α catalytic inhibitors. The apoptotic effect in the G1/S phase.	[9]
 <p>C2</p>	Cervix cancer cells (HeLa) Breast cancer cells (MCF-7) Lung cancer cells (A549) Liver cancer cells (HepG2)	IC₅₀ (μM) 2.88 ± 0.15 0.75 ± 0.07 4.37 ± 0.07 2.91 ± 0.11	Induce apoptosis. Bind with DNA through intercalation.	[10]
 <p>C3</p>	Brain (U87-MG) Colon (HCT-116) Breast (MDA-MB-231) Prostate (PC-3)	IC₅₀ (μM) 0.37±0.04 0.31±0.18 1.08±0.45 1.27±0.28	Focal Adhesion Kinase (FAK) inhibition. Arrest cells in the G2/M phase of the cell cycle. Retard cell growth	[12]
 <p>C4</p>	Cervical (HeLa)	IC₅₀ (μM) 25.3	Decreases angiogenesis in the peritoneum of EAT bearing mice.	[14]
 <p>C5</p>	Colon (HCT-116) Breast (MCF-7)	GI₅₀ (μM) 3.0 40	Cyclin-dependent kinase 6 (CDK6) inhibition and induce cancer cell apoptosis.	[15]
 <p>C6</p>	Human umbilical vein endothelial cells (HUVECs) Smooth muscle cells (SMCs)	IC₅₀ (μM) 0.4 5.5	Selective inhibitor for the HUVECs. Activation of p38 signaling pathway.	[16]





(Continued)

Table I (Continued).

Chemical Structure	Cell Line	Activity	Mechanism	Reference
 <p style="text-align: center;">C7</p>	Human myeloid leukemia cells (HL-60) Human myeloid leukemia cells (K562) Human myeloid leukemia cells (K562R) (multidrug-resistant cells) Human prostate carcinoma cells (PC-3) Human breast carcinoma cells (MCF-7) Human esophageal carcinoma cells (ECA-109) Human hepatocarcinoma cells (BEL-7402) Human non-small lung cancer cells (A549)	IC₅₀ (μM) 0.2 1 0.9 3.1 10.6 3.8 1.2 1.3	Exhibited excellent inhibitory activity on tumor growth in vivo	[17]
 <p style="text-align: center;">C8</p>	MCF-7 H1299 HeLa B16-F10 HUVEC	IC₅₀ (μM) 9.450±0.292 7.652±0.215 6.862±0.144 4.912±0.088 >50	Inhibit tubulin polymerization and arrest cell cycle at G2/M phase	[21]
 <p style="text-align: center;">C9</p>	HeLa	IC₅₀ (μM) 82.83±1.37	Arrest cell cycle at G1 phase and induce apoptosis	[22]





(Continued)

Table I (Continued).

Chemical Structure	Cell Line	Activity	Mechanism	Reference
 <p style="text-align: center;">C10</p>	Cervical (HeLa) Breast (MDA-MB-231) Renal cancer (ACHN)	GI₅₀ (μM) 0.36 0.30 0.38	ND Possibly due to dimerized estrogen receptor	[18]
 <p style="text-align: center;">C11</p>	Nasopharyngeal carcinoma (CNE-1)	IC₅₀ (μM) 1.1 ± 0.1	c-MYC G-quadruplex DNA stabilizers	[19]
 <p style="text-align: center;">C12</p>	breast cancer (MDA-MB-231) prostate cancer (PC3) neuroblastoma (SH-SY5Y)	IC₅₀ (μM) 0.38±0.08 1.09 ±0.24 0.77±0.12	Aurora kinase inhibition	[20]
 <p style="text-align: center;">C13</p>	NCI 60 LOX IMVI MALME-3M M14 MDA-MB-435 SK-MEL-2 SK-MEL-28 SK-MEL-5 UACC-257 UACC-62	IC₅₀ (μM) 2.4 1.75 1.69 1.76 1.97 2.21 2.13 1.60 2.08 1.85	Inhibit BRAF ^{V600E} kinase	[23]

(Continued)

Table I (Continued).

Chemical Structure	Cell Line	Activity	Mechanism	Reference
 <p style="text-align: center;">C14</p>	Caco-2 HCT-116 HeLa MCF-7	IC₅₀ (μM) 4.67 ± 0.11 16.78 ± 0.59 6.87 ± 0.32 0.38 ± 0.04	Inhibit GSK-3β	[24]
 <p style="text-align: center;">C15</p>	Melanoma (human A375) Melanoma (mouse B16)	IC₅₀ (μM) 16.1±0.5 31.6±1.1	Anti-proliferative	[11]
 <p style="text-align: center;">C16</p>	Cervical (HeLa) Colon (LS180) Breast (MCF-7) Jurkat cells	IC₅₀ (μM) >25 for all the cell lines	ND	[13]
 <p style="text-align: center;">C17</p>	A549 PC3 PANC ASPC-1	IC₅₀ (μM) 0.93 ± 0.34 0.83 ± 0.27 0.063 ± 0.09 0.062 ± 0.013	ND	[25]

Note: Values are means of three experiments.

Abbreviation: LC₅₀, The concentration required to kill half the cells in the cell culture; IC₅₀, Compound concentration required to inhibit tumor cell proliferation by 50%; GI₅₀, The concentration of drug causing 50% inhibition of cell growth; ND, Not determined.

immunocompromised people.²⁶ Infectious diseases were increasingly recognized as a serious worldwide public health concern and for this reason, tracking and reporting on antibiotic resistance in a recorded number of nations, making

a significant step forward in the global battle against drug resistance and the evidence showing that the medications that were used for treatment are increasingly resistant to a various number of bacterial infections.^{27,28}

The wide use of antimicrobial agents in the treatment of common infectious diseases such as urinary tract infections and abdominal tract infections suggests that the world is running out of effective strategies to deal with these infectious diseases. For example, in 33 countries, the resistance rate of ciprofloxacin ranged from 8.4% to 92.9%. Currently, in 2021 the world health organization is worried that the excessive use of antibiotics during the COVID-19 pandemic will further intensify the rise in antimicrobial resistance.²⁸ In this century, the growth of antimicrobial resistance expanded, and new antimicrobial agents that are selective, potent, and less harmful compared to current drugs will be needed to eliminate the life-threatening invasive infections in clinical care.²⁹ One of the key aspects to discover new agents is to determine the Structure–Activity Relationships (SAR) of a chemical scaffold, followed by structural modifications to optimize some of the biological activities at a specific target. Herein, we summarized the most potent imidazole derivatives as potential antimicrobial agents based on their SAR and essential structural features.

As reported by Ansari et al, the authors synthesized twenty-two imidazole derivatives to investigate and determine the antibacterial activity. All the synthesized compounds displayed good activity against Gram-positive but low activity against Gram-negative bacteria with the MIC ranging from 4 to 64 μM compared to Ampicillin and Ciprofloxacin as standard drugs. Among the twenty-two compound, M18 illustrated remarkable antibacterial activity with MIC 2 μM against *Staphylococcus aureus* as shown in Table 2. Besides, some of the synthesized compounds demonstrated good to moderate antifungal activity. However, compound M18 was found to be equipotent as Amphotericin B against *Candida Albicans* and *A. Flavus*. The author concluded that as the number of carbon atoms in the side chain at the 2-position of the oxadiazole heterocyclic ring increases, the antibacterial activity against *Candida Albicans* increases.³⁰

An additional study was conducted in 2009, screened twenty compounds against two Gram-positive, two Gram-negative bacteria, and five fungus strains. The antibacterial screening demonstrated that compound M19 showed good inhibitory activity against *Escherichia coli*, *S. typhimurium*, *B. subtilis*, and *Staphylococcus aureus* with a diameter of zone of inhibition 19, 17, 20, and 21 mm, respectively, compared to Chloramphenicol as a standard drug. Furthermore, it seemed that the antibacterial and antifungal activity both depend on the heterocyclic moiety, not only that but also, the activity was enhanced by the presence of nitro and chlorine substituents on the benzimidazole moiety. Besides, the twenty compounds were also tested for antifungal activity

in another series of experiments. Moreover, compound M20 showed good antifungal activity against *C. Albicans*, *H. oryzae*, *A. Niger*, *T. viridae*, and *Penicillium* species with zone inhibition diameter of 18, 15, 15, 10, and 19 mm, respectively. Nystatin was used as the standard drug. Unfortunately, the diameter of the inhibition zone against *T. viridae* was considerably lower than the other fungal strain in this study.³¹

In this regard, Pieczonka et al reported in vitro activity of 3-oxido-1H-imidazole-4-carbohydrazides against 4 Gram-positive strains, 4 Gram-negative strains, and *Candida albicans* as fungal species. The authors used vancomycin and oxacillin as reference agents for Gram-positive, and chloramphenicol for *Escherichia coli*. The results showed potent antibacterial activity of compound M21 against some strains, indicate that the 5-nitrofuryl group was essential for antibacterial activity. Eventually, compound M21 had antimicrobial properties due to nitro-substituted furan ring which can be considered as an essential moiety in antibacterial activity.³²






Knowing the mechanism of action of new potential anti-microbial agents can provide more information regarding the safety of compounds by understanding their effect on the human body however, two studies reported the antibacterial activity of imidazole derivatives without investigating the mechanism of action.

A good example of this is the newly synthesized imidazole derivatives benzimidazole-1,2,3-triazole-sulfonamide hybrids that were tested for antimicrobial activity. In this study, four bacterial strains *Bacillus cereus*, *Staphylococcus aureus*, *E. coli*, and *P. aeruginosa* were utilized with ciprofloxacin as a reference agent. The result demonstrated that compound M22 was the most potent antimicrobial derivative against the mentioned strains with MIC range from 32 to 64 μM .³³ A recent report was done by Daraji et al, described the antibacterial efficacy of imidazole derivatives against *E. coli*, *P. aeruginosa*, *S. aureus*, and *S. pyogenes*. Among all the compounds, compound M23 was found to be active against all the strains with broad-spectrum antibacterial activity against the Extended-spectrum beta-lactamase (ESBL), Vancomycin-resistant Enterococcus (VRE), and Methicillin-resistant *Staphylococcus aureus* (MRSA) strains.³⁴ All the discussed Imidazole derivatives with antimicrobial properties are summarized in Table 2.

The Anti-Fungal Properties of Imidazole Derivatives






Imidazole derivatives possessed several pharmacological activities, one of these is antifungal property which was also

Table 2 The Anti-Microbial Activities of Imidazole Derivatives Using Several Microorganisms

Chemical Structure	Microorganism	Activity	Mechanism	Reference
 <p style="text-align: center;">M18</p>	Bacteria <i>S. aureus</i> <i>B. subtilis</i> <i>S. mutans</i> <i>E. coli</i> <i>P. aeruginosa</i> <i>S. typhi</i> Fungi <i>C. Albicans</i> <i>A. niger</i> <i>A. flavus</i>	MIC (μM) 2 4 4 >128 NT NT Zone diameter of inhibition (mm) 22–28 10–15 22–28	Lipophilic cross membrane in the biological system	[30]
 <p style="text-align: center;">M19</p>	Bacteria <i>S. aureus</i> <i>B. subtilis</i> <i>S. typhimurium</i> <i>E. coli</i>	Zone diameter of inhibition (mm) 21 20 17 19	ND	[31]
 <p style="text-align: center;">M20</p>	Fungi <i>C. albicans</i> <i>H. oryzae</i> <i>A. niger</i> <i>T. viridae</i> <i>Penicillium sp.</i>	Zone diameter of inhibition (mm) 18 15 15 10 19	ND	[31]
 <p style="text-align: center;">M21</p>	Bacteria <i>E. coli</i> <i>S. aureus</i> <i>S. aureus</i> <i>P. vulgaris</i> <i>P. mirabilis</i> <i>E. faecalis</i> <i>S. epidermidis</i>	MIC (μM) 11 11 11 >600 >600 >600 4	Double strand breaks in DNA helix	[32]
 <p style="text-align: center;">M22</p>	Bacteria <i>Bacillus cereus</i> <i>S. aureus</i> <i>E. coli</i> <i>P. aeruginosa</i>	MIC (μM) 32 32 64 64	ND	[33]



(Continued)

Table 2 (Continued).

Chemical Structure	Microorganism	Activity	Mechanism	Reference
 <p>M23</p>	Bacteria <i>E. coli</i> <i>P. aeruginosa</i> <i>S. aureus</i> <i>S. pyogenes</i> ESBL VRE MRSA	MIC (μM) 12.5 12.5 25 50 62.5 100 100	ND	[34]
 <p>M24</p>	Fungi <i>Candida albicans</i> <i>Rhizopus oryza</i> <i>Penicillium citrum</i> <i>Aspergillus niger</i>	MIC (μM) 12.5 for all fungi	ND	[35]
 <p>M25</p>	Fungi <i>Candida albicans</i> <i>Rhizopus oryza</i> <i>Penicillium citrum</i> <i>Aspergillus niger</i>	MIC (μM) 12.5 for all fungi	ND	[35]
 <p>M26</p>	Protozoal <i>T. b. rhodesiense</i> <i>T. cruzi</i> <i>L. donovani</i> <i>P. falciparum</i>	IC₅₀ (μM) 34.43 0.203 15.01 0.059	Target-based an assay using recombinant <i>T. cruzi</i> CYP51.	[36]
 <p>M27</p>	Protozoal <i>L. donovani</i> <i>promastigotes</i>	IC₅₀ (μM) 10	Potential kinase inhibitor	[37]

(Continued)

Table 2 (Continued).

Chemical Structure	Microorganism	Activity	Mechanism	Reference
 <p style="text-align: center;">M28</p>	Protozoal <i>T. cruzi</i> <i>T. vaginalis</i>	Percentage of inhibition 61.0 87.5	ND	[38]
 <p style="text-align: center;">M29</p>	Protozoal <i>T. gondii</i>	EC₅₀ (μM) 0.003	ND	[39]

Note: The prefix M letter in the compound's number referred to Anti-Microbial activity.

Abbreviations: MIC, Minimal inhibitory concentration; IC₅₀/EC₅₀, Compound concentration required to decrease parasite viability by 50%; ESBL, Extended spectrum beta-lactamases; VRE, Vancomycin-resistant enterococci; MRSA, Methicillin-resistance staphylococcus aureus; NT, Not tested; ND, Not determined.

reported in 2012 by Hussain et al, that investigated the activity of imidazole derivatives towards *C. Albicans*, *R. Oryza*, *P. Citrum*, and *A. Niger* fungal strains using Ketoconazole as a control. The minimum inhibitory concentration (MIC) was tested for M24 and M25 which was found to be 12.5 μM for these two compounds among the other twenty compounds. The authors concluded that the two of the synthesized twenty imidazole derivatives, M24 and M25 were lead compounds with a strong antifungal activity that could be utilized for further development.³⁵

The Anti-Protozoal Properties of Imidazole Derivatives

With the lack of effective treatment in addition to the resistant infections worldwide, researchers began to

investigate new drugs against protozoan parasites. Herein, we reviewed the reported studies of imidazole derivatives based on their activities as anti-plasmodial, anti-leishmanial, and anti-protozoal.

According to Saccoliti et al, twenty-four Imidazole-derivatives were synthesized and biologically evaluated as antiprotozoal agents, unfortunately, none of the new derivatives exhibited higher potency compared to Artemisinin (ART) as a standard drug. However, compound M26 was three times more powerful than the reference drug Chloroquine (CHQ) as well as it possessed high selectivity index (SI = 253) that demonstrated the best anti-plasmodial activity.³⁶

Another study done by Ramu et al has tested three imidazolidinones as a potential antileishmanial agent

with a potential mechanism of action as a kinase inhibitor. Surprisingly, all three compounds were unable to inhibit human kinome but were potent kinome inhibitors on parasite cells. However, compound M27 had the highest IC_{50} of $10\mu M$. The authors concluded that compound M27 demonstrated significant antileishmanial activity via kinase inhibition.³⁷

Regarding the antiprotozoal activity of the (benz)imidazole derivatives, Aguirre et al examined the antiprotozoal activity in seven compounds and found that lipophilicity was essential for the potency of these compounds. At 100 mM, seven derivatives showed good activity against *T. cruzi*, while compound M28 was one of the best derivatives with a lipophilic substituent in the NO moiety. Benzimidazole derivatives showed high activity against *T. vaginalis* at 100 $\mu g/mL$ and 24 h of contact, which was sustained for three derivatives up to 48 h.³⁸

Additionally, Adeyme et al synthesized a total of twenty-six new imidazole derivatives through a comparative study to Pyrimethamine, a reference drug for the treatment of Toxoplasmosis. Among the five imidazole derivatives, compound M29 demonstrated an excellent selectivity against *T. gondii* versus the host cells.³⁹

Anti-Inflammatory Activity of Imidazole Derivatives

Inflammation is the predominant way of responding to a pathogen or injurious stimuli. This response includes the release of pro-inflammatory chemicals as a consequence, these chemicals increase blood vessel permeability and cause leakage of fluid into the tissues that result in pain, heat, redness, swelling, and loss of function.⁴⁰ The main two types of inflammation are acute and chronic inflammation.⁴¹ Medications that are commonly used to treat inflammation are nonsteroidal anti-inflammatory drugs (NSAID) and corticosteroids. The latter is known to cause both local and systemic side effects.⁴² Moreover, several undesirable adverse effects could be present due to the chronic usage of nonsteroidal anti-inflammatory drugs (NSAID).⁴³ Thus, in the pharmaceutical industry, benzimidazole is being investigated and the substituted forms of benzimidazole have also been used in various therapeutic applications. Several effective drugs currently on market, including proton pump inhibitors such as Omeprazole and Esomeprazole, and anthelmintics such as Mebendazole and Albendazole, which resulted in the optimization of benzimidazole derivatives based on their structures.⁴⁴ On

investigating imidazole derivatives, the current available studies reported to have promising anti-inflammatory activity based on their mechanism of action and important structure–activity relationship.

Shankar et al synthesized ten imidazole derivatives and screened their anti-inflammatory activities using cyclooxygenase-1 (COX-1) and cyclooxygenase-2 (COX-2) enzymes in vitro. Compound I30 demonstrated potent COX-2 inhibition with a percentage of 78.68 that is greater than the standard drug Ibuprofen (COX-2 inhibition percentage of 29.67). Additionally, in-vitro results were well-translated to the in-vivo environment in which compound I30 showed good anti-inflammatory activity against COX-2 enzyme with an inhibition percentage of 54.314 compared to Ibuprofen (52.986).⁴⁵

Using in vitro Ellman's method-based assay investigated by Bukhari et al, several anti-inflammatory agents were studied as potential inhibitors of COXs, phospholipase A2, lipoxygenase, TNF- α , and IL-6 using Indomethacin as a standard. Between thirty-five compounds, compound I31 was the most potent compound as an inhibitor of COX-1, COX-2, 5-LOX, and sPLA2-V with IC_{50} of 0.98, 11.56, 09.51, and 5.21, respectively.⁵⁰ Furthermore, Katikireddy et al investigated five compounds for their anti-inflammatory activity in vivo based on their potential antioxidant activity. The selected compounds showed comparable anti-inflammatory activity, decreased paw volume, relative to the standard drug Indomethacin. Moreover, compound I32 was the most effective anti-inflammatory agent with a percent inhibition of 90.30 at 6 hours. Compound I32 had a dichloride substitution at ortho and para positions which are responsible for the potent anti-inflammatory activity. Additional docking studies revealed that compound I32 binds to the active site of cyclooxygenase-2 (COX-2) confirming the in-vitro results.⁴⁷

Nascimento et al have synthesized eight novel imidazole derivatives to be tested as a potential anti-inflammatory agent in vitro. Compound I33 was the most potent inhibitor of nuclear factor kappa B (NF- κ B) transcription factor transmigration. Furthermore, compound I33 effectively reduced the pro-inflammatory mediators in addition to inhibition of nitric oxide release in J774 macrophage. The author concluded that compound I33 demonstrated similar efficacy in both in vitro and in vivo studies suggesting it could serve as a potential anti-inflammatory agent.⁴⁹

In this regard, Rocha et al have investigated two imidazole alkaloids to study the anti-inflammatory activity on human neutrophils with Indomethacin as a reference agent. The results of this study showed that I34 has good activity against neutrophils degranulation, reactive oxygen species (ROS), and IL-6 production. These results were confirmed using in vivo model that decreased inflammatory hypernociception and myeloperoxidase (MPO) release in mice.⁴⁸

Moreover, some authors reported SAR studies based on applying a medicinal chemistry approach from available anti-inflammatory drugs. For example, Husain et al believed that safe and effective NSAIDs would still be needed. Therefore, the author investigated Di- and tri-substituted imidazole derivatives using carrageenan-induced rat paw edema. The study results showed that six compounds displayed remarkable anti-inflammatory activity from 49.58 to 58.02% edema inhibition with minimal GI discomfort and severity index from 0.17 to 0.34 in vivo. Husain et al noted that compounds I35 and I36 are biologically active lead compounds with a promising anti-inflammatory activity that could be utilized as a novel anti-inflammatory candidate.³⁵

Furthermore, another study investigated in-vivo analgesic and anti-inflammatory activities of eleven synthesized compounds. The results showed that compound I37 demonstrated the strongest anti-inflammatory activity at 100 mg/kg compared to Nimesulide. However, the other nine compounds demonstrated a moderate anti-inflammatory activity. The initial in-vivo studies of these compounds showed that the chlorine group in meta position on the aniline ring improved both analgesic and anti-inflammatory properties, which could serve as new models for synthesizing more powerful immunomodulatory therapies.⁴⁴

An additional study in which authors evaluated twenty-three compounds against human erythrocytes. Among the synthesized compounds, compound I38 demonstrated excellent activity with an IC_{50} of 44 μ M that is higher than standard Aspirin (IC_{50} of 200 μ M) and Indomethacin (IC_{50} of 112 μ M). Furthermore, as the number of electrons withdrawing moiety in the benzene ring increases the anti-inflammatory activity also increases. The authors concluded that compounds I38 with two NO_2 substitutions on the benzene ring exhibited an excellent anti-inflammatory activity compared to aspirin and indomethacin.⁴⁶

Moreover, Gaba investigated the anti-inflammatory activity of nine imidazole compounds using a carrageenan-induced rat paw edema method, and results demonstrated

that three compounds showed anti-inflammatory activity ranging from 20.90 to 46.27% of inhibition.⁵¹ It is anticipated that the three electron-donating substituents in the three compounds showed a highly significant reduction in edema, the highest reduction was in compound I39 with 46.27% compared to the standard drug Indomethacin with 47.76%. The authors stated that compound I39 could be utilized as a lead compound for the discovery of novel anti-inflammatory drugs. All the anti-inflammatory studies are summarized in Table 3.

Pharmacokinetics ADME Properties of Selected Imidazole Derivatives

The ADME analysis was calculated using the SwissADME webserver. To determine whether the imidazole derivatives fulfill the drug-likeness properties, we have selected the most potent imidazole derivatives that demonstrated high potency as anti-cancer, anti-microbial, and anti-inflammatory effects from the above-discussed studies. The ADME properties are present in Tables 5 Tables 4–6 and the SMILE for each compound are summarized in Table 1S.







Molecular Weight (MW)

Considering Lipinski's rule limit of molecular weight (MW) of 500, it is apparent that all imidazole compounds were within the recommended range which improves their chances to be absorbed orally in the gastrointestinal tract, except for compound C2, C15, C16, M22, and M23 with an MW of 739.04, 607.65, 586.66, 834.95, and 521.61g/mol, respectively.⁵² The compounds with high MW do not pass through the cell membrane as readily as small MW, not only that but also low MW molecules are likely to be absorbed, diffused, transported across the cell membrane, and be more soluble compared to high MW.^{53,54} Importantly, compound I39 showed great promise as a lead compound to develop a novel class of orally active anti-inflammatory agents with fewer GI side effects.⁵¹

Lipophilicity (Log P) and Gastrointestinal (GI) Absorption





In drug discovery and design, lipophilicity, which is measured by the octanol/water partition coefficient, is an important physicochemical property that enhances compound GI absorption, safety, and efficacy.⁵⁵ All the imidazole derivatives demonstrated promising results in lipophilicity as leading compounds to generate a novel

Table 3 The Anti-Inflammatory Activities of Imidazole Derivatives in Multiple in vitro/ In vivo Models

Chemical Structure	Method/Target	Activity	Reported Target	Reference
 <p>I30</p>	In vitro COX-1 and COX-2 enzyme inhibition assay In vivo Carrageenan-induced rat paw edema assay after 24h	% of Inhibition COX-1 21.19 COX-2 78.68% of inhibition 54.314	Interacts with COX-2 enzyme	[45]
 <p>I31</p>	Ellman's method-based assay	IC₅₀ (μM) COX-1 0.98 ± 0.97 COX-2 11.56 ± 2.42 5-LOX 09.51 ± 3.40 sPLA2-V 5.21 ± 1.12	COX-1,2 5-LOX, sPLA2-V Inhibition	[50]
 <p>I32</p>	Carrageenan-induced rat paw edema assay after 6h	% of inhibition 90.30	Interacts with COX-2 enzyme	[47]
 <p>I33</p>	Leukocyte migration	Cytotoxicity Profile (CC₁₀) 51±1.7	Reducing leukocyte migration, Cg-Induced Oxidative Stress suppressor, and formation of exudate.	[31]
 <p>I34</p>	MTT assay on human neutrophils	% of inhibition MPO release 58.3 ± 3.8% ROS production 73.2 ± 0.9% IL-6 production 81.7%	Inhibition of neutrophils degranulation and reactive oxygen species (ROS) production	[49]
 <p>I35</p>	Carrageenan-induced rat paw edema method	(% of inhibition) ± SEM 37.78 ± 2.377	ND	[35]

(Continued)

Table 3 (Continued).

Chemical Structure	Method/Target	Activity	Reported Target	Reference
 <p style="text-align: center;">I36</p>	Carrageenan-induced rat paw edema method	(% of inhibition) ± SEM 24.67 ± 1.213	ND	[35]
 <p style="text-align: center;">I37</p>	Carrageenan-induced rat paw edema method after 3h	% of oedema inhibition 44.4	ND	[44]
 <p style="text-align: center;">I38</p>	Human erythrocyte	IC ₅₀ (µM) 44 ± 0.17	ND	[46]
 <p style="text-align: center;">I39</p>	Carrageenan-induced rat paw edema at 3 h	% of inhibition 53.73	ND	[51]

Notes: The prefix "I" letter in the compound's number referred to Anti-inflammatory activity. The cytotoxic concentration that killed 10% of the cells (CC₁₀) was calculated through nonlinear regression analysis of the logarithm of concentration in the function of the normalized response (percentage of cell viability). Results are expressed as mean ± E.P.M.

Abbreviations: IC₅₀, The concentration that inhibits enzyme activity by 50%; ND, Not determined.

class of orally active agents in the recommended range of -2.0–6.5 except for compounds C2, C10, C13, M19 M22, M26, and M29.⁵⁵ Moreover, all the compounds showed high gastrointestinal (GI) absorption except for compounds C2, C3, C14-16, M19, M22, M23, M26, M29, and I34 that could be due to their high MW.

Blood-Brain Barrier (BBB) Permeability

Chemical structures can be modified to increase or decrease BBB permeability by structural design. Since lipophilicity is a key determinant in BBB permeability, it is useful to use this property in brain tumors. All the imidazole compounds cannot cross BBB except for

compounds C1, C7-9, C11, C17, M18, M25, I31-33, and I36-37 due to the balanced lipophilicity/solubility properties that could greatly help to treat some central nervous system diseases which require BBB crossing ability.^{56,57}

Solubility (Log S)

Log S which predicts the aqueous solubility, with a recommended range of -6.5 to 0.5, of a compound has a direct effect on its absorption.⁵⁶ Therefore, low water solubility is associated with poor GI absorption, and thus the general goal is to avoid insoluble compounds such as C2, C3, C10, C15, C16, M22, M26, and M2. The rest of the imidazole derivatives displayed high to moderate water solubility based on the ADME predictions.⁵⁶

Table 4 The Pharmacokinetics ADME Properties of Selected Imidazole Derivatives as Anti-Cancer Agents

Compounds Codes	Molecular Weight	HB Donor	HB Acceptor	Log Po/w (WLOGP)	Log S (SILICO S-IT)	BBB Permeant	GI Absorption	Rule of Five (ROF)
Imidazole Derivatives as Anti-Cancer								
C1	360.84 g/mol	2	3	4.18	-6.12 Poorly soluble	Yes	High	Yes; 0 violation
C2	739.04 g/mol	2	5	7.36	-12.75 Insoluble	No	Low	Yes; 1 violation: MW>500
C3	388.43 g/mol	4	5	2.28	-6.84 Poorly soluble	No	High	Yes; 0 violation
C4	372.93 g/mol	1	2	6.39	-8.42 Poorly soluble	No	High	Yes; 0 violation
C5	269.17 g/mol	0	1	4.77	-4.86 Moderately soluble	Yes	High	Yes; 0 violation
C6	366.60 g/mol	2	3	3.23	-6.46 Poorly soluble	Yes	High	Yes; 0 violation
C7	447.28 g/mol	0	5	3.75	-5.97 Moderately soluble	Yes	High	Yes; 0 violation
C8	408.27 g/mol	0	5	4.32	-6.59 Poorly soluble	No	High	Yes; 0 violation
C9	294.31 g/mol	4	6	-1.48	-2.02 Soluble	No	Low	Yes; 0 violation
C10	490.60 g/mol	0	2	7.50	-12.48 Insoluble	No	Low	Yes; 0 violation
C11	365.22 g/mol	1	3	5.63	-9.22 Poorly soluble	Yes	High	Yes; 0 violation
C12	460.53 g/mol	2	6	2.44	-7.50 Poorly soluble	No	High	Yes; 0 violation
C13	607.65 g/mol	3	9	8.66	-12.88 insoluble	No	Low	Yes; 1 violation: MW>500
C14	586.66 g/mol	2	6	5.87	-10.64 insoluble	No	Low	No; 1 violation: MW>500
C15	437.66 g/mol	2	2	7.84	-10.19 Insoluble	No	Low	Yes; 0 violation
C16	390.39 g/mol	1	5	2.50	-5.09 Moderately soluble	No	High	Yes; 0 violation
C17	365.81 g/mol	1	2	3.13	-6.33 Poorly soluble	Yes	High	Yes; 0 violation

Hydrogen Bond Donor and Acceptor (HBD/HBA)

The hydrogen bond donors (HBD) increase acidity while hydrogen bond acceptor (HBA) increases basicity. Using

ADME predictions, all compounds obeyed Lipinski's rule regarding the number of hydrogen bond donors from 0 to 6 and hydrogen bond acceptors from 2 to 10 except for compounds M22 which has 13 hydrogen bond acceptors.^{54,58,59}

Table 5 The Pharmacokinetics ADME Properties of Selected Imidazole Derivatives as Anti-Microbial Agents

Compounds Codes	Molecular Weight	HB Donor	HB Acceptor	Log Po/w (WLOGP)	Log S (SILICO S-IT)	BBB Permeant	GI Absorption	Rule of Five (ROF)
Imidazole Derivatives as Anti-microbial								
M18	324.76 g/mol	0	4	4.10	-6.97 Poorly soluble	Yes	High	Yes; 0 violation
M19	390.99 g/mol	2	2	6.99	-8.26 Poorly soluble	No	Low	Yes; 0 violation
M20	304.86 g/mol	1	1	6.26	-7.23 Poorly soluble	No	High	Yes; 0 violation
M21	373.36 g/mol	1	6	0.98	-2.3 Soluble	No	High	Yes; 0 violation
M22	834.95 g/mol	2	13	6.58	-12.74 Insoluble	No	Low	No; 2 violations: MW>500, NorO>10
M23	521.61 g/mol	1	6	5.21	-8.08 Poorly soluble	No	Low	Yes; 1 violation: MW>500
M24	371.39 g/mol	0	4	5.12	-7.41 Poorly soluble	No	High	Yes; 0 violation
M25	356.42 g/mol	0	3	5.22	-8.17 Poorly soluble	Yes	High	Yes; 0 violation
M26	444.36 g/mol	1	1	7.49	-10.85 Insoluble	No	Low	Yes; 0 violation
M27	189.17 g/mol	2	3	-0.61	-2.77 Soluble	No	High	Yes; 0 violation
M28	351.31 g/mol	0	8	2.34	-2.56 Soluble	No	High	Yes; 0 violation
M29	423.55 g/mol	1	3	8.10	-10.53 Insoluble	No	Low	Yes; 0 violation

Target Predictions of Selected Imidazole Derivatives

The bioactivity scores of selected imidazole derivatives were calculated using Molinspiration virtual screening webserver.⁶⁰ In Table 7, the larger the bioactivity score correlates with the higher probability that a specific molecule will be active at that target.² Results showed that compound C4 may work by five different mechanisms that give a better chance to be a promising novel anticancer agent. Interestingly, the

computational bioavailability scores correlated well with the previously reported studies for some of the proposed targets. For example, C4 was reported to act through inhibition of focal adhesion kinase (FAK) and computational results demonstrated that C4 exhibited a high bioactivity score as a kinase inhibitor.¹² In similar manner, the anti-cancer mechanisms of compounds C1, C12, C13, and C15 were reported to inhibit Topoisomerase IIR, aurora kinase, tubulin polymerization, and BRAF^{V600E} kinase, respectively. These

Table 6 The Pharmacokinetics ADME Properties of Selected Imidazole Derivatives as Anti-Inflammatory Agents

Compounds Codes	Molecular Weight	HB Donor	HB Acceptor	Log Po/w (WLOGP)	Log S (SILICO S-IT)	BBB Permeant	GI Absorption	Rule of Five (ROF)
Imidazole Derivatives as Anti-inflammatory								
I30	356.42 g/mol	1	3	2.58	-6.94 Poorly soluble	Yes	High	Yes; 0 violation
I31	267.28 g/mol	2	3	2.34	-5.15 Moderately soluble	No	High	Yes; 0 violation
I32	389.28 g/mol	2	3	4.89	-8.27 Poorly soluble	No	High	Yes; 0 violation
I33	350.39 g/mol	0	4	4.68	-7.02 Poorly soluble	Yes	High	Yes; 0 violation
I34	286.33 g/mol	1	4	1.16	-3.46 Soluble	Yes	High	Yes; 0 violation
I35	371.39 g/mol	0	4	5.12	-7.41 Poorly soluble	No	High	Yes; 0 violation
I36	342.39 g/mol	1	3	4.92	-7.48 Poorly soluble	Yes	High	Yes; 0 violation
I37	257.72 g/mol	2	1	3.49	-6.50 Poorly soluble	Yes	High	Yes; 0 violation
I38	404.38 g/mol	3	8	2.06	-3.97 Moderately soluble	No	Low	Yes; 1 violation: NorO>10
I39	407.49 g/mol	1	4	4.94	-7.93 Poorly soluble	No	High	Yes; 0 violation

four compounds exhibited high bioactivity scores as kinase inhibitors that provide additional confirmation to the reported studies.^{14,20,21} Moreover, compound C10 could act via four mechanisms one of which is GPCR which is consistent with the previously reported study about C10 and its action through dimerization of estrogen receptor.¹⁸ Compound C11 was suggested to stabilize DNA c-MYC G-quadruplex by modulating ion-channel and this was confirmed with our target prediction score of 0.30.¹⁹ However, compound C14 was reported to work on GSK-3 β which is one of the kinase enzymes but our target prediction score of -0.34 did not correlate with Al-Blewi et al study.²⁴ All the proposed mechanisms for imidazole derivatives are summarized in Figure 1.

The results shown in Table 8, indicate that compound M21 was the most promising compound as a GPCR ligand and kinase inhibitor. Moreover, enzyme inhibition tends to be a crucial factor for selecting lead candidates for *T. species*

therapy. Moreover, M27 was proposed to be a kinase inhibitor, however, the computational results did not suggest that with a score of -0.30.³⁷ Furthermore, compound M26 exhibited the highest bioactivity against all the investigated targets, one of which is enzymes (a score of 0.6) that is consistent with Saccoliti et al study which mentioned the compound works by inhibiting the CYP51 enzyme as a possible mechanism of action.³⁶

The findings shown in Table 9, demonstrated that some anti-inflammatory mechanisms of imidazole derivatives could be due to the interactions with GPCRs, nuclear receptors, ion channels, proteases, kinases, and enzymes. According to the bioactivity scores, the most promising compounds were found to be I30, I31, I33, and I34 with scores of 0.24, 0.32, 0.21, and 0.72, respectively, as an enzyme inhibitor suggesting COX enzyme may be involved in mediating this anti-inflammatory effect that

Table 7 The Target Predictions of Imidazole Derivatives as Anti-Cancer

Compounds Codes	GPCR Ligand	Ion Channel Modulator	Kinase Inhibitor	Nuclear Receptor Ligand	Protease Inhibitor	Enzyme Inhibitor	Reported Target	Reference
Imidazole Derivatives as Anti-Cancer								
C1	-0.10	-0.20	0.08	-0.41	-0.57	-0.33	Topoisomerase IIR catalytic inhibitors.	[9]
C2	-0.89	-2.07	-1.81	-1.61	-0.81	-1.39	ND	[10]
C3	0.23	0.16	0.61	-0.87	0.21	0.12	Focal Adhesion Kinase (FAK) inhibition.	[12]
C4	0.53	0.34	0.28	-0.05	0.10	0.41	ND	[14]
C5	-0.09	-0.14	-0.12	-0.62	-0.62	-0.02	Cyclin-dependent kinase 6 (CDK6) inhibition	[15]
C6	-0.20	-0.23	0.41	-0.59	-0.48	0.25	ND	[16]
C7	-0.13	-0.64	0.03	-0.46	-0.52	-0.18	ND	[17]
C8	-0.16	-0.27	0.13	-0.48	-0.33	0.06	Inhibit tubulin polymerization	[21]
C9	-0.35	-1.07	-0.69	-1.19	-0.48	-0.14	ND	[22]
C10	0.13	0.11	0.28	-0.03	-0.09	0.09	ND Possibly due to dimerized estrogen receptor	[18]
C11	0.32	0.30	0.51	-0.19	-0.24	0.31	c-MYC G-quadruplex DNA stabilizers	[19]
C12	0.18	-0.16	0.37	-0.45	-0.11	0.04	Aurora kinase inhibition	[20]
C13	0.12	-0.47	0.13	-0.30	-0.07	-0.07	Inhibit BRAF ^{V600E} kinase	[23]
C14	-0.28	-0.85	-0.34	-0.62	-0.34	-0.20	Inhibit GSK-3 β	[24]
C15	0.35	0.16	0.20	-0.04	0.20	0.29	ND	[11]
C16	-0.86	-1.08	-1.15	-1.02	-1.16	-0.60	ND	[13]
C17	0.12	0.04	-0.15	-0.38	0.03	0.00	ND	[25]

correlated well with Shankar et al, Bukhari et al, Nascimento et al and Rocha et al studies.^{45,48–50} On the contrary, Katikireddy et al believed that compound I32 works by inhibiting the COX-2 enzyme, however, the target prediction did not show a promising bioactivity score at this target suggesting computational studies should not be used solely for target predictions.⁴⁷

Future Direction

The current review illustrates that imidazole derivatives have promising pharmacological effects particularly as

anticancer, anti-microbial, and anti-inflammatory with a great potential for treating various human diseases. Computational studies were performed to encapsulate some of the important parameters of lead optimization such as ADME and target predictions for the highly potent candidates. Our results showed that several imidazole derivatives demonstrated an acceptable pharmacokinetic profile with promising bioavailability scores that correlated well with reported targets. Moreover, several considerations could be taken into account for the development of imidazole derivatives such as the

Proposed Mechanisms for Imidazole Derivatives




Figure 1 Proposed mechanisms for imidazole derivatives.

Table 8 Target Prediction of Imidazole Derivatives as Anti-Microbial

Compounds Codes	GPCR Ligand	Ion Channel Modulator	Kinase Inhibitor	Nuclear Receptor Ligand	Protease Inhibitor	Enzyme Inhibitor	Reported Target	Reference
Imidazole Derivatives as Anti-microbial								
M18	-0.10	-0.48	-0.21	-0.46	-0.27	-0.19	ND	[30]
M19	0.34	0.22	0.03	0.06	0.08	0.29	ND	[31]
M20	0.36	0.15	0.22	0.04	-0.14	0.22	ND	[31]
M21	0.01	-0.53	-0.35	-0.54	-0.25	0.04	Double strand DNA helix breakage	[32]
M22	-2.04	-3.48	-2.85	-3.16	-1.60	-2.52	ND	[33]
M23	-0.62	-0.89	-0.62	-0.77	-0.62	-0.57	ND	[34]
M24	0.05	0.07	0.04	-0.28	-0.41	0.00	ND	[35]
M25	0.20	0.13	0.18	-0.19	-0.28	0.11	ND	[37]
M26	0.31	0.12	0.17	0.12	-0.02	0.61	CYP51	[36]
M27	-0.89	-1.48	-0.30	-1.93	-1.54	-0.67	Kinase inhibitor	[37]
M28	-0.27	0.08	-0.43	-0.51	-0.28	-0.18	ND	[38]
M29	-0.12	-0.26	-0.11	0.27	-0.44	0.10	ND	[39]

Table 9 The Target Prediction of Imidazole Derivatives as Anti-Inflammatory

Compounds Codes	GPCR Ligand	Ion Channel Modulator	Kinase Inhibitor	Nuclear Receptor Ligand	Protease Inhibitor	Enzyme Inhibitor	Reported Target	Reference
Imidazole Derivatives as Anti-inflammatory								
I30	0.42	0.32	0.67	-0.10	0.12	0.24	COX-2 enzyme	[45]
I31	0.17	0.08	0.48	-0.23	-0.25	0.32	COX-1,2 5-LOX, sPLA2-V Inhibition	[50]
I32	-0.12	-0.55	-0.27	-0.56	-0.61	-0.24	COX-2 enzyme	[47]
I33	0.05	-0.06	-0.19	0.02	-0.24	0.21	p65 NF-κB	[49]
I34	0.64	0.15	0.26	0.03	0.28	0.72	NF-κB	[48]
I35	0.05	0.07	0.04	-0.28	-0.41	0.00	ND	[35]
I36	0.21	0.12	0.18	-0.15	-0.29	0.16	ND	[35]
I37	-0.13	0.09	0.08	-0.69	-0.44	-0.07	ND	[44]
I38	-0.10	-0.57	-0.58	-0.54	-0.50	-0.23	ND	[46]
I39	-0.11	-0.19	0.07	-0.36	-0.23	-0.14	ND	[51]

in vitro testing using murine cell lines which could greatly influence the translation of data into the human biological system. The use of human cells, in-vivo models, and/or more complex in vitro 3D cultures that mimic the biological system could accelerate the development of these agents into the clinic. In summary, the current reported literature in imidazole core scaffold demonstrated potential therapeutic values that suggest further development and optimization in this area is required to discover novel and potent drug candidates.

Abbreviations

A375, Human Melanoma Cells; A549, Lung Cancer Cells; ADME, Absorption, Distribution, Metabolism, and Excretion; ASPC-1, pancreatic cancer cells; B16, Mouse Melanoma Cells; BBB, Blood-Brain Barrier; CDK6, Cyclin-Dependent Kinase 6; CNE-1, Nasopharyngeal Carcinoma; COX-1, cyclooxygenase-1; COX-2, cyclooxygenase-2; CTX, Cyclophosphamide; CYP2C19, Cytochrome P 450 2C19; CYP3A4, Cytochrome P 450 3A4; EAT, Ehrlich Ascites Tumor; EAT, Ehrlich Ascites Tumor; ERs, Estrogen Receptors; ESBL, Extended Spectrum Beta-Lactamases; FAK, Focal Adhesion Kinase; *Flavus*, *Aspergillus flavus*; GI, Gastrointestinal; GPCR, G protein Coupled Receptors; GSK-3 β , Glycogen Synthase Kinase 3 beta; *H. oryzae*, *Hirschiella oryzae*; HBA, Hydrogen Bond Acceptor; HBD, Hydrogen Bond Donor; HCT-116, Colon Cancer Cells; HEK 293, Kidney Cancer Cells; HL-60, Human Myeloid Leukemia Cells; HUVECS, Human Umbilical Vein Endothelial Cells; K562, Human Myeloid Leukemia Cells; Log P, Lipophilicity; Log S, Solubility; MCF-7, Breast Cancer Cells; Mcl-1, Myeloid Cell Leukemia 1; MDA-MB-231, Breast; MIC, Minimum Inhibitory Concentration; MPO, myeloperoxidase; MRSA, Methicillin-Resistance *Staphylococcus Aureus*; MW, Molecular Weight; NCI-60, National Cancer Institute- 60; NF- κ B, Nuclear Factor Kappa B N β , *Aspergillus Niger*; NSAID, Nonsteroidal Anti-inflammatory Drugs; PANC-1, human pancreatic cancer cell line; PC-3, Prostate Cancer Cells; ROF, Rule of five; ROS, reactive oxygen species; *S. Typhimurium*, *Salmonella typhimurium*; SH-SY5Y, Neuroblastoma Cells; SMCS, Smooth Muscle Cells; *subtilis*, *Bacillus subtilis*; *T. cruzi*, *Trypanosoma cruzi*; *T. vaginalis*, *Trichomonas vaginalis*; *T. viridae*, *Trichoderma viridae*; U87-MG, Brain Cancer Cells; VRE, Vancomycin-Resistant Enterococci.

Acknowledgments

The authors want to express their sincerest gratitude to the College of Pharmacy (COP) at King Saud bin Abdulaziz University for Health Sciences (KSAU-HS) for their continued support.

Funding

The authors acknowledge financial support from King Abdullah International Medical Research Center (KAIMRC), Ministry of National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia. Grant # (SP20.441.R).

Disclosure

The authors declare no conflicts of interest for this work.

References

1. Sherer C, Snape TJ. Heterocyclic scaffolds as promising anticancer agents against tumours of the central nervous system: exploring the scope of indole and carbazole derivatives. *Eur J Med Chem.* 2015;97:552–560. doi:10.1016/j.ejmech.2014.11.007
2. Molina P, Tárraga A, Otón F. Imidazole derivatives: a comprehensive survey of their recognition properties. *Org Biomol Chem.* 2012;10(9):1711–1724. doi:10.1039/c2ob06808g
3. Gaba M, Mohan C. Development of drugs based on imidazole and benzimidazole bioactive heterocycles: recent advances and future directions. *Med Chem Res.* 2016;25(2):173–210.
4. Institute NC. Cancer statistics; 2020. Available from: <https://www.cancer.gov/about-cancer/understanding/statistics>. Accessed July 20, 2021.
5. UK CR. Worldwide cancer statistics; 2018. Available from: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/worldwide-cancer#heading-One>. Accessed July 20, 2021.
6. Institute NC. What is cancer?; 2015. Available from: <https://www.cancer.gov/about-cancer/understanding/what-is-cancer>. Accessed July 20, 2021.
7. Pérez-Herrero E, Fernández-Medarde A. Advanced targeted therapies in cancer: drug nanocarriers, the future of chemotherapy. *Eur J Pharm Biopharm.* 2015;93:52–79.
8. Ali R, Mirza Z, Ashraf GM, et al. New anticancer agents: recent developments in tumor therapy. *Anticancer Res.* 2012;32(7):2999–3005.
9. Baviskar AT, Madaan C, Preet R, et al. N-fused imidazoles as novel anticancer agents that inhibit catalytic activity of topoisomerase α and induce apoptosis in G1/S phase. *J Med Chem.* 2011;54(14):5013–5030. doi:10.1021/jm200235u
10. Zhao F, Lu W, Su F, et al. Synthesis and potential antineoplastic activity of dehydroabietylamine imidazole derivatives. *MedChemComm.* 2018;9(12):2091–2099. doi:10.1039/C8MD00487K
11. Chen J, Wang Z, Lu Y, Dalton JT, Miller DD, Li W. Synthesis and antiproliferative activity of imidazole and imidazoline analogs for melanoma. *Bioorg Med Chem Lett.* 2008;18(11):3183–3187. doi:10.1016/j.bmcl.2008.04.073
12. Dao P, Smith N, Tomkiewicz-Rault C, et al. Design, synthesis, and evaluation of novel imidazo[1,2-a][1,3,5]triazines and their derivatives as focal adhesion kinase inhibitors with antitumor activity. *J Med Chem.* 2015;58(1):237–251. doi:10.1021/jm500784e
13. Sarkarzadeh H, Miri R, Firuzi O, et al. Synthesis and antiproliferative activity evaluation of imidazole-based indeno[1,2-b]quinoline-9,11-dione derivatives. *Arch Pharm Res.* 2013;36(4):436–447. doi:10.1007/s12272-013-0032-7

14. Roopashree R, Mohan CD, Swaroop TR, Jagadish S, Rangappa KS. Synthesis, characterization and in vivo biological evaluation of novel benzimidazoles as potential anticancer agents. *Asian J Pharm Clin Res.* 2014;7(5):309–313.
15. Alkahtani HM, Abbas AY, Wang S. Synthesis and biological evaluation of benzo[d]imidazole derivatives as potential anti-cancer agents. *Bioorg Med Chem Lett.* 2012;22(3):1317–1321. doi:10.1016/j.bmcl.2011.12.088
16. Chung K-H, Hong S-Y, You H-J, Park R-E, Ryu C-K. Synthesis and biological evaluation of 5-arylamino-1H-benzo[d]imidazole-4,7-diones as inhibitor of endothelial cell proliferation. *Bioorg Med Chem.* 2006;14(17):5795–5801. doi:10.1016/j.bmc.2006.05.059
17. Xue N, Yang X, Wu R, et al. Synthesis and biological evaluation of imidazol-2-one derivatives as potential antitumor agents. *Bioorg Med Chem.* 2008;16(5):2550–2557. doi:10.1016/j.bmc.2007.11.048
18. Meenakshisundaram S, Manickam M, Pillaiyar T. Exploration of imidazole and imidazopyridine dimers as anticancer agents: design, synthesis, and structure-activity relationship study. *Arch Pharm (Weinheim).* 2019;352(12):e1900011. doi:10.1002/ardp.201900011
19. Wu Q, Song Y, Liu R, et al. Synthesis, docking studies and antitumor activity of phenanthroimidazole derivatives as promising c-myc G-quadruplex DNA stabilizers. *Bioorg Chem.* 2020;102:104074.
20. Fan C, Zhong T, Yang H, et al. Design, synthesis, biological evaluation of 6-(2-amino-1H-benzo [d] imidazole-6-yl) quinazolin-4 (3H)-one derivatives as novel anticancer agents with Aurora kinase inhibition. *Eur J Med Chem.* 2020;190:112108.
21. Komuraiah B, Ren Y, Xue M, et al. Design, synthesis and biological evaluation of benz-fused five-membered heterocyclic compounds as tubulin polymerization inhibitors with anticancer activities. *Chem Biol Drug Des.* 2021;97(5):1109–1116.
22. Hassan AY, El-Sebaey SA, El Deeb MA, Elzoghbi MS. Potential antiviral and anticancer effect of imidazoles and bridgehead imidazoles generated by HPV-Induced cervical carcinomas via reactivating the P53/pRb pathway and inhibition of CA IX. *J Mol Struct.* 2021;1230:129865.
23. Ali EM, Abdel-Maksoud MS, Ammar UM, et al. Design, synthesis, and biological evaluation of novel imidazole derivatives possessing terminal sulphonamides as potential BRAFV600E inhibitors. *Bioorg Chem.* 2021;106:104508.
24. Al-Blewi F, Shaikh SA, Naqvi A, et al. Design and synthesis of novel imidazole derivatives possessing triazole pharmacophore with potent anticancer activity, and in silico ADMET with GSK-3 β molecular docking investigations. *Int J Mol Sci.* 2021;22(3):1162.
25. Zhang M, Ding Y, Qin H-X, et al. One-pot synthesis of substituted pyrrole-imidazole derivatives with anticancer activity. *Mol Divers.* 2020;24:1177–1184.
26. Badiie P, Hashemizadeh Z. Opportunistic invasive fungal infections: diagnosis & clinical management. *Indian J Med Res.* 2014;139(2):195–204.
27. US NIOH. Understanding emerging and re-emerging infectious diseases; 2007. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK20370/>. Accessed July 20, 2021.
28. World Health Organization. Record number of countries contribute data revealing disturbing rates of antimicrobial resistance; 2020. Available from: <https://www.who.int/news/item/01-06-2020-record-number-of-countries-contribute-data-revealing-disturbing-rates-of-antimicrobial-resistance>.
29. World Health Organization. *Antimicrobial Resistance and Primary Health Care.* World Health Organization; 2018.
30. Ansari KF, Lal C. Synthesis, physicochemical properties and antimicrobial activity of some new benzimidazole derivatives. *Eur J Med Chem.* 2009;44(10):4028–4033. doi:10.1016/j.ejmech.2009.04.037
31. Sharma S, Gangal S, Rauf A. Convenient one-pot synthesis of novel 2-substituted benzimidazoles, tetrahydrobenzimidazoles and imidazoles and evaluation of their in vitro antibacterial and antifungal activities. *Eur J Med Chem.* 2009;44(4):1751–1757. doi:10.1016/j.ejmech.2008.03.026
32. Pieczonka AM, Strzelczyk A, Sadowska B, Młostoń G, Stączek P. Synthesis and evaluation of antimicrobial activity of hydrazones derived from 3-oxido-1H-imidazole-4-carbohydrazides. *Eur J Med Chem.* 2013;64:389–395. doi:10.1016/j.ejmech.2013.04.023
33. Al-Blewi FF, Almeahadi MA, Aouad MR, et al. Design, synthesis, ADME prediction and pharmacological evaluation of novel benzimidazole-1,2,3-triazole-sulfonamide hybrids as antimicrobial and antiproliferative agents. *Chem Cent J.* 2018;12(1):110. doi:10.1186/s13065-018-0479-1
34. Daraji DG, Rajani DP, Rajani SD, et al. Structure based design, synthesis, and biological evaluation of imidazole derivatives targeting dihydropteroate synthase enzyme. *Bioorg Med Chem Lett.* 2021;36:127819.
35. Husain A, Drabu S, Kumar N, Alam MM, Bawa S. Synthesis and biological evaluation of di- and tri-substituted imidazoles as safer anti-inflammatory-antifungal agents. *J Pharm Bioallied Sci.* 2013;5(2):154–161. doi:10.4103/0975-7406.111822
36. Saccoliti F, Madia VN, Tudino V, et al. Design, synthesis, and biological evaluation of new 1-(Aryl-1 H-pyrrolyl)(phenyl)methyl-1 H-imidazole derivatives as antiprotozoal agents. *J Med Chem.* 2019;62(3):1330–1347. doi:10.1021/acs.jmedchem.8b01464
37. Ramu D, Jain R, Kumar RR, et al. Design and synthesis of imidazolidinone derivatives as potent anti-leishmanial agents by bioisosterism. *Arch Pharm.* 2019;352(4):e1800290. doi:10.1002/ardp.201800290
38. Aguirre G, Boiani M, Cerecetto H, et al. Novel antiprotozoal products: imidazole and benzimidazole N-Oxide derivatives and related compounds. *Arch Pharm.* 2004;337(5):259–270. doi:10.1002/ardp.200300840
39. Adeyemi OS, Eseola AO, Plass W, et al. Imidazole derivatives as antiparasitic agents and use of molecular modeling to investigate the structure-activity relationship. *Parasitol Res.* 2020;119(6):1925–1941.
40. Chen L, Deng H, Cui H, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget.* 2018;9(6):7204–7218. doi:10.18632/oncotarget.23208
41. Silva VG, Silva RO, Damasceno SRB, et al. Anti-inflammatory and antinociceptive activity of episopiloturine, an imidazole alkaloid isolated from *Pilocarpus microphyllus*. *J Nat Prod.* 2013;76(6):1071–1077. doi:10.1021/np400099m
42. Youssef J, Novosad SA, Winthrop KL. Infection risk and safety of corticosteroid use. *Rheum Dis Clin North Am.* 2016;42(1):157–x. doi:10.1016/j.rdc.2015.08.004
43. Sostres C, Gargallo CJ, Arroyo MT, Lanás A. Adverse effects of non-steroidal anti-inflammatory drugs (NSAIDs, aspirin and coxibs) on upper gastrointestinal tract. *Best Pract Res Clin Gastroenterol.* 2010;24(2):121–132. doi:10.1016/j.bpg.2009.11.005
44. Achar KC, Hosamani KM, Seetharamareddy HR. In-vivo analgesic and anti-inflammatory activities of newly synthesized benzimidazole derivatives. *Eur J Med Chem.* 2010;45(5):2048–2054. doi:10.1016/j.ejmech.2010.01.029
45. Shankar B, Jalapathi P, Valeru A, Kishor Kumar A, Saikrishna B, Kudle K. Synthesis and biological evaluation of new 2-(6-alkylpyrazin-2-yl)-1H-benz[d]imidazoles as potent anti-inflammatory and antioxidant agents. *Med Chem Res.* 2017;26(9):1835–1846. doi:10.1007/s00044-017-1897-7
46. Shantharam C. Synthesis and SAR studies of potent antioxidant and anti-inflammatory activities of imidazole Derived Schiff base analogues. *Biochem Anal Biochem.* 2017;06:314.

47. Katikireddy R, Kakkerla R, Krishna MPSM, Durgaiah G, Reddy YN, Satyanarayana M. Synthesis and biological evaluation of (E)-N'-benzylidene-7-methyl-2-propyl-1H-benzo[d]imidazole-5-carboximidazides as antioxidant, anti-inflammatory and analgesic agents. *Heterocycl Comm*. 2019;25(1):27–38. doi:10.1515/hc-2019-0009
48. Rocha TM, Machado NJ, de Sousa JAC, et al. Imidazole alkaloids inhibit the pro-inflammatory mechanisms of human neutrophils and exhibit anti-inflammatory properties in vivo. *J Pharm Pharmacol*. 2019;71(5):849–859. doi:10.1111/jphp.13068
49. Nascimento M, Munhoz ACM, Theindl LC, et al. A novel tetrasubstituted imidazole as a prototype for the development of anti-inflammatory drugs. *Inflammation*. 2018;41(4):1334–1348. doi:10.1007/s10753-018-0782-y
50. Bukhari SN, Lauro G, Jantan I, et al. Anti-inflammatory trends of new benzimidazole derivatives. *Future Med Chem*. 2016;8(16):1953–1967. doi:10.4155/fmc-2016-0062
51. Gaba M. Design, synthesis and biological evaluation of Novel 1, 2, 5-substituted benzimidazole derivatives as gastroprotective anti-inflammatory and analgesic agents. *Med Chem (Los Angeles)*. 2015;5(2). doi:10.4172/2161-0444.1000243
52. Lipinski C, Lipinski CA. Lead- and drug-like compounds: the rule-of-five revolution. *Drug Discov Today Technol*. 2004;1:337–341. doi:10.1016/j.ddtec.2004.11.007
53. Di L, Kerns EH. *Drug-Like Properties: Concepts, Structure, Design and Methods: From ADME to Toxicity Optimization*. Amsterdam [i pozostale]: Elsevier - Academic Press; 2019.
54. Sharom FJ. ABC multidrug transporters: structure, function and role in chemoresistance. *Pharmacogenomics*. 2008;9(1):105–127. doi:10.2217/14622416.9.1.105
55. Arnott JA, Planey SL. The influence of lipophilicity in drug discovery and design. *Expert Opin Drug Discov*. 2012;7(10):863–875. doi:10.1517/17460441.2012.714363
56. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep*. 2017;7(1):42717. doi:10.1038/srep42717
57. Noriega-Iribe E, Díaz-Rubio L, Estolano-Cobián A, et al. In vitro and in silico screening of 2,4,5-trisubstituted imidazole derivatives as potential xanthine oxidase and acetylcholinesterase inhibitors, anti-oxidant, and antiproliferative agents. *Appl Sci*. 2020;10(8):2889. doi:10.3390/app10082889
58. Abraham MH, Chadha HS, Mitchell RC. Hydrogen bonding. 33. Factors that influence the distribution of solutes between blood and brain. *J Pharm Sci*. 1994;83(9):1257–1268. doi:10.1002/jps.2600830915
59. Pajouhesh H, Lenz GR. Medicinal chemical properties of successful central nervous system drugs. *NeuroRx*. 2005;2(4):541–553.
60. Cheminformatics M; 2020. Available from: <https://www.molinspiration.com>. Accessed July 20, 2021.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>