

Research article

Ecdysteroids and oocyte development in the black fly *Simulium vittatum*

Rafael Noriega², Frank B Ramberg¹ and Henry H Hagedorn*¹

Address: ¹Department of Entomology and Center for Insect Science University of Arizona, Tucson, AZ, USA and ²University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania 15216, USA

E-mail: Rafael Noriega - ranst23+@pitt.edu; Frank B Ramberg - framberg@U.Arizona.EDU; Henry H Hagedorn* - hagedorn@ag.arizona.edu

*Corresponding author

Published: 24 April 2002

Received: 6 February 2002

BMC Developmental Biology 2002, 2:6

Accepted: 24 April 2002

This article is available from: <http://www.biomedcentral.com/1471-213X/2/6>

© 2002 Noriega et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Oocyte development was studied in the autogenous black fly, *Simulium vittatum* (Diptera, Nematocera), a vector of *Onchocerca volvulus*, the causative agent of onchocerciasis.

Results: Oocyte growth was nearly linear between adult eclosion and was complete by 72 hours at 21°C. The oocyte became opaque at 14 hours after eclosion indicating the initiation of protein yolk deposition. The accumulation of vitellogenin was measured using SDS-PAGE. The density of the yolk protein bands at about 200 and 65 kDa increased during the first and second days after eclosion. The amount of protein in the 200 kDa band of vitellogenin, determined using densitometry, rapidly increased between 12 and 25 hours after eclosion. Ecdysteroid levels were measured using a competitive ELISA. Ecdysteroid levels increased rapidly and subsequently declined during the first day after eclosion.

Conclusion: These data show a correlation between the appearance of vitellogenin in the oocyte, and the rise in ecdysteroids. A possible relationship to molting of the nematode, *Onchocerca volvulus*, is discussed.

Background

Blackflies are major nuisance pests and are vectors of the nematode, *Onchocerca volvulus*, which causes the serious human disease, onchocerciasis, mainly in tropical Africa but also in Central and South America. Microfilariae, ingested by the black fly from the human host, invade the thoracic muscles and molt several times. The steroid hormone, 20-hydroxyecdysone, is known to control molting in insects [1] and has been implicated in the control of molting of nematodes [2]. Ecdysteroids have been isolated from several parasitic nematodes [3], and have been shown to fluctuate in titer during the molt [4]. Genes related to the 20-hydroxyecdysone receptor gene of *Dro-*

sophila melanogaster have been isolated from *O. volvulus*[5]. Interestingly, 20-hydroxyecdysone caused premature timing of the third stage molt of *Diriofilaria immitis*[6]. This raises the possibility that changes in ecdysteroid titer in the insect host could affect the timing of developmental molts of the parasitic nematode.

20-hydroxyecdysone is important in the control of reproduction in Diptera, for example in *Drosophila melanogaster*[7,8], the stable fly, *Stomoxys calcitrans*[9] the blowfly *Phormia regina*[10], and the mosquitoes, *Aedes aegypti*[11,12], *Aedes atropalpus*[13], and *Anopheles albimanus*[14]. Ecdysteroid titers rise and fall in these insects




Figure 1
Follicle and oocyte length after eclosion in the black fly, *Simulium vittatum*. Each time point represents the mean of 20 follicle measurements from each of 3–5 females, \pm standard deviation.

during the development of eggs. In the hematophagous Diptera, the titer of ecdysteroids rise in the adult female after the blood meal as eggs develop [10,15,16]. It is possible that ecdysteroid titers also increase during egg development in the adult female black fly.

Simulium vittatum is one of the few blackflies that can be reared in the laboratory [17] making it especially useful for physiological studies. *S. vittatum* can develop its first batch of eggs autogenously [i.e. without a blood meal] but requires a blood meal for subsequent cycles of egg development [18]. We studied the relationship between egg development and ecdysteroids during autogenous egg development in this species.

Results

To determine the basic characteristics of oogenesis, blackflies were collected and follicle and oocyte lengths were measured. Oocyte growth in *S. vittatum* begins after adult eclosion and is completed by 72 hours after eclosion at 21°C. From 2–72 hours the increase in follicle and oocyte length was nearly linear (Fig. 1). The oocyte became opaque at 14 hours after eclosion indicating the initiation of protein yolk deposition.

To correlate the growth of the oocyte with the appearance of yolk proteins, the accumulation of protein in ovaries was measured using SDS-PAGE. As shown in Figure 2, major protein bands are present at about 200 and 65 kDa. These bands were not seen in extracts of males (data not shown). Given their presence in large amounts in the

oocytes, their absence in males, and the presence of similar bands in *Simulium ornatum* that have been immunologically identified as vitellogenin [23], we conclude that they represent the vitellogenin proteins of *S. vittatum*. The density of the vitellogenin bands increased during the first day after eclosion. The amount of protein in the 200 kDa band, determined using densitometry, increased linearly for 36 hours after eclosion (Fig. 3).

Ecdysteroid levels were measured using a competitive ELISA. Ecdysteroid levels increased rapidly and subsequently declined during the first day after eclosion (Fig. 4).

Discussion

These experiments demonstrate a correlation between growth of the oocyte, the appearance of vitellogenin in the ovary and rising titers of ecdysteroids in the black fly *S. vittatum*.

Previtellogenic growth of the follicle in *S. vittatum* takes about 12 hr. Ovarian follicles were observed to enter the vitellogenic stage at 14 hr after eclosion as indicated by the opaque appearance of the oocyte that occurs as yolk proteins are taken up. These observations correlate with those of Liu and Davies, [24]. Follicle growth was linear after eclosion. The oocyte reached maximum size by 72 hr after eclosion.

The appearance of opaque oocytes after 14 hr was correlated with an increase in the amount of vitellogenin present in oocytes.

Ecdysteroid levels were found to increase during the first day after eclosion. The changing titers of ecdysteroids were correlated with the appearance of vitellogenin in the oocytes and to the vitellogenic phase of oocyte growth. Vitellogenic growth of oocytes and vitellogenin synthesis are known to be stimulated by ecdysteroids in several cyclorrhaphid flies and mosquitoes [7–10,12]. This suggests that egg development in *S. vittatum* may also be regulated by ecdysteroids. Confirmation that ecdysteroids regulate vitellogenin synthesis in *S. vittatum* would require more direct evidence.

These observations are similar to those of the autogenous mosquito, *Ae. atropalpus*[14]. The timing of the growth of the follicle, and the rise in vitellogenin and ecdysteroids are similar in both species. In contrast, vitellogenic growth does not occur until after a blood meal in the anautogenous black fly, *Simulium ochraceum*[25], or the anautogenous mosquito *Ae. aegypti*[26]. In *Ae. aegypti* the vitellogenic period, including the rise in ecdysteroids and the synthesis of vitellogenin, does not begin until after the blood meal. Examining ecdysteroid titers after a blood



Figure 2
 SDS-PAGE Separation of vitellogenin protein in *Simulium vittatum* on SDS gels. Lane 1, molecular weight markers (200, 116, 97, 66 and 45 kDa). Lane 2, ovaries with proteinase inhibitors; lane 3, ovary extract without proteinase inhibitors. Ovaries were removed 36 hours after eclosion.




Figure 3
 Vitellogenin accumulation in *Simulium vittatum* ovaries after eclosion. Ovaries were homogenized in SDS sample buffer. Proteins were separated on SDS-PAGE and stained with Coomassie blue. The intensity of the stain in the 200 kDa protein band was determined by densitometry. Data are reported as the mean \pm standard deviation relative to the density of the 200 kDa band at 12 hours to correct for differences in staining intensity between gels. The number of band measurements used to calculate the mean is shown next to the data points.




Figure 4
 Ecdysteroid levels in *Simulium vittatum* with time after eclosion. Ecdysteroid levels were measured using a competitive ELISA as described by Kingan (21). Each point represents the average of 9 groups of 10 females, \pm standard deviation.

meal in an anautogenous black fly species would be of interest.

Conclusions

Our data suggest that microfilariae of *O. volvulus* would be exposed to changing titers of ecdysteroids during their development within the thoracic muscles. Development of microfilariae to the infective third stage takes 11 days in *Simulium ochraceum* during which time the host could complete 3 gonotrophic cycles [27], and 6 to 8 days in *Simulium damnosum* [28]. Given that the complete development of eggs in *S. vittatum* takes 3–4 days [29], the microfilariae might be exposed to several peaks of ecdysteroids during their development. It is possible that development of microfilariae is cued by the fluctuating titers of ecdysteroids in the host.

Methods

Blackflies, *Simulium vittatum*, were reared using the method described by Cupp and Ramberg [19]. Females were collected within 1 hr intervals under simulated natural conditions and stored in an incubator at 21°C. The adult females were fed 15% dextrose containing 1% streptomycin and mycostatin.

Follicle growth measurement

Ovaries were removed in *Aedes* saline [20] and follicles were teased apart using minuten needles. The follicle of the black fly is polytrophic and therefore contains an oocyte and accompanying nurse cells. Follicle and oocyte lengths were measured at different times after eclosion using a compound microscope and an ocular micrometer.

Ecdysteroid ELISA

Females were collected at various time points after adult eclosion, frozen, and homogenized in 50% MeOH (10 females / 600 ml). The homogenate was centrifuged and the supernatant was stored at -20°C. Ecdysteroid titers were assayed using a competitive ELISA as described by Kingan [21] using 20-hydroxyecdysone as the standard. The assay is about equally sensitive to ecdysone and 20-hydroxyecdysone. The latter is considered to be the active form of the hormone. The primary anti-ecdysone antibody was a generous gift of T.M. Kingan, University of California at Riverside. The secondary antibody was anti-rabbit IgG alkaline phosphatase conjugate. Activity was detected using p-nitrophenyl phosphatase [pNPP, Immunopure tablet from Pierce]. The absorbance was measured using an ELISA plate reader at 405 nm. Data analysis was performed as described by Kingan [21] using "Softmax" data analysis program (Molecular Devices Corp., Palo Alto, CA).

SDS-polyacrylamide gel electrophoresis

Females to be assayed were dissected in *Aedes* saline. Ovaries from two females were removed and stored at -70°C.

Ovaries were disrupted in 50 µl of sample buffer (5 mM Tris Base, 4% SDS, 0.002% Bromophenol Blue, 20% glycerol, pH 6.8, 2% mercaptoethanol) for 90 seconds in a microwave oven on high setting and then placed in boiling water for 1 minute at 100°C as described by Horscroft and Roy [22]. The mixture was then centrifuged in a microfuge for 3 minutes. Including protease inhibitors (aprotinin, chymostatin, antipain, leupeptin, all at 5 µg/ml, plus 1 mM PMSF and 5 mM benzamidine) had no effect on results presumably because boiling in SDS rapidly inhibited proteases (see Fig. 2). Proteins in the supernatant (5–20 µl) were separated electrophoretically using a vertical slab unit (Hofer Scientific Instruments) on 8.25% SDS-polyacrylamide gels. The molecular weight standard (Biorad, high molecular weight) had markers at 200, 116, 97, 66 and 45 kDa. Gels were stained with Coomassie Brilliant Blue and analyzed using a densitometer (Ultrascan XL Enhanced Laser Densitometer, Bromma). The density of the 65 kDa and 200 kDa vitellogenin bands were measured.

Abbreviations used

ELISA: enzyme linked immunosorbant assay

Competing interests

None declared

Acknowledgements

This work was supported by NIH grant number HD24869.

References

1. Nijhout HF: **Insect Hormones**. Princeton, Princeton University Press; 1994
2. Lawrence PO: **Hormonal effects on insects and other endoparasites in vitro**. *In Vitro Cell Dev Biol* 1991, **27A**:487-494
3. Franke S, Käuser G: **Occurrence and hormonal role of ecdysteroids in non-arthropods**. In: *From Chemistry to Mode of Action, Ecdysone* (Edited by: Koolman J) New York, Thieme Medical Publishers, Inc 1989, 296-307
4. Fleming MW: **Ascaris suum: Role of ecdysteroids in molting**. *Exp. Parasitol* 1985, **60**:207-210
5. Yates RA, Tuan RS, Shepley KJ, Unnasch TR: **Characterization of genes encoding members of the nuclear hormone-receptor superfamily from *Onchocerca volvulus***. *Mol Biochem Parasitol* 1995, **70**:19-31
6. Warbrick EV, Barker GC, Rees HH, Howells RE: **The effect of invertebrate hormones and potential hormone inhibitors on the 3rd larval molt of the filarial nematode, *Diriofilaria immitis*, in vitro**. *Parasitology* 1993, **107**:459-463
7. Schwartz MB, Kelly TJ, Woods CW, Imberski RB: **Ecdysteroid fluctuations in adult *Drosophila melanogaster* caused by elimination of pupal reserves and synthesis by early vitellogenic ovarian follicles**. *Insect Biochem* 1989, **19**:243-249
8. Kozlova T, Thummel CS: **Steroid regulation of postembryonic development and reproduction in *Drosophila***. *Trends Endocrinol Metabol* 2000, **11**:276-80
9. Chen AC, Kelly TJ: **Correlation of ecdysteroids with ovarian development and yolk protein synthesis in the adult stable fly, *Stomoxys calcitrans***. *Comp Biochem Physiol A-Physiol* 1993, **104**:485-490
10. Yin C-M, Zou B-X, Yi S-X, Stoffalano JG: **Ecdysteroid activity during oogenesis in the black blowfly, *Phormia regina***. *J Insect Physiol* 1990, **36**:375-382
11. Kapitskaya MZ, Li C, Miura K, Segraves W, Raikhel A: **Expression of the early-late gene encoding the nuclear receptor HR3 sug-**

- gests its involvement in regulating the vitellogenic response to ecdysone in the adult mosquito. *Molec Cell Endocrinol* 2000, **160**:25-37
12. Hagedorn HH: **The endocrinology of the adult female mosquito**, *Advan Disease Vector Res* 1994, **10**:109-148
 13. Masler EP, Fuchs MS, Sage B, O'Connor JD: **Endocrine regulation of ovarian development in the autogenous mosquito, *Aedes atropalpus***. *Gen Comp Endocrinol* 1980, **41**:250-259
 14. Lu YH, Hagedorn HH: **Egg development in the mosquito *Anopheles albimanus***. *Internat J Invert Reprod Develop* 1986, **9**:79-94
 15. Kelly TJ, Chen AC: **Cycling of ecdysteroid levels in adult female stable flies, *Stomoxys calcitrans* in relation to blood feeding**. *J Insect Physiol* 1997, **43**:789-794
 16. Hagedorn HH, O'Connor JD, Fuchs MS, Sage B, Schlaeger DA, Bohm MK: **The ovary as a source of alpha-ecdysone in an adult mosquito**. *Proc Natl Acad Sci* 1975, **72**:3255-3259
 17. Brenner RJ, Cupp EW: **Rearing blackflies (Diptera: Simuliidae) in a closed system of water circulation**. *Trop Med Parasitol* 1980, **31**:247-258
 18. Davies DM: **Ecology and behavior of adult blackflies (Simuliidae): a review**. *Quaestiones entomologicae* 1978, **14**:3-12
 19. Cupp EW, Ramberg FB: **are and maintenance of black fly colonies**. In: *Molecular Biology of Insect Disease Vectors: A Methods Manual* (Edited by: Crampton JM, Beard CB, Lewis C) New York, Chapman and Hall 1997, 31-40
 20. Hagedorn HH, Turner S, Hagedorn EA, Pontecorvo F, Greenbaum P, Pfeiffer D, Flanagan TR, Wheelock G: **Postemergence growth of the ovarian follicles of *Aedes aegypti***. *J Insect Physiol* 1977, **23**:203-206
 21. Kingan TG: **A competitive enzyme-linked immunosorbent assay: Applications in the assay of peptides, steroids and cyclic nucleotides**. *Anal Biochem* 1989, **183**:283-289
 22. Horscroft NJ, Roy P: **Thermal denaturation of proteins for SDS-PAGE analysis by microwave irradiation**. *BioTech* 1997, **22**:224-225
 23. Renshaw M, Hurd H: **Vitellogenin sequestration by *Simulium* oocytes: the effect of *Onchocerca* infection**. *Physiol Ento* 1994, **19**:70-74
 24. Liu TP, Davies DM: **Intramitochondrial transformations during lipid vitellogenesis in oocytes of a black fly, *Simulium vittatum***. *Inter J Insect Morph Embryol* 1973, **2**:233-245
 25. Cupp EW, Collins RC: **The gonotrophic cycle in *Simulium ochraceum***. *Am J Trop Med Hyg* 1979, **28**:422-426
 26. Yonge C, Hagedorn HH: **Dynamics of vitellogenin uptake in *Aedes aegypti* as demonstrated by trypan blue**. *J Insect Physiol* 1977, **23**:1199-1203
 27. Porter CH, Collins RC: **The gonotrophic cycle of wild *Simulium ochraceum* and the associated development of *Onchocerca volvulus***. *Trop Med Hyg* 1985, **34**:302-309
 28. Duke BOL: **Studies on factors influencing the transmission of onchocerciasis. II. The intake of microfilariae by *Simulium damnosum* and the survival of the parasite in the fly under laboratory conditions**. *Annal Trop Med Parasitol* 1962, **56**:255-263
 29. Edman JD, Simmons KR: **Rearing and colonization of blackflies (Diptera: Simuliidae)**. *J Med Entomol* 1985, **22**:1-17

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com