

ORIGINAL ARTICLE

Presenteeism in a Dutch hand eczema population— a cross-sectional survey

Jart A. F. Oosterhaven¹  | Peter A. Flach² | Ute Bültmann² | Marie L. A. Schuttelaar¹ 

¹Department of Dermatology, University of Groningen, University Medical Centre, Groningen, The Netherlands

²Department of Health Sciences, Community and Occupational Medicine, University of Groningen, University Medical Centre, Groningen, The Netherlands

Correspondence

Marie-Louise Anna Schuttelaar, Department of Dermatology, University Medical Centre Groningen, P.O. Box 30.001, 9700 RB Groningen, The Netherlands.
Email: m.l.a.schuttelaar@umcg.nl

Background: Presenteeism (attending work despite complaints and ill health, which should prompt rest and absence) has been overlooked in the field of hand eczema.

Objectives: To examine the 1-year prevalence of presenteeism related to hand eczema in a population of hand eczema patients who visited a tertiary referral centre. Secondary objectives: to identify intrinsic/extrinsic reasons for presenteeism and to evaluate associated factors.

Methods: This was a cross-sectional questionnaire study. Presenteeism was defined as “going to work despite feeling you should have taken sick leave because of hand eczema”. Respondents answered questions about socio-demographic factors, clinical features, occupational characteristics, and hand eczema related to occupational exposure.

Results: Forty-one per cent (141/346) of patients who had both worked and had hand eczema during the past 12 months reported presenteeism. The most often reported reasons were: “Because I do not want to give in to my impairment/weakness” (46%) and “Because I enjoy my work” (40%). Presenteeism was associated with: mean hand eczema severity; absenteeism because of hand eczema; improvement of hand eczema when away from work; and high-risk occupations.

Conclusions: In this study, presenteeism was common and predominantly observed in patients with more severe hand eczema and occupational exposure. The most frequently reported reasons for presenteeism were of an intrinsic nature.

KEYWORDS

absenteeism, hand eczema, occupational, presenteeism

1 | INTRODUCTION

Hand eczema is one of the most prevalent occupational skin diseases in Europe.^{1,2} In Germany, it is even the most frequently reported occupational disease.³ Hand eczema can lead to sickness absenteeism and eventually to job loss and change of profession.^{4–6} From other medical conditions, such as allergic rhinitis and arthritis, it is known that sickness absenteeism is often preceded by a phase in which workers try to continue their working activities, while their disease actually hampers their productivity and recovery.^{7–9} This phenomenon is called presenteeism: attending work despite complaints and ill health that should prompt rest and absence from work.¹⁰ Presenteeism can be regarded as a positive concept by workers with chronic conditions who are able to keep working.¹¹ However, in both the

medical and economic literature, presenteeism is mostly regarded as a negative and counterproductive phenomenon. Presenteeism received little attention for years, but has been increasingly studied in occupational medicine since the start of the 21st century. A recent review highlighted presenteeism as a risk factor for future sickness absence and decreased self-rated health.¹² Furthermore, it has been shown that presenteeism may be related to more productivity loss and higher costs than sickness absence in the long term.^{13,14}

Despite the fact that hand eczema is frequently caused or aggravated by occupational exposures,¹⁵ hand eczema-related presenteeism has hitherto not received much attention. Although a review from 2010 showed a significant impact of occupational contact dermatitis on work activities,¹⁶ to date only 1 study among patients with hand eczema has addressed presenteeism. Van der Meer et al studied

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. Contact Dermatitis published by John Wiley & Sons Ltd.

Dutch healthcare professionals with self-reported hand eczema. They considered presenteeism to be “lost time at work” (in terms of amount and quality of work performed). The 1-year prevalence of hand eczema in the healthcare professionals was relatively low (12%); of those with hand eczema, 3.1% reported presenteeism and 1.7% reported sickness absence because of hand eczema.¹⁷ To date, little is known about presenteeism in patients with more severe hand eczema, working in various occupations. Therefore, the aim of this study was to examine the 1-year prevalence of presenteeism related to hand eczema in a population of hand eczema patients who visited a tertiary referral centre. Secondary objectives were to identify intrinsic or extrinsic reasons for presenteeism, and to evaluate factors associated with the prevalence of presenteeism in hand eczema patients.

2 | METHODS

2.1 | Study design

This was a cross-sectional study carried out at the Department of Dermatology of the University Medical Centre Groningen, a tertiary referral centre for hand eczema. The population of hand eczema patients that visits the department predominantly lives in the 5 northern provinces of The Netherlands (population approximately 3.2 million). Patients were identified by searching electronic patient records from visits between January 1, 2011 and December 31, 2015. Identified patients received a postal questionnaire. In parallel, patients who visited our outpatient clinic and were diagnosed with hand eczema were also recruited. These patients completed the questionnaire digitally on-site. The questionnaire was developed to assess the prevalence of presenteeism, intrinsic or extrinsic reasons for presenteeism, and factors associated with presenteeism prevalence. Before the start of the study, a pilot study was conducted in 5 hand eczema patients to finalize the questionnaire. The study was reviewed and approved by the Medical Ethical Review Board of the University Medical Centre Groningen (reference METc 2016/169).

2.2 | Study population and recruitment

Patients of working age, between 20 and 67 years, with hand eczema diagnosed by a dermatologist in the past 5 years were included. The diagnosis was made in accordance with guidelines by Menné et al and Diepgen et al.^{18,19}

Patients were identified from electronic records by use of the International Classification of Disease (ICD, 10th edition), according to the diagnoses L20 (Atopic dermatitis), L23 (Allergic contact dermatitis), L24 (Irritant contact dermatitis), L25 (Unspecified contact dermatitis), and L30 (Other dermatitis). This yielded a total of 1168 patients. One author (J.O.) manually screened these files and identified a total of 789 patients in whom the diagnosis of hand eczema could be confirmed. The other 379 patients were excluded, mainly because the ICD codes were not specific for hand eczema and the dermatitis occurred on body areas other than the hands in these patients. For a study flow diagram, see Figure 1.

In June 2016, an invitation letter, a questionnaire and a prepaid return envelope were sent to the 789 eligible patients. A reminder was sent after 5 weeks. The recruitment of the on-site patients took place between June 2016 and March 2017. A total of 115 patients were approached to complete the questionnaire. For data entry by participants on-site and data entry of the postal questionnaires, the online survey software QUALTRICS was used (Qualtrics, Provo, Utah; <http://www.qualtrics.com>). To improve the response rate, 10 gift coupons of €50 were raffled among the participants who responded to the postal questionnaire.

2.3 | Measurements

All concepts are briefly described below. For a comprehensive overview of the definitions and categorization for the analyses, see Appendix S1. All variables concern the past 12 months unless otherwise indicated.

Presenteeism. Patients were asked whether they had both worked and had hand eczema during the past 12 months. In these patients, presenteeism was assessed with the question: “During the past 12 months, did you go to work despite feeling that you should have taken sick leave because of your hand eczema? Yes/no.” The duration of presenteeism was also measured.²⁰

Reasons for presenteeism. Intrinsic and/or extrinsic reasons for presenteeism were measured with the following question: “You indicated that during the past 12 months you went to work despite feeling that you should have taken sick leave because of your hand eczema. What was the reason for this? (multiple answers possible).” Answer categories were assembled from Johansen et al,²¹ Johns et al,²² and Aronsson et al.¹⁰ Following the pilot study, 2 answer categories were added: “Because I think it is expected of me” and “Because I don’t want to give in to my affliction/weakness”.

Socio-demographic factors. Sex; age at questionnaire completion; and education (low/middle or high).

Clinical features. First episode of hand eczema ≤ 18 years;²³ atopic dermatitis ever;²³ mean hand eczema severity, which was determined with the photographic guide developed by Coenraads et al,^{24,25} and other longstanding diseases.

Occupational characteristics. Type of employment (paid employed/self-employed);²³ hours per week; sufficient time at work to perform tasks satisfactorily;²⁶ sufficient resources at work to perform tasks satisfactorily;²⁶ number of employees; supervising tasks ([non]-management);²⁷ shift work;²⁷ high-risk occupation;^{28–33} and monthly income.²¹

Hand eczema related to occupational exposure. Absenteeism because of hand eczema; improvement of hand eczema when away from work;²³ hand eczema related to occupational exposure;²³ and wet work, which was determined according to the German “Technische Regeln für Gefahrstoffe” (TRGS) 401 criteria³⁴ and work by Behroozy et al.³⁵

Covariables. Frequency of hand eczema ([nearly] all the time or more than once);²³ months worked; and job loss or early retirement because of hand eczema.




FIGURE 1 Study flow diagram. HE, hand eczema; ICD, International Classification of Disease

2.4 | Statistical analysis

Before the analyses were performed, 3 preparatory steps were taken. First, to handle missing values, each completed postal questionnaire was screened. When missing values were found, the sender was contacted by telephone or email to obtain an answer. In this way, all missing data were retrieved. The design of the digital questionnaire did not allow for missing data. Second, respondents and non-respondents to the postal questionnaire were compared in a non-response analysis. Third, respondents who completed the postal questionnaire were compared with respondents who completed the questionnaire digitally on-site. Descriptive statistics were reported by the use of mean and standard deviation for normally distributed

variables, and median and interquartile range (IQR) for non-normally distributed variables. For comparison of normally and non-normally distributed variables, respectively, Student's *t* test and the Mann-Whitney *U* test were used. Proportions were compared by use of the χ^2 test.

Logistic regression analysis was used to evaluate associations with presenteeism prevalence. All variables that were significant in the univariable analysis were entered into a multivariable model. A post hoc sample size calculation for logistic model building was performed. The method based on the work by Peduzzi et al³⁶ states that: $N = 10 k/p$. Here, p is the smallest of the proportions of negative or positive cases in the population, and k is the number of covariates (the number of independent variables). To build a model with 12 independent variables

in a sample with a presenteeism proportion of 0.40, the necessary minimal sample size is $N = 300$. Odds ratios (ORs) with 95% confidence intervals were calculated. Analyses were performed with IBM SPSS STATISTICS for Windows, Version 23.0 (IBM Corp., Armonk, New York). A P value of $<.05$ was regarded as being statistically significant.

A sensitivity analysis was conducted to examine the influence of income on presenteeism in the multivariable model. Information on income was not provided by all respondents (respondents choosing the options "I don't know" or "I would rather not answer this question"), so, for the analysis, these response options were regarded as missing. As logistic regression analysis only handles complete cases, income was left out of the main multivariable model. In a subgroup analysis, reasons for presenteeism were assessed for the digital questionnaire on-site group to check whether reasons in this group differed from those in the total group.

3 | RESULTS

3.1 | Sample characteristics

Data were collected from 500 participants, aged between 20 and 67 years. The response to the postal questionnaire was 48.8%. The non-response analysis showed no sex difference, but non-respondents were significantly younger than respondents: 40 years (IQR 29.0-52.0) vs 49 years (IQR 36.0-58.0); $P < .01$. Respondents to the digital questionnaire on-site were significantly younger than respondents to the postal questionnaire. Also, during the past 12 months, their hand eczema prevalence was higher, they had more severe hand eczema, they worked during fewer months, and they called in sick more often because of their hand eczema (Table 1).

3.2 | Presenteeism prevalence and reasons for presenteeism

In the analyses of presenteeism prevalence, 346 respondents were included because they had both worked and had hand eczema during the past 12 months. A total of 141 (40.8%) respondents reported that they had experienced presenteeism during the past year. No significant difference in presenteeism prevalence was observed between the respondents to the digital and postal questionnaires.

Table 2 shows the reported reasons for presenteeism with a differentiation between 5 intrinsic and 11 extrinsic reasons. In total, 505 answers were provided by the 141 respondents who reported presenteeism (175 intrinsic; 308 extrinsic; 22 other reasons that were not categorized). The median number of reasons reported was 3 (IQR 2-5). Two intrinsic reasons for presenteeism were mentioned most often: "I do not want to give in to my impairment/weakness" (46.1% of respondents) and "I enjoy my work" (39.7% of respondents). Notably, 22.7% of respondents went to work because they were "afraid of losing their job". Self-employed respondents ($n = 29$) were more inclined to choose extrinsic reasons, with financial motives and irreplaceable responsibilities being the most often reported reasons (total reasons 85; 28 intrinsic reasons [33%], and 57 extrinsic reasons

[67%]). Paid employed respondents ($n = 112$) chose intrinsic reasons more often than self-employed individuals (total reasons 398; 147 intrinsic reasons [37%], and 251 extrinsic reasons [63%]). See Appendix S2 for all reported reasons for both groups.

3.3 | Factors associated with presenteeism

In a univariable logistic regression analysis (Table 3), significant associations were found between presenteeism and variables from all 4 domains (socio-demographics, clinical features, occupational characteristics, and hand eczema related to occupational exposure). Significant variables were included in the multivariable model. For income, see sensitivity analyses.

In the multivariable regression model (Table 4), more severe hand eczema; absence because of hand eczema in the past 12 months, hand eczema that improved when away from work; and working in a high-risk occupation were significantly associated with presenteeism prevalence.

3.4 | Sensitivity and subgroup analyses

A higher income was significantly associated with a lower presenteeism prevalence in the univariable analysis. Eighty-eight respondents chose not to disclose information about their income or could not answer this question. When income was added to the multivariable model in Table 4, it was no longer significantly associated with presenteeism. The effect of having a high-risk occupation attenuated and became non-significant ($P = .07$, $n = 258$). The other significant independent variables remained significant; see Appendix S3.

In a subgroup analysis, the distribution of reasons for presenteeism in the digital questionnaire on-site group was assessed. This showed a very similar ranking to that in the whole group. The same 2 intrinsic reasons for presenteeism were most frequently reported: "Because I do not want to give in to my impairment/weakness" (57.7%) and "Because I enjoy my work" (40.9%). See Appendix S3.

4 | DISCUSSION

In this study, we found a 1-year prevalence of 41% for presenteeism because of hand eczema. Intrinsic reasons for presenteeism were common, and the phenomenon was strongly associated with severity and occupational characteristics.

More than 40% of the study population indicated that they had experienced presenteeism because of hand eczema in the past 12 months. This proportion is strikingly higher than the low percentage (3%) that was found by van der Meer et al in their study in Dutch healthcare workers.¹⁷ The discrepancy could partly be explained by the probably mild hand eczema severity of their homogeneous working population, which might have minimized the interference with work, as compared with our occupationally heterogeneous patient population with much more severe hand eczema. Another explanation could be the different definition of presenteeism that was used. We focused on whether respondents felt that they should have called in sick although they did not (subjectively), instead of asking

TABLE 1 Basic characteristics of the total study population (N = 500) and a comparison between respondents to the postal questionnaire and respondents to the digital questionnaire on-site

	Total, N = 500	Respondents to postal questionnaire, n = 385	Respondents to digital questionnaire on-site, n = 115	P value
Socio-demographics				
Female sex, % (n/n _{total})	55.8 (279/500)	57.7 (222/385)	49.6 (57/115)	.13
Age (years), median (IQR)	46.5 (34.0-57.0)	49.0 (36.0-58.0)	40.0 (27.0-53.0)	<.001
Education, % (n/n_{total})				
Low/middle	75.4 (377/500)	73.8 (284/385)	80.9 (93/115)	.14
High	24.6 (123/500)	26.2 (101/385)	19.1 (22/115)	
Clinical features, % (n/n_{total})				
First episode of HE ≤18 years	74.8 (374/500)	76.4 (294/385)	69.6 (80/115)	.14
Atopic dermatitis, ever	47.6 (238/500)	45.5 (175/385)	54.8 (63/115)	.09
HE during the past 12 months	86.2 (431/500)	82.3 (317/385)	99.1 (114/115) ^a	<.001
HE (nearly) all the time or more than once	86.3 (372/431)	84.5 (268/317)	91.2 (104/114)	.08
Mean HE severity				
				<.001
No HE during past 12 months	11.2 (56/500)	14.5 (56/385) ^a	0 (0/115) ^a	
Almost clear	25.4 (127/500)	29.9 (115/385)	10.4 (12/115)	
Moderate	38.2 (191/500)	35.6 (137/385)	47.0 (54/115)	
Severe	20.6 (103/500)	17.1 (66/385)	32.2 (37/115)	
Very severe	4.6 (23/500)	2.9 (11/385)	10.4 (12/115)	
Occupational characteristics				
Not employed during past 12 months, % (n/n _{total})	19.7 (85/431)	19.9 (63/317)	19.3 (22/114)	.81
Paid employed, % (n/n_{total})				
Months worked, median (IQR)	12 (12-12)	12 (12-12)	12 (11-12)	.03
Work hours per week, mean (±SD)	31.0 (±12.5)	30.8 (±12.3)	31.5 (±12.9)	.67
Self-employed, % (n/n_{total})				
Months worked, median (IQR)	12 (10-12)	12 (10-12)	12 (10.5-12)	.33
Work hours per week, mean (±SD)	37.5 (±20.3)	37.4 (±19.2)	37.6 (±24.2)	.98
High-risk occupation, % (n/n _{total}) ^b	43.6 (151/346)	42.9 (109/254)	45.7 (42/92)	.71
Monthly income, % (n/n_{total})				
Low	45.7 (118/258)	46.2 (86/186)	44.4 (32/72)	.89
Middle/high	54.3 (140/258)	53.8 (100/186)	55.6 (40/72)	
HE related to occupational exposure, % (n/n_{total})				
Worked during the past 12 months while having had HE during the past 12 months	80.3 (346/431)	80.1 (254/317)	80.7 (92/114)	.89
Absenteeism because of HE during the past 12 months	16.2 (56/346)	12.2 (31/254)	27.2 (25/92)	.001
Improvement of HE when away from work	45.4 (157/346)	48.0 (122/254)	38.0 (35/92)	.11
HE related to occupational exposure				
Wet work	50.3 (174/346)	52.4 (133/254)	44.6 (41/92)	.22
Job loss or early retirement because of HE during the past 12 months	46.2 (160/346)	45.7 (116/254)	47.8 (44/92)	.81
	4.0 (20/500)	3.1 (12/385)	7.0 (8/115)	.10
Presenteeism, % (n/n_{total})				
Prevalence during the past 12 months	40.8 (141/346)	38.2 (97/254)	47.8 (44/92)	.11
Duration of >42 days	17.0 (24/141)	17.5 (17/97)	15.9 (7/44)	.81

HE, hand eczema; IQR, interquartile range; SD, standard deviation.

^a Thirteen respondents responded inconsistently to the questions on the presence of hand eczema in the past 12 months and mean severity over the past 12 months. These patients were not included in further analyses.

^b See Appendix S4 for an overview of the occupations in the sample.

for lost time at work in terms of amount and quality of work performed (a somewhat more objective measure). However, the difference is large, indicating that the phenomenon is indeed quite

common in our patient population. A comparison with presenteeism in patients with other chronic diseases is difficult, mainly because many studies measure presenteeism as a decline in productivity or

TABLE 2 Intrinsic and extrinsic reasons for presenteeism in 141 workers with hand eczema

Reasons for presenteeism	Intrinsic/extrinsic motivation	n (%)
Because ...		
... I do not want to give in to my impairment/weakness	Intrinsic	65 (46.1)
... I enjoy my work	Intrinsic	56 (39.7)
... I think it is expected of me	Extrinsic	52 (36.9)
... I do not want to burden my colleagues	Extrinsic	41 (29.1)
... I do not want to be considered lazy or unproductive	Extrinsic	37 (26.2)
... financially I cannot afford taking sick leave	Extrinsic	32 (22.7)
... I am afraid of losing my job	Extrinsic	32 (22.7)
... my pride keeps me from calling in sick	Intrinsic	28 (19.9)
... my employer expects it of me	Extrinsic	25 (17.7)
... no one else can take over my responsibilities	Extrinsic	25 (17.7)
... I have appointments with clients/patients	Extrinsic	23 (16.3)
... I need to catch up on a lot of work if I have been sick	Extrinsic	17 (12.1)
... going to work is good for my health	Intrinsic	14 (9.9)
... I do not want to be suspected of cheating	Extrinsic	13 (9.2)
... I want to maintain my social network	Intrinsic	12 (8.5)
... I feel ashamed to call in sick	Extrinsic	11 (7.8)
Other reasons ^a		22 (15.6)

The total percentage exceeds 100% because subjects were permitted to choose multiple reasons.

^a Other reported reasons were: "I don't 'feel sick'" (6); "I am self-employed" (2); "other work could (temporarily) replace my normal work" (2); "it never occurred to me to call in sick for hand eczema" (2); "I work with livestock" (2); "I didn't want to get in trouble over calling in sick; I could easily work from home; I can't sit still; I don't have enough insurance and can't afford employees; re-organization at work; I don't consider hand eczema a reason for calling in sick; people don't take eczema seriously so I don't want to be considered a fraud; it doesn't match my character to call in sick".

quality of work, using various measurement instruments. The problem here is that there are still no generally accepted measurement instruments with which to assess presenteeism. A large range of existing instruments were found to be insufficiently validated.³⁷ With these instruments, a wide range of presenteeism (19%–79%) was found in studies in several chronic diseases, including systemic lupus erythematosus, spondyloarthritis, inflammatory bowel disease, low back pain, and rheumatic arthritis.^{38–44} A recent study with a definition more comparable to that used in the present study was performed by d'Errico et al in a sample of the European working population. The authors asked >30 000 workers the question "Over the past 12 months did you work when you were sick?". For 6 health problems (insomnia, stomach pain, headache, upper arm pain, back pain, and wounds), they found rather similar percentages of presenteeism, ranging between 42.3% (back pain) and 52.2% (insomnia).⁴⁵ The

percentage found in our population is slightly lower. This could be explained by the fact that we did not mark every respondent who worked with hand eczema as having presenteeism. We incorporated an additional aspect in our definition of presenteeism by including the phrase "... despite feeling you should have taken sick leave because of your hand eczema". This could be regarded by respondents as a proxy for expected productivity loss ("I am going to work although I think I will be less productive because of my hand eczema today"). It could also be regarded as resulting from fear that their hands would become much worse if they worked, regardless of whether this was at the expense of their productivity ("I am going to work although I think my hand eczema will get worse due to my working activities today"). These 2 explanations might influence each other greatly and may very often coexist. New and better-validated instruments are needed to assess productivity loss caused by presenteeism.^{37,46} Once these reliable and valid instruments become available, future studies should focus on determining whether presenteeism in patients with hand eczema actually causes productivity loss.

A notable finding in our study is the intrinsic nature of the most often reported reasons for presenteeism. This was also found by Robertson et al⁴⁷ and partly by Johansen et al ("enjoying work" was the second most reported reason in their study, after "don't want to burden my colleagues").²¹ It seems that presenteeism in our patients is often self-imposed. This is an important finding, especially in patients with occupational hand eczema. It is easily assumed that the tendency to attend work, although being detrimental to hand eczema, is mainly financially driven or forced by third parties. Our study, however, shows that, although feelings of external pressure along with financial motives exist, intrinsic reasons are even more important for many individuals. This does not apply to self-employed individuals, in whom extrinsic reasons are more common than in paid employees. Patients should be made aware of the fact that presenteeism can lead to deterioration of hand eczema. Along with their occupational physician, patients need to determine their occupational exposure and the feasibility of using adequate protection against hazardous exposure in their work. To enable adequate assessment of this, the dermatologist should provide the occupational physician with information about the sensitization profile from patch testing and about the tolerance of the skin of the hands to irritants, and its repair capacity after exposure to irritants.³⁵

Although intrinsic reasons are most often reported, it is important to note that more than 1 in 4 respondents reported "Because I am afraid of losing my job" as a reason for presenteeism. This figure is much higher than the 4% reported by Johansen et al in a sample of the general working population with unspecified medical conditions in Norway and Sweden.²¹ It seems that having hand eczema causes patients much worry about being able to keep their jobs.

Strong associations were found between presenteeism prevalence and mean hand eczema severity, absenteeism because of hand eczema, improvement of hand eczema when away from work, and working in a high-risk occupation. The association of these factors provides a picture of a predominance of patients who experience hand eczema-related presenteeism because of more severe hand eczema that is at least partly caused or aggravated by work. Many of

TABLE 3 Univariable logistic regression analysis for the association of factors within 4 domains with presenteeism

	Total % (n/n _{total})	Presenteeism % (n/n _{total})	Crude OR (95%CI)	P value
Socio-demographics				
Sex				
Male	48.3 (167/346)	40.1 (67/167)	1.00 (ref.)	.82
Female	51.7 (179/356)	41.3 (74/179)	1.05 (0.69-1.62)	
Age group (y)				
20-35	33.2 (115/346)	46.1 (53/115)	1.00 (ref.)	.17
36-50	32.4 (112/346)	33.9 (38/112)	0.60 (0.35-1.03)	
51-67	34.4 (119/346)	42.0 (50/119)	0.85 (0.51-1.42)	
Education				
Low/middle	74.6 (258/346)	44.6 (115/258)	1.00 (ref.)	.01
High	25.4 (88/346)	29.5 (26/88)	0.52 (0.31-0.87)	
Clinical features				
First episode of HE				
≤18 years	29.5 (102/346)	39.2 (40/102)	1.00 (ref.)	.71
>18 years	70.5 (244/346)	41.4 (101/244)	1.10 (0.68-1.76)	
Atopic dermatitis ever				
No	50.0 (173/346)	34.7 (60/173)	1.00 (ref.)	.02
Yes	50.0 (173/346)	46.8 (81/173)	1.66 (1.08-2.56)	
Mean HE severity				
Almost clear	30.3 (105/346)	13.3 (14/105)	1.00 (ref.)	<.001
Moderate	43.1 (149/346)	42.3 (63/149)	4.76 (2.49-9.12)	
Severe	22.5 (78/346)	70.5 (55/78)	15.54 (7.39-32.71)	
Very severe	4.0 (14/346)	64.3 (9/14)	11.70 (3.42-40.01)	
Other longstanding diseases				
No	68.5 (237/346)	41.8 (99/237)	1.00 (ref.)	.57
Yes	31.5 (109/346)	38.5 (42/109)	0.87 (0.55-1.39)	
Occupational characteristics				
Type of employment				
Paid employed	79.2 (274/346)	40.9 (112/274)	1.00 (ref.)	.93
Self-employed	20.8 (72/346)	40.3 (29/72)	0.98 (0.58-1.66)	
Mean weekly working hours				
≤23	24.6 (85/346)	47.1 (40/85)	1.00 (ref.)	.39
24-35	31.2 (108/346)	38.0 (41/108)	0.69 (0.39-1.23)	
≥36	44.2 (153/346)	39.2 (60/153)	0.73 (0.43-1.24)	
Sufficient time at work				
No	9.8 (34/346)	64.7 (22/34)	1.00 (ref.)	.004
Yes	90.2 (312/346)	38.1 (119/312)	0.34 (0.16-0.71)	
Sufficient resources at work				
No	4.6 (16/346)	75.0 (12/16)	1.00 (ref.)	.009
Yes	95.4 (330/346)	39.1 (129/330)	0.21 (0.07-0.68)	
Number of employees				
Self-employed	11.8 (41/346)	39.0 (16/41)	1.00 (ref.)	.61
1-9	15.3 (53/346)	49.1 (26/53)	1.51 (0.66-3.44)	
10-99	29.5 (102/346)	40.2 (41/102)	1.05 (0.50-2.21)	
≥99	43.4 (150/346)	38.7 (58/150)	0.99 (0.49-2.00)	
Supervising tasks				
Non-management	81.2 (281/346)	40.2 (113/281)	1.00 (ref.)	.67
Middle management/executive	18.8 (65/346)	43.1 (28/65)	1.13 (0.66-1.94)	
Shift work				
No	79.2 (274/346)	40.1 (110/274)	1.00 (ref.)	.66
Yes	20.8 (72/346)	43.1 (31/72)	1.13 (0.67-1.91)	
High-risk occupation				
No	56.4 (195/346)	33.3 (65/195)	1.00 (ref.)	.002

TABLE 3 (Continued)

	Total % (n/n _{total})	Presenteeism % (n/n _{total})	Crude OR (95%CI)	P value
Yes	43.6 (151/346)	50.3 (76/151)	2.02 (1.31-3.14)	
Income				.04
Low	45.7 (118/258)	50.0 (59/118)	1.00 (ref.)	
Middle/high	54.3 (140/258)	37.1 (52/140)	0.59 (0.36-0.97)	
HE related to occupational exposure				<.001
Absenteeism because of HE				<.001
No	83.8 (290/346)	32.1 (93/290)	1.00 (ref.)	
Yes	16.2 (56/346)	85.7 (48/56)	12.7 (5.78-28.0)	
Improvement of HE when away from work				<.001
No	54.6 (189/346)	31.7 (60/189)	1.00 (ref.)	
Yes	45.4 (157/346)	51.6 (81/157)	2.29 (1.48-3.55)	
HE related to occupational exposure				.002
No	49.7 (172/346)	32.6 (56/172)	1.00 (ref.)	
Yes	50.3 (174/346)	48.9 (85/174)	1.98 (1.28-3.06)	
Wet work				.05
No	53.8 (186/346)	36.0 (67/186)	1.00 (ref.)	
Yes	46.3 (160/346)	46.3 (74/160)	1.53 (0.99-2.35)	

CI, confidence interval; HE, hand eczema; OR, odds ratio.

TABLE 4 Multivariable logistic regression model for presenteeism

	Mutually adjusted OR (95%CI)	P value
Mean HE severity		<.001
Almost clear	1.00 (ref.)	
Moderate	5.52 (2.63-11.6)	
Severe	17.6 (7.32-42.4)	
Very severe	6.80 (1.57-29.4)	
Absenteeism because of HE		<.001
No	1.00 (ref.)	
Yes	9.96 (4.06-24.5)	
Improvement of HE when away from work		.009
No	1.00 (ref.)	
Yes	2.20 (1.22-3.96)	
High-risk occupation		.009
No	1.00 (ref.)	
Yes	2.14 (1.21-3.78)	
Sufficient time at work		.15
No	1.00 (ref.)	
Yes	0.44 (0.15-1.33)	
Sufficient resources at work		.19
No	1.00 (ref.)	
Yes	0.34 (0.07-1.71)	
HE related to exposure at work		.24
No	1.00 (ref.)	
Yes	1.42 (0.79-2.53)	
Atopic dermatitis ever		.53
No	1.00 (ref.)	
Yes	1.19 (0.69-2.06)	
Education		.60
Low/middle	1.00 (ref.)	
High	1.20 (0.61-2.32)	

CI, confidence interval; HE, hand eczema; OR, odds ratio.

these patients had already called in sick on several occasions because of hand eczema, but probably only to give their hands a little rest. From other diseases, we know that this kind of behaviour can eventually cause long-term absence if it is continued for some time.¹² Interestingly, in the group of patients with very severe hand eczema, the adjusted OR for presenteeism prevalence is lower than in the severe group. This is possibly because the patients with such very severe disease eventually do call in sick for a longer time (data not shown). Notably, self-employment was not associated with presenteeism. This is probably related to the importance of intrinsic motivations that drive workers to attend work while having hand eczema, regardless of employment status.

A limitation of our study is that we chose not to incorporate psychosocial factors, mainly because this would further increase the amount of items in the questionnaire. By asking our question about presenteeism specifically in relationship to hand eczema, we mainly addressed the health state aspect of presenteeism. However, the question can be raised of whether to look at presenteeism as a health state phenomenon (originating mainly from the medical condition) or rather as behaviour (implying that a choice is made, supported by psychosocial characteristics). A recent meta-analysis found evidence for both.⁴⁸ This is in agreement with Brooks et al, who stated that presenteeism should be approached as a complex system, with incorporation of variables from both the health state and a behavioural point of view.⁴⁹

Another possible limitation can be found in the definition of high-risk occupations. For certain occupations, working activities are quite well known and similar between different workers in this branch (eg, hairdressers and bakers). For other occupations, much more variation exists (eg, in healthcare workers). To precisely determine whether an individual is working in a high-risk occupation, the job content, working process and exposure levels are more important than the job title.

Methodologically, a possible limitation could be common method variance; we measured the outcome and all associated variables with the same self-report questionnaire. Furthermore, it is possible that non-response bias in the postal questionnaire group could have influenced the results of this study. Unfortunately, we only had sex and age of the postal non-respondents available, so it is not possible to draw conclusions about the presence of non-response bias.

Data were collected with 2 different methods (postal and digital). The postal and digital on-site respondents were combined for analysis. Age, mean severity of hand eczema, absenteeism because of hand eczema and months worked during the past 12 months significantly differed between the postal and digital on-site respondents. This was expected regarding severity and absenteeism, as the digital respondents were all patients who sought care from a dermatologist when completing the questionnaire, whereas the postal respondents had visited the department at some point during the past 5 years. Nevertheless, presenteeism prevalence was not significantly different between the groups. This, along with similarities in several other variables (eg, sex, education, occupational characteristics, and, especially, whether respondents had both worked and had hand eczema during the past 12 months), led us to combine the respondent groups. Also, reasons for presenteeism in the digital respondent group showed a very similar ranking to those of the whole group in the subgroup analysis.

Finally, we did not control for a possible influence of certain lifestyle factors, such as smoking, high body mass index, or alcohol use, which have been identified as possible risk factors for presenteeism.⁵⁰

In this study, we have shown that presenteeism is a common phenomenon in patients with more severe hand eczema. The most frequently reported reasons for presenteeism were of an intrinsic nature. Dermatologists and occupational physicians should pay attention to presenteeism to provide more individually targeted care for hand eczema patients.

CONFLICT OF INTEREST

There was no funding and the authors report no conflicts of interest, either actual or perceived.

ORCID

Jart A. F. Oosterhaven  <http://orcid.org/0000-0001-7214-7528>

Marie L. A. Schuttelaar  <http://orcid.org/0000-0002-0766-4382>

REFERENCES

- European Commission Employment SA and EO. *Eurostat Statistical Books. Health and Safety at Work in Europe (1999-2007). A Statistical Portrait*. Luxembourg: Publications Office of the European Union, 2010, p. 58 [Internet]. <http://ec.europa.eu/eurostat/documents/3217494/5718905/KS-31-09-290-EN.PDF/88eef9f7-c229-40de-b1cd-43126bc4a946> Accessed March 12, 2018.
- Mahler V, Aalto-Korte K, Alfonso JH, et al. Occupational skin diseases: actual state analysis of patient management pathways in 28 European countries. *J Eur Acad Dermatol Venereol*. 2017;31(suppl 4):12-30.
- Deutsche Gesetzliche Unfallversicherung. DGUV Statistics 2016. <http://publikationen.dguv.de/dguv/pdf/10002/12640neu.pdf>.
- Cvetkovski RS, Rothman KJ, Olsen J, et al. Relation between diagnoses on severity, sick leave and loss of job among patients with occupational hand eczema. *Br J Dermatol*. 2005;152:93-98.
- Thyssen JP, Johansen JD, Linneberg A, Menné T. The epidemiology of hand eczema in the general population—prevalence and main findings. *Contact Dermatitis*. 2010;62:75-87.
- Petersen AH, Johansen JD, Hald M. Hand eczema—prognosis and consequences: a 7-year follow-up study. *Br J Dermatol*. 2014;171:1428-1433.
- Schultz AB, Edington DW. Employee health and presenteeism: a systematic review. *J Occup Rehabil*. 2007;17:547-579.
- Bergström G, Bodin L, Hagberg J, Aronsson G, Josephson M. Sickness presenteeism today, sickness absenteeism tomorrow? A prospective study on sickness presenteeism and future sickness absenteeism. *J Occup Environ Med*. 2009;51:629-638.
- Hansen CD, Andersen JH. Sick at work—a risk factor for long-term sickness absence at a later date? *J Epidemiol Community Health*. 2009;63:397-402.
- Aronsson G, Gustafsson K, Dallner M. Sick but yet at work. An empirical study of sickness presenteeism. *J Epidemiol Community Health*. 2000;54:502-509.
- Hansen CD. "Making a virtue" of going ill to work—reflections on the necessity of everyday workplace "suffering". *Tidsskr Forsk i Sygd og Samf*. 2010;13:69-88.
- Skagen K, Collins AM. The consequences of sickness presenteeism on health and wellbeing over time: a systematic review. *Soc Sci Med*. 2016;161:169-177.
- Goetzel RZ, Long SR, Ozminkowski RJ, Hawkins K, Wang S, Lynch W. Health, absence, disability, and presenteeism cost estimates of certain physical and mental health conditions affecting US employers. *J Occup Environ Med*. 2004;46:398-412.
- Collins JJ, Baase CM, Sharda CE, et al. The assessment of chronic health conditions on work performance, absence, and total economic impact for employers. *J Occup Environ Med*. 2005;47:547-557.
- Böhm D, Stock Gissendanner S, Finkeldey F, et al. Severe occupational hand eczema, job stress and cumulative sickness absence. *Occup Med (Lond)*. 2014;64:509-515.
- Lau MY, Burgess JA, Nixon R, Dharmage SC, Matheson MC. A review of the impact of occupational contact dermatitis on quality of life. *J Allergy (Cairo)*. 2011;2011:964509.
- van der Meer EWC, Boot CRL, van der Gulden JWW, Jungbauer FH, Coenraads PJ, Anema JR. Hand eczema among healthcare professionals in the Netherlands: prevalence, absenteeism, and presenteeism. *Contact Dermatitis*. 2013;69:164-171.
- Diepgen TL, Elsner P, Schliemann S, et al. Guideline on the management of hand eczema ICD-10 code: L20. L23. L24. L25. L30. *J Dtsch Dermatol Ges*. 2009;7(suppl 3):S1-S16.
- Menné T, Johansen JD, Sommerlund M, Veien NK. Hand eczema guidelines based on the Danish guidelines for the diagnosis and treatment of hand eczema. *Contact Dermatitis*. 2011;65:3-12.
- Taloyan M, Aronsson G, Leineweber C, Magnusson Hanson L, Alexanderson K, Westerlund H. Sickness presenteeism predicts sub-optimal self-rated health and sickness absence: a nationally representative study of the Swedish working population. *PLoS One*. 2012;7:e44721.
- Johansen V, Aronsson G, Marklund S. Positive and negative reasons for sickness presenteeism in Norway and Sweden: a cross-sectional survey. *BMJ Open*. 2014;4:e004123-2013-004123.
- Johns G. Presenteeism in the workplace: a review and research agenda. *J Organiz Behav*. 2010;31:519-542.
- Susitaival P, Flyvholm MA, Meding B, et al. Nordic occupational skin questionnaire (NOSQ-2002): a new tool for surveying occupational skin diseases and exposure. *Contact Dermatitis*. 2003;49:70-76.
- Coenraads PJ, van der Walle H, Thestrup-Pedersen K, et al. Construction and validation of a photographic guide for assessing severity of chronic hand dermatitis. *Br J Dermatol*. 2005;152:296-301.
- Hald M, Veien NK, Laurberg G, Johansen JD. Severity of hand eczema assessed by patients and dermatologist using a photographic guide. *Br J Dermatol*. 2007;156:77-80.

26. Hansen CD, Andersen JH. Going ill to work—what personal circumstances, attitudes and work-related factors are associated with sickness presenteeism? *Soc Sci Med*. 2008;67:956-964.
27. Centraal Bureau voor de Statistiek (CBS), Nederlandse Organisatie voor Toegepast Natuurwetenschappelijk Onderzoek (TNO). NEA: Nationale Enquête Arbeidsomstandigheden voor werknemers [Internet]. 2014. http://www.monitorarbeid.tno.nl/dynamics/modules/SFIL0100/view.php?fil_id=125 Accessed March 12, 2018.
28. Diepgen TL. Occupational skin diseases. *J Dtsch Dermatol Ges*. 2012;10:295-297.
29. Halkier-Sørensen L. Notified occupational skin diseases in Denmark. Important exposure sources, occupations and trades. The course from notification to compensation and socio-economical aspects. *Contact Dermatitis*. 1996;35(suppl 1):1-120.
30. Diepgen TL. Occupational skin-disease data in Europe. *Int Arch Occup Environ Health*. 2003;76:331-338.
31. Skoet R, Olsen J, Mathiesen B, Iversen L, Johansen JD, Agner T. A survey of occupational hand eczema in Denmark. *Contact Dermatitis*. 2004;51:159-166.
32. Schwensen JF, Friis UF, Menné T, Johansen JD. One thousand cases of severe occupational contact dermatitis. *Contact Dermatitis*. 2013;68:259-268.
33. Diepgen TL, Andersen KE, Chosidow O, et al. Guidelines for diagnosis, prevention and treatment of hand eczema. *J Dtsch Dermatol Ges*. 2015;13:e1-e22.
34. Bundesanstalt für Arbeitsschutz und Arbeitsmedizin. TRGS 401: Risks Resulting from Skin Contact—Identification, Assessment, Measures [Internet]. 2008. <https://www.baua.de/DE/Angebote/Rechtstexte-und-Technische-Regeln/Regelwerk/TRGS/TRGS-401.html> Accessed March 12, 2018.
35. Behroozy A, Keegel TGG. Wet-work exposure: a main risk factor for occupational hand dermatitis. *Saf Health Work*. 2014;5:175-180.
36. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol*. 1996;49:1373-1379.
37. Ospina MB, Dennett L, Wayne A, Jacobs P, Thompson AH. A systematic review of measurement properties of instruments assessing presenteeism. *Am J Manag Care*. 2015;21:e171-e185.
38. Boonen A, Brinkhuizen T, Landewe R, van der Heijde D, Severens JL. Impact of ankylosing spondylitis on sick leave, presenteeism and unpaid productivity, and estimation of the societal cost. *Ann Rheum Dis*. 2010;69:1123-1128.
39. Braakman-Jansen LM, Taal E, Kuper IH, van de Laar MA. Productivity loss due to absenteeism and presenteeism by different instruments in patients with RA and subjects without RA. *Rheumatology (Oxford)*. 2012;51:354-361.
40. d'Errico A, Viotti S, Baratti A, et al. Low back pain and associated presenteeism among hospital nursing staff. *J Occup Health*. 2013;55:276-283.
41. Haglund E, Petersson IF, Bremander A, Bergman S. Predictors of presenteeism and activity impairment outside work in patients with spondyloarthritis. *J Occup Rehabil*. 2015;25:288-295.
42. Tran-Duy A, Nguyen TT, Thijs H, et al. Longitudinal analyses of presenteeism and its role as a predictor of sick leave in patients with ankylosing spondylitis. *Arthritis Care Res (Hoboken)*. 2015;67:1578-1585.
43. Zand A, van Deen WK, Inserra EK, et al. Presenteeism in inflammatory bowel diseases: a hidden problem with significant economic impact. *Inflamm Bowel Dis*. 2015;21:1623-1630.
44. Cosatti MA, Muñoz S, Alba P, et al. Multicenter study to assess presenteeism in systemic lupus erythematosus and its relationship with clinical and sociodemographic features. *Lupus*. 2017;27:33-39. <https://doi.org/10.1177/0961203317701843>.
45. d'Errico A, Ardito C, Leombruni R. Work organization, exposure to workplace hazards and sickness presenteeism in the European employed population. *Am J Ind Med*. 2016;59:57-72.
46. Kigozi J, Jowett S, Lewis M, Barton P, Coast J. The estimation and inclusion of presenteeism costs in applied economic evaluation: a systematic review. *Value Health*. 2017;20:496-506.
47. Robertson I, Leach D, Doerner N, Smeed M. Poor health but not absent: prevalence, predictors, and outcomes of presenteeism. *J Occup Environ Med*. 2012;54:1344-1349.
48. Miraglia M, Johns G. Going to work ill: a meta-analysis of the correlates of presenteeism and a dual-path model. *J Occup Health Psychol*. 2016;21:261-283.
49. Brooks A, Hagen SE, Sathyanarayanan S, Schultz AB, Edington DW. Presenteeism: critical issues. *J Occup Environ Med*. 2010;52:1055-1067.
50. Burton WN, Chen CY, Conti DJ, Schultz AB, Edington DW. The association between health risk change and presenteeism change. *J Occup Environ Med*. 2006;48:252-263.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Oosterhaven JAF, Flach PA, Bültmann U, Schuttelaar MLA. Presenteeism in a Dutch hand eczema population—a cross-sectional survey. *Contact Dermatitis*. 2018;79:10–19. <https://doi.org/10.1111/cod.12993>