

The quest of finding the perfect spermatozoon

Cristian O'Flaherty

Departments of Surgery (Division of Urology) and Pharmacology and Therapeutics, McGill University and the Research Institute, McGill University Health Centre (MUHC), Montréal, Canada

Correspondence to: Dr. Cristian O'Flaherty, DVM, PhD. Departments of Surgery (Division of Urology) and Pharmacology and Therapeutics, McGill University and the Research Institute, McGill University Health Centre (MUHC), Montréal, Canada. Email: cristian.oflaherty@mcgill.ca.

Comments on: Agarwal A, Majzoub A, Esteves SC, *et al.* Clinical utility of sperm DNA fragmentation testing: practice recommendations based on clinical scenarios. *Transl Androl Urol* 2016;5:935-50.

Submitted May 05, 2017. Accepted for publication May 14, 2017.

doi: 10.21037/tau.2017.05.38

View this article at: <http://dx.doi.org/10.21037/tau.2017.05.38>

Infertility is a health problem that touches around 15% of couples worldwide and male infertility is the sole cause in half of these cases (1,2). Since the development of assisted reproductive techniques (ART) particularly intracytoplasmic sperm injection (ICSI), it has been a challenge to select the very best spermatozoon to inject into the mature oocyte. This quest has not been easy and up to now there is no diagnostic tool to assure the health of the selected sperm.

Due to the lack of high percentages of success rate for ART it is mandatory to develop new strategies to select and improve the quality of the sperm sample to be used either for *in vitro* fertilization (IVF) or ICSI. There is increasing number of studies that raise a red flag in terms of the safety of the sperm sample to be used in ART (3,4). In this review, Agarwal *et al.* (5) aim to deliver useful guidelines for sperm DNA fragmentation testing based on commonly encountered clinical scenarios and considering what the different test measure and their feasibility in terms of costs and practicality for the clinic.

Studies using animal models and human spermatozoa are providing increasing evidence that sperm DNA fragmentation is a major culprit for abnormal reproductive outcomes (6-9). What is important to address is that the sperm chromatin is a complex structure with all its components being susceptible to damage. Spermatozoa from cancer survivors have a variety of sperm chromatin damage from single and double DNA strand breaks to different levels of DNA compaction due to either low levels of protamination, loss of disulfides bridges between protamines or in some cases due to both. Particularly in the case of levels of DNA compaction, it has been reported

that an overoxidation of thiol groups is associated with male infertility (10,11), thus caution must be taken at the time of analyzing this characteristic of the sperm chromatin and highlights the importance of an appropriate balance in the redox status of the sperm nuclear thiol groups. These findings indicate that sperm chromatin quality should be defined by analyzing separate components. Moreover, the way these men recovered their sperm chromatin integrity varied among individuals and with time (12,13).

It is now evident that the standard semen analysis does not help clinicians to decide, in some cases, what therapeutic path to follow to help infertile men. Thus, it is imperative to find new alternatives that will provide sufficient information to better understand male infertility. The inclusion of sperm chromatin structure assays as those indicated by Agarwal *et al.* (5) can be of help in the treatment of male infertility. However, there is still not enough evidence that these tests can predict the reproductive outcome. There are two important issues that need to be addressed in order to support or not the introduction of these techniques in clinical practice: (I) standardization of these assays using unified protocols and (II) development of randomized controlled studies with sufficient number of participants. However, standardization can be difficult to implement and efforts from large institutions and recognized research groups in the field must come together to accomplish these goals. It is also time for governments to get involved by funding these studies as the outcome of this research may help to design new diagnostic and treatment strategies to maximize subsidized reproduction assisted programs.

Acknowledgements

Funding: Supported by a grant from the Canadian Institutes of Health Research (MOP 133661 to C O'Flaherty).

Footnote

Conflicts of Interest: C O'Flaherty is the recipient of the Chercher Boursier Junior 2 salary award from the Fonds de la Recherche en Santé du Québec.

References

1. World Health Organization. Towards More Objectivity in Diagnosis and Management of Male Infertility: Results of a World Health Organization Multicentre Study. *Int J Androl* 1987;1:53.
2. De Kretser DM, Baker HW. Infertility in men: recent advances and continuing controversies. *J Clin Endocrinol Metab* 1999;84:3443-50.
3. Agarwal P, Loh SK, Lim SB, et al. Two-year neurodevelopmental outcome in children conceived by intracytoplasmic sperm injection: prospective cohort study. *BJOG* 2005;112:1376-83.
4. Hansen M, Bower C, Milne E, et al. Assisted reproductive technologies and the risk of birth defects--a systematic review. *Hum Reprod* 2005;20:328-38.
5. Agarwal A, Majzoub A, Esteves SC, et al. Clinical utility of sperm DNA fragmentation testing: practice recommendations based on clinical scenarios. *Transl Androl Urol* 2016;5:935-50.
6. Evenson D, Wixon R. Meta-analysis of sperm DNA fragmentation using the sperm chromatin structure assay. *Reprod Biomed Online* 2006;12:466-72.
7. Lewis SE, John Aitken R, Conner SJ, et al. The impact of sperm DNA damage in assisted conception and beyond: recent advances in diagnosis and treatment. *Reprod Biomed Online* 2013;27:325-37.
8. Ozkosem B, Feinstein SI, Fisher AB, et al. Absence of Peroxiredoxin 6 Amplifies the Effect of Oxidant Stress on Mobility and SCSA/CMA3 Defined Chromatin Quality and Impairs Fertilizing Ability of Mouse Spermatozoa. *Biol Reprod* 2016;94:68.
9. Fatehi AN, Bevers MM, Schoevers E, et al. DNA damage in bovine sperm does not block fertilization and early embryonic development but induces apoptosis after the first cleavages. *J Androl* 2006;27:176-88.
10. O'Flaherty C, Vaisheva F, Hales BF, et al. Characterization of sperm chromatin quality in testicular cancer and Hodgkin's lymphoma patients prior to chemotherapy. *Hum Reprod* 2008;23:1044-52.
11. Seligman J, Kosower NS, Weissenberg R, et al. Thiol-disulfide status of human sperm proteins. *J Reprod Fertil* 1994;101:435-43.
12. O'Flaherty C, Hales BF, Chan P, et al. Impact of chemotherapeutics and advanced testicular cancer or Hodgkin lymphoma on sperm deoxyribonucleic acid integrity. *Fertil Steril* 2010;94:1374-9.
13. O'Flaherty CM, Chan PT, Hales BF, et al. Sperm chromatin structure components are differentially repaired in cancer survivors. *J Androl* 2012;33:629-36.

Cite this article as: O'Flaherty C. The quest of finding the perfect spermatozoon. *Transl Androl Urol* 2017;6(Suppl 4):S491-S492. doi: 10.21037/tau.2017.05.38