Poster presentation

Open Access

Two-compartment models of spasticity in spinal motor neurons following spinal cord injury Mini Kurian^{*1} and Sharon Crook^{1,2}

Address: ¹Center for Adaptive Neural Systems, Biodesign Institute & Department of Mathematics and Statistics, Arizona State University, Tempe, AZ 85287, USA and ²School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA

Email: Mini Kurian* - kurian@mathpost.la.asu.edu * Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007 Toronto, Canada. 7–12 July 2007

Published: 6 July 2007 BMC Neuroscience 2007, 8(Suppl 2):P101 doi:10.1186/1471-2202-8-S2-P101

© 2007 Kurian and Crook; licensee BioMed Central Ltd.

Spasticity is characterized by hyperreflexia, clonus and hypertonic musculature and is an impediment to functional locomotor recovery after spinal cord injury (SCI). It is known that the structure and biophysical properties of motoneurons are altered following SCI; however, the physiological mechanisms underlying spasticity are not well understood. Under control conditions motoneurons can produce plateau potentials, which are sustained depolarizations triggered by brief synaptic inputs. These plateau potentials are mediated by L-type calcium currents and are known to cause bistable behavior in the motoneurons of brain-stem intact animals and humans. This bistable behavior endows motoneurons with a mechanism for translating short lasting synaptic inputs into long lasting motor output [1]. During the acute stage following SCI, rat motoneurons lose the endogenous ability to generate plateau potentials but at chronic stages the plateau potentials reappear [2]. Voltage gated persistent sodium and calcium currents (PICs) have been identified as the cause of plateau potentials in the chronic stage following SCI [3]. In the presence of PICs, a brief stimulus can produce self sustained firing.

Two-compartment models with Hodgkin-Huxley type channel kinetics were constructed to mimic motoneuron dynamics for the control case and for the acute and chronic stages following SCI. Table 1 shows the channels used in constructing the motoneuron models.

Experimental data for injected ramp currents and injected pulse currents were used to constrain the model parameters [2]. Computational studies were used to systematically investigate the mechanisms underlying the generation of plateau potentials and the firing properties in the three cases including: the influences of the specific ionic currents, the coupling strength between the somatic and dendritic compartments, and the relative sizes of the two compartments.

Table I: Motoneuron model currents.

Motoneuron	Soma Channels	Dendritic Channels
Control	I _{Na} , I _{K-dr} , I _{Ca-N} , I _{K(Ca)} , I _{Leak}	I _{Ca-N} , I _{Ca-L} , I _{K(Ca)} , I _{Leak}
Acute	I _{Na} , I _{K-dr} , I _{Ca-N} , I _{K(Ca)} , I _{Leak}	I _{Ca-N} , I _{K(Ca)} , I _{Leak}
Chronic	I _{Na} , I _{K-dr} , I _{Ca-N} , I _{K(Ca)} , I _{Leak}	$I_{Ca-N}, I_{Ca-P} I_{Na-P}, I_{K(Ca)}, I_{Leak}$

References

- Heckman CJ, Gorassini MA, Bennett DJ: Persistent inward currents in motoneurons dendrites: implications for motor output. *Muscle Nerve* 2004, 31:135-156.
- Bennett DJ, Li Y, Siu M: Plateau potentials in sacrocaudal motoneurons of chronic spinal rats recorded in vitro. J Neurophysiol 2001, 86:1955-1971.
- Li Y, Bennett DJ: Persistent sodium and calcium currents cause plateau potentials in motoneurons of chronic spinal rats. J Neurophysiol 2003, 90:857-869.

