

RESEARCH ARTICLE

Retinoic Acid Can Exacerbate T Cell Intrinsic TLR2 Activation to Promote Tolerance

Vivien Nguyen¹, Kandyce Pearson², Jee-Hyun Kim², Karishma Kamdar², R. William DePaolo^{2*}

1 Department of Pediatric Gastroenterology and Nutrition, Children's Hospital of Los Angeles, Los Angeles, California, United States of America, **2** Department of Molecular Microbiology and Immunology, Keck School of Medicine and University of Southern California, Los Angeles, California, United States of America

* depaolo@usc.edu



OPEN ACCESS

Citation: Nguyen V, Pearson K, Kim J-H, Kamdar K, DePaolo RW (2015) Retinoic Acid Can Exacerbate T Cell Intrinsic TLR2 Activation to Promote Tolerance. *PLoS ONE* 10(3): e0118875. doi:10.1371/journal.pone.0118875

Academic Editor: Lena Alexopoulou, Centre d'Immunologie de Marseille-Luminy, CNRS-Inserm, FRANCE

Received: December 12, 2013

Accepted: January 13, 2015

Published: March 31, 2015

Copyright: © 2015 Nguyen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Research reported in this publication was supported by National Institute of Digestive Disease and Kidney Research of the National Institutes of Health under award number K01DK082725. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

The contribution of vitamin A to immune health has been well established. However, recent evidence indicates that its active metabolite, retinoic acid (RA), has the ability to promote both tolerogenic and inflammatory responses. While the outcome of RA-mediated immunity is dependent upon the immunological status of the tissue, the contribution of specific innate signals influencing this response have yet to be delineated. Here, we found that treatment with RA can dampen inflammation during intestinal injury. Importantly, we report a novel and unexpected requirement for TLR2 in RA-mediated suppression. Our data demonstrate that RA treatment enhances TLR2-dependent IL-10 production from T cells and this, in turn, potentiates T regulatory cell (T_{REG}) generation without the need for activation of antigen presenting cells. These data also suggest that combinatorial therapy using RA and TLR2 ligands may be advantageous in the design of therapies to treat autoimmune or inflammatory disease.

Introduction

The general concept that Vitamin A (VA) contributes to immunity dates as far back as Hippocrates [1], and recent advances have demonstrated specific roles for VA in many different types of disease. For instance, VA deficiency (VAD) increases mortality during gastrointestinal, respiratory and HIV infections [2–5] which can be reversed by VA supplementation [6–7]. Despite these observations the role of VA is still not well understood in the context of intestinal inflammation even though more than 15% of children with inflammatory bowel disease (IBD) have low serum levels of VA at the time of diagnosis [8].

VA mediates its metabolic and immune effects via conversion to its active metabolite, RA, via retinaldehyde dehydrogenase (RALDH) enzymes [9–11]. In the last decade, many studies have provided insight into the nature of RA-mediated responses, especially its role in innate and adaptive immunity within the gut associated lymphoid tissues (GALT). Most notably, RA promotes T cell trafficking to the GALT via $\alpha 4\beta 7$ and CCR9 expression [12–14] and contributes to the polarization of Foxp3⁺ T_{REG} by RALDH-expressing CD103⁺ GALT DC [15–20].

These effects are dependent on TGF- β mediated T cell expression of retinoic acid receptor (RAR) and repression of the IL-6R, respectively [21–23]. Corroborating these *in vitro* findings, the generation of induced T_{REG} (iT_{REG}) in response to ingested antigens is abrogated in VAD mice [24].

iT_{REG} and IL-17-producing CD4 helper T cells (T_H17) have a reciprocal relationship [19, 25, 26], leading one to expect an inhibitory effect of RA on T_H17 differentiation and maintenance. A number of studies have shown that direct activation of RA on T cells can suppress T_H17 differentiation through the inhibition of IL-6R and IL-23R [13, 19, 23]. However, antigen-presenting cells activated via MyD88-dependent innate signals and treated with RA have been shown to potentiate T_H17 differentiation [27]. These data suggest that RA, in concert with microbial-driven signals, may help to promote T_H17 cell differentiation further suggesting that RA may have a dual nature imparting it with the ability to both promote and inhibit iT_{REG} generation via the regulation T_H17 cells [28].

Pathogens crossing the epithelial barrier during infection or exposure of the tissue to commensal bacteria during injury can provide the microbial signals needed to impact RA-mediated immunity. While tissue-derived homeostatic factors may promote the expression of RALDH in CD103⁺ DC in order to potentiate iT_{REG} cell numbers [29, 30, 31], inflammation and exposure to microbes may have an opposite effect. This has been observed in models of experimental colitis in which the expression of RALDH in CD103⁺ DC is reduced resulting in fewer iT_{REG} and worse inflammation [32]. In an IL-15-enriched microenvironment, RA was shown to increase the production of IL-12 and IL-23 by gut CD103⁺ DC, diminishing their capacity to promote iT_{REG} and restrain T_H1 and T_H17 responses to dietary gluten [33]. These data align with clinical reports linking pharmacological retinoid therapy to the development of IBD in a subset of patients and point to RA as a potential instigator of inflammation in the appropriate milieu [24, 34].

TLR2 is a member of the Toll-like receptor (TLR) family of pattern recognition receptors, and detects tri- [35] and di-acylated [36] bacterial lipoproteins by forming heterodimers with TLR1 or TLR6, respectively. TLR2 signaling in splenic DC induces RALDH activity [30] and IL-10 [37], imparting them with gut-specific imprinting and iT_{REG}-promoting functions. In contrast, others have demonstrated preservation of RALDH activity in MyD88-deficient DC [38] and promotion of T_H17 cells [27] and RALDH [32] during microbial stimulation. Studies examining the relationship between TLR2 and RA have focused on the DC, despite reports that TLR2 is expressed on T_{REG} [39] and may influence T_{REG} expansion and function [40–42]. Here we show that exogenous RA can suppress inflammation during intestinal injury and that this ability is lost in a TLR2-deficient environment. Further, we show that RA potentiates TLR2-induced IL-10 production directly from T cells and promotes iT_{REG} differentiation. These findings further demonstrate the ability of RA to act as an adjuvant to promote signals from the local tissue microenvironment and suggest a potential benefit of combining TLR2 ligands and RA to suppress inflammatory disease and promote tolerance.

Materials and Methods

Mice

This study was performed in strict accordance and compliance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health, the Animal Welfare Act and U.S. Government Principles for The Utilization and Care of Vertebrate Animals Used in Testing Research and Training. All studies were approved and in accordance with the Institutional Biosafety Committee (IBC) and the Institutional Care and Use

Committee (IACUC) of the University of Southern California. Mice were housed in specific pathogen free (SPF) conditions.

C57Bl/6 (WT) mice were purchased from The Jackson Laboratory (Bar Harbor, ME) and TLR2KO mice on a B6 background were bred in-house. Male mice aged 6–8 weeks were used for all experiments and the numbers of mice used for each experimental group are listed in the figure legends.

Induction of colitis

2.5% (weight/volume) dextran sodium sulfate (DSS; molecular weight 36,000–50,000; lot NM4241; MP Biomedicals; Santa Ana, CA) was added to drinking water for 7 days, followed by a 7-day period of recovery with untreated drinking water.

Clinical and histological evaluation of colitis

Body weight, presence of occult or gross blood per rectum (scoring: 0, no blood; 1, occult blood positive; 2, visible blood in fecal pellet; 4, blood at anus) were determined daily. Colonic specimens were opened longitudinally and made into “Swiss-roll” preparations. Hematoxylin and eosin staining was performed on 5 μ m paraffin-embedded sections. Slides were viewed with a Leica DM 750 microscope with PLAN 10x/0.22 and PLAN 40x/0.65 objectives. Histologic severity was assessed using a scoring system developed at The Jackson Laboratory [42]. All slides were evaluated in a blinded manner by independently two gastroenterologist/pathologists.

Oral tolerance induction

WT and TLR2KO mice were orally gavaged as previously described [33]. Fecal occult blood tests (Hemocult Sensa II; Beckman Coulter; Brea, CA) were performed daily and upon conversion from negative to positive, 1 μ M of RA dissolved in 100 μ l of corn oil (Sigma-Aldrich; St. Louis, MO) was administered to each mouse via oral gavage with an 18 gauge round-tipped needle (Kent Scientific Corp; Torrington, CT) for the duration of the experiment. Controls received the same volume of DMSO diluted in 100 μ l of vehicle (corn oil). In some experiments the mice received 100 μ l of sterile PBS containing 100 μ g of ovalbumin or PBS alone every other day for a total of five feedings. Mice were euthanized two days after the last feeding. In some experiments 1×10^5 syngeneic ovalbumin-specific T cells from RAG-OT2-Tg mice were transferred to naïve TLR2KO mice prior to antigen feeding via retro-orbital injection [33].

Protein extraction from tissues and cecal lysates

Total protein was extracted from whole colon specimens by flash freezing with liquid nitrogen and manually homogenizing with a mortar and pestle. For cecal lysates, total contents were placed in 500 μ l sterile PBS and mixed with 500 μ l 0.1 mm glass beads and mechanically homogenized for 3–4 minutes using a Bead Beater (BioSpec). After beads and debris settle to the bottom of the tube, the supernatant is removed. Protein concentrations of supernatants were determined via Bradford assay per manufacturer’s protocol (Bio-Rad; Hercules, CA) and homogenates were diluted to a concentration of 1 mg/mL with sterile PBS (Sigma-Aldrich).

Quantitative real-time RT-PCR

Quantitative real-time RT-PCR was performed using the iCycler iQ real-time PCR detection system (Bio-Rad Laboratories) and a SYBR green amplification kit (PE Biosystems). Each PCR reaction was performed and normalized using primers for GAPDH. $\Delta\Delta$ CT was calculated using Δ CT values of naïve WT or naïve TLR2KO samples as the reference.

In vitro assays

For CD4⁺ T cell isolation, spleens were mechanically disrupted through a 70 µm cell strainer (BD Biosciences; San Jose, CA). CD4⁺ cells were isolated by positive immunoselection using CD4 (L3T4) microbeads (Miltenyi Biotec; Bergisch Gladbach, DE). Purified CD4⁺ T cells were stained with anti-CD62L and anti-CD44 and sorted on a FACS Aria II resulting in a 98% pure population of naïve T cells. The purified T cells were stimulated with 1 µg/mL plate-bound anti-CD3ε (eBioscience; San Diego, CA), 0.1 µg/mL TLR ligands, 10 µg/mL cecal lysate, and 2 ng/mL recombinant TGF-β (R&D Systems; Minneapolis, MN), in the presence or absence of 10 nM RA. In some experiments 50 µg/ml anti-IL-10Rα antibody (clone 1B1.3a) or isotype rat IgG1 (BD Biosciences) control was added to the wells.

For dendritic cell isolation, spleens were digested with 400 units/mL collagenase type IV (Sigma-Aldrich). Cells were filtered, re-suspended in 22.5% Optiprep (Sigma-Aldrich), overlaid with Hank's Buffered Saline (HBS; Sigma-Aldrich) and centrifuged at 670g for 30 minutes. Dendritic cells were sorted from the interphase using magnetic CD11c Microbeads (Miltenyi Biotec; Bergisch Gladbach, DE) and stimulated with 10 µg/mL cecal lysate and 2 ng/mL TGF-β in the presence or absence of 10 nM RA.

Antibodies and flow cytometry

The following conjugated antibodies were purchased from eBioscience: CD4 (GK1.5), CD45.1 (A20), CD45.2 (104), IFN-γ (XMG1.2), Foxp3 (FJK-16a), IL-10 (JES5-16E3) and isotype controls. A fixation and permeabilization kit was used for intranuclear detection of Foxp3 followed by an intracellular staining protocol for IL-10 (eBioscience). Flow cytometry analysis was performed with a FACS Canto (BD Biosciences) and analyzed using FlowJo software.

Detection of cytokines by ELISA

Cell supernatants (50 µL per well) and tissue homogenates (50 µg per well) were evaluated for IL-12p40 (BD Biosciences), IL-10, (BD Biosciences), TNF-α (BD Biosciences), IFN-γ (BD Biosciences) and IL-17 (R&D).

Statistical analysis

Data are expressed as means ± standard error of the mean (SEM) of 2 or more independent experiments. Differences between means are evaluated using 2-tailed, unpaired Student's *t*-tests (GraphPad Prism version 6; San Diego, CA) where appropriate. *p* values of < 0.05 are considered significant.

Results

RA-mediated repair against epithelial injury is dependent on TLR2-signaling in vivo

Due to the prevalence of vitamin and micronutrient deficiencies in patients with IBD, enteral repletion is often initiated at the time of diagnosis [8]. Given the potential for RA to promote either regulatory [13,19,23] or inflammatory [24,33] immune responses depending on the status of the tissue microenvironment, we assessed the effects of RA supplementation during acute colitis and evaluated the role of TLR2 in modulating these effects. DSS was administered to WT and TLR2 deficient mice to induce epithelial injury for seven days followed by a repair phase in which the mice received regular drinking water. In order to mirror the clinical presentation and course of individuals with IBD, we delayed enteral supplementation of RA until

overt signs of colitis (i.e. loose stool and fecal occult blood) were noted. In line with prior studies demonstrating that the absence of TLR2 signaling exacerbates chemically-induced and spontaneous colitis in mice, we found that TLR2KO mice lost more body weight than WT counterparts during induction of colitis as well as during recovery [43,44] (Fig 1A, closed and open circles). TLR2KO mice also had significantly worse fecal occult blood scores (Fig 1B, closed and open circles), and worse histopathology (Fig 1C) compared to WT mice, with larger areas of denuded, ulcerated epithelium (Fig 1D and 1E, white bars, closed and open circles) and shorter colon lengths (Fig 1F), indicative of inflammation. WT mice treated with RA showed significant improvement in fecal occult bleeding (Fig 1B, closed squares), histopathology (Fig 1C, bottom left), less epithelial damage (Fig 1D and 1E, black bar and closed squares) and significantly longer colons (Fig 1F). In contrast, RA supplementation had adverse effects in TLR2KO mice, which demonstrated severe rectal bleeding (Fig 1B, open squares); worse histopathology (Fig 1C and 1D, bottom right, black bars), and shorter colons (Fig 1F) compared to both WT mice receiving RA and TLR2KO mice receiving vehicle control. Together these data demonstrate that treatment with RA can protect against epithelial injury during acute colitis however, in the absence of TLR2, treatment with RA exacerbates disease.

RA mediated suppression of colonic cytokine production in vivo is dependent on TLR2

To determine if RA supplementation affects inflammatory cytokine production, we examined the levels of cytokines in the colonic mucosa at day 10. RA treatment suppressed IFN- γ (Fig 2A) and TNF- α (data not shown) production in WT mice and exacerbated their production in TLR2KO mice (Fig 2A). In contrast, RA treatment increased IL-10 production in the WT mice, an effect that was not observed in the TLR2KO mice (Fig 2A), corroborating earlier observations that TLR2 directly promotes IL-10 responses [48]. RA had no effect on IL-17 production in either condition (Fig 2A).

The colonic mucosa was analyzed for mRNA expression of T-helper associated transcription factors at day 14. Complimenting the cytokine data described above, RA treatment exacerbated the expression of *TBET* in TLR2KO but not WT mice. Furthermore, although RA increased *FOXP3* expression in both conditions, its expression was significantly lower in TLR2KO animals (Fig 2B). As seen with the cytokine analysis, RA treatment had no impact on the T_H17 -associated factor *RORC*. The increased IL-10 and *FOXP3* levels lead us to speculate that the protective effect of RA observed in WT mice may be due to the induction of anti-inflammatory responses via the induction of IL-10 and $Foxp3^+$ T regulatory cells. Examination of colonic LP cells 10 days after initial DSS treatment demonstrated an increase in the frequency of $CD4^+$ T cells expressing $Foxp3^+$ and IL-10 in WT mice treated with RA, an effect that was not in TLR2KO mice (Fig 2C). Taken together, these data suggest that *in vivo* RA treatment affects both cytokine production and T cell polarization in a manner dependent upon TLR2 signaling.

The suppressive effect of RA and TLR2 is not mediated by dendritic cells

Both TLR2 and RA have been shown to have effects on DC as well as T cells [21,23,27,37]. In order to further dissect the role of RA and TLR2 in regulating cytokine production, splenic DC were purified from the spleens of WT and TLR2KO mice. To evaluate the specific effects mediated by RA and TLR2 on DC in a gut environment [33] TGF- β and the TLR2/6-agonist Pam2-Cysk4 (Pam2) were added to cultures containing RA. RA treatment suppressed Pam2 mediated IL-10 production by WT DC, and enhanced IL-12p40 production (Fig 3A). RA




Fig 1. RA potentiates colonic injury and inflammation in the absence of TLR2 signaling. WT and TLR2KO mice were given 2.5% DSS for seven days to induce colonic damage and then placed on normal drinking water for seven days to allow for tissue repair. Upon first presence of fecal occult blood positivity for each, mice were supplemented with RA or vehicle control, which continued for the rest of the disease course. Weight loss (A) and fecal occult blood (B) (n = 10–14 mice per group) were monitored daily. (C) H&E staining of colonic sections of mice at day 14 (after 7 days of DSS followed by 7 days of water). (D) Histology scoring of H&E-stained colonic slides (n = 4 mice per group). (E) The percent of surface area containing ulcerated/denuded epithelium quantified using J Image software. (F) Colon lengths from mice at day 10 (n = 5–6 mice per group). *, p < 0.05, **, p < 0.01 using Students t-test.

doi:10.1371/journal.pone.0118875.g001

treatment had no effect on the production of these cytokines by WT DC in the absence of Pam2, or in TLR2KO DC (data not shown).

To determine whether specific, endogenous signals within the microbiota or DC-modulators produced by the epithelium can alter the response to RA in a TLR2 dependent manner, we treated splenic DC from WT and TLR2KO mice with lysates of their own cecal contents. RA exacerbated the production of IL-12p40 by both WT and TLR2KO DC stimulated with cecal lysates, however there was no impact of RA on IL-10 in these conditions (Fig 3B). The intestinal micro-environment differs greatly from peripheral tissues such as the spleen, to determine whether the effects of TLR2 and RA would also occur in mucosal derived cells we sorted and stimulated CD103⁺ DC from the colonic LP and spleen as previously described [33]. CD103⁺ LP DC will produce RA *ex vivo* [18, 20] and the addition a TLR2 agonist produced levels of IL-10 equal to splenic CD103⁺ DC plus TLR2 agonist and RA (Fig 3C) suggesting that the ability of TLR2 to induce IL-10 is not tissue specific. Previous studies have also indicated that TLR2 signaling in splenic DC induces RALDH which converting vitamin A to RA [30]. We evaluated the transcript




Fig 2. RA treatment suppresses IFN- γ during DSS in WT mice but enhances their secretion in TLR2KO mice. (A) Mucosal scrapings from mice receiving the DSS and water were harvested on day 10 and analyzed by ELISA for cytokine levels. (B) Quantitative RT-PCR for transcription factors associated with T helper subsets were also performed on colonic lamina propria samples taken at day 14. Data shown are the fold increase of Vehicle- and RA-treated mice compared to naive controls. Data are the mean \pm SEM of 5–8 mice per group pooled from two independent experiments. (C) Expression of Foxp3 and IL-10 in colonic LP CD4⁺ T cells (n = 3 per group), one representative facs plot is shown. *, p < 0.05, **, p < 0.01 using Student's t-test.

doi:10.1371/journal.pone.0118875.g002

levels of *aldha1a2* in splenic and LP DC to determine whether TLR2KO mice had an intrinsic defect in RA levels accounting for the phenotype we observed. However, we found similar levels of *aldha1a2* regardless of TLR2 expression (Fig 3D). Overall, RA was unable to suppress *ex vivo* IL-12p40 production by WT splenic DC stimulated with either a TLR2-specific agonist or commensal lysate and had little effect on cytokine production by TLR2KO DC.

RA enhances TLR2-induced IL-10 production from CD4⁺ T cells

Direct signaling via TLR2 on iT_{REG} has been shown to affect their expansion and function [40, 41] and the role of RA in iT_{REG} function has been clearly established [21,23,45]. We




Fig 3. RA potentiates cytokine responses in both WT and TLR2KO DC. (A) WT splenic DC were cultured with 100 ng/ml of TLR2 ligand Pam2CysK4 and (B) WT and TLR2KO splenic DC were cultured with 10 µg/ml cecal lysate in the presence or absence of RA, TGF-β or RA/TGF-β. Supernatants were analyzed after 24 hours for cytokine production by ELISA. (C) Comparison of IL-10 IL- production from CD103⁺ LP DC and CD103⁺ SP DC. (D) Transcript levels of *aldh1a2* in spleen and LP of naïve WT and TLR2KO mice determined by quantitative PCR. For (A-C), data are the mean ± SEM of three independent experiments, for (D) 5 individual mice were analyzed. *, p < 0.05, **, p < 0.01 using Student's t-test.

doi:10.1371/journal.pone.0118875.g003

questioned whether the ability of RA to promote iT_{REG} responses could be influenced by TLR2 signaling, which may explain the marked difference in response to RA we observed *in vivo*. To address this question, naïve splenic CD4⁺ T cells were stimulated polyclonally in the presence of Pam2. Engagement of TLR2 by Pam2 directly on T cells suppressed IFN-γ and increased IL-10 production in WT T cells. Interestingly, RA exacerbated both the IFN-γ suppression and IL-10 production (Fig 4A) suggesting an adjuvant effect of RA on TLR2-mediated T cell cytokine production. There was no effect by either TLR2 activation or RA treatment on IL-17 production (Fig 4A).

To determine whether RA could alter T cell responses in a TLR2-dependent manner in the presence of endogenous microbial ligands, we stimulated WT and TLR2KO CD4 T cells in the presence of cecal lysate, TGF-β as well as RA. Unlike the pure TLR2 agonist, Pam2, cecal lysate increased IFN-γ production by CD4 T cells, which is likely the result of the presence of alternative TLR-ligands in the culture (Fig 4B). Importantly, however, RA dramatically suppressed IFN-γ production by CD4 T cells in the presence of cecal lysate and this was not observed in TLR2KO cells (Fig 4B). Cecal lysate also increased IL-10 production by WT CD4 T cells and similar to the Pam2 stimulated cells, RA enhanced this phenotype in a TLR2-dependent manner (Fig 4B). IL-17 production was increased by cecal lysate in the presence of exogenous TGF-




Fig 4. RA and TLR2 signals suppress pro-inflammatory cytokines from T cells. Cytokine production from purified splenic CD4⁺ T cells stimulated with anti-CD3 ϵ and (A) Pam2CysK4, (B) cecal lysate, or (C) indicated TLR agonists in the presence or absence RA, TGF- β or RA/TGF- β . Data are the mean \pm SEM of 3–4 independent experiments. *, $p < 0.05$, **, $p < 0.01$ using Student's t-test.

doi:10.1371/journal.pone.0118875.g004

β , but this was not altered by either RA treatment or TLR2 deficiency. Because of the abundance of potential TLR ligands in the cecal lysate and because a recent study has shown a relationship between yeast zymosan, TLR2/dectin-1 signaling and IL-10 in DC [49], we checked to see if other TLR ligands induced IL-10 from T cells and if there was an adjuvant effect mediated by RA. The TLR5 ligand, flagellin, was the only TLR agonist screened that induced IL-10 from T cells, however RA had no effect on IL-10 levels (Fig 4C). Zymosan, which is a Dectin-1 and TLR2 agonist, was unable to induce IL-10 from purified, polyclonally stimulated T cells *in vitro*, and has no role in RA-mediated immune-regulation. Taken together, we conclude that in the presence of specific TLR2 ligands, RA suppresses IFN- γ and enhances IL-10 production by CD4 T cells and this is dependent on CD4 T cell intrinsic TLR2 signaling.

RA and TLR2 synergize to promote the generation of Foxp3⁺ T cells in an IL-10, T cell intrinsic manner

To investigate the synergy of TLR2 and RA in IL-10 production by CD4 T cells, we analyzed intracellular expression of IL-10 and Foxp3 in polyclonally stimulated WT and TLR2KO CD4⁺ T cells treated with Pam2, RA and/or TGF- β . In the presence of Pam2, RA treatment caused a robust increase in IL-10⁺ CD4 T cells, which were Foxp3⁻ and this was not observed in the absence of Pam2 or in TLR2KO CD4 T cells (Fig 5A). Furthermore, in the presence of only




Fig 5. TLR2 and RA act in concert to potentiate iT_{REG} generation via IL-10 production. Foxp3 and IL-10 expression by WT and TLR2KO T cells stimulated with Pam2CysK4 (A) or cecal lysate (B) in the presence or absence RA, TGF- β or RA/TGF- β . Data are representative plots from three independent experiments for (A) and mean and frequency of Foxp3⁺IL-10⁺ and IL-10⁺CD4⁺ T cells from WT (white bars) and TLR2KO (gray bars) treated with cecal lysate (n = 5–7 mice) for (B). (C) Frequency of Foxp3 and IL-10 expression in WT T cells cultured with Pam2CysK4, RA and TGF- β and treated with anti-IL-10R α antibody or isotype control. Data are representative FACS plots from six independent experiments shown in the bar graphs. (D) Cytokine production from re-stimulated *in vitro* MLN cells of WT and TLR2KO mice fed OVA. Data are the mean \pm SEM of two independent experiments with three mice per group. *, p < 0.05, **, p < 0.01.

doi:10.1371/journal.pone.0118875.g005

Pam2, RA treatment caused a robust increase in non-IL-10 producing Foxp3⁺ CD4⁺ T cells and this same Pam2-mediated increase was also observed after Foxp3⁺ CD4⁺ T cells were enhanced with exogenous TGF- β (Fig 5A). Importantly, the effect of Pam2 and RA to increase Foxp3⁺ CD4⁺ T cells was not observed when TLR2KO cells were used (Fig 5A, bottom panels). Similar data was also obtained with T cells stimulated with cecal lysates instead of Pam2 alone and summarized in Fig 5B.

We hypothesized that the synergy of TLR2 and RA in increasing IL-10 production was itself driving subsequent Foxp3⁺ and IL-10⁺ CD4⁺ T cell expansion in these cultures. To test this we added IL-10R α blocking antibodies to CD4⁺ T cell cultures stimulated with Pam2, TGF- β and RA. Anti-IL-10R α blockade significantly ablated both IL-10⁺ and Foxp3⁺ CD4⁺ T cell production, showing that the effect of Pam2 and RA to promote regulatory T cell responses is IL-10 dependent (Fig 5C).

Endogenous TLR2 signals promote oral tolerance via T cells

The importance of RA in the induction of oral tolerance via generation of iT_{REG} is well established [21–23]. However, the function of endogenous microbial signals in the context of food-derived metabolic signals in this process is not well understood. Based on the above observation that TLR2 signaling on CD4 T cells synergizes with RA to promote the generation of iT_{REG} *in vitro*, we hypothesized that the absence of TLR2 may impair tolerance against orally administered proteins. To test whether RA and TLR2 work *in vivo* to enhance iT_{REG} numbers, WT or TLR2KO mice were fed OVA with or without concomitant RA supplementation. Two days after the last feeding, mesenteric lymph node (MLN) cells were re-stimulated *in vitro* to assess antigen-specific immune polarization. As expected, RA enhanced the normal tolerogenic response to fed OVA in WT mice by significantly increasing IL-10 and suppressing IFN- γ and IL-2 (Fig 5D, data not shown) in a TLR2 dependent manner. This was further demonstrated by the elevated pro-inflammatory IFN- γ from TLR2KO and the inability of these mice to produce IL-10 (Fig 5D). To assess whether the phenotype observed is due to intrinsic TLR2 signaling on T cells we transferred naïve OVA-specific CD4 OT2 T cells from WT mice to WT and TLR2KO mice. When these mice were fed OVA to induce tolerance, we found a reduction in IFN- γ and an induction of IL-10, similar to WT tolerized controls (Fig 5D). These data confirm that TLR2 signaling on T cells is required for oral tolerance and enhancement thereof by RA.

Discussion

Numerous studies have delineated the role of RA in lymphocyte trafficking and differentiation [10,14,16,46]. However, the innate signaling pathways that modulate these effects are not well understood. There is evidence that RA promotes divergent immune responses depending on the environmental milieu [24,33] and that TLR2 is involved in propagating these responses [30]. Our findings illustrate a novel relationship between dietary metabolites and specific signals from the microbiome to generate tolerogenic responses. Importantly, we have shown that RA is capable of inhibiting colonic disease and this is dependent upon TLR2 as seen by decreased pro-inflammatory cytokine production, better histology scores and less bleeding. It is noteworthy that in the absence of TLR2 signaling, RA is altered from a tolerogenic adjuvant to a promoter of hyper-inflammation, drawing a close tie between RA and TLR2 signaling. In the context of DSS-induced tissue injury, our data illustrate a protective role for TLR2 signaling as TLR2-deficient mice displayed shorter colons, more weight loss and worse histological features. The addition of RA during DSS had a protective anti-inflammatory response associated with an increase in Foxp3⁺ and Foxp3⁺ IL-10⁺ expressing cells in WT mice. The effects of RA were lost in the TLR2KO mice and in fact worsened disease. These data lead us to propose a model in which IL-10 is produced during DSS-induced injury via commensal activation of TLR2 on mucosal T cells. The addition of RA increases IL-10 production by TLR2-stimulated T cells, which favors an environment that promotes the generation of Foxp3⁺ Treg cells. These data align with our previous work that suggests RA can act as an adjuvant to promote dominant responses in the mucosal tissue [33]. However, It is possible that RA may promote TLR2-induced IL-10 through another one of its signaling mechanisms.

These data illustrate that RA and TLR2 signal directly via the T cell to increase the production of IL-10. Further, we demonstrate that the IL-10 produced by RA and TLR2 increases the generation of Foxp3⁺ T cells. TLR2 has been shown to be expressed and impact the function of differentiated, activated T_{REG} [40, 41]. In fact, human T cells have been shown to induce TLR2 upon activation and Pam3Cysk4 acts as a co-stimulatory molecule inducing IL-10, TNF- α , IL-2 and IFN- γ production [47]. Our data demonstrates an important role for RA in TLR2-mediated *de novo* generation of Foxp3⁺ iT_{REG} from naïve T cells independent of TGF- β . However, when TGF- β is added to the cultures the numbers of Foxp3⁺ T cells are increased even further by the addition of TLR2 agonists. These data lead us to hypothesize that the absence of TLR2 signaling would negatively affect oral tolerance, as it appears to result in reduced numbers of iT_{REG} against endogenous or commensal antigens. We found that TLR2-deficient mice have a defect in the ability to be tolerized to oral antigen and RA administration promotes production of inflammatory responses in these mice. However, tolerance was restored in TLR2KO animals if they were given a population of TLR2-expressing, antigen-specific, naïve T cells prior to oral antigen and RA treatment.

In summary, our studies reveal that RA and TLR2 signaling synergize to promote tolerogenic responses in a T cell-dependent manner to promote tolerogenic responses via the production of IL-10 and ultimately the potentiation of iT_{REG}. This not only has implications during homeostasis for the generation of tolerogenic responses against dietary antigens, but also during inflammation as a way to temper the inflammatory responses and potentially reduce damage to the tissue. Our observations indicate that enteral vitamin A supplementation or retinoid therapy at the time of diagnosis or during disease relapse may be detrimental to patients with IBD. Our findings also suggest that people with aberrant TLR2 signaling, which has been associated with the single nucleotide polymorphism R753Q, may be predisposed to inflammatory or autoimmune conditions due to impaired induction of iT_{REG}, particularly in the setting of VAD [44]. Finally, given the potential of RA to transition from an essential to pathological mediator of immune responses, further investigation of how vitamin A metabolism affects disease is warranted and the molecular mechanism of TLR2 and RA interactions must be assessed.

Author Contributions

Conceived and designed the experiments: RWD. Performed the experiments: VN JHK KK KP RWD. Analyzed the data: VN RWD. Wrote the paper: VN RWD.

References

1. Wolf G. A history of vitamin A and retinoids. *FASEB J.* 1996; 10: 1102–1107. PMID: [8801174](#)
2. McDermott MR, Mark DA, Befus AD, Baliga BS, Suskind RM, Bienenstock J. Impaired intestinal localization of mesenteric lymphoblasts associated with vitamin A deficiency and protein-calorie malnutrition. *Immunology.* 1982; 45: 1–5. PMID: [6799390](#)
3. Sirisinha S, Darip MD, Moongkarndi P, Ongsakul M, Lamb AJ. Impaired local immune response in vitamin A-deficient rats. *Clin Exp Immunol* 1980; 40: 127–135. PMID: [7389210](#)
4. Sommer A, Tarwotjo I, Hussaini G, Susanto D. Increased mortality in children with mild vitamin A deficiency. *Lancet* 1983; 2: 585–588. PMID: [6136744](#)
5. Wiedermann U, Hanson LA, Kahu H, Dahlgren UI. Aberrant T-cell function in vitro and impaired T-cell dependent antibody response in vivo in vitamin A-deficient rats. *Immunology* 1993; 80: 581–586. PMID: [8307607](#)
6. Semba RD, Ndugwa C, Perry RT, Clark TD, Jackson JB, Melikian G, et al. Effect of periodic vitamin A supplementation on mortality and morbidity of human immunodeficiency virus-infected children in Uganda: A controlled clinical trial. *Nutrition* 2005; 21: 25–31. PMID: [15661475](#)
7. Sommer A, Tarwotjo I, Djunaedi E, West KP, Loeden AA, Tilden R, et al. Impact of vitamin A supplementation on childhood mortality. A randomised controlled community trial. *Lancet* 1986; 1: 1169–1173. PMID: [2871418](#)

8. Alkhouri RH, Hashmi H, Baker RD, Gelfond D, Baker SS. Vitamin and mineral status in patients with inflammatory bowel disease. *J Pediatr Gastroenterol Nutr.* 2013; 56: 89–92. doi: [10.1097/MPG.0b013e31826a105d](https://doi.org/10.1097/MPG.0b013e31826a105d) PMID: [22832510](https://pubmed.ncbi.nlm.nih.gov/22832510/)
9. Hall JA, Grainger JR, Spencer SP, Belkaid Y. The role of retinoic acid in tolerance and immunity. *Immunity* 2011; 35: 13–22. doi: [10.1016/j.immuni.2011.07.002](https://doi.org/10.1016/j.immuni.2011.07.002) PMID: [21777796](https://pubmed.ncbi.nlm.nih.gov/21777796/)
10. Mora JR, Iwata M, von Andrian UH. Vitamin effects on the immune system: vitamins A and D take centre stage. *Nat Rev Immunol.* 2008; 8: 685–698. doi: [10.1038/nri2378](https://doi.org/10.1038/nri2378) PMID: [19172691](https://pubmed.ncbi.nlm.nih.gov/19172691/)
11. Chambon P. A decade of molecular biology of retinoic acid receptors. *FASEB J.* 1996; 10: 940–954. PMID: [8801176](https://pubmed.ncbi.nlm.nih.gov/8801176/)
12. Johansson-Lindbom B, Svensson M, Wurbel MA, Malissen B, Marquez G, et al. Selective generation of gut tropic T cells in gut-associated lymphoid tissue (GALT): requirement for GALT dendritic cells and adjuvant. *J Exp Med.* 2003; 198: 963–969. PMID: [12963696](https://pubmed.ncbi.nlm.nih.gov/12963696/)
13. Kang SG, Park J, Cho JY, Ulrich B, Kim CH. Complementary roles of retinoic acid and TGF-beta1 in coordinated expression of mucosal integrins by T cells. *Mucosal Immunol.* 2011; 4: 66–82. doi: [10.1038/mi.2010.42](https://doi.org/10.1038/mi.2010.42) PMID: [20664575](https://pubmed.ncbi.nlm.nih.gov/20664575/)
14. Ohoka Y, Yokota A, Takeuchi H, Maeda N, Iwata M. Retinoic acid-induced CCR9 expression requires transient TCR stimulation and cooperativity between NFATc2 and the retinoic acid receptor/retinoid X receptor complex. *J Immunol.* 2011; 186: 733–744. doi: [10.4049/jimmunol.1000913](https://doi.org/10.4049/jimmunol.1000913) PMID: [21148038](https://pubmed.ncbi.nlm.nih.gov/21148038/)
15. Kim JM, Rasmussen JP, Rudensky AY. (2007) Regulatory T cells prevent catastrophic autoimmunity throughout the lifespan of mice. *Nat Immunol.* 2007; 8: 191–197. PMID: [17136045](https://pubmed.ncbi.nlm.nih.gov/17136045/)
16. Iwata M, Hirakiyama A, Eshima Y, Kagechika H, Kato C, et al. Retinoic acid imprints gut-homing specificity on T cells. *Immunity.* 2004; 21: 527–538. PMID: [15485630](https://pubmed.ncbi.nlm.nih.gov/15485630/)
17. Mora JR, Bono MR, Manjunath N, Weninger W, Cavanagh LL, Roseblatt M, et al. Selective imprinting of gut-homing T cells by Peyer's patch dendritic cells. *Nature.* 2003; 424: 88–93. PMID: [12840763](https://pubmed.ncbi.nlm.nih.gov/12840763/)
18. Coombes JL, Siddiqui KR, Arancibia-Carcamo CV, Hall J, Sun CM, Belkaid Y, et al. A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. *J Exp Med.* 2007; 204: 1757–1764. PMID: [17620361](https://pubmed.ncbi.nlm.nih.gov/17620361/)
19. Mucida D, Park Y, Kim G, Turovskaya O, Scott I, Krienenberg M, et al. Reciprocal TH17 and regulatory T cell differentiation mediated by retinoic acid. *Science* 2007; 317: 256–260. PMID: [17569825](https://pubmed.ncbi.nlm.nih.gov/17569825/)
20. Sun CM, Hall JA, Blank RB, Bouladoux N, Oukka M, Mora JR, et al. Small intestine lamina propria dendritic cells promote de novo generation of Foxp3 Treg cells via retinoic acid. *J Exp Med.* 2007; 204: 1775–1785. PMID: [17620362](https://pubmed.ncbi.nlm.nih.gov/17620362/)
21. Benson MJ, Pino-Lagos K, Roseblatt M, Noelle RJ. All-trans retinoic acid mediates enhanced T reg cell growth, differentiation, and gut homing in the face of high levels of co-stimulation. *J Exp Med.* 2007; 204: 1765–1774. PMID: [17620363](https://pubmed.ncbi.nlm.nih.gov/17620363/)
22. Nolting J, Daniel C, Reuter S, Stuelten C, Li P, Sucov H, et al. Retinoic acid can enhance conversion of naive into regulatory T cells independently of secreted cytokines. *J Exp Med.* 2009; 206: 2131–2139. doi: [10.1084/jem.20090639](https://doi.org/10.1084/jem.20090639) PMID: [19737861](https://pubmed.ncbi.nlm.nih.gov/19737861/)
23. Xiao S, Jin H, Korn T, Liu SM, Oukka M, Lim B, et al. Retinoic acid increases Foxp3+ regulatory T cells and inhibits development of Th17 cells by enhancing TGF-beta-driven Smad3 signaling and inhibiting IL-6 and IL-23 receptor expression. *J Immunol* 2008; 181: 2277–2284. PMID: [18684916](https://pubmed.ncbi.nlm.nih.gov/18684916/)
24. Hall JA, Cannons JL, Grainger JR, Dos Santos LM, Hand TW, Naik S, et al. Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor alpha. *Immunity.* 2011; 34: 435–447. doi: [10.1016/j.immuni.2011.03.003](https://doi.org/10.1016/j.immuni.2011.03.003) PMID: [21419664](https://pubmed.ncbi.nlm.nih.gov/21419664/)
25. Manel N, Unutmaz D, Littman DR. The differentiation of human T(H)-17 cells requires transforming growth factor-beta and induction of the nuclear receptor RORgamma. *Nat Immunol.* 2008; 9: 641–649. doi: [10.1038/ni.1610](https://doi.org/10.1038/ni.1610) PMID: [18454151](https://pubmed.ncbi.nlm.nih.gov/18454151/)
26. Iwakura Y, Ishigame H, Saijo S, Nakae S. Functional specialization of interleukin-17 family members. *Immunity.* 2011; 34: 149–162. doi: [10.1016/j.immuni.2011.02.012](https://doi.org/10.1016/j.immuni.2011.02.012) PMID: [21349428](https://pubmed.ncbi.nlm.nih.gov/21349428/)
27. Uematsu S, Fujimoto K, Jang MH, Yang BG, Jung YJ, Nishiyama M, et al. Regulation of humoral and cellular gut immunity by lamina propria dendritic cells expressing Toll-like receptor 5. *Nat Immunol.* 2008; 9: 769–776. doi: [10.1038/ni.1622](https://doi.org/10.1038/ni.1622) PMID: [18516037](https://pubmed.ncbi.nlm.nih.gov/18516037/)
28. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, et al. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature.* 2006; 441: 235–238. PMID: [16648838](https://pubmed.ncbi.nlm.nih.gov/16648838/)
29. Manicassamy S, Reizis B, Ravindran R, Nakaya H, Salazar-Gonzalez RM, Wang YV, et al. Activation of beta-catenin in dendritic cells regulates immunity versus tolerance in the intestine. *Science.* 2010; 329: 849–853. doi: [10.1126/science.1188510](https://doi.org/10.1126/science.1188510) PMID: [20705860](https://pubmed.ncbi.nlm.nih.gov/20705860/)

30. Wang S, Villablanca EJ, De Calisto J, Gomes DC, Nguyen DD, Mizoguchi E, et al. MyD88-dependent TLR1/2 signals educate dendritic cells with gut-specific imprinting properties. *J Immunol.* 2011; 187: 141–150. doi: [10.4049/jimmunol.1003740](https://doi.org/10.4049/jimmunol.1003740) PMID: [21646294](https://pubmed.ncbi.nlm.nih.gov/21646294/)
31. Yokota A, Takeuchi H, Maeda N, Ohoka Y, Kato C, Song SY, et al. GM-CSF and IL-4 synergistically trigger dendritic cells to acquire retinoic acid-producing capacity. *Int Immunol.* 2009; 21: 361–377. doi: [10.1093/intimm/dxp003](https://doi.org/10.1093/intimm/dxp003) PMID: [19190084](https://pubmed.ncbi.nlm.nih.gov/19190084/)
32. Laffont S, Siddiqui KR, Powrie F. Intestinal inflammation abrogates the tolerogenic properties of MLN CD103+ dendritic cells. *Eur J Immunol.* 2010; 40: 1877–1883. doi: [10.1002/eji.200939957](https://doi.org/10.1002/eji.200939957) PMID: [20432234](https://pubmed.ncbi.nlm.nih.gov/20432234/)
33. DePaolo RW, Abadie V, Tang F, Fehlner-Peach H, Hall JA, Wang W, et al. Co-adjutant effects of retinoic acid and IL-15 induce inflammatory immunity to dietary antigens. *Nature.* 2011; 471: 220–224. doi: [10.1038/nature09849](https://doi.org/10.1038/nature09849) PMID: [21307853](https://pubmed.ncbi.nlm.nih.gov/21307853/)
34. Crockett SD, Porter CQ, Martin CF, Sandler RS, Kappelman MD. Isotretinoin use and the risk of inflammatory bowel disease: a case-control study. *Am J Gastroenterol.* 2010; 105: 1986–1993. doi: [10.1038/ajg.2010.124](https://doi.org/10.1038/ajg.2010.124) PMID: [20354506](https://pubmed.ncbi.nlm.nih.gov/20354506/)
35. Jin MS, Kim SE, Heo JY, Lee ME, Kim HM, Paik SG, et al. Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. *Cell.* 2007; 130: 1071–1082. PMID: [17889651](https://pubmed.ncbi.nlm.nih.gov/17889651/)
36. Kang JY, Nan X, Jin MS, Youn SJ, Ryu YH, Mah S, et al. Recognition of lipopeptide patterns by Toll-like receptor 2-Toll-like receptor 6 heterodimer. *Immunity.* 2009; 31: 873–884. doi: [10.1016/j.immuni.2009.09.018](https://doi.org/10.1016/j.immuni.2009.09.018) PMID: [19931471](https://pubmed.ncbi.nlm.nih.gov/19931471/)
37. Manicassamy S, Ravindran R, Deng J, Oluoch H, Denning TL, Kasturi SP, et al. Toll-like receptor 2-dependent induction of vitamin A-metabolizing enzymes in dendritic cells promotes T regulatory responses and inhibits autoimmunity. *Nat Med.* 2009; 15: 401–409. doi: [10.1038/nm.1925](https://doi.org/10.1038/nm.1925) PMID: [19252500](https://pubmed.ncbi.nlm.nih.gov/19252500/)
38. Molenaar R, Knippenberg M, Goverse G, Olivier BJ, de Vos AF, O'Toole T, et al. Expression of retinaldehyde dehydrogenase enzymes in mucosal dendritic cells and gut-draining lymph node stromal cells is controlled by dietary vitamin A. *J Immunol.* 2011; 186: 1934–1942. doi: [10.4049/jimmunol.1001672](https://doi.org/10.4049/jimmunol.1001672) PMID: [21220692](https://pubmed.ncbi.nlm.nih.gov/21220692/)
39. Liu H, Komai-Koma M, Xu D, Liew FY. Toll-like receptor 2 signaling modulates the functions of CD4+CD25+ regulatory T cells. *PNAS.* 2006; 103: 7048–7053. PMID: [16632602](https://pubmed.ncbi.nlm.nih.gov/16632602/)
40. Netea MG, Suttmuller R, Hermann C, Van der Graaf CA, Van der Meer JW, van Krieken JH, et al. Toll-like receptor 2 suppresses immunity against *Candida albicans* through induction of IL-10 and regulatory T cells. *J Immunol.* 2004; 172: 3712–8. PMID: [15004175](https://pubmed.ncbi.nlm.nih.gov/15004175/)
41. Suttmuller RP, den Brok MH, Kramer M, Bennink EJ, Toonen LW, Kullberg BJ, et al. Toll-like receptor 2 controls expansion and function of regulatory T cells. *J Clin Invest.* 2006; 116: 485–494. PMID: [16424940](https://pubmed.ncbi.nlm.nih.gov/16424940/)
42. Bleich A, Mahler M, Most C, Leiter EH, Liebler-Tenorio E, Elson CO, et al. Refined histopathologic scoring system improves power to detect colitis QTL in mice. *Mamm Genome.* 2004; 15: 865–871. PMID: [15672590](https://pubmed.ncbi.nlm.nih.gov/15672590/)
43. Cario E, Gerken G, Podolsky DK. Toll-like receptor 2 controls mucosal inflammation by regulating epithelial barrier function. *Gastroenterology.* 2007; 132: 1359–1374. PMID: [17408640](https://pubmed.ncbi.nlm.nih.gov/17408640/)
44. Ey B, Eysing A, Klepak M, Salzman NH, Gothert JR, Runzi M, et al. Loss of TLR2 worsens spontaneous colitis in MDR1A deficiency through commensally induced pyroptosis. *J Immunol.* 2013; 190: 5676–5688. doi: [10.4049/jimmunol.1201592](https://doi.org/10.4049/jimmunol.1201592) PMID: [23636052](https://pubmed.ncbi.nlm.nih.gov/23636052/)
45. Round JL, Lee SM, Li J, Tran G, Jabri B, Chatila TA, et al. The Toll-like receptor 2 pathway establishes colonization by a commensal of the human microbiota. *Science.* 2011; 332: 974–977. doi: [10.1126/science.1206095](https://doi.org/10.1126/science.1206095) PMID: [21512004](https://pubmed.ncbi.nlm.nih.gov/21512004/)
46. Eksteen B, Mora JR, Haughton EL, Henderson NC, Lee-Turner L, Villablanca EJ, et al. Gut homing receptors on CD8 T cells are retinoic acid dependent and not maintained by liver dendritic or stellate cells. *Gastroenterology.* 2009; 137: 320–329. doi: [10.1053/j.gastro.2009.02.046](https://doi.org/10.1053/j.gastro.2009.02.046) PMID: [19233184](https://pubmed.ncbi.nlm.nih.gov/19233184/)
47. Komai-Koma M, Jones L, Ogg GS, Xu D, Liew FY. TLR2 is expressed on activated T cells as a co-stimulatory receptor. *Proc Natl Acad Sci.* 2004; 101: 3029–3034 PMID: [14981245](https://pubmed.ncbi.nlm.nih.gov/14981245/)
48. DePaolo RW, Tang F, Kim I, Han M, Levin N, Ciletti N, et al. Toll-like receptor 6 drives differentiation of tolerogenic dendritic cells and contributes to LcrV-mediated plague pathogenesis. *Cell Host & Microbe.* 2008; 4: 350–361.
49. Manoharan I, Hong Y, Suryawanshi A, Angus-Hill ML, Sun Z, Mellor AL, et al. TLR2-Dependent Activation of β -Catenin Pathway in Dendritic Cells Induces Regulatory Responses and Attenuates Autoimmune Inflammation. *The Journal of Immunology.* 2014; 193: 4203–4213. doi: [10.4049/jimmunol.1400614](https://doi.org/10.4049/jimmunol.1400614) PMID: [25210120](https://pubmed.ncbi.nlm.nih.gov/25210120/)