



RESEARCH ARTICLE

REVISED **Delivering Behaviour Change Interventions: Development of a Mode of Delivery Ontology [version 2; peer review: 2 approved]**

Marta M. Marques ^{1-3*}, Rachel N. Carey^{1*}, Emma Norris ¹, Fiona Evans¹, Ailbhe N. Finnerty¹, Janna Hastings ¹, Ella Jenkins ¹, Marie Johnston ⁴, Robert West ⁵, Susan Michie ¹

¹Centre for Behaviour Change, University College London, London, UK

²ADAPT SFI Research Centre, Trinity College Dublin, Dublin, Ireland

³Trinity Centre for Healthcare and Practice Innovation, Trinity College Dublin, Dublin, Ireland

⁴Aberdeen Health Psychology Group, University of Aberdeen, Aberdeen, Scotland, UK

⁵Research Department of Epidemiology & Public Health, University College London, London, UK

* Equal contributors

V2 **First published:** 10 Jun 2020, **5**:125
<https://doi.org/10.12688/wellcomeopenres.15906.1>

Latest published: 26 Feb 2021, **5**:125
<https://doi.org/10.12688/wellcomeopenres.15906.2>

Abstract

Background: Investigating and improving the effects of behaviour change interventions requires detailed and consistent specification of all aspects of interventions. An important feature of interventions is the way in which these are delivered, i.e. their mode of delivery. This paper describes an ontology for specifying the mode of delivery of interventions, which forms part of the Behaviour Change Intervention Ontology, currently being developed in the Wellcome Trust funded Human Behaviour-Change Project.

Methods: The Mode of Delivery Ontology was developed in an iterative process of annotating behaviour change interventions evaluation reports, and consulting with expert stakeholders. It consisted of seven steps: 1) annotation of 110 intervention reports to develop a preliminary classification of modes of delivery; 2) open review from international experts (n=25); 3) second round of annotations with 55 reports to test inter-rater reliability and identify limitations; 4) second round of expert review feedback (n=16); 5) final round of testing of the refined ontology by two annotators familiar and two annotators unfamiliar with the ontology; 6) specification of ontological relationships between entities; and 7) transformation into a machine-readable format using the Web Ontology Language (OWL) and publishing online.

Results: The resulting ontology is a four-level hierarchical structure comprising 65 unique modes of delivery, organised by 15 upper-level classes: Informational, Environmental change, Somatic, Somatic

Open Peer Review

Reviewer Status

	Invited Reviewers	
	1	2
version 2		
(revision)		
26 Feb 2021	report	report
	↑	↑
version 1		
10 Jun 2020		
	report	report

1. **Lucie Byrne-Davis** , University of Manchester, Manchester, UK

2. **Ann DeSmet** , Université Libre de Bruxelles, Brussels, Belgium

Any reports and responses or comments on the article can be found at the end of the article.

alteration, Individual-based/ Pair-based /Group-based, Uni-directional/Interactional, Synchronous/ Asynchronous, Push/ Pull, Gamification, Arts feature. Relationships between entities consist of *is_a*. Inter-rater reliability of the Mode of Delivery Ontology for annotating intervention evaluation reports was $\alpha=0.80$ (very good) for those familiar with the ontology and $\alpha=0.58$ (acceptable) for those unfamiliar with it.

Conclusion: The ontology can be used for both annotating and writing behaviour change intervention evaluation reports in a consistent and coherent manner, thereby improving evidence comparison, synthesis, replication, and implementation of effective interventions.

Keywords

ontology, intervention, behaviour, reporting, expert feedback, evidence synthesis, delivery



This article is included in the [Human Behaviour-Change Project](#) collection.

Corresponding authors: Marta M. Marques (MMOREIRA@tcd.ie), Rachel N. Carey (rachel.carey@ucl.ac.uk), Susan Michie (s.michie@ucl.ac.uk)

Author roles: **Marques MM:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Carey RN:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Norris E:** Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing – Review & Editing; **Evans F:** Investigation, Writing – Review & Editing; **Finnerty AN:** Formal Analysis, Investigation, Software, Visualization, Writing – Review & Editing; **Hastings J:** Investigation, Software, Writing – Review & Editing; **Jenkins E:** Formal Analysis, Investigation, Writing – Review & Editing; **Johnston M:** Conceptualization, Funding Acquisition, Investigation, Writing – Review & Editing; **West R:** Conceptualization, Methodology, Writing – Review & Editing; **Michie S:** Conceptualization, Funding Acquisition, Investigation, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work is supported by Wellcome through a collaborative award to The Human Behaviour-Change Project [201524]. MMM is funded by a Marie-Sklodowska-Curie fellowship [EU H2020 EDGE program grant agreement No. 713567].

Copyright: © 2021 Marques MM *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Marques MM, Carey RN, Norris E *et al.* **Delivering Behaviour Change Interventions: Development of a Mode of Delivery Ontology [version 2; peer review: 2 approved]** Wellcome Open Research 2021, 5:125 <https://doi.org/10.12688/wellcomeopenres.15906.2>

First published: 10 Jun 2020, 5:125 <https://doi.org/10.12688/wellcomeopenres.15906.1>

REVISED Amendments from Version 1

This version of the manuscript includes the changes made in response to the two reviewers. It provides more description of the peer-review process and how the Mode of Delivery Ontology can be used for different purposes. Two minor corrections were made to the definitions in the Ontology on the entities *video game mode of delivery* and *somatic alteration mode of delivery*.

Any further responses from the reviewers can be found at the end of the article

Introduction

Patterns of human behaviour contribute significantly to the global disease burden, as well as to a wide range of environmental and social problems (e.g. [Gakidou et al., 2017](#); [Watts et al., 2017](#)). The development of behaviour change interventions, defined as coordinated sets of activities designed to change specified behaviour patterns ([Michie et al., 2011](#)), can be an effective and cost-effective solution to such global problems. Research investigating the development, evaluation and implementation of behaviour change interventions, as well as evidence syntheses, demonstrate striking variability in effectiveness across different studies (see Cochrane database, e.g. [Flodgren et al., 2017](#); [Ussher et al., 2012](#)). Understanding this variability is difficult given the complexity of interventions, with variations in content and delivery potentially interacting with each other and with the intervention setting, population and target behaviour.

Being able to specify intervention characteristics in a way that facilitates replication and evidence synthesis is an important step in building evidence efficiently and cumulatively. This requires conceptual frameworks that organise knowledge using clear, coherent, and shared terminology ([Michie et al., 2017](#)). Such frameworks promote communication and collaboration across disciplines and research groups, and can be helpful in advancing knowledge generation to inform intervention development, implementation, evaluation, and reporting ([Craig et al., 2008](#); [Hoffmann et al., 2014](#); [Moher et al., 2001](#)). Another benefit of using conceptual frameworks is that they can enhance researchers' ability to examine associations between specific intervention components and outcomes ([Sheeran et al., 2017](#)). This allows for a more thorough understanding of interventions and how they bring about their effects which, in turn, can inform the development of more effective interventions.

Previously published classification systems for describing behaviour change interventions include the widely used Behaviour Change Techniques Taxonomy v1 (BCTTv1 ([Michie et al., 2013](#))), covering intervention content (e.g. [Newbury-Birch et al., 2014](#); [Zebis et al., 2016](#)). The BCTTv1 is a hierarchical taxonomy used to classify the potentially 'active ingredients' of behaviour change interventions, known as behaviour change techniques (BCTs) ([Michie et al., 2019a](#); [Michie et al., 2013](#); [Michie et al., 2015](#)). BCTTv1, which includes 93 discrete BCTs, has been used to identify and define

BCTs in intervention research ([Newbury-Birch et al., 2014](#); [Paul et al., 2017](#); [Young et al., 2014](#)) and to categorise intervention content in evidence syntheses ([Arnott et al., 2014](#); [Jones et al., 2014](#)). By providing a common language with which to describe interventions, BCTTv1 has facilitated a level of rigour and specificity in reporting intervention content that was not previously commonplace ([Sheeran et al., 2017](#)). While BCTTv1 and other classification systems for intervention content (e.g. [Hollands et al., 2017](#)) have provided a shared language for specifying intervention content, there are other crucial aspects of behaviour change interventions that have received comparatively little attention, including *how* such content is delivered ([Dombrowski et al., 2016](#)).

Ontologies

BCTTv1 is an example of a taxonomy, a knowledge representation structure in which a controlled vocabulary of agreed-upon terms is arranged hierarchically. An *ontology* is a more expressive structure for organising knowledge (see glossary of italicised terms, [Table 1](#)). It includes a controlled vocabulary, unambiguous identifiers for each entity, and additional information such as synonyms and examples of usage. It includes relationships between entities, usually beyond the hierarchical class-subclass relationship as well as a formal, logic-based encoding of domain knowledge where possible ([Arp et al., 2015](#); [Hastings, 2017](#); [Larsen et al., 2017](#); [Michie & Johnston, 2017](#); [Norris et al., 2019](#)). Ontologies enable entities to be compared and integrated across fields of study and allow large datasets to be synthesised efficiently using computational tools (e.g. in biology, the [Gene Ontology](#) ([Ashburner et al., 2000](#))).

The potential for ontologies to facilitate knowledge synthesis in behaviour change is being developed in the [Human Behaviour-Change Project](#) ([Michie et al., 2018](#); [Michie et al., 2020a](#); [Michie et al., 2020b](#)). This collaboration between behavioural scientists, computer scientists and systems architects is building a database and platform for researchers, practitioners and policy-makers to address variants of the 'big question' of behaviour change: "What works, compared with what, how well, with what exposure, with what behaviours (for how long), for whom, in what settings and why?" Answering this involves extending previous work to classify all entities of behaviour change interventions and the relationships between them, i.e. a *Behaviour change intervention ontology* (BCIO), specified by a controlled vocabulary that by the upper level of the BCIO ([Michie et al., 2020b](#)) contains 42 entities. The *Behaviour change intervention delivery* entity of the ontology (i.e. the means by which BCI content is provided), comprises (a) *BCI Source* (i.e., a role played by a person, population or organisation that provides a behaviour change intervention), (b) *BCI Schedule of delivery* (an attribute of a behaviour change intervention that involves its temporal organisation), (c) *BCI Style of delivery* (an attribute of a BCI delivery that encompasses the characteristics of how a behaviour change intervention is communicated), and (d) *BCI Mode of delivery* (an attribute of a BCI delivery that is the physical or informational medium through which a behaviour change intervention is provided).

Table 1. Glossary.

Term	Definition	Source
Annotation	Process of coding selected parts of documents or other resources to identify the presence of ontology entities.	Michie et al., 2018.
Annotation guidance manual	Written guidance on how to identify and tag pieces of text from intervention evaluation reports with specific codes relating to entities in the ontology.	
Basic Formal Ontology (BFO)	An upper level ontology consisting of continuants and occurrents developed to support integration, especially of data obtained through scientific research.	Arp et al., 2015.
Entity	Anything that exists, that can be a continuant or an occurrent as defined in the Basic Formal Ontology.	Arp et al., 2015.
EPPI-Reviewer	A web-based software program for managing and analysing data in all types of systematic review (meta-analysis, framework synthesis, thematic synthesis etc). It manages references, stores PDF files and facilitates qualitative and quantitative analyses such as meta-analysis and thematic synthesis. It also has a facility to annotate published papers.	Thomas et al., 2010; <i>EPPI-Reviewer 4:</i> http://eppi.ioe.ac.uk/eppireviewer4/ <i>EPPI-Reviewer Web Version:</i> https://eppi.ioe.ac.uk/eppireviewer-web/
GitHub	A web-based platform used as a repository for sharing code, allowing version control.	https://github.com/
Inter-rater reliability	Statistical assessment of similarity and dissimilarity of coding between two or more coders. If inter-rater reliability is high this suggests that ontology entity definitions and labels are being interpreted similarly by the coders.	Gwet, 2014. <i>Handbook of inter-rater reliability: The definitive guide to measuring the extent of agreement among raters.</i> Gaithersburg, Advanced Analytics.
Interoperability	Two systems are interoperable if data coming from each system can be used by the other system. Note: An ontology is interoperable with another ontology if it can be used together with or re-uses parts from the other ontology	http://www.obofoundry.org/principles/fp-010-collaboration.html
Issue tracker	An online log for problems identified by users accessing and using an ontology.	BCIO Issue Tracker: https://github.com/HumanBehaviourChangeProject/ontologies/issues
OBO Foundry	The Open Biological and Biomedical Ontology (OBO) Foundry is a collective of ontology developers that are committed to collaboration and adherence to shared principles. The mission of the OBO Foundry is to develop a family of interoperable ontologies that are both logically well-formed and scientifically accurate.	Smith et al., 2007; www.obofoundry.org/
Ontology	A standardised framework providing a set of terms that can be used for the consistent annotation (or "tagging") of data and information across disciplinary and research community boundaries.	Arp et al., 2015.
Parent class	A class within an ontology that is hierarchically related to one or more child (subsumed) classes such that all members of the child class are also members of the parent class and all properties of the parent class are also properties of the child class.	Arp et al., 2015.
Reconciliation	The process of discussing differences between the annotations of two paired annotators on the same papers. Differences are discussed before a final reconciled version of coding for each paper is produced.	Stan et al., 2014.
ROBOT	An automated command line tool for ontology workflows.	Jackson et al., 2019, http://robot.obolibrary.org
Unique resource identifier (URI)	A string of characters that unambiguously identifies an ontology or an individual entity within an ontology. Having URI identifiers is one of the OBO Foundry principles.	http://www.obofoundry.org/principles/fp-003-uris.html
Web Ontology Language (OWL)	A formal language for describing ontologies. It provides methods to model classes of "things", how they relate to each other and the properties they have. OWL is designed to be interpreted by computer programs and is extensively used in the Semantic Web where rich knowledge about web documents and the relationships between them are represented using OWL syntax.	https://www.w3.org/TR/owl2-quick-reference/

Delivery of Behaviour Change Interventions

An important characteristic of behaviour change interventions is the method or methods by which the content (e.g. BCTs) is brought to its target population (i.e. the intervention's mode of delivery; MoD). MoDs can act synergistically or antagonistically with the intervention techniques in influencing intervention outcomes and effects. An example of this is a meta-analysis of evidence about the effectiveness of smoking cessation interventions, which found effectiveness to be higher with increasing numbers of intervention techniques but only if delivered in person and not when delivered in written form (Black *et al.*, 2020).

Several systematic reviews have extracted information about MoDs (Bock *et al.*, 2014; van Genugten *et al.*, 2016; MacDonald *et al.*, 2016), and an annotation scheme for MoD within internet-based interventions has been developed (Webb *et al.*, 2010). However, MoD has received comparatively little attention in intervention research (Dombrowski *et al.*, 2016), and there is a lack of clarity and consensus across behavioural intervention research regarding how MoD is defined, what it includes, and how it should be reported. This is in contrast to the reporting of BCTs as the content of behaviour change interventions, for which there is now wide shared understanding, for example, featuring in the Encyclopaedia of Behavioural Medicine (Michie *et al.*, 2019a) and in many hundreds of publications. The various conceptualisations of MoD, and the lack of a shared language or framework with which to describe it has made the study of interactions between it and other intervention entities difficult to analyse systematically (Dombrowski *et al.*, 2016). Here, we define MoD as the attribute of BCI delivery that is the informational or physical medium through which a behaviour change intervention is provided (Michie *et al.*, 2020b). For example, providing someone with information about the health consequences of performing a particular behaviour could be conducted face-to-face (e.g. by a GP), through a poster or leaflet, or through a digital device (e.g. an app). 'Item 6: How' of the TiDieR framework highlights the need for researchers to clearly specify the MoD of the intervention. An example of a classification that briefly addresses MoD is the Effective Practice and Organisation of Care (EPOC) Taxonomy (Effective Practice and Organisation of Care (EPOC), 2015) which includes a category for "How and when care is delivered" and "Information and Communication Technology" including some elements of MoD such as group vs individual care. EPOC was developed specifically to classify delivery of health systems interventions. A framework that systematically describes a vast range of MoD entities that can be implemented in behaviour change interventions for any domain of human behaviour was needed. An ontology provides a mechanism for doing this. The development of an MoDontology that can be linked to other ontologies relevant to behaviour change interventions would advance scientific understanding, the development and evaluation of interventions and methods for evidence synthesis.

Aim

The aim of the MoD Ontology is to provide a clear, usable and reliable classification system to specify the MoDs of behaviour change interventions, including single BCTs. The

development of an ontology with clear and unambiguously defined terms enables precision of reporting, which in turn promotes evidence synthesis, replication and analyses of associations between MoDs, other intervention characteristics and intervention outcomes.

Methods

The ontology was developed in seven iterative steps (detailed below), involving reviewing existing classification systems, *annotation* of behaviour change intervention reports (including testing of *inter-rater reliability*) and feedback from international expert stakeholders (outlined in Table 2).

Step 1: Development of the preliminary ontology and piloting

Initial descriptions of MoD *entities* were extracted from 20 published behaviour change intervention evaluation reports, randomly selected using a random number generator from a larger *database of reports* annotated by behaviour change techniques and mechanisms of action (Michie *et al.*, 2018), covering a range of health behaviours. Next, two researchers independently piloted the preliminary MoD ontology with another set of intervention reports, taken from the same database and using the same selection method. Guidance on how to annotate papers for MoD was developed by the research team, providing clear instructions on how to code each entity, including definitions and examples for each. Reports were annotated in batches of 10 until a satisfactory and stable criterion of inter-rater reliability was achieved. Inter-rater reliability of the extent to which researchers capture the same information from a report was measured in two ways. The first was percentage agreement of instances where both researchers had annotated an MoD. The second was the proportion of times annotators agreed on a code when both of them captured the same information from a report. This was calculated at every level of the hierarchy, and it was performed using Cohen's Kappa (Cohen, 1960), in Microsoft Excel 365. Kappa values >.61 were deemed as 'substantial' and values >.81 as 'strong' (Landis & Koch, 1977). The preliminary ontology was revised and updated iteratively throughout the annotation process. Where there were discrepancies between the two annotators, these were discussed, and amendments were made to the ontology if both annotators judged that these changes would improve clarity. In the case of disagreement, a senior member of the research team was consulted.

Step 2: Stakeholder review (Round 1)

Nine international behavioural scientists with experience in behaviour change interventions, across a range of behavioural domains, were invited to provide feedback on the structure, content and terminology of the preliminary MoD Ontology. Following small adjustments based on this feedback, the MoD Ontology was published online, and a wider international research community was invited through mailing lists to submit feedback using an *Qualtrics* form presenting the preliminary MoD structure, and entity labels and definitions (see <https://osf.io/eyn3b/> (West *et al.*, 2020)). Twenty-five behavioural scientists responded to indicate whether 1) there were any entities missing, 2) the structure was coherent, 3) there were changes needed

Table 2. Steps for developing the Mode of Delivery Ontology.

Phase	Step	Methods
Initial development	1. Developing and piloting a preliminary ontology	Data extraction from 120 BCI reports; - 20 reports for initial draft + 100 for improvements and inter-rater reliability calculations; Group discussions
	2. Requesting feedback on preliminary ontology from expert stakeholders	Open peer review from 25 experts; Group discussions
Testing and refinement	3. Testing & refining ontology through second round of data annotations	Data annotations from 55 BCI reports; inter-rater reliability calculations; inter-rater reliability; Group discussions
	4. Requesting feedback on refined ontology from experts	Open peer-review from 16 experts; consultation with an ontology expert; Group discussions
Consolidation of changes and agreement on final version	5. Testing & finalising ontology through final round of data annotations	Data annotations from 150 BCI reports; inter-rater reliability calculations
	6. Specifying the relationships between entities	Group discussions
	7. Transforming into machine-readable format	Ontology content was transformed automatically into an OWL ontology using the ROBOT library's template functionality

MoD, mode of delivery; BCI, Behaviour change intervention.

in the terminology of the labels and definitions, and 4) there were additional suggestions for improvement.

Step 3: Inter-rater reliability testing (Round 2)

The revised version was used to annotate MoD entities in a set of 55 published reports, randomly selected using a random number generator from the database mentioned in Step 1 (Michie *et al.*, 2018). These papers covered the behavioural domains of physical activity, diet and smoking. Annotation of the reports was conducted independently by two researchers. The annotation process was carried out in batches of five papers. After every batch, annotations were compared, and discrepancies discussed. Inter-rater reliability was calculated using the same procedure as in Step 1. Where there were discrepancies, consensus was reached through discussion.

Step 4: Stakeholder review (Round 2)

Experts who provided feedback in Step 2 were invited to submit feedback on the revised ontology. Experts were sent an email with a request to review the structure, labels and definitions of each entity, and indicate whether the structure was coherent and whether there was anything missing and provide suggestions for improved terminology. During this step, an ontology expert (JH) was consulted regarding the structure and definitions.

Step 5: Inter-rater reliability testing (Round 3)

To test the range of applicability of this revised version of the MoD Ontology (as well as the *annotation guidance manual*), we conducted a final round of annotations as part of the annotations being conducted in the Human Behaviour-Change Project. First, two developers of the MoD ontology annotated reports that were selected from a database of reports used in the Human Behaviour-Change Project (Michie *et al.*, 2017) (see <https://osf.io/myje6/> (West *et al.*, 2020)). These annotations

were conducted using *EPPI reviewer 4* software (Thomas *et al.*, 2010). An open alternative to this software used for annotation is *PDFAnno* (Shindo *et al.*, 2018). All reports were randomised controlled trials from one of three datasets: Cochrane Reviews, *papers annotated for behaviour change techniques* and papers from the IC-SMOKE project (De Bruin *et al.*, 2016) (list of systematic reviews included as *Extended data* at <https://osf.io/myje6/> (West *et al.*, 2020)). There was a reconciliation process after the first batch of 10, followed by any necessary amendments to the annotation manual. These amendments mainly involved the inclusion of examples (e.g. illustrating when to code or not to code certain pieces of information as MoD).

To examine the usability of the MoD Ontology for researchers and intervention developers with no prior knowledge of the MoD Ontology, we conducted a final round of inter-rater reliability assessment by asking two researchers unfamiliar with the ontology and without specific expertise in modes of delivery to annotate a random sample of randomised controlled trials from a *database of papers annotated by BCTs*, with no restrictions on the outcome behaviour. Inter-rater reliability was assessed using Krippendorff's Alpha (Hayes & Krippendorff, 2007), using Python 3.6 (code available on *GitHub* (Finnerty & Moore, 2020)).

Step 6: Specifying relationships within the MoD Ontology

The research team developed relationships between ontology entities to formally capture the types of knowledge that are present in the ontology. The relationships were specified following best practices from Basic Formal Ontology (BFO) described in *Arp et al.* (2015) and Relation Ontology (Smith *et al.*, 2005). Relationships can be generic and shared across multiple ontologies (e.g. the "is a" relationship between classes

where one class is a subclass of another class, or the “part of” relationship which captures the relationship between wholes and their parts) or they can be domain specific, which are introduced when needed to formally capture relationships unique to a given domain.

Step 7: Making the MoD Ontology machine-readable and available online

The MoD Ontology was initially developed as a table of entities, with separate rows for each entity annotated in columns for different types of annotation, including a primary label, definition, synonyms and relationships. When the MoD Ontology was at a stable level of development for initial release, it was converted into the *Web Ontology Language (OWL)* (Antoniou & van Harmelen, 2004) format, enabling it to be viewed and visualised using ontology software such as *Protégé* and to be compatible with other ontologies and software tools. The conversion to OWL used the *ROBOT* ontology toolkit library (Jackson *et al.*, 2019), which provides a facility to create well-formatted ontologies from templates. A ROBOT template can be prepared easily in common spreadsheet software, annotated with instructions for translation from spreadsheet columns to OWL language and metadata entities. Within the input template spreadsheet, separate columns represent the entity ID (e.g. BCIO:011004), name, definition, relationship with other entities, examples and synonyms.

This OWL version of the MoD Ontology was then stored on the [project GitHub repository](#) (Finnerty & Moore, 2020), as *GitHub* has an *issue tracker*, which allows feedback to be submitted by members of the community that can be responded to, and if necessary, addressed in subsequent releases. When the full BCIO has been finalised, it will be submitted to the *OBO Foundry* (Smith *et al.*, 2007).

Results

Step 1: Development of the preliminary ontology and piloting

The data extracted from the behaviour change intervention reports led to the identification of 160 unique entities, which were represented in a four-level hierarchical structure, as well as two ‘cross-cutting’ entities (a description of the preliminary version is available as *Extended data* at <https://osf.io/gu5ke/> (West *et al.*, 2020)). A hundred reports were annotated, with adjustments made to the ontology as a result of the first 70; the ontology was stable for the final 30 reports. Average agreement between annotators for each batch of 10 reports varied between 72% and 95%. Inter-rater reliability was calculated for each level of the hierarchy separately and considered to be ‘good’ for all levels (% agreement 86.6 to 97.8; Kappa 0.68 to 0.97). Reliability was also calculated for each of the cross-cutting entities (Kappa = .55 and .75). Further details on the inter-rater reliability and changes made to the MoD Ontology in this step can be found as *Extended data* at: <https://osf.io/r3wn2/> (West *et al.*, 2020).

Step 2: Stakeholder review (Round 1)

Feedback on the MoD ontology through the open review feedback form was received by 25 experts, of which 18 were from

universities, 5 were from commercial sector organisations, 1 from public sector organisations and 1 from third sector. Twelve experts were from the United Kingdom, 2 from the United States of America, 3 from Ireland, 1 from Canada, 1 from the Netherlands, 1 from New Zealand, and we have no information about the country for the remaining 5 experts. These data were collated, synthesised, and discussed among the research team. This led to further amendments to the structure, content and terminology (full details on the feedback and corresponding changes made to the MoD Ontology are available as *Extended data* at <https://osf.io/95n3a/> (West *et al.*, 2020)).

Step 3: Inter-rater reliability testing (Round 2)

For the 55 papers annotated in this round, agreement for whether a particular entity was considered an MoD was 61%; and agreement on the specific MoD code assigned was 87.9% (Kappa = .857) (inter-rater reliability results are available as *Extended data* at <https://osf.io/sw2jv/> (West *et al.*, 2020)).

Step 4: Stakeholder review (Round 2)

Feedback was received from 16 of the 25 experts invited. Based on this, the following changes were made: 1) the entities “other” and “unclear” were removed, as all entities represented in an ontology need to be fully specified; and (2) increased clarity was provided on how the cross-cutting entities related to the other upper-level classes (see <https://osf.io/3zhbc/> (West *et al.*, 2020) for more details”).

For the revised version, definitions were developed using pre-specified guidance, with the standard format of definitions being: A is a B that C, or involves or relates to C in some way, where A is the class being defined, B is a *parent class* and C describes a set of properties of A that distinguish it from other members of B (Michie *et al.*, 2019b).

Step 5: Inter-rater reliability testing (Round 3)

For the annotations conducted by researchers familiar with the MoD ontology, a very good agreement ($\alpha=0.80$) was achieved after annotating 50 reports (25 smoking and 25 physical activity). For the annotations conducted by researchers unfamiliar with the ontology, acceptable agreement ($\alpha=0.58$) was achieved after annotating 96 papers, targeting various behaviours (26 physical activity; 22 diet; 13 alcohol; 11 treatment adherence; nine sexual behaviours; seven multiple health behaviours; two for prescription, smoking, and screening, respectively; and one paper for organ donation and one for oral health) (Hayes & Krippendorff, 2007) (inter-rater reliability results are available as *Extended data* at <https://osf.io/efp4x/> (West *et al.*, 2020)).

Step 6: Specifying relationships within the MoD Ontology

Currently, the only relationship used in the ontology represent its hierarchical structure, i.e. “subclass of” (is_a) relationships (e.g. *face to face MoD* “is_a” *human interactional MoD*). Formal representations of knowledge using explicit logical relationships allow computational tools to perform additional checks and inferences to enhance the resulting consistency of reporting for complex interventions.

Step 7 - Making the MoD Ontology machine-readable and available online

A downloadable version of the final MoD Ontology can be found on [GitHub](#) (Finnerty & Moore, 2020). The hierarchical structure, labels, *uniform resource identifiers (URIs)* and definitions for all entities are described in [Table 3](#). The ontology is accompanied by an annotation manual that provides guidance on how to annotate for these entities in reports of behaviour change interventions (available as *Extended data* at <https://osf.io/4j2xh/> (West et al., 2020)). The final MoD Ontology presents a four-level hierarchical structure comprising 65 entities. There are 15 upper-level classes: 1.1. Informational MoD; 1.2. Environmental change MoD; 1.3. Somatic MoD; 1.4. Somatic alteration MoD; 1.5. Individual-based MoD vs 1.6. Pair-based MoD, vs 1.7. Group-based MoD; 1.8. Uni-directional MoD vs. 1.9. Interactional MoD; 1.10. Synchronous MoD vs. 1.11. Asynchronous MoD; 1.12. Push MoD vs. 1.13. Pull MoD; 1.14. Gamification MoD; 1.15. Arts feature MoD. The first upper-level classes include lower level entities (sub-classes). For example, *Informational MoD* includes *Printed material MoD*, which includes sub-classes of *Letter MoD*, *Public notice MoD*, *Printed publication MoD*, and *Labelling MoD*. Entities from 1.5 to 1.15 correspond to entities that can be present at the same time as at least one of the other MoD. For example, an intervention that is *delivered through face to face* (sub-class of *Human interactional MoD*), can also be classified as an *Individual-based* or *Group-based MoD*. It is worth noting that, given the exponential growth in new technologies, this MoD Ontology captures a specific moment in time, and will need updating as technologies and methods develop.

Discussion

Given the lack of classification systems providing comprehensive coverage of how behaviour change interventions and techniques are delivered, we developed the first ontology of modes of delivery (MoD). This ontology consists of 65 entities organised in 15 upper-level entities. Inter-rater reliability was found to be 0.80 (very good) for those familiar with the ontology and 0.58 (acceptable) for those unfamiliar with it, as assessed by Krippendorff's alpha. Together with *Source*, *Schedule* and *Style* it represents the characteristics of *Delivery* of a behaviour change intervention. Ontologies aim to be dynamic representations that are updated according to new evidence on entities and relationships. As with other lower level ontologies that form part of the BCIO (Michie et al., 2020b), the MoD Ontology will be improved upon and refined through application and feedback by users.

The MoD Ontology contributes to the growing number of methodological resources now freely available to intervention researchers (e.g. Bartholomew et al., 2011; Hoffmann et al., 2014; Hollands et al., 2017; Michie et al., 2013). For example, a [Theory and Techniques Tool](#) available for free online, provides an interactive dataset of links between BCTs and their mechanisms of action (i.e. the processes through which BCTs have their effects). The tool was informed by data from evidence synthesis (Carey et al., 2019) and expert consensus

(Connell et al., 2019), which were triangulated (Johnston et al., 2018); all three sets of data are available in the tool.

The MoD Ontology contributes to a larger programme of work developing ontologies for other intervention components, the Human Behaviour-Change Project (Michie et al., 2018; Michie et al., 2020a). Within this project, lower level ontologies are being developed for intervention-related entities of content, delivery, tailoring, context, engagement, mechanism of action, and outcome behaviour within the BCIO (Michie et al., 2020b). These ontologies have been developed using an explicit, standardised, and tested method for ontology development created within the Human Behaviour-Change Project (Wright et al., 2020). As the development of the MoD ontology started prior to the development of the BCIO, the process of development was slightly different from the one described in this collection (Wright et al., 2020), containing more rounds of expert feedback and inter-rater reliability testing.

The MoD ontology provides a crucial contribution to the much needed body of research examining the links between MoDs and the content of behaviour change interventions, using the BCTTv1 or other classification systems of techniques (e.g. Knittle et al., 2020; Kok et al., 2016). For example, coding existing behaviour change interventions for their modes of delivery and BCTs can increase our understanding of which mode(s) of delivery are the most effective in delivering a given BCT. Further, by linking with other HBCP ontologies characterising behaviour change interventions, it will be possible to go a step further and identify which MoD(s) are more appropriate for different behaviours, populations, contexts, if they need to be tailored, and their potential for reach and engagement.

Strengths and limitations

These ontologies provide a framework for applying machine learning and reasoning algorithms to synthesise and interpret evidence, as well as predict outcome. This allows real-time up-to-date evidence to be interrogated by users such as policy-makers, planners and intervention designers to answer variants of the "big question": "What works, compared with what, how well, with what exposure, with what behaviours (for how long), for whom, in what settings and why?", across a wide range of contexts. This body of work has the potential to have far-reaching use by and implications for policy-makers, practitioners and researchers - for example, by informing evidence-based guidelines and identifying knowledge gaps.

Further, the use of entity IDs for each entity in the ontology provides a machine-readable identifier for integration in future systems and also allows *interoperability* between existing ontologies.

Several limitations should be noted about the development process, and the resulting MoD Ontology. Given the rapid growth in new technologies and the fast-moving pace of behavioural science research, the MoD Ontology will need updating and refining as existing methods develop and new methods emerge. However, this is common to all ontologies and indeed considered 'best practice' in ontology development (Arp et al., 2015).

Table 3. Entity labels, definitions, URIs and examples of usage for Mode of Delivery Ontology entities.

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
Informational mode of delivery BCIO:011001				Mode of delivery that involves intentional transmission of a representation of the world to an intervention recipient with the aim of changing that person's representation of the world.	This includes delivery of rewards, prompts, and cues that result in learning and information about the environment and environmental contingencies.
	Human interactional mode of delivery BCIO:011002			Informational mode of delivery that involves a person as intervention source who interacts with an intervention recipient in real time.	
		Face to face mode of delivery BCIO:011003		Human interactional mode of delivery that involves an intervention source and recipient being together in the same location and communicating directly.	
		At-a-distance mode of delivery BCIO:011004		Human interactional mode of delivery that involves an intervention source and recipient being in different locations and communicating through a communication channel.	
	Printed material mode of delivery BCIO:011005			Informational mode of delivery that involves use of printed material.	Can include paper, acetate, text, diagrams and photographic images.
		Letter mode of delivery BCIO:011006		Printed material mode of delivery that involves a letter or postcard that can be sent through the post or handed directly to the recipient.	
		Public notice mode of delivery BCIO:011007		Printed material mode of delivery that involves display of a poster, sign or notice in a public location.	
		Printed publication mode of delivery BCIO:011008		Printed material mode of delivery that involves use of a printed publication.	Includes leaflets, brochures, newspapers, newsletter, booklets, magazines, manuals or worksheets.
		Labelling mode of delivery BCIO:011009		Printed material mode of delivery that involves information printed on a product or its packaging, or a label attached to or included with, a product or its packaging, and aims to convey information about that product.	
	Electronic mode of delivery BCIO:011010			Informational mode of delivery that involves electronic technology in the presentation of information to an intervention recipient.	

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
		Television mode of delivery BCIO:011011		Electronic mode of delivery that involves presentation of information that is broadcast and displayed by television.	Includes internet and satellite television.
		Mobile digital device mode of delivery BCIO:011012		Electronic mode of delivery that involves presentation of information by a handheld mobile digital device that can store, retrieve and process data.	
		Computer mode of delivery BCIO:011013		Electronic mode of delivery that involves presentation of information by a desktop or laptop computer.	
		Electronic billboard mode of delivery BCIO:011014		Electronic mode of delivery that involves presentation of information by an electronic screen positioned in a public location.	
		Wearable electronic device mode of delivery BCIO:011015		Electronic mode of delivery that involves presentation of information by an electronic screen positioned in a public location.	Includes a watch, clip-on device, spectacles, in-ear device, vibrating device.
		Electronic environmental object mode of delivery BCIO:011016		Electronic mode of delivery that involves an electronic device positioned in the environment of the intervention recipient that can gather information and respond to commands.	Includes robots, and internet of things.
		3-D projection mode of delivery BCIO:011017		Electronic mode of delivery that involves presentation of a 3-D image.	Includes hologram but does not include virtual reality headsets.
		Virtual reality mode of delivery BCIO:011018		Electronic mode of delivery that involves use of virtual reality through a virtual reality headset and optionally body movement sensors.	
		Playable electronic storage mode of delivery BCIO:011019		Electronic mode of delivery that involves presentation of information stored on an object that is inserted into a playing device.	Includes cassettes, video tapes, DVDs, CDs.
		Radio broadcast mode of delivery BCIO:011020		Electronic mode of delivery that involves presentation of audio information that is broadcast and received by a radio receiver.	Includes cassettes, video tapes, DVDs, CDs.
		Call mode of delivery BCIO:011021		Electronic mode of delivery that involves a communication process in which a signal is sent by a caller to a recipient to alert them of the communication intent, giving the recipient the opportunity to engage with the communication.	Includes automated calls and audio messaging.

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
			Audio call mode of delivery BCIO:011022	Call mode of delivery that involves only audio information in the communication.	
			Video call mode of delivery BCIO:011023	Call mode of delivery that involves video and audio information in the communication.	
			Messaging mode of delivery BCIO:011024	Call mode of delivery that involves textual information in the communication.	Text message can include emojis, and additional audio and pictorial material. Includes SMS, WhatsApp and other messaging services.
		Email mode of delivery BCIO:011025		Electronic mode of delivery that involves communication by email.	
		Video game mode of delivery BCIO:011026		Electronic mode of delivery that involves the intervention recipient playing a computer game	
		Website mode of delivery BCIO:011027		Electronic mode of delivery that involves the intervention recipient interacting with a website.	
		Mobile application mode of delivery BCIO:011028		Electronic mode of delivery that involves the intervention recipient interacting with a mobile application.	
		E-book mode of delivery BCIO:011029		Electronic mode of delivery that involves the intervention recipient being given access to an e-book.	
	Audio informational mode of delivery BCIO:011030			Informational mode of delivery that involves sound.	
	Visual informational mode of delivery BCIO:011031			Informational mode of delivery that involves visual images.	
	Textual mode of delivery BCIO:011032			Informational mode of delivery that involves written text.	
Environmental change mode of delivery BCIO:011033				Mode of delivery that involves changing the physical shape, size, structure or appearance of objects in the environment of the intervention recipient.	This does not include use of textual or pictorial information. It includes lighting, speed humps, use of music, shape and size of containers of consumables.
Somatic mode of delivery BCIO:011034				Mode of delivery that involves devices or substances that alter bodily processes or structure.	
	Ingestion mode of delivery BCIO:011035			Somatic mode of delivery that involves ingestion of a chemical into the body.	

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
		Transdermal mode of delivery BCIO:011036		Ingestion mode of delivery that involves ingestion of a chemical through the skin.	
		Alimentary mode of delivery BCIO:011037		Somatic mode of delivery that involves ingestion of a chemical through the stomach or intestine.	
			Pill mode of delivery BCIO:011038	Alimentary mode of delivery that involves swallowing of a pill or oral capsule.	
			Ingestible liquid mode of delivery BCIO:011039	Alimentary mode of delivery that involves swallowing of a liquid.	
		Buccal mode of delivery BCIO:011040		Ingestion mode of delivery that involves absorption of a chemical through the lining of the buccal cavity.	
		Inhalation mode of delivery BCIO:011041		Ingestion mode of delivery that involves absorption of a chemical through the upper airways or lungs by inspiration.	
		Injection mode of delivery BCIO:011042		Ingestion mode of delivery that involves a chemical being introduced into body tissue through a hollow needle that punctures the skin.	
			Subcutaneous injection mode of delivery BCIO:011043	Injection mode of delivery in which the tissue receiving the chemical is subcutaneous tissue.	
			Intravenous injection mode of delivery BCIO:011044	Injection mode of delivery in which the tissue receiving the chemical is subcutaneous tissue.	
			Intramuscular injection mode of delivery BCIO:011045	Injection mode of delivery in which the tissue receiving the chemical is muscle.	
		Wearable ingestion mode of delivery BCIO:011046		Injection mode of delivery in which the tissue receiving the chemical is muscle.	Includes insulin pump.
	Chewable substance mode of delivery BCIO:011047			Ingestion mode of delivery that involves chewing of a soft material.	This includes chewing gum. Often involves ingestion of a chemical that is released by chewing and absorbed through the lining of the buccal cavity.

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
	Physical stimulus mode of delivery BCIO:011048			A mode of delivery that involves application of a physical stimulus to the body.	
		Light exposure mode of delivery BCIO:011049		Physical stimulus mode of delivery that involves exposure of light to the body.	
		Temperature mode of delivery BCIO:011050		Physical stimulus mode of delivery that involves application of heat or cold to the body.	
		Electrical stimulation mode of delivery BCIO:011051		Physical stimulus mode of delivery that involves application of electrical stimulation to the body.	
		Physical pressure mode of delivery BCIO:011052		Physical stimulus mode of delivery that involves application of physical pressure to the outside of the body.	Includes massage.
		Wearable stimulus mode of delivery BCIO:011053		Physical stimulus mode of delivery that involves a device that is worn on the body.	
Somatic alteration mode of delivery BCIO:011054				Mode of delivery that involves modifying the structure of the body of the recipient of the intervention.	Includes surgery.
Individual-based mode of delivery *BCIO:011055				Mode of delivery that involves one recipient in the location where the intervention is delivered.	
Pair-based mode of delivery *BCIO:011056				Mode of delivery that involves two recipients in the location where the intervention is delivered who have an interpersonal relationship.	
Group-based mode of delivery *BCIO:011057				Mode of delivery that involves three or more people in the location where the intervention is delivered.	
Uni-directional mode of delivery **BCIO:011058				Mode of delivery in which the only causal influence is from the intervention source to the recipient.	
Interactional mode of delivery **BCIO:011059				Mode of delivery in which there is causal influence from the intervention source to the recipient and from the recipient to the source.	
Synchronous mode of delivery ***BCIO:011060				Mode of delivery that involves delivery and receipt of the intervention or its components occurring at the same time or very close in time.	

Upper-Level	Sub-Level 1	Sub-Level 2	Sub-Level 3	Definition	Examples of usage
Asynchronous mode of delivery ***BCIO:011061				Mode of delivery that involves receipt of the intervention or its components taking place a significant period of time after delivery.	
Push mode of delivery ***BCIO:011062				Mode of delivery that is not dependent on actions on the part of the intervention recipient.	
Pull mode of delivery ***BCIO:011063				Mode of delivery that requires some action on the part of the recipient.	
Gamification mode of delivery BCIO:011064				Mode of delivery that involves application of typical elements of game playing to other areas of activity, typically as an online marketing technique to encourage engagement with a product or service.	Includes point scoring, competition with others, and rules of play.
Arts feature mode of delivery BCIO:011065				Mode of delivery that involves application of creativity on the part of the intervention recipient.	Includes art therapy, music therapy, dance and acting.

Note. Entity IDs correspond to Behaviour Change Intervention Ontology (BCIO):* Only one of individual-based, group-based or pair-based mode of delivery will apply, **only one of uni-directional or interactional mode of delivery will apply; ***only one of synchronous or asynchronous mode of delivery will apply; ****only one of push or pull mode of delivery will apply.

Secondly, the intervention reports included in the annotation process were from two larger projects, the Theory and Techniques Project (Michie *et al.*, 2018) and the Human Behaviour-Change Project (Michie *et al.*, 2017). The intervention reports annotated within the ontology development mainly addressed two health-related behaviours, smoking cessation and physical activity; there is always the possibility that other literature within and outside the health domain may indicate modes of delivery not captured in our set of papers or by our group of experts. However, external inter-rater reliability was tested across diverse behaviours and found to be acceptable. Future applications of the ontologies to a wider collection of non-health related behaviours and contexts is likely to extend and improve the ontology. The inter-rater reliability of the annotations conducted by coders unfamiliar with the ontology was lower than that found in other ontologies of the BCIO such as the Intervention Setting Ontology (Norris *et al.*, 2020), a result that can be explained by the complexity of this ontology. Nonetheless, the coding guidelines were refined throughout the process and the level of reliability increased considerably between the first and second sets of 50 papers. It is our recommendation that anyone interested in using the MoD ontology should first familiarise themselves with the MoD entities (labels, definitions and examples) and their relationships, read the coding manual, and conduct some trial annotations and assessment of reliability.

Conclusions

The MoD Ontology provides a foundation on which future research can build, and its development is intended to be an ongoing and collaborative process. By providing greater clarity about *how* an intervention and its components are delivered, researchers can add to knowledge as to how MoDs influence intervention effectiveness, both directly and in interaction with other intervention-related entities. This will inform the selection of appropriate MoDs for interventions.

Ethics

Ethical approval was granted by University College London's ethics committee (CEHP/2016/555). Participant consent was gained from the first page of the online Qualtrics survey.

Data availability

Underlying data

The BCIO is available from: <https://github.com/HumanBehaviourChangeProject/ontologies>.

Archived ontology as at time of publication: <https://doi.org/10.5281/zenodo.4476603> (Norris *et al.*, 2021).

License: [CC-BY 4.0](#).

Extended data

Open Science Framework: Human Behaviour-Change Project. <https://doi.org/10.17605/OSF.IO/UXWDB> (West *et al.*, 2020).

This project contains the following extended data related to this method:

- Copy of feedback form (PDF)
- Papers used in HBCP annotations (PDF)
- Description of the preliminary version of the MoD Ontology (PDF)
- Step 1 - Inter-Rater Reliability of the preliminary version of the Mode of Delivery Ontology (PDF)
- Feedback Report feedback and corresponding changes made to the Ontology (PDF)
- Step 3 - Inter-Rater Reliability of the preliminary version of the Mode of Delivery Ontology (PDF)
- General guidance for Mode of Delivery Ontology (PDF)

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

Code used to calculate alpha for IRR: <https://github.com/HumanBehaviourChangeProject/Automation-InterRater-Reliability>.

Archived code as at time of publication: <https://doi.org/10.5281/zenodo.3833816> (Finnerty & Moore, 2020).

License: [GNU General Public License v3.0](#)

Acknowledgements

We would like to express our gratitude to the experts who contributed to the open peer-review stages of this study and to Kirsty Atha for the support in annotating papers.

References

- Antoniu G, van Harmelen F: **Web Ontology Language: OWL**. In Staab S, Studer R (Eds.): *Handbook on Ontologies. International Handbooks on Information Systems*. Berlin: Springer. 2004; 91–110. [Publisher Full Text](#)
- Arnott B, Rehackova L, Errington L, *et al.*: **Efficacy of behavioural interventions for transport behaviour change: systematic review, meta-analysis and intervention coding**. *Int J Behav Nutr Phys Act*. 2014; 11(1): 133. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Arp R, Smith B, Spear AD: **Building Ontologies with Basic Formal Ontology**.

Massachusetts: MIT Press. 2015. [Publisher Full Text](#)

Ashburner M, Ball C, Blake J, *et al.*: **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium**. *Nat Genet*. 2000; 25(1): 25–29. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Bartholomew Eldredge LK, Parcel GS, Kok G, *et al.*: **Planning Health Promotion Programs: An Intervention Mapping Approach**. Jossey-Bass. 2011; 3. [Reference Source](#)

Black N, Eisma M, Viechtbauer W, *et al.*: **Variability and Effectiveness of**

Comparator Group Interventions in Smoking Cessation Trials: A Systematic Review and Meta-Analysis. *Addiction*. 2020; **115**(9): 1607–1617.

[PubMed Abstract](#) | [Publisher Full Text](#)

Bock C, Jarczok M, Litaker D: **Community-based Efforts to Promote Physical Activity: A Systematic Review of Interventions Considering Mode of Delivery, Study Quality and Population Subgroups.** *J Sci Med Sport*. 2014; **17**(3): 276–282.

[PubMed Abstract](#) | [Publisher Full Text](#)

Carey R, Connell L, Johnston M, *et al.*: **Behaviour change techniques and their mechanisms of action: a synthesis of links described in published intervention literature.** *Ann Behav Med*. 2019; **53**(8): 693–707.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Cohen J: **A Coefficient of Agreement for Nominal Scales.** *Educ Psychol Meas*. 1960; **20**(1): 37–46.

[Publisher Full Text](#)

Connell L, Carey R, Johnston M, *et al.*: **Links between behavior change techniques and mechanisms of action: an expert consensus study.** *Ann Behav Med*. 2019; **53**(8): 708–720.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Craig P, Dieppe P, Macintyre S, *et al.*: **Developing and evaluating complex interventions: the new Medical Research Council guidance.** *BMJ*. 2008; **337**(7676): a1655.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

De Bruin M, Viechtbauer W, Eisma MC, *et al.*: **Identifying effective behavioural components of Intervention and Comparison group support provided in SMOKing cEssation (IC-SMOKE) interventions: a systematic review protocol.** *Syst Rev*. 2016; **5**(1): 77.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Dombrowski SU, O'Carroll RE, Williams B: **Form of delivery as a key 'active ingredient' in behaviour change interventions.** *Br J Health Psychol*. 2016; **21**(4): 733–740.

[PubMed Abstract](#) | [Publisher Full Text](#)

Effective Practice and Organisation of Care (EPOC): **EPOC Taxonomy**. 2015. (accessed 10 December 2020).

[Reference Source](#)

Finnerty A, Moore C: **Human BehaviourChange Project/Automation-InterRater-Reliability: Release of HBCP inter-rater reliability code v1.0.0 (Version v1.0.0).** *Zenodo*. 2020.

<http://www.doi.org/10.5281/zenodo.3833816>

Flodgren G, Gonçalves-Bradley DC, Summerbell CD: **Interventions to change the behaviour of health professionals and the organisation of care to promote weight reduction in children and adults with overweight or obesity.** *Cochrane Database Syst Rev*. 2017; **11**(11): CD000984.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Gakidou E, Afshin A, Abajobir A, *et al.*: **Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016.** *Lancet*. 2017; **390**(10100): 1345–1422.

[Publisher Full Text](#)

van Genugten L, Dusseldorp E, Webb TL, *et al.*: **Which Combinations of Techniques and Modes of Delivery in Internet-Based Interventions Effectively Change Health Behavior? A Meta-Analysis.** *J Med Internet Res*. 2016; **18**(6): e155.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Gwet KL: **Handbook of Inter-Rater Reliability: The definitive guide to measuring the extent of agreement among raters.** Advanced Analytics, LLC. 2014.

[Reference Source](#)

Hastings J: **Primer on Ontologies.** *Methods Mol Biol*. 2017; **1446**: 3–13.

[PubMed Abstract](#) | [Publisher Full Text](#)

Hayes AF, Krippendorff K: **Answering the Call for a Standard Reliability Measure for Coding Data.** *Communication Methods and Measures*. 2007; **1**(1): 77–8.

[Publisher Full Text](#)

Hoffmann T, Glasziou P, Boutron I, *et al.*: **Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide.** *BMJ*. 2014; **348**: g1687.

[PubMed Abstract](#) | [Publisher Full Text](#)

Hollands G, Bignardi G, Johnston M, *et al.*: **The TIPME intervention typology for changing environments to change behaviour.** *Nat Hum Behav*. 2017; **1**(0140).

[Publisher Full Text](#)

Jackson R, Balhoff J, Douglass E, *et al.*: **ROBOT: A tool for automating ontology workflows.** *BMC Bioinformatics*. 2019; **20**(1): 407.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Jones CJ, Smith H, Llewellyn C: **Evaluating the effectiveness of health belief model interventions in improving adherence: a systematic review.** *Health Psychol Rev*. 2014; **8**(3): 253–269.

[PubMed Abstract](#) | [Publisher Full Text](#)

Johnston M, Carey RN, Connell Bohlen L, *et al.*: **Linking behavior change techniques and mechanisms of action: Triangulation of findings from**

literature synthesis and expert consensus. 2018.

[Publisher Full Text](#)

Kok G, Gottlieb NH, Peters GJY, *et al.*: **A taxonomy of behaviour change methods: an Intervention Mapping approach.** *Health Psychol Rev*. 2016; **10**(3): 297–312.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Knittle K, Heino M, Marques MM, *et al.*: **The compendium of self-enactable techniques to change and self-manage motivation and behaviour v.1.0.** *Nat Hum Be*. 2020; **4**(2): 215–223.

[PubMed Abstract](#) | [Publisher Full Text](#)

Landis J, Koch G: **The measurement of observer agreement for categorical data.** *Biometrics*. 1977; **33**(1): 159–174.

[PubMed Abstract](#) | [Publisher Full Text](#)

Larsen KR, Michie S, Hekler EB, *et al.*: **Behavior change interventions: the potential of ontologies for advancing science and practice.** *J Behav Med*. 2017; **40**(1): 6–22.

[PubMed Abstract](#) | [Publisher Full Text](#)

MacDonald J, Lorimer K, Knussen C, *et al.*: **Interventions to increase condom use among middle-aged and older adults: A systematic review of theoretical bases, behaviour change techniques, modes of delivery and treatment fidelity.** *J Health Psychol*. 2016; **21**(11): 2477–2492.

[PubMed Abstract](#) | [Publisher Full Text](#)

Michie S, Thomas J, Mac Aonghusa P, *et al.*: **The Human Behaviour-Change Project: An artificial intelligence system to answer questions about changing behaviour [version 1; peer review: not peer reviewed].** *Wellcome Open Res*. 2020a.

[Publisher Full Text](#)

Michie S, West R, Finnerty AN, *et al.*: **Representation of behaviour change interventions and their evaluation: Development of the Upper Level of the Behaviour Change Intervention Ontology [version 1; peer review: 1 approved, 1 approved with reservations].** *Wellcome Open Res*. 2020b.

[Publisher Full Text](#)

Michie S, Carey R, Johnston M, *et al.*: **From theory-inspired to theory-based interventions: A protocol for developing and testing a methodology for linking behaviour change techniques to theoretical mechanisms of action.** *Ann Behav Med*. 2018; **52**(6): 501–512.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Michie S, Johnston M: **Optimising the value of the evidence generated in implementation science: the use of ontologies to address the challenges.** *Implement Sci*. 2017; **12**(1): 131.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Michie S, Johnston M, Carey R: **Behavior Change Techniques.** In G. M. (Ed.), *Encyclopedia of Behavioral Medicine*. New York, NY: Springer. 2019a.

[Publisher Full Text](#)

Michie S, West R, Hastings J: **Creating ontological definitions for use in science.** *Qeios*. 2019b.

[Publisher Full Text](#)

Michie S, Richardson M, Johnston M, *et al.*: **The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions.** *Ann Behav Med*. 2013; **46**(1): 81–95.

[PubMed Abstract](#) | [Publisher Full Text](#)

Michie S, Thomas J, Johnston M, *et al.*: **The Human Behaviour-Change Project: Harnessing the power of Artificial Intelligence and Machine Learning for evidence synthesis and interpretation.** *Implement Sci*. 2017; **12**(1): 121.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Michie S, van Stralen MM, West R: **The behaviour change wheel: A new method for characterising and designing behaviour change interventions.** *Implement Sci*. 2011; **6**: 42.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Michie S, Wood C, Johnston M, *et al.*: **Behaviour Change Techniques: The Development and Evaluation of a Taxonomic Method for Reporting and Describing Behaviour Change Interventions (A Suite of Five Studies Involving Consensus Methods, Randomised Controlled Trials and Analysis of Qualitative Data).** *Health Technol Assess*. 2015; **19**(99): 1–188.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Moher D, Schulz KF, Altman DG: **The CONSORT statement: revised recommendations for improving the quality of reports of parallel group randomized trials.** *BMC Med Res Methodol*. 2001; **1**(1): 2.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Newbury-Birch D, Coulton S, Bland M, *et al.*: **Alcohol screening and brief interventions for offenders in the probation setting (SIPS trial): a pragmatic multicentre cluster randomized controlled trial.** *Alcohol Alcohol*. 2014; **49**(5): 540–548.

[PubMed Abstract](#) | [Publisher Full Text](#)

Norris E, Finnerty AN, Hastings J, *et al.*: **A scoping review of ontologies related to human behaviour change.** *Nat Hum Behav*. 2019; **3**(2): 164–172.

[PubMed Abstract](#) | [Publisher Full Text](#)

Norris E, Hastings J, Ailbhe F: **HumanBehaviourChangeProject/ontologies: Upper-Level, Setting & MoD papers Submitted.** *Zenodo*. 2021.

<http://www.doi.org/10.5281/zenodo.4476603>

Norris E, Marques MM, Finnerty AN, *et al.*: **Development of an Intervention Setting Ontology for behaviour change: Specifying where interventions**

take place [version 1; peer review: 2 approved]. *Wellcome Open Res.* 2020.

[Publisher Full Text](#)

Paul L, Brewster S, Wyke S, *et al.*: **Increasing physical activity in older adults using STARFISH, an interactive smartphone application (app); a pilot study.** *J Rehabil Assist Technol Eng.* 2017; **4**: 2055668317696236.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sheeran P, Klein WMP, Rothman AJ: **Health behavior change: moving from observation to intervention.** *Annu Rev Psychol.* 2017; **68**: 573–600.

[PubMed Abstract](#) | [Publisher Full Text](#)

Shindo H, Munesada Y, Matsumoto Y: **PDFAnno: a web-based linguistic annotation tool for pdf documents.** In: *Proceedings of the Eleventh International Conference on Language Resources and Evaluation (LREC 2018)*. 2018. [Reference Source](#)

Smith B, Ceusters W, Klagges B, *et al.*: **Relations in biomedical ontologies.** *Genome Biol.* 2005; **6**(5): R46.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Smith B, Ashburner M, Rosse C, *et al.*: **The OBO Foundry: coordinated evolution of ontologies to support biomedical data integration.** *Nat Biotechnol.* 2007; **25**(11): 1251–1255.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Stan J, Demner-Fushman D, Fung KW, *et al.*: **Facilitating reconciliation of inter-annotator disagreements.** In: *AMIA Annual Symposium Proceedings*. 2014; 1596.

[Reference Source](#)

Thomas J, Brunton J, Graziosi S: **EPPI-Reviewer 4: software for research synthesis.** EPPI-Centre Software. London: Social Science Research Unit, UCL Institute of Education. 2010.

[Reference Source](#)

Ussher MH, Taylor A, Faulkner G: **Exercise interventions for smoking cessation.** *Cochrane Database Syst Rev.* 2012; **1**(8): CD002295.

[PubMed Abstract](#) | [Publisher Full Text](#)

Watts N, Adger WN, Ayeb-Karlsson S, *et al.*: **The Lancet Countdown: tracking progress on health and climate change.** *Lancet.* 2017; **389**(10074): 1151–1164.

[PubMed Abstract](#) | [Publisher Full Text](#)

Webb TL, Joseph J, Yardley L, *et al.*: **Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy.** *J Med Internet Res.* 2010; **12**(1): e4.

[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

West R, Michie S, Shawe-Taylor J, *et al.*: **Human Behaviour-Change Project.** 2020.

<http://www.doi.org/10.17605/OSF.IO/UXWDB>

Wright AJ, Norris E, Finnerty, *et al.*: **Ontologies relevant to behaviour change interventions: a method for their development [version 1; peer review: 1 not approved].** *Wellcome Open Res.* 2020.

[Publisher Full Text](#)

Young MD, Collins CE, Callister R, *et al.*: **The SHED-IT Weight Loss Maintenance trial protocol: A randomised controlled trial of a weight loss maintenance program for overweight and obese men.** *Contemp Clin Trials.* 2014; **37**(1): 84–97.

[PubMed Abstract](#) | [Publisher Full Text](#)

Zebis MK, Andersen LL, Brandt B, *et al.*: **Effects of evidence-based prevention training on neuromuscular and biomechanical risk factors for ACL injury in adolescent female athletes: a randomised controlled trial.** *Br J Sports Med.* 2016; **50**(9): 552–7.

[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 23 March 2021

<https://doi.org/10.21956/wellcomeopenres.18343.r42863>

© 2021 Byrne-Davis L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Lucie Byrne-Davis 

Division of Medical Education, University of Manchester, Manchester, UK

I have no further comments to make. The authors have responded fully to each of my initial comments. This is a very interesting paper that provides insight into the development of this much needed ontology.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Behaviour change in health settings, particularly health worker practice change; health worker education

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 08 March 2021

<https://doi.org/10.21956/wellcomeopenres.18343.r42862>

© 2021 DeSmet A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ann DeSmet 

Université Libre de Bruxelles, Brussels, Belgium

All previous comments were sufficiently addressed.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: health psychology, behaviour change, digital health interventions, serious games

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 14 September 2020

<https://doi.org/10.21956/wellcomeopenres.17447.r39974>

© 2020 DeSmet A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ann DeSmet 

Université Libre de Bruxelles, Brussels, Belgium

The authors describe their approach and results in building an ontology of mode of delivery of interventions in this paper. The paper is well-written, clearly structured and methodologically sound. I have listed a few suggestions and minor comments below:

- I personally appreciate the initiative the authors have taken here. Having created and evaluated serious game interventions, I have noticed how certain techniques may be recommended or effective in one type of intervention, but may not work so well when delivered in a game format. I agree that a detailed description of delivery modes and their structure (what belongs to which group) is necessary, but I was wondering if the authors could also provide more detailed suggestions for future use in the discussion/conclusion part.
For example, the BCTv1 is mentioned in the introduction, but how do the authors plan to make the connection between this taxonomy and the ontology? Could they shed more light on future initiatives to clarify the importance of this work to the reader?
The authors also refer to several taxonomies of techniques that exist (it may be useful to also refer to the Intervention Mapping protocol list that is relatively well-known - Kok, G., Gottlieb, N. H., Peters, G. J. Y., Mullen, P. D., Parcel, G. S., Ruiters, R. A., ... & Bartholomew, L. K. (2016). A taxonomy of behaviour change methods: an intervention mapping approach. *Health psychology review*, 10(3), 297-312¹). Could they clarify where this ontology is inherently linked to the taxonomy or could in the future also be used with other BCT taxonomies, as in a type of open platform communication?
- In the method part it was sometimes difficult to see the link between the text and Table 2. The text, for example, mentions 20 pilot reports in step 1, and then another 'set of interventions'. Table 2 then shows 120 BCI reports were extracted. Why 120? How did the authors decide this was an appropriate number? Same for step 3 (55 reports). Could more information be provided on the database? Are these reports that maybe

already follow a certain protocol of annotation, could this create some bias?

The authors mention 'mailing lists' as a way to recruit the experts. Could they provide more information on the mailing lists, or characteristics of experts?

- Could the authors elaborate more on the potential reasons for discrepancies in interrater reliability 'whether a particular entity was considered an MoD was 61%; and agreement on the specific MoD code assigned was 87.9%' in round 2?
- Step 5: could it be that the lower agreement between raters was not related to the fact that they were less familiar with the ontology, but by the fact that there were was a wider variety in target behaviors in this selection of reports? Taxonomies are also mostly applied to diet, physical activity, addictive behaviours; could it be that the ontology does not fit as well with screening, infectious diseases etc?
- Table 2 mentions inter-rater reliability twice for step 3: typo?
- Table 3: definition of video game delivery seems to copy-pasted from the level above?
- Table 3: Somatic alteration mode of delivery - also typo (copy-paste above)?

References

1. Kok G, Gottlieb NH, Peters GJ, Mullen PD, et al.: A taxonomy of behaviour change methods: an Intervention Mapping approach. *Health Psychol Rev.* 2016; **10** (3): 297-312 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: health psychology, behaviour change, digital health interventions, serious

games

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 15 Feb 2021

Marta Marques, University College London, London, UK

The authors describe their approach and results in building an ontology of mode of delivery of interventions in this paper. The paper is well-written, clearly structured and methodologically sound. I have listed a few suggestions and minor comments below:

We appreciate the reviewer's positive feedback. We have addressed all comments.

Personally appreciate the initiative the authors have taken here. Having created and evaluated serious game interventions, I have noticed how certain techniques may be recommended or effective in one type of intervention, but may not work so well when delivered in a game format. I agree that a detailed description of delivery modes and their structure (what belongs to which group) is necessary, but I was wondering if the authors could also provide more detailed suggestions for future use in the discussion/conclusion part.

For example, the BCTv1 is mentioned in the introduction, but how do the authors plan to make the connection between this taxonomy and the ontology? Could they shed more light on future initiatives to clarify the importance of this work to the reader?

Thank you for this important remark. We have added a new paragraph to the discussion as follows: "The MoD ontology provides a crucial contribution to the much needed body of research examining the links between MoDs and the content of behaviour change interventions, using the BCTv1 or other classification systems of techniques (e.g. Knittle *et al.*, 2020; Kok *et al.*, 2016). For example, coding existing behaviour change interventions for their modes of delivery and BCTs can increase our understanding of which mode(s) of delivery are the most effective in delivering a given BCT. Further, by linking with other HBCP ontologies characterising different aspects of behaviour change interventions, it will be possible to go a step further and identify which MoD(s) are more appropriate for different behaviours, populations, contexts, if they need to be tailored, and their potential for reach and engagement."

Taxonomies of techniques that exist (it may be useful to also refer to the Intervention Mapping protocol list that is relatively well-known - Kok, G., Gottlieb, N. H., Peters, G. J. Y., Mullen, P. D., Parcel, G. S., Ruitter, R. A., ... & Bartholomew, L. K. (2016). A taxonomy of behaviour change methods: an intervention mapping approach. *Health psychology review*, 10(3), 297-312¹. Could they clarify where this ontology is inherently linked to the taxonomy or could in the future also be used with other BCT taxonomies, as in a type of open platform communication?

Thank you for this suggestion. We have added information related to this to the paragraph

presented in the previous comment.

In the method part it was sometimes difficult to see the link between the text and Table 2. The text, for example, mentions 20 pilot reports in step 1, and then another 'set of interventions'. Table 2 then shows 120 BCI reports were extracted. Why 120? How did the authors decide this was an appropriate number? Same for step 3 (55 reports).

Thank you for noticing this. In step 1 there was an initial extraction of 20 reports for the first skeleton of the ontology and then 100 papers more were annotated to improve the coverage and specificity of the ontology and test its reliability. We have amended the table as follows: "Data extraction from 120 BCI reports: 20 reports for initial draft + 100 for improvements and inter-rater reliability calculations". The number of papers was not pre defined, the coders kept reviewing until an adequate Kappa was reached. The same was true for the number of papers in step 3.

Could more information be provided on the database? Are these reports that maybe already follow a certain protocol of annotation, could this create some bias?

Thank you for pointing this out. The 55 reports came from a collection of articles assembled for a previous project in our research group (Michie et al., 2018). These are articles in which authors described links between behaviour change techniques and intervention mechanisms of action. Mode of delivery might be described in more detail in these papers where other aspects of interventions are also specified in detail. This greater level of nuance is likely to be a greater challenge to create ontology categories to fit, and so make achieving good inter-rater reliability more difficult.

The authors mention 'mailing lists' as a way to recruit the experts. Could they provide more information on the mailing lists, or characteristics of experts?

We thank the reviewer for this important point. Invitations to potential participants were sent out via third-party mailing lists (e.g. conference). We have some data on the characteristics of the experts who participated, such as the type of organisations reviewers were from and countries. We also have a list of the specific institutions they were from. We have added this information to the results section, step 2 as follows: "Feedback on the MoD ontology through the open review feedback form was received by 25 experts, of which 18 were from universities, 5 were from commercial sector organisations, 1 from public sector organisations and 1 from third sector. Twelve experts were from the United Kingdom, 2 from the United States of America, 3 from Ireland, 1 from Canada, 1 from the Netherlands, 1 from New Zealand, and we have no information about the country for the remaining 5 experts."

Could the authors elaborate more on the potential reasons for discrepancies in interrater reliability 'whether a particular entity was considered an MoD was 61%; and agreement on the specific MoD code assigned was 87.9%' in round 2?

The first element corresponds to recognizing that part of the text contains a description of a

mode of delivery. One of the reasons for this lower agreement can be due to the fact that many papers describe mode of delivery poorly and it is stated in the coding manual that MoD should be coded when it is clearly stated in the paper (similarly to BCTTv1). When both coders identified a segment of the text as stating a MoD there was higher agreement about which specific MoD was stated, which demonstrates the utility of the MoD in distinguishing between different MoDs and clearly defining them.

Step 5: could it be that the lower agreement between raters was not related to the fact that they were less familiar with the ontology, but by the fact that there were a wider variety in target behaviors in this selection of reports? Taxonomies are also mostly applied to diet, physical activity, addictive behaviours; could it be that the ontology does not fit as well with screening, infectious diseases etc?

This is an interesting point. The MoD ontology was designed to be applicable across behaviours, and MoD reporting or lack of it seems to be consistent across behaviours. We hope that future research using this ontology will provides the necessary data to explore this issue further, i.e. if lower agreement are related with familiarity and/or stability across behaviours.

Table 2 mentions inter-rater reliability twice for step 3: typo?

Yes, it was a typo. Thank you for pointing it out.

Table 3: definition of video game delivery seems to copy-pasted from the level above?

Thank you for noticing this. We have now changed the definition to "Electronic mode of delivery that involves the intervention recipient playing a computer game."

Table 3: Somatic alteration mode of delivery - also typo (copy-paste above)?

Again, thank you for spotting this typo. We have changed to "Mode of delivery that involves modifying the structure of the body of the recipient of the intervention"

Competing Interests: No competing interests were disclosed.

Reviewer Report 15 July 2020

<https://doi.org/10.21956/wellcomeopenres.17447.r39010>

© 2020 Byrne-Davis L. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Lucie Byrne-Davis

Division of Medical Education, University of Manchester, Manchester, UK

This paper presents the development of an ontology of 'modes of delivery' of behaviour change interventions. It is one of the studies from the Human Behaviour Change Project and is a welcome

addition to the literature. Overall, the paper is very well written and the studies sound. My only issue is about the extent to which the introduction includes references to other taxonomies/ontologies beyond the three that it does mention, and therefore how the paper is situated in the literature both in the introduction and discussion sections.

Abstract

I didn't understand the sentence "Relationships between entities consist of *is_a*."

Should the conclusion in the abstract recommend that people should be familiar with the ontology to ensure that it was used reliably, given that the reliability was only 0.58 when they were unfamiliar?

Introduction

You introduce three classification systems but then move straight into the BCTTv1. It is not clear why you focus on that one and so this paragraph seems to come from nowhere. Could you make the reason you are moving from the three systems to the BCTTv1 more obvious? Also, you start a new paragraph after introducing the three systems but that is a very short paragraph, so I would suggest this needs to be one paragraph together. I also expected in the introduction to see more reference to previous taxonomies and problematising these to establish why this ontology was so important. You don't, for example, mention the EPOC taxonomy and I was not sure why.

Methods and results

Step 1. This step specifies health behaviours. Previously, you have not specified that this relates to health behaviours specifically, in fact you introduce this as including environmental and social problems, and some of the earlier work is related to health worker behaviours. It would be good to have some clarity about whether this is all human behaviour (which I think it is) and to what extent the methods relied on interventions related to health behaviours and whether this is a limitation of the methods. I know you do state this as a limitation but it would be good to see this up front. in the methods and a rationale for why the study was conducted in this way.

Step 2. Can you report the response rate (either in methods or results) and where the raters were from. I'm particularly interested in whether all were from a particular part of the world, what institutions were included. Much of the work rests on these individuals being experts so I think it would be appropriate to include some further information in the text that summarises their credentials and any potential biases they might introduce into the initial ontology.

Discussion

As per the introduction, it would be useful to see how this ontology fits with previous attempts at classifying modes of delivery. If there are none (if the EPOC taxonomy is not an example of this) then it would be good to state that as part of the reason for developing this anew.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Behaviour change in health settings, particularly health worker practice change; health worker education

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 15 Feb 2021

Marta Marques, University College London, London, UK

This paper presents the development of an ontology of 'modes of delivery' of behaviour change interventions. It is one of the studies from the Human Behaviour Change Project and is a welcome addition to the literature. Overall, the paper is very well written and the studies sound. My only issue is about the extent to which the introduction includes references to other taxonomies/ontologies beyond the three that it does mention, and therefore how the paper is situated in the literature both in the introduction and discussion sections.

We appreciate the reviewer' positive feedback and addressed each suggestion and comment.

Abstract: I didn't understand the sentence "Relationships between entities consist of *is_a*."

We have changed the sentence to "Relationships between entities are hierarchical e.g. "Face-to-face mode of delivery *is_a* human interactional mode of delivery"

Abstract: Should the conclusion in the abstract recommend that people should be familiar with the ontology to ensure that it was used reliably, given that the reliability was only 0.58 when they were unfamiliar?

We understand the reviewer concern. The recommendation for this and the other HBCP ontologies is that anyone interested in using it, especially for formal coding exercises

should first familiarise themselves with it. We have added a sentence to the discussion section of the manuscript as follows: "It is our recommendation that anyone interested in using the MoD ontology should first familiarise themselves with the MoD entities (labels, definitions and examples) and their relationships, read the coding manual, and conduct some trial annotation and assessment of reliability."

Introduction

You introduce three classification systems but then move straight into the BCTTv1. It is not clear why you focus on that one and so this paragraph seems to come from nowhere. Could you make the reason you are moving from the three systems to the BCTTv1 more obvious? Also, you start a new paragraph after introducing the three systems but that is a very short paragraph, so I would suggest this needs to be one paragraph together. I also expected in the introduction to see more reference to previous taxonomies and problematising these to establish why this ontology was so important. You don't, for example, mention the EPOC taxonomy and I was not sure why.

Thank you for your comment. We have revised this section to reflect the BCTTv1 as an example of a taxonomy focusing on the content of interventions. In addition, we added information about the EPOC taxonomy in the "Delivery of Behaviour Change Interventions" section.

Methods and results

Step 1. This step specifies health behaviours. Previously, you have not specified that this relates to health behaviours specifically, in fact you introduce this as including environmental and social problems, and some of the earlier work is related to health worker behaviours. It would be good to have some clarity about whether this is all human behaviour (which I think it is) and to what extent the methods relied on interventions related to health behaviours and whether this is a limitation of the methods. I know you do state this as a limitation but it would be good to see this up front. in the methods and a rationale for why the study was conducted in this way.

Thank you for pointing this out. This is indeed intended as an ontology of modes of delivery for all domains of behaviour change interventions. The limitations section of the discussion addresses the limitations of having annotated mainly health-related behaviour papers within the ontology development stages, and we have now made this point clearer, as follows: "Secondly, the intervention reports included in the annotation process were from two larger projects, the Theory and Techniques Project ([Michie et al., 2018](#)) and the Human Behaviour-Change Project ([Michie et al., 2017](#)). The intervention reports annotated within the ontology development mainly addressed two health-related behaviours, smoking cessation and physical activity; there is always the possibility that other literature within and outside the health domain may indicate modes of delivery not captured in our set of papers or by our group of experts. However, external inter-rater reliability was tested across diverse behaviours and found to be acceptable. Future applications of the ontologies to a wider collection of non-health related behaviours and contexts is likely to extend and improve the ontology."

Step 2. Can you report the response rate (either in methods or results) and where the raters were from. I'm particularly interested in whether all were from a particular part of the world, what institutions were included. Much of the work rests on these individuals being experts so I think it would be appropriate to include some further information in the text that summarises their credentials and any potential biases they might introduce into the initial ontology.

We thank the reviewer for this important point. We have data on the type of organisations reviewers were from: 18 were from universities, 5 from commercial sector organisations, 1 from public sector organisations and 1 third sector; and the countries: 12 experts were from the United Kingdom, 2 from the United States of America, 3 from Ireland, 1 from Canada, 1 from the Netherlands, 1 from New Zealand, and for 5 of them we have no information about the country. We also have a list of the specific institutions they were from. We don't have response rate data as the invitations to participate were sent out via third-party mailing lists (e.g. conference) and so we do not know how many people were subscribed to each list. We have added the following information to the results, step 2: "Feedback on the MoD ontology through the open review feedback form was received by 25 experts, of which 18 were from universities, 5 were from commercial sector organisations, 1 from public sector organisations and 1 from third sector. Twelve experts were from the United Kingdom, 2 from the United States of America, 3 from Ireland, 1 from Canada, 1 from the Netherlands, 1 from New Zealand, and we have no information about the country for the remaining 5 experts."

Discussion

As per the introduction, it would be useful to see how this ontology fits with previous attempts at classifying modes of delivery. If there are none (if the EPOC taxonomy is not an example of this) then it would be good to state that as part of the reason for developing this.

Thank you for this comment. We have addressed this comment in the introduction in "Delivery of Behaviour change interventions". Further we added a sentence in the discussion to reflect this as follows "Given the lack of classification systems providing comprehensive coverage of how behaviour change interventions and techniques are delivered, we developed the first ontology of modes of delivery (MoD)."

Competing Interests: No competing interests were disclosed.