Hindawi Evidence-Based Complementary and Alternative Medicine Volume 2017, Article ID 9439682, 7 pages https://doi.org/10.1155/2017/9439682

Research Article

Association between Nine Types of TCM Constitution and Five Chronic Diseases: A Correspondence Analysis Based on a Sample of 2,660 Participants

Yanbo Zhu,¹ Huimei Shi,² Qi Wang,² Yangyang Wang,¹ Xiaohan Yu,² Jie Di,¹ Xiaomei Zhang,¹ Yanni Li,¹ Tong Li,¹ and Hui Yan¹

¹School of Management, Beijing University of Chinese Medicine, Beijing 100029, China

Correspondence should be addressed to Yanbo Zhu; yanbo0722@sina.com and Qi Wang; wangqi710@126.com

Received 17 January 2017; Revised 11 March 2017; Accepted 8 May 2017; Published 1 June 2017

Academic Editor: Yoshiki Mukudai

Copyright © 2017 Yanbo Zhu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. The purpose of this study was to explore the association of nine types of Traditional Chinese Medicine (TCM) constitution with the five chronic diseases: hypertension, hyperlipidemia, diabetes mellitus, heart disease, and obesity. *Methods*. Chi-squared test was performed to investigate the distribution characteristics of TCM constitutions in the participants with the five chronic diseases in questionnaire. Correspondence analysis was used to explore the correlation between them. *Results*. A total of 2,660 participants (1,400 males; 1,260 females) were included in this study. The mean age was 52.54 ± 13.92 . Of them, 600 were of gentleness type accounting for 22.56%. Proportions of gentleness type in the chronic diseases ($16.00\% \sim 23.70\%$) were less than that in general population (32.14%). The gentleness type and yin-deficiency type were significantly correlated with hypertension and diabetes mellitus, qi-deficiency type was correlated with heart disease, phlegm-dampness type was associated with obesity, and dampnessheat type was correlated with hyperlipidemia. *Conclusions*. The correlations between TCM constitution types and the five chronic diseases were different. This may have a significant implication for TCM practice, and even the people with gentleness type should not be ignored in health management.

1. Introduction

Chronic diseases such as cardiovascular disease-heart disease, hypertension, and stroke-cancer, diabetes, and chronic respiratory disease are the main causes of death [1]. The rapid growth of chronic diseases brings heavy burden on healthcare and society. The major risk factors of these diseases come from tobacco smoking, alcohol consumption, unhealthy diet, and lack of physical activity [1–3].

In terms of the theory of Traditional Chinese Medicine (TCM), TCM constitution, depending on the intrinsic characteristics of human body, is innate and influenced by the environment. It integrates the morphological structure and physiological function with psychological state. Although relatively stable, TCM constitution may also develop and change so as to adapt to the changes in the environment throughout

a human body's life [4]. The TCM constitutions are classified into nine categories: gentleness type, qi-deficiency type, yang-deficiency type, yin-deficiency type, phlegm-dampness type, dampness-heat type, blood-stasis type, qi-depression type, and special diathesis type. The classification is defined in terms of physical features, common manifestations, psychological characteristics, susceptibility to certain diseases, and adaptability to the environment [5].

Previous studies have shown that unhealthy status [6, 7] or diseases like diabetes mellitus and cardiovascular diseases were correlated with specific constitution types [8–12]. It has been found that more likely to suffer from angina pectoris are the constitutions of dampness-heat type, phlegm-dampness type, and blood-stasis type commonly found in Lingnan area in the southern part of China [12]. It also has been demonstrated that some health conditions and disease

²School of Preclinical Medicine, Beijing University of Chinese Medicine, Beijing 100029, China

status can be improved by the interventions according to TCM constitution [13–16]. These findings suggest that the constitution types found in people with chronic diseases may provide valuable information for disease prevention and treatment [17]. However, the correlation between TCM constitution types and chronic diseases was still unclear.

Number of cases and its proportion as well as binary logistic regression were often used to evaluate the correlation between TCM constitution types and some diseases [8, 10, 11]. However, when analyzing categorical variables like the nine types of TCM constitution and the five chronic diseases, these methods are difficult to reveal the correlation between them. In this study, a multivariate statistical method-correspondence analysis (CA) was applied to investigate the correlation of TCM constitution types in participants with the five chronic diseases.

2. Materials and Methods

2.1. Subjects. The data of this study was obtained from a cross-sectional survey of 21,218 subjects from December 2005 to January 2007 [18]. By using purposive sampling method, nine provinces and municipalities in China (Jiangsu, Anhui, Gansu, Qinghai, Fujian, Beijing, Jilin, Jiangxi, and Henan) were selected. In each province and municipality, the subjects were collected from communities, health examination centers, or colleges. Systemic sampling method was used to select subjects. All the participants signed an informed consent form.

In our study, the eligible subjects were the participants with self-reported chronic diseases: hypertension, hyperlipidemia, diabetes mellitus, heart disease, and obesity. A number of 2,660 participants (1,400 males and 1,260 females) were included in this study and 18,558 individuals who have none of the five chronic diseases were excluded.

2.2. Measurement of TCM Constitution. TCM constitutions were measured and classified with Constitution in Chinese Medicine Questionnaire (CCMQ) [19–21]. CCMQ is a self-rating scale with good reliability and validity. It has 60 items, which contains nine subscales: gentleness, qi-deficiency, yang-deficiency, yin-deficiency, phlegm-dampness, dampness-heat, blood-stasis, qi-depression, and special diathesis. Score of each subscale is standardized from 0 to 100. Gentleness type is a balanced one with higher score indicating a better constitution status, while the other eight types are pathological types with higher score indicating worse constitution status.

Discrimination analysis was applied to determine the constitution type of each subject based on the data from 542 subjects with typical constitutions diagnosed by TCM constitution experts [22].

2.3. Correspondence Analysis. It is a statistical method to analyze two-way and multiway category data that are transformed into cross tables [23–26]. In CA, the relationship between the categories of the same variable and the correspondence among different variables can be revealed. The results of CA are demonstrated graphically in a biplot with

rows and columns as points, and the interpretation is based upon proximities between points. The procedure of biplot analysis is as follows [27]. First, the proximity of each category of the same variable on horizontal and vertical axes of the biplot should be compared. If they are in close proximity on horizontal or vertical axis, it means there are little differences among them on that axis. Second, the proximity among categories of different variables should be compared. If the categories are in close proximity, they are associated with each other. The closer the proximity, the stronger the association.

2.4. Statistical Methods. Chi-squared test and CA were performed to identify the relationship between nine TCM constitution types and the five chronic diseases. Data analysis was conducted with SPSS 17.0 (SPSS Inc., released 2008, SPSS for Windows, Version 17.0, Chicago, IL, USA) and the critical value of statistical significance was set as 0.05.

3. Results

3.1. Clinical Characteristics of the Participants. Totally 2,660 participants (52.63% males and 47.37% females) were included in this study. The mean age was 52.54 ± 13.92 years, and 1,188 (44.66%) participants were 55 years and above. Of the 2660 participants, 600 were in gentleness type, accounting for 22.56%; the numbers of participants with hypertension, hyperlipidemia, diabetes mellitus, heart disease, and obesity were 1,466 (55.11%), 911 (34.25%), 412 (15.49%), 650 (24.44%), and 351 (13.20%), respectively (Table 1).

3.2. Distribution of TCM Constitution Types among the Participants with the Five Chronic Diseases. There were differences among the distributions of TCM constitution types in different chronic diseases ($\chi^2=138.31,\,P<0.001$). The proportions of gentleness type in the chronic diseases varied from 16.00% to 23.70%, which were less than that in general population (32.14%) [22]; the proportions of special diathesis type for each chronic disease were less than the others; and the proportions of qi-deficiency type were higher than the others in chronic diseases except obesity (Table 2).

The top three pathological constitution types in the five chronic diseases were qi-deficiency type (19.00%), phlegm-dampness type (10.90%), and yin-deficiency type (9.10%) in hypertension participants; qi-deficiency type (17.30%), phlegm-dampness type (14.70%), and dampnessheat type (10.20%) in hyperlipidemia participants; qi-deficiency type (19.20%), dampness-heat type (11.90%), and phlegm-dampness type (11.40%) in diabetes mellitus participants; qi-deficiency type (30.80%), yang-deficiency type (11.10%), and blood-stasis type (9.40%) in heart disease participants; phlegm-dampness type (19.90%), qi-deficiency type (15.70%), and dampness-heat type (11.10%) in obesity participants (Table 2).

3.3. Correspondence Analysis. The results of CA are presented in Tables 3 and 4 and Figure 1. From Table 3, the singular value indicates the correlation between the row and column profiles. The eigenvalues of dimension 1 and dimension 2

TABLE 1: Clinical characteristics of 2,660 patients.

Characteristic	Classification	n	Percentage (%)
Gender	Male	1,400	52.63
Gender	Female	1,260	47.37
	15-24	85	3.20
	25-34	190	7.14
Age (years)	35-44	495	18.61
rige (years)	45-54	702	26.39
	55-64	573	21.54
	≥65	615	23.12
	Primary school	324	12.18
Education	Middle school	1,267	47.63
	College degree and above	1,069	40.19
	Gentleness type	600	22.56
	Qi-deficiency type	508	19.10
	Yang-deficiency type	230	8.65
Constitution	Yin-deficiency type	225	8.46
types	Phlegm-dampness type	295	11.09
	Dampness-heat type	248	9.32
	Blood-stasis type	216	8.12
	Qi-depression type	194	7.29
	Special diathesis type	144	5.41
	Hypertension	1,466	55.11
Chronic	Hyperlipidemia	911	34.25
diseases	Diabetes mellitus	412	15.49
	Heart disease	650	24.44
	Obesity	351	13.20

were 0.778 and 0.139, respectively, and 91.70% of the variance was represented well in the first two dimensions (Table 3).

The factor loading matrix of constitution types and chronic diseases on dimension 1 and dimension 2 was showed in Table 4, and it was also graphically showed in a biplot (Figure 1). Since dimension 1 (77.80%) revealed much more information than dimension 2 (13.90%), we can only see the results on dimension 1 (horizontal axis of Figure 1). On dimension 1 of Figure 1, the nine TCM constitution types fell into two categories: (1) 1 (gentleness type), 5 (phlegm-dampness type), 6 (dampness-heat type), 8 (qi-depression type), and 9 (special diathesis type); (2) 2 (qi-deficiency type), 3 (yang-deficiency type), 4 (yin-deficiency type), and 7 (blood-stasis type). The five chronic diseases were also divided into two categories: (1) A (hypertension), B (hyperlipidemia), C (diabetes mellitus), and E (obesity); (2) D (heart disease). According to the spatial distribution of variables, gentleness type and yin-deficiency type were correlated with hypertension and diabetes mellitus; qi-deficiency type was associated with heart disease; phlegmdampness type was correlated with obesity; and dampnessheat type was correlated with hyperlipidemia.

4. Discussion

Different chronic diseases have different distribution of TCM constitutions according to the constituent ratios. But it is

difficult to clearly explain the distribution of TCM constitution types in chronic diseases. CA is a multivariate statistical analysis method used to explore the relationship between several variables. It is based on the analysis of the contingency table through the row and column profiles to present a unique graphical display showing the relationship among variables [28]. Therefore, on the basis of constituent ratios, CA was used to reveal the relationship between chronic diseases and TCM constitutions.

Results showed that hypertension and diabetes mellitus were related to yin-deficiency type. Hypertension is recognized as dizziness, headache, liver yang, and liver wind in TCM. The main characteristics of yin-deficiency type are the consumption of body fluids, essence and blood, and the deficiency of internal heat [5]. It is recognized in *Internal Canon of Huangdi* that deficiency of kidney essence and hyperactivity of liver yang may lead to dizziness. Meanwhile, some study [29] has found that, in hypertension participants with yin-deficiency and yang hyperactivity, the renin, angiotensin II, and serum lipid peroxidation significantly increased, and plasma nitric oxide and atrial natriuretic factor significantly decreased, which may relate to the development of hypertension.

The diabetes mellitus is recognized as consumptive thirst in TCM. It is documented in *Jingyue Quanshu* • *Zazhengmo* • *Xuezheng* that all blood syndromes result from insufficiency of fluids leading to fire flaring up regardless of zang-organs. It is more likely to suffer from consumptive thirst with excessive fire flaring up due to insufficiency of yin for a long time.

Interestingly, we found that gentleness type was also related to hypertension. The characteristics of gentleness type are moderate posture, shiny and glossy complexion, and vigorous and good function of zang-fu organs. In this research, the mean age of the participants was 52.54, and about half of them were 55 years old and above. One possible explanation was that hypertensive participants with gentleness type may have a better survival rate, which may justify the association of gentleness type with hypertension. Another possible explanation was that people with gentleness type believe that they are in good physical condition and pay no attention to their unhealthy behavior and habits like drinking and smoking [30], which increases hypertension prevalence in the gentleness type participants.

In addition, we found that obesity was related to phlegm-dampness type, and hyperlipidemia was related to dampness-heat type. Phlegm-dampness type resulted from accumulation of phlegm due to internal retention of dampness. Previous study has indicated that phlegm-dampness type and qi-deficiency type are the main constitution risk factors of overweight or obesity [31]. Dampness-heat type is a constitution type of internal retention of dampness-heat, which is an important pathogenesis of hyperlipidemia according to both TCM traditional literatures and modern studies [32]. Studies have also found that dampness-heat type is associated with metabolic syndrome [33, 34].

Our findings also show that qi-deficiency was related to heart disease. In TCM, heart disease is recognized as heart pain, which is due to qi blockage in the chest. Throughout the process of qi blockage in the chest, heart qi is deficient [35].

Table 2: Distribution of nine types of TCM constitution in five chronic diseases.

Chronic diseases	и	Gentleness	deficiency	deficiency	deficiency	dampness	heat	Blood-stasis	depression	diathesis
		2d/v	type	type	type	type	type	277	type	type
A	1466	348 (23.70)	279 (19.00)	117 (8.00)	134 (9.10)	160 (10.90)	127 (8.70)	121 (8.30)	94 (6.40)	86 (5.90)
В	911	206 (22.60)	158 (17.30)	65 (7.10)	61(6.70)	134 (14.70)	93 (10.20)	73 (8.00)	70 (7.70)	51(5.60)
C	412	85 (20.60)	79 (19.20)	43 (10.40)	34 (8.30)	47 (11.40)	49 (11.90)	30 (7.30)	24 (5.80)	21 (5.10)
D	650	104 (16.00)	200 (30.80)	72 (11.10)	52 (8.00)	43 (6.60)	40 (6.20)	61(9.40)	49 (7.50)	29 (4.50)
田	351	74 (21.10)	55 (15.70)	21 (6.00)	28 (8.00)	70 (19.90)	39 (11.10)	18 (5.10)	29 (8.30)	17 (4.80)

TABLE 3: Statistics summary of correspondence analysis.

	Singular value Inertia		Proportion of inertia	
	Siligular value	mertia	Accounted for	Cumulative
Dimensio	on			
1	0.168	0.028	0.778	0.778
2	0.071	0.005	0.139	0.917
3	0.046	0.002	0.057	0.974
4	0.031	0.001	0.026	1.000
	Total	0.036	1.000	1.000

Table 4: Factor loading matrix of correspondence analysis.

Variable	Dimension 1	Dimension 2
Constitution types		
Gentleness type	-0.230	0.310
Qi-deficiency type	0.566	-0.204
Yang-deficiency type	0.423	-0.011
Yin-deficiency type	0.059	0.286
Phlegm-dampness type	-0.685	-0.404
Dampness-heat type	-0.391	-0.060
Blood-stasis type	0.264	0.181
Qi-depression type	-0.032	-0.343
Special diathesis type	-0.127	0.295
Chronic diseases		
Hypertension	-0.024	0.293
Hyperlipidemia	-0.289	-0.111
Diabetes mellitus	-0.021	0.031
Heart disease	0.809	-0.249
Obesity	-0.623	-0.513

Qi acts as the commander of blood while blood is the mother of qi, which means that qi manages the generation and circulation of blood and blood is the carrier of qi. The deficiency of heart qi can result in heart disease due to slow or impeded flow of blood.

There are some limitations in this study. First, this study was based solely on cross-sectional data. Longitudinal studies should be conducted to confirm the findings. Second, the chronic diseases were self-reported and only five chronic diseases were involved in the study. More chronic diseases are recommended to be evaluated in further studies. Third, TCM constitution is also correlated with other factors like eating habits, geological locations, and climates [36]. It would be better to recruit more participants in different areas and to further analyze the correlations between TCM constitution types and chronic diseases in the future.

5. Conclusions

In summary, the associations of TCM constitution types differ in the participants with the five different chronic diseases. Yin-deficiency type and gentleness type were associated with hypertension and diabetes mellitus; phlegm-dampness type was correlated with obesity; dampness-heat




FIGURE 1: Correspondence analysis biplot of TCM constitution types and five chronic diseases. ○: TCM constitution types, 1: gentleness type, 2: qi-deficiency type, 3: yang-deficiency type, 4: yin-deficiency type, 5: phlegm-dampness type, 6: dampness-heat type, 7: bloodstasis type, 8: qi-depression type, and 9: special diathesis type; ▲: chronic diseases, A: hypertension, B: hyperlipidemia, C: diabetes mellitus, D: heart disease, and E: obesity.

type was associated with hyperlipidemia; and qi-deficiency type was correlated with heart disease. Therefore, different interventions based on the TCM constitution should be conducted for different chronic diseases preventions and interventions. At the same time, people with gentleness type should not be ignored in health management.

Disclosure

Huimei Shi is co-first author.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Authors' Contributions

Yanbo Zhu performed the study concept and design; Yanbo Zhu had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis; Yanbo Zhu and Qi Wang were responsible for the acquisition of data; Yanbo Zhu, Yangyang Wang, and Huimei Shi performed the analysis and interpretation of data; Yanbo Zhu, Huimei Shi, and Yangyang Wang drafted the manuscript; Yanbo Zhu, Huimei Shi, Qi Wang, Yangyang Wang, Xiaohan Yu, Jie Di, Xiaomei Zhang, Yanni Li, Tong Li, and Hui Yan performed critical revision of the manuscript for

important intellectual content. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by grants from the National Program on Key Basic Research Project of China (973 Program) (nos. 2011CB505403; 2005CB523501) and Science and Technology Basic Work of Science and Technology (2013FY114400-5).

References

- [1] R. Beaglehole, J. Epping-Jordan, V. Patel et al., "Improving the prevention and management of chronic disease in low-income and middle-income countries: a priority for primary health care," *The Lancet*, vol. 372, no. 964, pp. 940–949, 2008.
- [2] M. Deacon-Crouch, I. Skinner, M. Connelly, and J. Tucci, "Chronic disease, medications and lifestyle: perceptions from a regional Victorian Aboriginal community," *Pharmacy Practice*, vol. 14, no. 3, p. 798, 2016.
- [3] W. H. Dietz, C. E. Douglas, and R. C. Brownson, "Chronic disease prevention, and nutrition for a healthy start," *Journal of the American Medical Association*, vol. 316, no. 16, pp. 1–14, 2016.
- [4] Q. Wang, Constitution Science of Traditional Chinese Medicine, People's Medical Publishing House, Beijing, China, 2005.
- [5] Q. Wang, "Classification and diagnosis basis of nine basic constitutions in chinese medicine," *Journal of Beijing University of Traditional Chinese Medicine*, vol. 28, no. 4, pp. 1–8, 2005.
- [6] L. Yi, T. Pan, and W. Zou, "Relationship between traditional Chinese medicine constitution types with chemotherapy-induced nausea and vomiting in patients with breast cancer: an observational study," BMC Complementary & Alternative Medicine, vol. 16, no. 1, p. 451, 2016.
- [7] T. Wang, J. Chen, X. Sun et al., "Effects of TCMC on transformation of good health status to suboptimal health status: a nested case-control study," *Evidence-based Complementary and Alternative Medicine*, vol. 2015, Article ID 259727, 8 pages, 2015.
- [8] C. H. Lee, T. C. Li, and C. I. Tsai, "Yang deficiency body constitution acts as a predictor of diabetic retinopathy in patients with type 2 diabetes: taichung diabetic body constitution study," *Evidence-Based Complementary and Alternative Medicine*, vol. 2015, Article ID 940898, 8 pages, 2015.
- [9] X. Q. Wu, Y. Y. Luo, J. H. Xu et al., "Diabetes mellitus patients and its relationship with insulin resistance and insulin secretion," *Journal of Guangzhou University of Traditional Chinese Medicine*, vol. 30, no. 3, pp. 312–315, 2013.
- [10] J. H. Cheng, Z. X. Wu, Y. H. Zeng et al., "Study on the distribution of TCM constitution types in patients with primary dyslipidemia," *Journal of New Chinese Medicine*, vol. 46, no. 2, pp. 66–69, 2014.
- [11] X. H. Yu, Y. B. Zhu, Q. Wang et al., "Logistic regression analysis on relationships between metabolic syndrome and constitution type of traditional Chinese medicine and related risk factors," *China Journal of Traditional Chinese Medicine and Pharmacy*, vol. 30, no. 10, pp. 3536–3539, 2015.
- [12] L. M. Li, S. N. Li, and D. L. Wei, "The constitution distribution of angina pectoris patients in lingnan area," *Journal of Traditional Chinese Medicine*, vol. 53, no. 15, pp. 1305–1307, 2012.

- [13] S. X. Zhang, C. Zhang, and X. Ma, "TCM constitution identification and intervention study in mild and moderate hypertension patients in community," *Chinese Archives of Traditional Chinese Medicine*, vol. 33, no. 1, pp. 110–112, 2015.
- [14] J. Wang and X. J. Wang, "Evaluation on the TCM constitution and intervention effect among middle-aged elderly residents in Ningbo," *Chinese Preventive Medicine*, vol. 16, no. 5, pp. 346– 350, 2015.
- [15] X. H. Yu, Y. B. Zhu, Q. Wang et al., "Study on intervention of hypertension patients by comprehensive methods about constitution of traditional Chinese medicine," *Tianjin Journal of Traditional Chinese Medicine*, vol. 32, no. 8, pp. 455–458, 2015.
- [16] J. H. Cheng, Z. X. Wu, Y. B. Zhu et al., "Evaluation of intervention effect of TCM constitution health education," *Journal of Beijing University of Traditional Chinese Medicine*, vol. 39, no. 6, pp. 511–515, 2016.
- [17] Q. Wang, "Individualized medicine, health medicine, and constitutional theory in chinese medicine," *Frontiers of Medicine in China*, vol. 6, no. 1, pp. 1–7, 2012.
- [18] Y. Zhu, Q. Wang, G. Pang et al., "Association between body mass index and health-related quality of life: the 'obesity paradox' in 21,218 adults of the chinese general population," *PLoS ONE*, vol. 10, no. 6, Article ID e0130613, pp. 1–13, 2015.
- [19] Q. Wang, Y. B. Zhu, H. S. Xue, and S. Li, "Primary compiling of constitution in chinese medicine questionnaire," *Chinese Journal of Clinical Rehabilitation*, vol. 10, no. 3, pp. 12–14, 2006.
- [20] Y. B. Zhu, Q. Wang, H. S. Xue et al., "Preliminary assessment on performance of constitution in chinese medicine questionnaire," *Chinese Journal of Clinical Rehabilitation*, vol. 10, no. 3, pp. 15–17, 2006.
- [21] Y. B. Zhu, Q. Wang, and H. Origasa, "Evaluation on reliability and validity of the constitution in chinese medicine questionnaire (CCMQ)," *Chinese Journal of Behavioral Medical Science*, vol. 16, no. 7, pp. 651–654, 2007.
- [22] Q. Wang and Y. B. Zhu, "Epidemiological investigation of constitution types of Chinese medicine in general population: base on 21, 948 epidemiological investigation data of nine provinces in China," *China Journal of Traditional Chinese Medicine and Pharmacy*, vol. 24, no. 1, pp. 7–12, 2009.
- [23] M. Zalewska, K. Furmańczyk, S. Jaworski, W. Niemiro, and B. Samoliński, "The prevalence of asthma and declared asthma in poland on the basis of ECAP survey using correspondence analysis," Computational and Mathematical Methods in Medicine, vol. 2013, Article ID 597845, 2013.
- [24] F. Tekaia, "Genome data exploration using correspondence analysis," *Bioinformatics and Biology Insights*, vol. 10, pp. 59–72, 2016.
- [25] M. Grassi and S. Visentin, "Correspondence analysis applied to grouped cohort data," *Statistics in Medicine*, vol. 13, no. 23-24, pp. 2407–2425, 1994.
- [26] E. S. Gelsema, M. Hunink, C. E. Queiros, and T. Timmers, "The use of correspondence analysis in the assessment of morphologic changes during carcinogenesis," *Cytometry*, vol. 5, no. 5, pp. 463–468, 1984.
- [27] Y. H. Li, "The application of the correspondence analysis technique in the market research," *Sci-Tech Information Development & Economy*, vol. 16, no. 21, pp. 164-165, 2006.
- [28] K. J. Li, A. Y. Wang, and L. Y. Feng, "Application of correspondence analysis in a study of the relationship between row and column variables in a cross table," *Chinese Journal of Health Statistics*, vol. 22, no. 5, pp. 281-282, 2005.

- [29] C. W. Dong and E. X. Gao, "Review of hypertension patients with Yin deficiency with yang hyperactivity," *Journal of Anhui TCM College*, vol. 17, no. 5, pp. 59–61, 1998.
- [30] Y. Zhu, Q. Wang, Z. Dai et al., "Case-control study on the associations between lifestyle-behavioral risk factors and phlegmwetness constitution," *Journal of Traditional Chinese Medicine*, vol. 34, no. 3, pp. 286–292, 2014.
- [31] Y. B. Zhu, Q. Wang, C. Y. Wu et al., "A logistic regression analysis: relationship between traditional chinese medicine body constitution type classification and oveweight and obesity in 18805 chinese adults," *Journal of Chinese Integrative Medicine*, vol. 8, no. 11, pp. 1023–1028, 2010.
- [32] M. Y. Lu, "Dampness-heat and coronary disease," *Journal of Shandong University of TCM*, vol. 27, no. 1, pp. 16–20, 2003.
- [33] Y. L. Feng, G. Y. Zheng, and C. Q. Ling, "The investigation of the correlation between metabolic syndrome and chinese medicine constitution types in senior retired military personnel of the people's liberation army," *Chinese journal of integrative medicine*, vol. 18, no. 7, pp. 485–489, 2012.
- [34] Y. L. Feng, Q. J. Lv, Y. Wang et al., "Relationship between phlegm-dampness and metabolic syndrome in aged people," *Chinese Journal of Clinical Healthcare*, vol. 13, no. 4, pp. 345-346, 2010.
- [35] M. Qu, M. X. Zhang, L. Zhang et al., "Concerning 'heart-qi deficiency' in the role of coronary heart disease (CHD)'s outbreak," *Chinese Archives of Traditional Chinese Medicine*, vol. 28, no. 2, pp. 22–24, 2010.
- [36] C. T. Low, P. C. Lai, and H. D. Li, "Neighbourhood effects on body constitution-a case study of Hong Kong," *Social Science & Medicine*, pp. 158–161, 2016.