

ORIGINAL RESEARCH ARTICLE

Glioblastoma cells express crucial enzymes involved in androgen synthesis: 3 β -hydroxysteroid dehydrogenase, 17-20 α -hydroxylase, 17 β -hydroxysteroid dehydrogenase and 5 α -reductase

Jose Antonio Mondragón Dr PhD, Professor^{1,2}  | Yesenia Serrano Lic Biologist, Student¹ | Andrea Torres¹ | Martin Orozco¹  | Jose Segovia Dr PhD, Professor¹  | Gabriel Manjarrez Dr M.D., Professor²  | Marta Catalina Romano¹ 

¹Departamento de Fisiología, Biofísica y Neurociencias, Centro de Investigación y de Estudios Avanzados del IPN, Ciudad de Mexico, Mexico

²Unidad de Enfermedades Neurológicas, Hospital de Especialidades, México City, México

Correspondence

Marta C. Romano, Departamento de Fisiología, Biofísica y Neurociencias, Centro de Investigación y de Estudios Avanzados del IPN, Avenida IPN # 2508, Ciudad de Mexico 07360, Mexico.
Email: mromano@fisio.cinvestav.mx

Funding information

CONACYT Fronteras to Dr. M.C. Romano, Grant/Award Number: 2015-2 1256

Abstract

Glioblastoma (GB) is the most common and aggressive primary brain tumour in adult humans. Therapeutic resistance and tumour recurrence after surgical removal contribute to poor prognosis for glioblastoma patients. Men are known to be more likely than women to develop an aggressive form of GB, and differences in sex steroids have emerged as a leading explanation for this finding. Studies indicate that the metabolism and proliferation of GB-derived cells are increased by sex steroids, the expression of androgen receptors (ARs) and the synthesis of androgens and oestrogens, suggesting that these hormones have a role in the tumour pathogenesis. The expression of aromatase, the enzyme that converts androgens to oestrogens, has been reported in glial cells and GB cell lines. Thus, it was necessary to test whether the steroidogenic enzymes involved in androgen synthesis are expressed in GB cells. Therefore, here, we investigated the expression of four key enzymes involved in androgen synthesis in human-derived GB cells. U87 cells were cultured in Dulbecco's modified Eagle medium plus foetal bovine serum and antibiotics on slides for immunocytochemistry or immunofluorescence. U87, LN229 and C6 cells were also cultured in multi-well chambers to obtain proteins for Western blotting. We used primary antibodies against 3 β -hydroxysteroid dehydrogenase (3 β -HSD), 17 α -hydroxylase/17,20-lyase (P450c17), 17 β -hydroxysteroid dehydrogenase (17 β -HSD) and 5 α -reductase. Immunocytochemistry, and immunofluorescence results revealed that glioblastoma cells express 3 β -HSD, P450c17, 17 β -HSD and 5 α -reductase proteins in their cytoplasm. Moreover, Western blot analyses revealed bands corresponding to the molecular weight of these four enzymes in the three GB cell lines. Thus, glioblastoma cells have the key enzymatic machinery necessary to synthesize androgens, and these enzymes might be useful targets for new therapeutic approaches.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Endocrinology, Diabetes & Metabolism* published by John Wiley & Sons Ltd.

KEYWORDS

cancer, glioblastoma, neurosteroids, steroidogenic enzymes

1 | INTRODUCTION

Among local brain tumour types, glioma is the most aggressive. The World Health Organization (WHO) has proposed classifying gliomas into four grades based on a set of morphological criteria that determine the cell lineage.¹ More recently, a molecular-based classification (WHO, 2016) for GB was proposed, where isocitrate dehydrogenase wild-type IDH that accounts for about 90% of cases, frequently corresponds with the clinically defined primary or *de novo* glioblastoma, and predominates in patients over 55 years old, and GB IDH-mutant (accounting for 10% of cases), so-called secondary glioblastoma with a history of prior lower grade diffuse glioma, occurs preferentially in younger patients.² The treatments for those types of neoplasms consist mainly of surgery for tumour ablation, radiotherapy and chemotherapy, mainly with temozolomide (TMZ). Unfortunately, patient survival is poor because of frequent recurrences.

Several studies have proposed that steroid hormones play an important role in developing gliomas.³ Preliminary evidence was provided by statistics from the United States of America, where the incidence in the adult population is 50% higher in men than in women^{4,5}). Likewise, the expression of receptors to oestrogen, progesterone and androgens in glioma cell lines and tumour biopsies in GB cell lines has been reported (Gandhari et al., 2006,^{6,7,40}, Chang et al. 2014,⁷

The synthesis of neurosteroids by the central nervous system (CNS) comprises several enzymes. Steroidogenesis in the CNS begins with cholesterol entry into the mitochondria, which is regulated by the protein StAR (steroidogenic acute regulatory protein), which captures cholesterol in the outer mitochondrial membrane and transfers it to the peripheral receptor to benzodiazepines (PBR) located in the mitochondrial membrane. This forms a channel that directs the cholesterol to the inner mitochondrial membrane, where the first steroidogenic enzyme is found, cholesterol desmolase, cytochrome P450_{scc} (side-chain cleavage), which synthesizes pregnenolone (P₅) in the endoplasmic reticulum. Starting from cholesterol, the first step of the pathway is to synthesize pregnenolone (P₅), which is then metabolized to various hormones. The 3 β -hydroxysteroid dehydrogenase delta 4-5 isomerase (3 β -HSD) is a key enzyme in these metabolic pathways and is necessary for producing the 3-oxo-group present in most steroid hormones. This enzyme is also expressed in SH-SY5Y cells⁸ [9]. CYP17 A1 (also termed 17 α -hydroxylase/C17-20-lyase) catalyses the hydroxylation of C21 steroids to C19 steroids. CYP17 A1 is also required to form dehydroepiandrosterone (DHEA) and androstenedione, weak androgens that are converted to testosterone (T) and estrone (E₁). The expression of CYP17A1 mRNAs in the nervous system is controversial, but has been reported in neurons and astrocytes of the

rat brain and SH-SY5Y cells.^{8,9} The multifunctional enzyme 17 β -hydroxysteroid dehydrogenase catalyses the oxidation of neuroactive steroids in the brain, which are involved in brain development, neurogenesis, neural plasticity and neuroprotection and has an impact on brain diseases. The originally described functions of 17 β -hydroxysteroid dehydrogenases were to maintain the balance between the less potent (17-Keto) and more potent (17 β -hydroxy) forms of oestrogen and androgens, also, they are involved in the metabolism of fatty acids and sterols (for a review, please see.¹⁰

The expression of the steroidogenic enzymes P450 cholesterol side-chain-cleavage and P450-aromatase (P450arom) in glioma cell lines and tumour biopsies also accounts for the possible participation of steroid hormones in GBM. Papadopoulos et al.¹¹ confirmed the presence of cytochrome P450 cholesterol side-chain-cleavage enzyme by immunoblotting in the mitochondria of the rat GB C6 cell line. The expression of the enzyme responsible for oestrogen biosynthesis, P450-aromatase (P450arom), in human and rat glioblastoma cell lines was described by Yague et al.¹² Furthermore, Dueñas Jiménez et al.¹³ proposed this enzyme as a prognostic biomarker in astrocytoma patients.

The ability of GB-derived cells to synthesize steroids has been reported by Papadopoulos et al.¹¹, who demonstrated that rat C6 cells transformed 22-hydroxy cholesterol to pregnenolone, and by Melcangi et al.¹⁴, who reported that C6 and 1321 N1 human astrocytoma-derived cells converted testosterone and progesterone into 5 α -reduced derivatives DHT and dihydroprogesterone (DHP). We have recently shown that U87 cells synthesize several androgens, and that the production can be inhibited by the 5 α -reductase inhibitors finasteride and dutasteride.¹⁵ Those results strongly suggest the presence of a panoply of enzymes involved in androgen synthesis.

The final products of steroidogenesis in each tissue depend on the enzymes that synthesize these hormones, such as P450-cytochromes, steroid dehydrogenases and reductases.¹⁶ Baulieu¹⁷ described the CNS as a steroidogenic tissue and showed that steroids (eg pregnenolone (P₅) and DHEA) and their sulphated forms (P₄S and DHEAS) were found in higher concentrations in the brain than in the peripheral circulation of rodents Robel and Baulieu.¹⁸

The central nervous system (CNS) expresses most of the synthesizing enzymes or modifiers of steroids; most enzymes involved in the steroid synthesis have been documented in the rat brain and regions of the human brain.¹⁹⁻²² The enzymatic pathways involved in neurosteroid synthesis and regulation have been extensively studied.²³ Oligodendrocytes, astrocytes and neurons present steroidogenic capacity. However, it is important to note that these enzymes have distinct expression profiles and activities in neurons and glial cells.⁹ The conversion of androgens like testosterone

to DHT, the potent AR agonist, results from 5α -reductase activities. Therefore, these enzymes are essential for androgen synthesis in several classic steroidogenic tissues.²⁴ Three types of this enzyme have been described, they are expressed in the normal prostate and are associated with prostate cancer initiation and progression²⁵, the prostate 5α -reductase mRNA levels of types 1 and 2 are regulated by testosterone and DHT.²⁵

The 5α -reductase isoforms 1 and 2 are expressed in the human brain where they participate in synthesis of neurosteroids.²² The 5α -reductase type 1 is constitutively expressed in the rat brain, its expression is negatively regulated by T and DHT, while gene expression of 5α -reductase type 2 is under the positive control of T and DHT.²⁶

The physiological importance of 5α -reductases in the brain derives from its ability to convert testosterone (T) to the potent androgen, dihydrotestosterone (DHT) and progesterone and deoxycorticosterone (DOC) to their respective 5α -reduced derivatives, precursors of allopregnanolone and tetrahydro DOC, which are both strong allosteric modulators of the aminobutyric acid receptor (GABA_AR).²⁷ The implication of 5α -reductases and aromatase in the growth of androgen and oestrogen-sensitive tumours, such as prostate, breast and GB cancers, suggested that these steroidogenic enzymes are expressed in GB-derived cells. Using RT-qPCR, Zamora-Sanchez et al.²⁸ have shown 5α -reductase mRNA expression in U87 cells.

We have recently shown that the androgens T₄ and DHT, but not androstenedione, significantly increase GB metabolism, and that the effect of these androgens is inhibited by using the 5α -reductase inhibitor dutasteride or the androgen receptor blocker finasteride.²⁹ Since androgens have a role in GB development, it is important to have a deeper insight into the tumour steroidogenic ability, focusing on androgen synthesis. Here, we aimed to describe the expression of key steroidogenic enzymes involved in the androgen synthesis by GB-derived cells.

2 | MATERIALS AND METHODS

2.1 | Cell culture

Human GB-derived U87 cells were purchased from American Type Culture Collection (ATCC, Manassas, VA, USA). Cells (5×10^4) were seeded in culture slides (Falcon, New York, USA), cultured in high-glucose Dulbecco's modified Eagle medium (GIBCO BRL) supplemented with 10% foetal bovine serum (FBS, GIBCO BRL), 2 mM glutamine (Sigma) and 100 units- μ g penicillin-streptomycin (GIBCO BRL) and maintained at 37°C in 95% air, 5% CO₂. For the experiments, the complete medium was removed, and the cells were washed with phosphate-buffered-saline solution, pH 7.2 (PBS). For the Western blot analysis, the cells were seeded in multi-well plates and cultured in the same conditions described above. LN 229 and C6 cells were cultured with the same methodology described for U87 cells.

2.2 | Immunocytochemistry

The cultured cells were fixed in 4% paraformaldehyde and ethanol dehydrated. An antigenic unmask was carried using a 0.1 M citrate buffer solution, pH 6.0 (DECLERE, Cell Mark, Rocklin, CA, USA), followed by the inhibition of endoperoxidase activity with 5% H₂O₂. After washing the slides, they were incubated with specific primary antibodies for CYP17 (Rabbit Polyclonal CYP17A1 antibody, GeneTex), for 5α -reductase Type2 (rabbit monoclonal [EPR6281(B)] to SRD5A2, Abcam) and antisera against 17 β HSD type I (Raised in rabbit using purified type 1 from the human placenta) and 3 β -HSD (Raised in rabbit using purified type I human placental 3 β -HSD) at a dilution of 1:1000 in a commercial antibody diluent solution (Cell Marque) and incubated for 18 h at 4°C. Next, the sections were washed with PBS and incubated with the secondary antibody diluted in PBS and Triton X 100 (Biotinylated anti-mouse-rabbit IgG antibody) for 30 min. The cells were then incubated with avidin-biotin complex for 15 min at room temperature and washed. Immunoreactivity was determined with a commercial kit (3,3-diaminobenzidine and H₂O₂) according to Vector Laboratories' protocol³⁰ and mounted in ENTELLAN mounting media for microscopy. Photomicrographs were taken with a digital camera (Infinity 1-Lumenera) with a 40 \times objective, using an Infinity 1-Lumenera camera equipped with a 10 \times objective, aided by an Olympus microscope.

2.3 | Immunofluorescence

For immunofluorescence observations, we fixed the slides as described for immunocytochemistry and exposed them to the same primary antibodies. After 24 h, the slides were washed with 0.1% PBS-Tween and then incubated with Alexa Fluor 546 anti-rabbit diluted to 1:100. The slides were mounted with ProLong Diamond Antifade Mountant for immunofluorescence and observed under a Leica TCS SP8 Confocal Laser Scanning Microscope (Leica Microsystems) with a 40 \times objective.

2.4 | Western Blotting

Cells were scrapped with a cell homogenizer and mixed for 30 s at 4°C in a solution of 50 mM Tris-HCl, pH 7.4, plus protease inhibitors (Protease Inhibitor Cocktail, Sigma-Aldrich Quimica SA de CV, Mexico). The samples were then centrifuged at 29,000 g for 15 min at 4°C. Protein concentration was determined by the bicinchoninic acid method. Then, 30 μ g of protein were placed in each of the 1-mm thick channels of an SDS-polyacrylamide gel at 12%. The electrophoretic run was performed at 100 V for 2 h (Mini-Protean Tetra System, BioRad). For protein electrotransference, the gels were mounted on nitrocellulose membranes, and the run was performed at 25 V for 45 min (Trans-Blot SD Semi-Dry Transfer Cell, BioRad). Nitrocellulose membranes with the transferred proteins were placed in a blocking solution (Millipore Chemiluminescent Blocker)

at 50% for 30 min. The membranes were incubated with the same primary antibodies specific for 3β -HSD, CYP17, 17β HSD and 5α reductase used for immunocytochemistry described above, at a dilution of 1:1000 in the same blocking solution. On the following day, membranes were incubated with a secondary anti-rabbit antibody (BioRad) at a dilution of 1:5000 in blocking solution. Membranes were revealed with chemiluminescence (Millipore, corporation, Billerica, U.S.A) and exposed to a film (Hiperfilm). The internal control used was glyceraldehyde-3-phosphate dehydrogenase (GADPH) or tubulin. The bands obtained were analysed and quantified by densitometry using Image Studio Software (LI-COR Biosciences).

3 | RESULTS

Figure 1 shows the abundant immunostaining for 3β -HSD detected in the cytoplasm of human GB-derived cells U87 using an immunocytochemistry technique (Figure 1B). The negative control is shown in Figure 1A. Confocal images obtained by immunofluorescent also detected the enzyme in the cytoplasm (Figure 1C), and 3D images revealed the lack of immunoreactivity in the nucleus (not shown).

The expression of U87 protein (42 KDa MW) corresponding to 3β -HSD can be observed in the Western blot analysis in Figure 1D. This figure also shows the expression of the 3β -HSD protein in another GB cell lines, LN229 and C6.

By immunocytochemistry, we detected the expression of 17α -hydroxylase (CYP17), the enzyme that converts progesterone to 17OH -progesterone and androstenedione in U87 cells (Figure 2B). Confocal images confirmed the cytoplasmic distribution of the enzyme in U87 cells (Figure 2C). Western blot analysis showed a band corresponding to the molecular weight of the of CYP17 protein (57 KDa MW, Figure 2D). This figure also shows the expression of the 3β -HSD protein in another GB cell lines, LN229 and C6.

The 17β -hydroxysteroid dehydrogenase expression was found in the cytoplasm of U87 cells exposed to a specific antibody by immunocytochemistry (Figure 3B). Figure 3C displays the immunofluorescent images obtained by confocal microscopy. Both techniques showed a cytoplasmic distribution of the enzyme. Western blotting showed a band corresponding to the molecular weight of the protein (35 KDa MW, Figure 3D). This figure also shows that the expression of the 3β -HSD protein is also present in another GB cell lines, LN229 and C6.




FIGURE 1 Micrographs illustrating immunostaining and protein expression for 3β -hydroxysteroid dehydrogenase in GB-derived U87 cells. Immunoreactivity by immunocytochemistry: (A) Negative control, (B) expression of the enzyme. (C) expression of the protein by immunofluorescence (400X). Western blot expression of the protein in U87, LN229 and C6 cells, and corresponding densitometric analysis. 400x




FIGURE 2 17 α -hydroxylase (CYP17) immunoreactivity in glioblastoma cells. (A) Negative control. (B) expression of the enzyme. (C) expression of the protein by immunofluorescence. Western blot expression of the protein by, and corresponding densitometry analysis. (400X). Western blot expression of the protein in U87, LN299 and C6 cells, and corresponding densitometric analysis

The expression of 5 α -reductase type 2, an enzyme that converts T to the potent androgen DHT, is present in the cytoplasm of U 87 cells, as shown by immunocytochemistry (Figure 4B). The enzyme was also observed in confocal microscopy images (Figure 4C). Western blot analysis detected the expression of the 5 α -reductase protein (25 KDa MW) that can be observed in Figure 4D. This figure also shows that L229 and C6 cells express the protein.

Figure 5 shows the densitometric analysis of the expression of the four steroidogenic enzymes included in our study in U87, LN229 and C6 cells lines. The two human and the rat GB cell lines expressed the four enzymes, although their expression level was different, the prevalence of the investigated 5 α -reductase in U87 cells was evidenced by the statistical analysis.

4 | DISCUSSION

As described in the Introduction, several authors have shown that normal glial cells synthesize androgens and oestrogens and express steroidogenic enzymes (P450c17, 3 β -HSD, 17 β -HSD and P450 aromatase).⁹ These findings suggest that the expression of these enzymes can be found in transformed glial cells. Supporting this

possibility, our own studies and those of other authors have found that GB-derived cells synthesize androgens and oestrogens, for the synthesis of these sex steroids, the action of several steroidogenic enzymes is essential, including 3 β -HSD, 17-20 α -lyase, 17 β -hydroxysteroid dehydrogenase and 5 α -reductase. The inhibition of androgen synthesis caused by inhibitory drugs such as finasteride and dutasteride, strongly suggested the presence of specific enzymes in U87 GB-derived cells.¹⁵ Following these findings, we have investigated the expression and distribution of the key enzymes directly involved in androgen synthesis in cultured GB-derived cells using different techniques and confirmed their presence in two other cell lines by Western blot.

The expression of 3 β -HSD found in U87 cells has been reported in cultured normal glial cells³¹, and the results of our immunocytochemical techniques show that this enzymes are preserved in GB-derived cells. This finding supports the reported steroidogenic capacity of C6 and U87 cells to provide steroids by transforming pregnenolone, DHEA and androstenediol to its metabolites.¹⁵ Our Western blotting analyses further demonstrated the presence of the enzyme protein in U87, LN229 and C6 GB-derived cells. Two isoforms of 3 β -HSD have been characterized in humans, type 1 is mainly expressed in the placenta, and type 2 in the adrenal cortex


17 β -Hydroxysteroid Dehydrogenase (HSD17B)

FIGURE 3 Light microscope micrograph illustrating 17 β -hydroxysteroid dehydrogenase expression as studied by immunocytochemistry, confocal microscopy, and Western blot. (A) Negative control. (B) expression of the enzyme. (C) expression of the protein by immunofluorescence. (400X). Western blot expression of the protein in U87, LN299 and C6 cells, and corresponding densitometric analysis

and gonads.³²⁻³⁴ Immunostaining using an antibody against 3 β -HSD type 1 shows that U87, LN229 and C6 cells express this isoform in the cytoplasm.

In the brain, DHEA and androstenedione are synthesized from pregnenolone and progesterone, respectively. The expression of the immunoreactive protein P450-Cyp 17 has been found in neurons of the hippocampus, where it localizes in pyramidal neurons, granule cells of the dentate gyrus (Hojo et al., 2004), hypothalamic periventricular neurons and Purkinje cells.³⁵ Furthermore, the immunocytochemical expression of P450-Cyp17 in astrocytes was found in the rat brain by Zwain and Yen.⁹ We have previously reported that U87 cells synthesize DHEA and androstenedione,¹⁵ a finding that suggested the presence of P450-Cyp 17 in these cells. The subcellular expression of this enzyme is now confirmed by immunocytochemical techniques and by the expression of the protein by Western blot analysis.

The activity of 17 β -HSD has been suspected based on brain tissue capacity to synthesize androgens and oestrogens^{18,23}. We have previously found the transformation of A₄ to testosterone, and DHEA to androstenediol by U87-derived GB cells¹⁵, therefore, the enzyme should be necessarily present in these cells, as we show in this study using immunohistochemical procedures. The expression

of the enzyme found by Western blot further confirms the presence of the protein in GB cells.

Type 1 17 β -HSD also participates in the synthesis of oestrogens, transforming estrone to estradiol and vice-versa.³⁶ Our study showed that the type 1 17 β -HSD antibody recognized the GB enzyme in the immunocytochemical and Western blot studies presented here, confirming that type 1 is present in the cells. This type facilitates the reduction in DHEA to androstenediol, and DHT to 3 α and 3 β -diol, and reactions that other authors and we have shown to occur in GB cells.³⁷

We and others have previously shown that the potent androgens testosterone and DHT increased the metabolism and proliferation of GB-derived cells^{29,38}, and these steroids positively control the gene expression of 5 α -reductase type 2.²⁶ The 5 α -reductase protein was highly expressed in glioblastoma-derived cells if compared with the expression of the three other enzymes investigated in this study, and a finding that could be explained by the positive control of the enzyme by the abundant androgens synthesized by these cells.

As we have shown studying the effect of androgens on U87 cells by MTT, testosterone and the strong androgen DHT, but not androstenedione, significantly enhance the metabolism, proliferation and


5 α -Reductase (SRD5A2)

FIGURE 4 Micrographs illustrating immunostaining and protein expression for 5 α -reductase Type 1 in GB-derived U87 cells. Immunoreactivity by immunocytochemistry: (A) Negative control, (B) expression of the enzyme. (C) expression of the protein by immunofluorescence. Western blot expression of the protein in U87, LN299 and C6 cells, and corresponding densitometric analysis




FIGURE 5 Densitometry corresponding to the expression of 3 β -hydroxysteroid dehydrogenase, 17 α -hydroxylase (CYP17), 17 β -hydroxysteroid dehydrogenase and 5 α -reductase in U87, LN229 and C6 GB-derived cells. Data are presented as means \pm SD. $N = 3$. ANOVA one way was followed by Kruskal-Wallis and Dunn's Multiple comparison tests. GraphPad Prism 5.01. * $p < .05$

invasion of U87 GB-derived cells.²⁹ Rodriguez-Lozano et al.³⁹ have shown that testosterone, promotes cell proliferation, migration and invasion of GB-derived cells through the androgen receptors. Therefore, the expression of 5 α -reductase would be essential to provide the cells with potent androgens necessary for cell metabolism. We recently showed that dutasteride, or the combination of this 5 α -reductase inhibitor with the AR blockers, cyproterone and flutamide, significantly reduced the metabolism, cell proliferation

and migration capacity of U87,²⁹ findings that strongly support the presence of 5 α -reductases found in these cells.

The results presented here indicate that U87 cells maintain the steroidogenic enzymes present in glial cells. Therefore, U87 cells have the machinery necessary to synthesize androgens and other neurosteroids that might have a role in the GB development. These findings were also shown in LN229 and C6 cells. These enzymes might be considered as therapeutic targets for the treatment of human glioblastomas.

ACKNOWLEDGEMENTS

Supported by CONACYT Fronteras 2015-2 1256 to Dr. M.C. Romano.

CONFLICT OF INTEREST

We declare that there is not conflict of interest


AUTHOR CONTRIBUTIONS

Jose Antonio Mondragón: formalAnalysis (equal); investigation (equal); methodology (equal). Yesenia C Serrano: investigation (equal); methodology (equal). Andrea C Torres: methodology (equal). Martin C Orozco: investigation (equal); methodology (equal). Jose C Segovia: conceptualization (equal); supervision (equal). Gabriel Manjarrez: conceptualization (equal); investigation (equal); methodology (equal). Marta Catalina Romano Pardo: conceptualization (lead); dataCuration (lead); formalAnalysis (equal); fundingAcquisition (lead); investigation (equal); writingOriginalDraft (lead); writingReviewEditing (lead).

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in [repository name e.g. "figshare"] at <https://www.doi.org/> [doi], reference number [reference number].


ORCID

Jose Antonio Mondragón  <https://orcid.org/0000-0002-4817-3598>

Martin Orozco  <https://orcid.org/0000-0002-6264-4026>

Jose Segovia  <https://orcid.org/0000-0001-8215-4772>

Gabriel Manjarrez  <https://orcid.org/0000-0003-0101-6875>

Marta Catalina Romano  <https://orcid.org/0000-0003-2932-0801>

REFERENCES

- Louis DN, Ohgaki H, Wiestler OD, et al. The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol.* 2007;2007(114):97-109g.
- Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health organization classification of tumors of the central nervous system: A summary. *Acta Neuropathol.* 2016;131:803-820. <https://doi.org/10.1007/s00401-016-1545-13>.
- Kabat GC, Etgen AM, Rohan TE. Do steroid hormones play a role in the etiology of glioma? *Cancer Epidemiol, Biomarkers Prev.* 2010;19:2421-2427. <https://doi.org/10.1158/1055-9965.EPI-10-0658>.
- McKinley BP, Michalek AM, Fenstermaker RA, Plunkett RJ. The impact of age and sex on the incidence of glial tumors in New York State from 1976 to 1995. *J Neurosurg.* 2000;93:932-939.
- Tamimi AF. Epidemiology and Outcome of Glioblastoma, Glioblastoma, , Vleschouwer S (Eds.), *Glioblastoma*. Brisbane (AU): Codon Publications; 2017.
- González-Agüero G, Ondarza R, Gamboa-Dominguez A, Cerbon MA, Camacho-Arroyo I. Progesterone receptor isoforms expression pattern in human astrocytomas. *Brain Res Bull.* 2001;56:43-48. [https://doi.org/10.1016/S0361-9230\(01\)00590-1](https://doi.org/10.1016/S0361-9230(01)00590-1).
- Yin Y, Zhang X, Li Z, et al. Glucocorticoid receptor β regulates injury-mediated astrocyte activation and contributes to glioma pathogenesis via modulation of β -catenin/TCF transcriptional activity. *Neurobiol Dis.* 2013;59:165-176.
- Emanuelsson I, Almokhtar M, Wikvall K, et al. Expression and regulation of 3β -hydroxysteroid dehydrogenase in cells of the nervous system: Potential effects of vitamin D on brain steroidogenesis. *Neurochem Int.* 2018;113:46-55. <https://doi.org/10.1016/j.neuint.2017.11.007>.
- Zwain IH, Yen SS. Neurosteroidogenesis in astrocytes, oligodendrocytes, and neurons of cerebral cortex of rat brain. *Endocrinology.* 1999;140:3843-3852.
- Poutanen M, Penning TM. Biology and clinical relevance of hydroxysteroid (17 β) dehydrogenase enzymes. *Mol Cell Endocrinol.* 2019;489:1-2.
- Papadopoulos V. Structure and function of the peripheral-type benzodiazepine receptor in steroidogenic cells. *Proc Soc Exp Biol Med.* 1998;217:130-142. <https://doi.org/10.3181/00379727-217-44215>
- Yague JG, Lavaque E, Carretero J, Azcoitia I, Garcia-Segura LM. Aromatase, the enzyme responsible for estrogen biosynthesis, is expressed by human and rat glioblastomas. *Neurosci Lett.* 2004;368:279-284.
- Dueñas-Jiménez JM, Candanedo-Arellano A, Santerre A, et al. Aromatase and estrogen receptor alpha mRNA expression as prognostic biomarkers in patients with astrocytomas. *J Neurooncol.* 2014;119:275-284.
- Melcangi RC, Cavarretta I, Magnaghi V, Ballabio M, Martini L, Motta M. Crosstalk between normal and tumoral brain cells. Effect on sex steroid metabolism. *Endocrine.* 1998;8:65-71. <https://doi.org/10.1385/ENDO:8:1:65>.
- Pinacho-García LM, Valdez RA, Navarrete A, Cabeza M, Segovia J, Romano MC. The effect of finasteride and dutasteride on the synthesis of neurosteroids by glioblastoma cells. *Steroids.* 2020;155:108556.
- Sanderson T. The steroid biosynthesis pathway as a target for endocrine-disrupting chemicals. *Toxicol Sci.* 2006;94:3-21.
- Baulieu EE. Neurosteroids: A novel function of the brain. *Psychoneuroendocrinology.* 1998;23:963-987.
- Robel P, Baulieu E-E. Neurosteroids: Biosynthesis and function. *Trends Endocrinol Metab.* 1994;5:1-8.
- Mellon SH, Deschepper CF. Neurosteroid biosynthesis: Genes for adrenal steroidogenic enzymes are expressed in the brain. *Brain Res.* 1993;629:283-292.
- Gomez-Sanchez EP, Ahmad N, Romero DG, Gomez-Sanchez CE. Is aldosterone synthesized within the rat brain? *Am J Physiol Endocrinol Metab.* 2005;288:E342-E346. <https://doi.org/10.1152/ajpendo.00355.2004>.
- Yu L, Romero DG, Gomez-Sanchez CE, Gomez-Sanchez EP. Steroidogenic enzyme gene expression in the human brain. *Mol Cell Endocrinol.* 2002;190:9-17.
- Pelletier G. Steroidogenic enzymes in the brain. In Martini L (Eds.), *Progress in Brain Research*. Elsevier. Netherlands: 2010;181:193-207. ISSN: 0079-6123.
- Do Rego JL, Seong JY, Burel D, et al. Neurosteroid biosynthesis: enzymatic pathways and neuroendocrine regulation by neurotransmitters and neuropeptides. *Front Neuroendocrinol.* 2009;30(3):259-301. <https://doi.org/10.1016/j.yfrne.2009.05.006>.
- Azzouni F, Godoy A, Li Y, Mohler J. The 5 alpha-reductase isozyme family: A review of basic biology and their role in human diseases. *Adv Urol.* 2012;2012:530121. <https://doi.org/10.1155/2012/530121>.
- Li J, Wang Z, et al. Androgen regulation of 5 α -reductase in prostate cancer: Implications for prostate cancer prevention. *PLoS One.* 2011;6(12):e28840.
- Torres JM, Ortega E. Differential regulation of steroid 5 α -reductase isoenzymes expression by androgens in the adult brain. *FASEB J.* 2020;17:1428-1433.
- García-Segura LM, Melcangi RC. Steroids and glial cell function. *Glia.* 2006;54:485-498.

28. Zamora-Sanchez CJ, Hansberg-Pastor V, Salido-Guadarrama I, Rodríguez-Dorantes M, Camacho-Arroyo I. Allopregnanolone promotes proliferation and differential gene expression in human glioblastoma cells. *Steroids*. 2017;119:36-42.
29. Orozco M, Valdez RA, Ramos L, Cabeza M, Segovia J, Romano MC. Dutasteride combined with androgen receptor antagonists inhibit glioblastoma U87 cell metabolism, proliferation, and invasion capacity: Androgen regulation. *Steroids*. 2020;164:108733.
30. Naish JS, Boenisch T, Farmilo AJ, Stead RH. *Handbook of Immunohistochemical staining methods*. Ed. Dako Corp. Carpinteria, California USA: 1989:1-41.
31. Brown RC, Cascio C, Papadopoulos V. Pathways of neurosteroid biosynthesis in cell lines from human brain: regulation of dehydropiandrosterone formation by oxidative stress and beta-amyloid peptide. *J Neurochem*. 2000;74:847-859.
32. Luu The V, Lachance Y, Labrie C, et al. Full length cDNA structure and deduced amino acid sequence of human 3beta-hydroxy-5-enesteroiddehydrogenase. *Mol Endocrinol*. 1989;3:1310-1312.
33. Rhéaume E, Lachance Y, Zhao HF, et al. Structure and expression of a new complementary DNA encoding the almost exclusive 3 beta-hydroxysteroid dehydrogenase/delta 5-delta 4-isomerase in human adrenals and gonads. *Mol Endocrinol*. 1991;5:1147-1157. <https://doi.org/10.1210/mend-5-8-1147>.
34. Simard J, Ricketts ML, Gingras S, Soucy P, Feltus FA, Melner MH. Molecular biology of the 3β-hydroxysteroid dehydrogenase/Δ⁵-Δ⁴ isomerase gene family. *Endocr Rev*. 2005;26:525-582. <https://doi.org/10.1210/er.2002-0050>.
35. Yamada H, Kominami S, Takemori S, Kitawaki J, Kataoka Y. Immunohistochemical localization of cytochrome P450 enzymes in the rat brain, considering the steroid synthesis in the neurons. *AH Cytochem*. 1997;30:609-616.
36. Mensah-Nyagan AG, Do-Rego JL, Beaujean D, Luu-The V, Pelletier G, Vaudry H. Neurosteroids: Expression of steroidogenic enzymes and regulation of steroid biosynthesis in the central nervous system. *Pharmacol Rev*. 1999;51:63-82.
37. Hilborn E, Stal O, Jansson A. Estrogen and androgen-converting enzymes 17β-hydroxysteroid dehydrogenase and their involvement in cancer: with a special focus on 17β- hydroxysteroid dehydrogenase type 1, 2, and breast cancer. *Oncotarget*. 2017;8:30552-30562.
38. Rodríguez-Lozano DC, Velázquez-Vázquez D, Del Moral-Morales A, Camacho-Arroyo I. Dihydrotestosterone induces proliferation, migration and invasion of human glioblastoma cell lines. *Onco Targets Ther*. 2020;13:8813-8823. <https://doi.org/10.2147/OTT.S262359>.
39. Rodríguez-Lozano DC, Piña-Medina AG, Hansberg-Pastor V, Bello-Álvarez C, Camacho-Arroyo I. Testosterone promotes glioblastoma cell proliferation, migration and invasion through androgen receptor activation. *Front Endocrinol*. 2019;10:16. <https://doi.org/10.3389/fendo.2019.00016>.
40. Gandhari MK, Frazier CR, Hartenstein JS, Cloix JF, Bernier M, Wainer IW. Identification and characterization of estrogen receptor-related receptor alpha and gamma in human glioma and astrocytoma cells. *Mol Cell Endocrinol*. 2010;5(315):314-318.
41. Hojo Y, Hattori T, Enami T, Furukawa A, Suzuki K, Ishii H, Mukai H, Morrison JH, Janssen WGM, Kominami S, Kominami S, Harada N, Kimoto T, Kawato S. Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017α and P450 aromatase localized in neurons. *PNAS*. 2004;101(3):865-870.

How to cite this article: Mondragón JA, Serrano YC, Torres AC, et al. Glioblastoma cells express crucial enzymes involved in androgen synthesis: 3β-hydroxysteroid dehydrogenase, 17-20α-hydroxylase, 17β-hydroxysteroid dehydrogenase and 5α-reductase. *Endocrinol Diab Metab*. 2021;4:e00289. <https://doi.org/10.1002/edm2.289>