

Research Article

Effect of Eye Movement Training on Sleep Quality of Patients with Advanced Lung Cancer Based on Pittsburgh Sleep Quality Index

Haiping Hu,¹ Wenying Yang,² Zhimin Liu,¹ Xiaona Zhang,¹ Junmei Shi,¹ and Huixia Xu ¹

¹Department of Oncology, The Fourth Hospital of Hebei Medical University, Hebei Cancer Hospital, Shijiazhuang 050011, Hebei, China

²Department of Blood Purification Center, Wuhan No. 1 Hospital, Wuhan Hospital of Traditional Chinese and Western Medicine, Wuhan 430022, Hubei, China

Correspondence should be addressed to Huixia Xu; xuhuixia@hbydsy.org.cn

Received 29 October 2021; Revised 11 November 2021; Accepted 22 November 2021; Published 16 December 2021

Academic Editor: Kalidoss Rajakani

Copyright © 2021 Haiping Hu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. To explore the effect of eye movement training on sleep quality of patients with advanced lung cancer based on the Pittsburgh Sleep Quality Index (PSQI). **Methods.** 120 advanced lung cancer patients admitted to our hospital from January 2019 to January 2020 were selected as the research object and divided into group A (PSQI scores ≥ 10 points, $n = 60$) and group B (PSQI < 10 points, $n = 60$). Routine nursing was performed to both groups, and patients in group A received the eye movement training additionally, so as to compare their PSQI scores, negative emotion scores, adverse reaction rate (ARR), Cancer Coping Modes Questionnaire (CCMQ) scores, and pain scores. **Results.** After training, group A obtained significantly better sleep quality ($P < 0.05$), lower negative emotion scores ($P < 0.001$), lower ARR ($P < 0.05$), better CCMQ scores ($P < 0.05$), and lower pain scores ($P < 0.001$) than group B. **Conclusion.** Eye movement training should be promoted in practice because it can reduce negative emotions, alleviate pain sensation, improve sleep quality and body condition, and lower the ARR for advanced lung cancer patients.

1. Introduction

Lung cancer is a common clinical condition. The latest annual report from the National Cancer Registry in 2018 showed that lung cancer is the malignant tumor with the highest incidence (11.6%) and mortality (18.4%) globally [1]. The causing factors include the environment, pulmonary history, and gene. Lung cancer patients usually present with cough, phlegm, and other clinical symptoms, and some patients even experience the phenomenon of coughing up blood and hoarseness, which greatly affect their life health in a negative way [2, 3]. In the middle to late stage of lung cancer, patients will feel intense cancerous pain that not only affects body condition but also aggravates psychological stress, hence increasing the possibility of sleep disorders. Sleep science is an emerging borderline science, and there are few theoretical studies on this discipline in China currently, not to mention its application, but the relationship

between sleep science and cancer has been confirmed by many literatures [4–7]. As patients with advanced lung cancer suffer from various symptoms, more emphasis is placed on treating their somatic symptoms in the clinic with assignment-oriented nursing. Even though some nursing plans add the content of psychological care, it is not targeted enough in managing sleep disorders, so the sleep condition is not well-improved yet [8–10]. With the increasing focus on cancer patients, there are more and more reports involving the quality of life in advanced lung cancer patients, but studies related to their sleep quality are extremely rare. In recent years, it has been reported that oculomotor training can ameliorate the condition of patients with diabetic retinopathy and improve their sleep quality [11], so such intervention measure may also be applied to cancer patients. To explore the effect of eye movement training on the sleep quality of advanced lung cancer patients based on PSQI, 120 patients with advanced lung cancer admitted to

our hospital from January 2019 to January 2020 were selected as the research object for the study, with the results summarized as follows.

2. Materials and Methods

2.1. General Information. 120 advanced lung cancer patients treated in our hospital from January 2019 to January 2020 were selected as the research object and divided into group A (PSQI scores ≥ 10 points, $n=60$) and group B (PSQI scores < 10 points, $n=60$), with no statistical difference in their general information ($P > 0.05$), see Table 1. The study was approved by the Hospital Ethics Committee.

2.2. Inclusion Criteria. The inclusion criteria of the study were as follows. (1) The patients or their family members fully understood the study process and signed the informed consent; (2) the patients met the diagnosis criteria for advanced lung cancer after examination [12] and were accepting treatment in our hospital without chemotherapy yet; (3) the estimated survival time of the patients was more than 3 months; and (4) the patients had sleep disorders [13].

2.3. Exclusion Criteria. The exclusion criteria for the patients of the study were as follows. (1) Presence of mental problems or inability to communicate with others; (2) suffering from other organic diseases [14]; (3) having the need for analgesic drugs [15]; (4) with sleep disorders not caused by cancer.

2.4. Methods. Both groups accepted the routine nursing with the following specific steps. (1) The nursing personnel provided medication intervention and recorded the drug usage according to medical advice. (2) The nursing personnel customized a healthy diet for the patients according to their actual situation and asked their sport preference to guide them to do exercise reasonably. (3) The nursing personnel carried out health education to the patients according to their actual situation, informed the patients and their family members of correct home nursing methods, and adjusted the nursing plan at any time based on the patients' condition, so as to improve the effect and quality of nursing.

Eye movement training was given to patients in group A additionally, with the following specific steps. (1) The nursing personnel informed the patients of the function of eye movement training to gain their trust and increase their compliance. (2) The patients were kept in a quiet and relaxed condition, gazed with both eyes into the distance horizontally for 0.5 h, during which the eyeballs were moved inwards (with medial rectus), upwards/inwards (with superior rectus), downwards/inwards (with inferior rectus), upwards/outwards (with inferior oblique), and downwards/outwards (with superior oblique) for 36 times each and moved annular from left to right to the maximum extent. (3) The patients did the eye movement training once a day and 5 days a week for 1 month. (4) The sleep index table was put at the foot of the patients' beds for dynamic observation; their sleep indexes were evaluated daily during training, and

outpatient evaluation and guidance were provided at the time of discharge from the hospital.

2.5. Observation Criteria

- (1) Pittsburgh Sleep Quality Index (PSQI): six factors in the PSQI scale (sleep quality, sleep latency, sleep efficiency, use of sleeping medication, and daytime dysfunction) were selected and compared before and after training. The total score was 18 points, with 0–3 points for each item, and lower scores indicated better sleep [16].
- (2) Negative emotion scores: the negative emotions were evaluated by the Self-Rating Anxiety Scale (SAS) and Self-Rating Depression Scale (SDS) before and after training, and each scale contained 20 items and has a total score of 100 points. Higher scores indicated that the negative emotions of patients were more serious [17].
- (3) Adverse reaction rate (ARR): the adverse reactions included digestive tract reaction, allergic reaction, injury of skin and mucous, and myelosuppression, and the number of patients with adverse reactions was counted.
- (4) Cancer Coping Modes Questionnaire (CCMQ): it included 26 items and 5 dimensions, of which the higher scores on confrontation and avoidance and depression indicated a more positive attitude towards the condition, while the higher scores on acceptance-resignation, imagination, and abreaction indicated a more negative attitude towards the condition [18].
- (5) Pain scores: the Chinese version of the Brief Pain Inventory (BPI-C) scale was used to evaluate the pain degree of patients in both groups before and after training, which covered five dimensions such as the most serious pain to the slightest pain within 24 hours. On a scale of 0–10 points, higher scores indicated more serious pain [19].

2.6. Statistical Processing. In this study, the data processing software was SPSS20.0, the picture drawing software was GraphPad Prism 7 (GraphPad Software, San Diego, USA), items included were enumeration data and measurement data, methods used were X^2 test and t -test, and differences were considered statistically significant at $P < 0.05$.

3. Results

3.1. Comparison of PSQI Scores. After training, the sleep quality of group A was significantly better than that of group B ($P < 0.05$), see Table 2.

3.2. Comparison of Negative Emotion Scores. After training, the negative emotion scores of group A were significantly lower than those of group B ($P < 0.001$), see Table 3.

TABLE 1: Comparison of general information.

Group	Group A (n = 60)	Group B (n = 60)	χ^2/t	P value
Gender			0.034	0.853
Male	35	34		
Female	25	26		
Age (years old)				
Range	32–74	33–74		
Mean age	61.21 ± 6.20	61.23 ± 6.21	0.018	0.986
TNM stage			0.034	0.854
IIIb	34	33		
IV	26	27		
Distant metastasis				
Lymphatic metastasis	30	30	0.000	1.000
Bone metastasis	16	17	0.042	0.838
Liver metastasis	10	11	0.058	0.810
Metastasis to other parts	4	3	0.152	0.697
Clinical history				
Diabetes	8	9	0.069	0.793
Hypertension	12	13	0.051	0.822
Coronary heart disease	7	6	0.086	0.769
Lung disease	10	11	0.058	0.810
Educational level				
Junior high school and below	12	13	0.051	0.822
Senior high school	30	30	0.000	1.000
College and above	18	17	0.040	0.841
Monthly income (yuan)			0.035	0.852
≥3000	36	37		
<3000	24	23		

TABLE 2: Comparison of PSQI scores after training ($\bar{x} \pm s$, points).

Group	Sleep latency	Sleep duration	Sleep efficiency	Sleep disorders	Use of sleeping medication	Daily dysfunction	Sleep quality
Group A	1.68 ± 0.65	1.18 ± 0.24	0.59 ± 0.35	1.00 ± 0.32	1.02 ± 0.24	1.45 ± 0.65	1.75 ± 0.65
Group B	2.01 ± 0.54	1.52 ± 0.28	1.20 ± 0.39	1.23 ± 0.65	1.38 ± 0.34	2.10 ± 0.54	2.00 ± 0.54
t	3.025	7.141	9.017	2.459	6.700	5.958	2.292
P value	0.003	0.000	0.000	0.015	0.000	0.000	0.024

TABLE 3: Comparison of negative emotion scores ($\bar{x} \pm s$, points).

Group	SDS before and after		SAS before and after	
Group A	58.65 ± 6.68	42.11 ± 5.98	60.11 ± 10.65	40.65 ± 5.68
Group B	58.84 ± 6.57	48.98 ± 6.95	61.20 ± 10.54	47.11 ± 5.65
t	0.157	5.804	0.563	6.246
P value	0.876	0.000	0.574	0.000

3.3. Comparison of ARR. The ARR of group A was significantly lower than that of group B ($P < 0.05$), see Figure 1.

Note: in Figure 1, the black areas indicate digestive tract reaction, the dark gray areas indicate allergic reaction, the light gray areas indicate injury of skin and mucous, and the white areas indicate myelosuppression; the left image indicates group A, and the right image indicates group B.

The number of patients with digestive tract reaction in group A and group B was 30 and 40, respectively ($\chi^2 = 11.868$, $P < 0.05$).

The number of patients with allergic reaction in group A and group B was 24 and 36, respectively ($\chi^2 = 4.800$, $P < 0.05$).

The number of patients with injury of skin and mucous in group A and group B was 24 and 42, respectively ($\chi^2 = 10.909$, $P < 0.05$).

The number of patients with myelosuppression in group A and group B was 25 and 45, respectively ($\chi^2 = 13.714$, $P < 0.05$).

3.4. Comparison of CCMQ Scores. After training, the CCMQ scores of group A were significantly better than those of group B ($P < 0.05$), see Table 4.

3.5. Comparison of Pain Scores. After training, the pain scores of group A were significantly lower than those of group B ($P < 0.001$), see Figure 2.

Note: in Figure 2, the horizontal axis indicates before and after training, and the vertical axis indicates the pain scores (points); the black areas indicate group A, and the gray areas indicate group B; and * indicates $P < 0.001$.

The data before training indicated that the results of comparing the pain scores between group A and group B




FIGURE 1: Comparison of ARR. (a) Group A = 103. (b) Group B = 171.

TABLE 4: Comparison of CCMQ scores after training ($\bar{x} \pm s$, points).

Group	Confrontation	Avoidance and depression	Acceptance-resignation	Imagination	Abreaction
Group A	3.25 ± 0.54	1.68 ± 0.65	2.18 ± 0.34	1.45 ± 0.25	1.54 ± 0.54
Group B	2.54 ± 0.65	2.20 ± 0.48	2.65 ± 0.54	1.87 ± 0.60	1.89 ± 0.68
<i>t</i>	6.508	4.985	5.705	5.005	3.122
<i>P</i> value	0.000	0.000	0.000	0.000	0.002




FIGURE 2: Comparison of pain scores before and after training ($\bar{x} \pm s$, points).

were not statistically different (4.98 ± 0.54 vs 4.87 ± 0.58 , $P > 0.001$).

The data after training indicated that the pain scores of group A were significantly lower than those of group B (3.65 ± 0.35 vs 4.25 ± 0.48 , $P < 0.001$).

4. Discussion

At present, the number of patients suffering from malignant tumors is increasing year by year worldwide, expanding the group of cancer patients continuously. Reports indicated that there are more than 2 million new lung cancer patients every year with poor prognosis. The academia has gradually

recognized that the purpose of treating cancer is not only limited to resection of tumors but also to improve the life health of patients and their quality of life. Lung cancer is one of the most common cancers in China, and 47% of lung cancer patients experience sleep disorders [20] that seriously compromise the treatment efficacy. Patients present severe nausea and decreased appetite from long-term treatment and cancer-induced fatigue after progressing to the end stage of the disease; at this time, their depression is more pronounced and their sleep quality decreases even further, raising the possibility of sleep disorders to over 50%. As sleep quality is closely related to life quality and treatment effect, and clinical practice has regarded life quality as one of the first indicators to evaluate the prognosis of patients, it is crucial to increase the efforts in conducting corresponding research.

Sleep science is an emerging science in China, but it is not well studied in academia at present nor fully applied in practice, so there is still a lack of well-established measures for sleep quality improvement in China. The oculomotor training selected in this study is often used in the treatment of diseases such as cognitive impairment. As an aerobic exercise of the brain, it is able to improve the cognitive ability of patients and reduce their physical and mental burden, and Secinti et al. stated that oculomotor training can help the body of patients to enter a state of complete relaxation, hence it is positive in weakening the anxiety of patients [21]. The study showed that the negative emotion scores after training of group A were significantly lower than those of group B ($P < 0.001$), indicating that the oculomotor training successfully brought the patients into a quiet and relaxed

atmosphere to enable them to take control of themselves again, greatly improve their self-efficacy, and alleviate their anxiety. Since anxiety and depression are the major factors that trigger sleep disorders in advanced lung cancer patients [22], the sleep quality will be improved with these emotions alleviated, as indicated by shorter sleep latency and lower likelihood of using sleep medication.

It is indisputable that sleep quality is of positive relevance with life health [23]. For instance, the study by Pongthavornkamol et al. showed that high-quality sleep can improve the psychological and body state and accelerate the rehabilitation progress, hence it is in favor of body recovery [24], while poor sleep quality affects the regular work and rest, disrupts the circadian rhythm, and negatively influences the quality of life and treatment effect of patients. Better sleep quality will improve the physical and psychological conditions in patients. Therefore, group A obtained significantly better CCMQ scores ($P < 0.05$) and lower pain scores ($P < 0.001$) than group B after training, indicating that because of a good sleep, patients in group A promoted their self-awareness, greatly enhanced their confidence in treatment, kept a more positive mind, and improved their body tolerance and that the sleep quality can affect the prognosis of advanced lung cancer patients.

In addition, the study also concluded that the ARR of group A was remarkably lower than that of group B ($P < 0.05$), which was consistent with the findings of Kizilgz and other scholars. In their study, patients in the experimental group received routine nursing combined with eye movement treatment, and patients in the control group were given the routine nursing only, and it was concluded that after treatment, the probabilities of patients in the experimental group having digestive tract reaction, allergic reaction, injury of the skin, and mucous and myelosuppression were 48%, 50%, 48% and 45%, respectively, which were obviously lower than those in the control group ($P < 0.05$) [25], indicating that the eye movement training could promote the body status by improving the sleep quality, thereby enhancing the tolerance and lowering the possibility of adverse reactions in patients.

In conclusion, eye movement training should be promoted in practice because it can reduce negative emotions, alleviate pain sensation, improve sleep quality and body condition, and lower the ARR for advanced lung cancer patients.

Data Availability

Data that supported the findings of this study are available on reasonable request from the corresponding author.

Disclosure

The authors received no financial support for the research, authorship, and/or publication of this article.

Conflicts of Interest

The authors have no conflicts of interest to declare.

Authors' Contributions

Haiping Hu and Wenying Yang contributed equally to this article.

References

- [1] N. Belloumi, S. B. Maalej, I. Bachouche, F. B. A. Chermiti, and S. Fenniche, "Comparison of sleep quality before and after chemotherapy in locally advanced nonsmall cell lung cancer patients: a prospective study," *Sleep Disorders*, vol. 2020, Article ID 8235238, 8 pages, 2020.
- [2] G. Jakobsen, M. Engstrøm, M. Thronæs et al., "Sleep quality in hospitalized patients with advanced cancer: an observational study using self-reports of sleep and actigraphy," *Supportive Care Cancer*, vol. 28, 2019.
- [3] A. Perkins and K. Cham, "Sleep in patients living with mesothelioma: a scoping review," *BMJ Open Respiratory Research*, vol. 6, no. 1, 2019.
- [4] D. Papadopoulos, A. Papadoudis, M. Kiagia, and K. Syrigos, "Nonpharmacologic interventions for improving sleep disturbances in patients with lung cancer: a systematic review and meta-analysis," *Journal of Pain and Symptom Management*, vol. 55, 2018.
- [5] C. Dsab, B. Mla, B. RCA et al., "Adjuvant melatonin for the prevention of recurrence and mortality following lung cancer resection (AMPLCaRe): a randomized placebo controlled clinical trial," *EclinicalMedicine*, vol. 33, 2021.
- [6] M. T. Gentry, P. J. Atherton, M. Lapid et al., "Effects of a multidisciplinary quality of life intervention on sleep quality in patients with advanced cancer receiving radiation therapy," *Palliative and Supportive Care*, vol. 18, no. 3, pp. 1–7, 2019.
- [7] J. H. Check, D. Check, and T. Poretta, "Mifepristone extends both length and quality of life in a patient with advanced non-small cell lung cancer that has progressed despite chemotherapy and a check-point inhibitor," *Anticancer Research*, vol. 39, no. 4, pp. 1923–1926, 2019.
- [8] Z. Matjaz, "Toxicity and quality of life in published clinical trials for advanced lung cancer," *Supportive Care in Cancer*, vol. 26, no. 10, pp. 3453–3459, 2018.
- [9] J. H. Check, D. Check, and T. Poretta, "Abstract 4715: mifepristone extends both length and quality of life in a patient with advanced non-small cell lung cancer that progressed despite chemotherapy and check-point inhibitors," *Cancer Research*, vol. 78, no. 13, 2018.
- [10] G. Demiray, S. Degirmencioglu, E. Ugurlu et al., "Relationship between serum ghrelin levels, weight loss and quality of life in patients with advanced non-small cell lung cancer," *International Journal of Cell Science & Molecular Biology*, vol. 4, 2018.
- [11] C. Weiss, S. Dickerson, M. L. Dubocovich, and G. E. Dean, "Bright light therapy to reduce fatigue and improve sleep quality in lung cancer survivors," *Sleep*, vol. 41, no. 1, 2018.
- [12] P. Horvat, C. Gray, A. Lambova et al., "Comparing findings from a friends of cancer research exploratory analysis of real-world end points with the cancer analysis system in England," *JCO Clinical Cancer Informatics*, vol. 5, pp. 1155–1168, 2012.
- [13] M. Mahdieh, F. Hossein, F. Fatemeh, and M. Mohammad, "Sleep quality and cancer-related fatigue in patients with cancer," *International Journal of Caring Sciences*, vol. 3, pp. 145–152, 2021.
- [14] S. J. Rice and C. P. Belani, "Diversity and heterogeneity of immune states in non-small cell lung cancer and small cell lung cancer," *PLoS One*, vol. 16, no. 12, Article ID e0260988, 2021.

- [15] Y. Bulbul, T. Ozlü, and S. Arinc, "Sleep disturbances in patients with lung cancer in Turkey," *Tuberkuloz Ve Toraks*, vol. 66, no. 4, pp. 297–303, 2018.
- [16] N. A. Arrato, S. B. Lo, M. Grogan et al., "Improving bio-behavioral outcomes with progressive muscle relaxation in patients with advanced lung cancer," *Journal of Clinical Oncology*, vol. 37, no. 27, p. 292, 2019.
- [17] D. Ge, P. Ziegler, H. Chen, L. M. Steinbrenner, and S. S. Dickerson, "Trajectory of insomnia symptoms in older adults with lung cancer: using mixed methods," *Supportive Care in Cancer*, vol. 27, no. 1, 2019.
- [18] M. Ida, H. Onodera, M. Yamauchi, and M. Kawaguchi, "Preoperative sleep disruption and postoperative functional disability in lung surgery patients: a prospective observational study," *Journal of Anesthesia*, vol. 33, no. 4, pp. 501–508, 2019.
- [19] T. R. Mendoza, K. L. Kehl, O. Bamidele et al., "Assessment of baseline symptom burden in treatment-naïve patients with lung cancer: an observational study," *Supportive Care Cancer*, vol. 27, 2019.
- [20] D. Kizilirmak, T. Goktalay, Y. Havlucu, Ö. Gültekin, and P. Çelik, "Effects of chemotherapy in lung cancer on sleep and quality of life," *Turkish Thoracic Journal*, vol. 20, no. 1, p. 191, 2019.
- [21] E. Secinti, D. Tometich, S. A. Johns, and C. Mosher, "Symptom experiences in advanced stage cancer: relationships to acceptance and commitment therapy constructs," in *Proceedings of the Society of Behavioral Medicine*, Washington, DC, USA, March 2019.
- [22] D. Papadopoulos, M. Kiagia, A. Charpidou, I. Gkiozos, and E. Syrigos, "Psychological correlates of sleep quality in lung cancer patients under chemotherapy: a single-center cross-sectional study," *Psycho-Oncology*, vol. 28, no. 9, 2019.
- [23] M. Dómine, B. Massuti, J. Puente et al., "Observational prospective study to determine the evolution of the symptomatic profile of metastatic non-small cell lung cancer (NSCLC) patients and its relation to the control of the disease," *Advances in Therapy*, vol. 36, 2019.
- [24] K. Pongthavornkamol, P. Lekdamrongkul, P. Pinsuntorn, and A. Molassiotis, "Physical symptoms, unmet needs, and quality of life in Thai cancer survivors after the completion of primary treatment," *Asia-Pacific Journal of Oncology Nursing*, vol. 6, no. 4, p. 363, 2019.
- [25] D. Kizilgz, H. Yeildal, and A. Kabalak, "Effect of eye movement training on sleep quality of patients with advanced lung cancer," *Eurasian Journal of Pulmonology*, vol. 21, no. 2, p. 93, 2019.