

Research Article

Pressure-dependent NOS activation contributes to endothelial hyperpermeability in a model of acute heart failure

Andreia Z. Chignalia¹, Ayman Isbatan¹, Milan Patel¹, Richard Ripper^{1,2}, Jordan Sharlin¹, Joelle Shosfy¹, Barry A. Borlaug³ and Randal O. Dull^{1,4}

¹Department of Anesthesiology, University of Illinois at Chicago, Chicago, IL 60612, U.S.A.; ²Research and Development Service, Jesse Brown Veterans Affairs Medical Center, 820 S Damen Ave., Chicago, IL 60612, U.S.A.; ³Department of Cardiovascular Medicine, Mayo Clinic and Foundation, 200 First St SW, Rochester, MN 55905, U.S.A.; ⁴Department of Anesthesiology, University of Arizona College of Medicine and Banner-University Medical Center, Tucson, AZ 85724, U.S.A.

Correspondence: Andreia Zago Chignalia (andreiac@uic.edu)



Aims: Acute increases in left ventricular end diastolic pressure (LVEDP) can induce pulmonary edema (PE). The mechanism(s) for this rapid onset edema may involve more than just increased fluid filtration. Lung endothelial cell permeability is regulated by pressure-dependent activation of nitric oxide synthase (NOS). Herein, we demonstrate that pressure-dependent NOS activation contributes to vascular failure and PE in a model of acute heart failure (AHF) caused by hypertension.

Methods and results: Male Sprague–Dawley rats were anesthetized and mechanically ventilated. Acute hypertension was induced by norepinephrine (NE) infusion and resulted in an increase in LVEDP and pulmonary artery pressure (P_{pa}) that were associated with a rapid fall in P_{aO_2} , and increases in lung wet/dry ratio and injury scores. Heart failure (HF) lungs showed increased nitrotyrosine content and ROS levels. L-NAME pretreatment mitigated the development of PE and reduced lung ROS concentrations to sham levels. Apocynin (Apo) pretreatment inhibited PE. Addition of tetrahydrobiopterin (BH4) to AHF rats lung lysates and pretreatment of AHF rats with folic acid (FA) prevented ROS production indicating endothelial NOS (eNOS) uncoupling.

Conclusion: Pressure-dependent NOS activation leads to acute endothelial hyperpermeability and rapid PE by an increase in NO and ROS in a model of AHF. Acute increases in pulmonary vascular pressure, without NOS activation, was insufficient to cause significant PE. These results suggest a clinically relevant role of endothelial mechanotransduction in the pathogenesis of AHF and further highlights the concept of active barrier failure in AHF. Therapies targetting the prevention or reversal of endothelial hyperpermeability may be a novel therapeutic strategy in AHF.

Introduction

Acute increases in left ventricular end diastolic pressure (LVEDP) and pulmonary capillary pressure (P_{pc}) can cause rapid and severe pulmonary edema (PE) [1]. Clinically, however, the severity of PE is often out of proportion to the increase in P_{pc} suggesting that other mechanisms may be operational. To understand this clinical disparity, we explored the relationship between sudden changes in hydrostatic pressure and acute changes in endothelial permeability. We have previously demonstrated that small changes in hydrostatic pressure produced large, rapid, and sustained increases in endothelial permeability, an effect mediated by nitric oxide (NO) [2] and reactive oxygen species (ROS) [3]. In fact, NO-mediated increases

Received: 12 April 2018
Revised: 11 October 2018
Accepted: 21 October 2018

Accepted Manuscript Online:
24 October 2018
Version of Record published:
23 November 2018

in endothelial permeability during increased hydrostatic pressure are common throughout the vascular system [4,5]. Thus, increases in both pressure and shear stress can activate nitric oxide synthase (NOS) and increase endothelial NO. Nitrosylation of adherens junction proteins is a trigger for junctional disassembly and results in acute increases in endothelial permeability [6]. The mechanisms whereby ROS contributes to endothelial hyperpermeability, however, are less understood. Reactive oxygen species are generated in a controlled and compartmentalized manner in the vasculature and although NADPH oxidases (Nox) are considered to be the main source of ROS in the vascular system, there is no evidence that Nox regulates barrier function; thus, other sources for ROS production, such as uncoupled endothelial NOS (eNOS) [7], may play a role in endothelial hyperpermeability and PE.

Interestingly, the majority of patients with chronic heart failure (HF) do not develop frank PE, even when P_{pc} is elevated [8,9] suggesting that alterations in endothelial signaling pathways, structural alteration of the vascular wall, or both, contribute to compensatory mechanism(s) that protect against sustained high pressure. We recognize that while NO causes acute changes in endothelial permeability, organic nitrites are used to treat pulmonary congestion and mitigate the symptoms of chronic HF. This paradox suggests, perhaps, that the physiological response of the pulmonary vasculature to NO and/or oxidative stress changes during the progression from acute HF (AHF) to chronic HF and, therefore, warrants further mechanistic characterization. Understanding the early mechanisms that regulate disease progression maybe an important step to better understand this paradox and improve therapeutic approaches to HF.

Herein, we describe a series of studies demonstrating the contribution of NOS activation to acute endothelial barrier failure in a model of AHF. The results demonstrate that eNOS has a dual role in AHF-related PE: it acts by increasing NO concentrations and second as a source of superoxide anion production leading to endothelial hyperpermeability and pulmonary vascular failure. Inhibition of NOS attenuates the physiological and histological pulmonary changes associated with AHF. The novel findings demonstrate that pressure-dependent NOS-activation results in enhancement of endothelial permeability and rapid PE. This represents a major paradigm shift in our understanding of hydrostatic PE and may lead to new therapeutic strategies to re-establish barrier integrity.

Methods

Reagents

NADPH, Norepinephrine (NE), Dihydroethidium (DHE), protease inhibitor cocktail, and Krebs Ringer Buffer were bought from Sigma Co, Ltd (St. Louis, MO); BSA was purchased from Proliant Biologicals (Boone, IA); antibodies were bought from EDM Millipore (Billerica, MA), Aviva Systems Biology (San Diego, CA) and Cell Signaling (Danvers, MA); Lucigenin and L-NAME were bought from Cayman Biochemicals; apocynin was bought from Calbiochem.

Animals

Animal studies were approved by the University of Illinois Institutional Animal Care and Use Committee. The investigation conforms to the *Guide for the Care and Use of Laboratory Animals* published by the U.S. National Institutes of Health. Male Sprague–Dawley rats were divided in the following groups: Sham, AHF; L-NAME + AHF (L-NAME + AHF); L-NAME; Apocynin (Apo); Apocynin + AHF (Apo + AHF); Ethanol (EtOH); Folic Acid (FA) and FA + AHF (FA + AHF).

AHF model

Rats received a NE infusion starting at 7 $\mu\text{g}/\text{Kg}/\text{min}$ and titrated to a mean arterial pressure (MAP) of 150 mm Hg for 2 h. Sham rats received a lactated ringers infusion (1.5 ml/h). For specific experiments, L-NAME or Apo (200 $\mu\text{mol}/\text{l}/\text{kg}$) was administered as a bolus over a 10-min period immediately before the induction of AHF. Additional control groups included solo administration of L-NAME and Apo to determine if it had any effects on the measured parameters independent of NE infusion. In order to assess the role of mechanotransduction in the development of PE during AHF, a non-hypertensive dose of L-NAME was chosen to be used in this model. With this approach, although increased mechanical forces and hemodynamics of HF are still present, we can specifically assess the contribution of NOS signaling to PE development.

Arterial blood gas analysis

Arterial blood gases, hematocrit (HCT) and pH were measured in 250 μl of whole blood using a GEM Premier 3000 machine (Instrumentation Laboratory, Orangeburg, NY), according to manufacturer's instructions.

Hemodynamics

Left ventricular end systolic pressure (LVESP), LVEDP, and pulmonary artery (PA) pressures were measured using a saline-filled PE tubing (PE 50) connected to an arterial pressure transducer interfaced to a TAM-A amplifier (Harvard Apparatus, Holliston, MA).

Lung wet-to-dry ratio

Wet-to-dry (W/D) ratios were determined after drying lungs for 24 h at 60°C.

Lung injury score

Lung injury was assessed by five blinded investigators and scored based on perivascular cuffing (PVC) and intra-alveolar hemorrhage (IAH). Total lung injury score (LIS) was determined by the weighted average of PVC and IAH.

Isolated perfused lung preparation

The rat isolated perfused lung preparation was used as previously described [2,10]. Briefly, rats were anesthetized and mechanically ventilated. The PA and left atria were cannulated and lungs were perfused with Krebs-Ringer bicarbonate solution containing 3% BSA. Lungs were exposed to low pressure (6 cm H₂O) or high pressure (12 cm H₂O) for 60 min.

To assess if NE could induce lung hyperpermeability, 10⁻⁶ mol/L of NE was added to the perfusate reservoir and circulated for 1 h at low pressure. Lung W/D ratios were determined.

To assess if NE had a direct effect on lung vasculature, a concentration-response curve for NE (10⁻⁸ to 10⁻³ mol/L) was performed and pulmonary artery pressure (P_{pa}) was recorded.

BAL albumin content

BAL albumin content was assessed using a fluorimetric detection kit (Active Motif, Carlsbad, CA) according to manufacturer's instructions.

MPO activity

MPO activity was measured in lung tissue and in BAL as previously described [11].

Immunoblotting

NOS, Nox, and NO production

Total eNOS, iNOS, Nox1, and Nox2 expression was assessed in whole lung lysates. eNOS activity was determined by p-Ser¹¹⁷⁷/eNOS ratio; NO production was assessed for nitrotyrosine content as a marker for mechanotransduction activation. Immunoblots were performed as previously described [12]. Signal was detected by chemiluminescence using Li-COR system and band intensities were measured using Image Studio software (Li-COR, Lincoln, NE).

eNOS uncoupling

To assess whether eNOS is uncoupled, we performed a low-temperature gradient gel following standard Western blot techniques as previously described [13]. Signal was detected by chemiluminescence using Li-COR system. Band intensities were measured using Image Studio software (Li-COR).

ROS measurements

Lucigenin Enhanced Chemiluminescence

ROS production was assessed by lucigenin enhanced chemiluminescence (ECL) as previously described [12]. To determine NOS-dependent superoxide generation, rats were treated with L-NAME and lucigenin ECL assay was performed. To confirm uncoupled eNOS as a source of ROS production in lung homogenates, we assessed ROS levels in lung lysates with exogenous tetrahydrobiopterin (BH₄) supplementation (*in vitro*). Addition of PEG-SOD *in vitro* to lung samples was done as an additional control to show superoxide anion is being detected in this assay.

Nox activity

Nox activity was assessed in lung membrane fractions as previously described [12]. Results were normalized by protein content in the samples.

Table 1 AHF model

	SHAM	AHF	L-NAME + AHF	L-NAME
MAP values (mm Hg)				
Baseline	109.8 ± 10.1	123.8 ± 19.3	117.2 ± 14.3	110.6 ± 15.0
Final (120 min)	105.7 ± 9.7	152.2 ± 22.2 ^{†‡}	149.9 ± 20.2 ^{*,§‡}	118.3 ± 25.4
P_{pa} (mm Hg)				
Baseline SBP	29.3 ± 3.2	26.6 ± 7.8	27.0 ± 2.6	27.0 ± 2.2
Baseline DBP	4.7 ± 1.1	10.1 ± 5.3	9.3 ± 5.5	11.3 ± 5.9
Final SBP (120 min)	30.3 ± 3.21	39.3 ± 5.9	38.3 ± 8.8	27.3 ± 2.87
Final DBP (120 min)	5.0 ± 1.7	18 ± 6.1 [‡]	14.2 ± 7.5 [‡]	13.0 ± 7.5
mPAP	13.4 ± 1.0	25.1 ± 6.3 [*]	22.23 ± 7.5	17.8 ± 5.6
Cardiac parameters				
HR (beats/min)	348.0 ± 8.2	361.2 ± 16.2	352.3 ± 12.4	344.0 ± 28.6
Baseline LVESP/LVEDP (mm Hg)	138.3 ± 7.6/6.3 ± 0.6	148.6 ± 21.9/9.67 ± 5.6	149.4 ± 19.0/10.4 ± 8.4	152.4 ± 16.62/7.8 ± 3.1
Final (120 min) LVESP/LVEDP (mm Hg)	138.0 ± 0.5/6.0 ± 0.1	229.6 ± 28.1/16.3 ± 9.5 [‡]	212.0 ± 35.1/15.2 ± 13.9 [‡]	165.6 ± 25.4/10.6 ± 5.7
Arterial blood gases				
pH	7.48 ± 0.1	7.159 ± 0.2 [*]	7.50 ± 0.3 [†]	7.49 ± 0.1
PaO ₂	109.6 ± 28.5	71.29 ± 11.6 [*]	93.0 ± 18.8	91.8 ± 10.6
PaCO ₂	34.78 ± 2.7	36.50 ± 8.9	39.80 ± 5.7 [§]	33.63 ± 10.1
HCT	33.63 ± 4.4	41.20 ± 4.9 [*]	36 ± 5.7	38.57 ± 3.4

Experimental groups: Sham; AHF; L-NAME + AHF (L-NAME bolus followed by NE infusion), and L-NAME (L-NAME bolus followed by lactated ringers infusion); $n \geq 3$ /group. Abbreviations: mPAP, mean P_{pa}; PaCO₂, arterial CO₂ pressure; PaO₂, arterial O₂ pressure.

* $P < 0.05$ compared with Sham.

[†] $P < 0.05$ compared with NE.

[‡] $P < 0.05$ compared with baseline from same group.

[§] $P < 0.05$ compared with L-NAME.

Statistical analysis

Data are presented as mean ± S.D. Groups were compared using one-way ANOVA or Student's *t* test as appropriate. Tukey's post-test was used to compensate for multiple test procedures. $P < 0.05$ was considered statistically significant.

Results AHF model

During general anesthesia, baseline MAP was 109.8 ± 10 mm Hg and heart rate (HR) averaged 348/min (Table 1). All hemodynamic variables were stable over the 2-h time period. Acute hypertension was induced by a NE infusion resulting in a rapid increase in MAP to 150 mm Hg and a significant increase in LVESP and LVEDP without changes in HR (see Table 1).

Acute hypertension caused a significant increase in pulmonary artery systolic pressure (PASP, 26 ± 8 compared with 39 ± 16 mm Hg) and a doubling of mean P_{pa} (mPAP; Sham = 13.14 ± 1.0 compared with AHF = 25.1 ± 6.3 mm Hg; Table 1). To characterize the direct effects NE on the pulmonary pressures, we used the *in situ* isolated perfused rat lung preparation and tested if NE infusion (10^{-8} to 10^{-3} mol/L) altered P_{pa}. NE had no direct effect on pulmonary vascular pressure (Supplementary Figure S1A), indicating that NE-induced pulmonary hypertension requires an intact circulation.

To clarify if NE *per se* could induce lung hyperpermeability, we used the isolated perfused lung preparation to test the effects of high pressure and NE on lung edema. While high pressure induced lung edema when compared with control group, NE did not alter lung W/D ratio when compared with control group (Supplementary Figure S1B). These results demonstrate that pressure, and not NE, is the main stimulus of endothelial hyperpermeability during AHF.

Elevations in LVEDP and PAP during AHF were associated with the development of PE. P_aO₂ dropped on an average to 50% (Figure 1A) and lung W/D ratio increased significantly (5.3 ± 0.52 compared with 6.44 ± 0.75 ; Figure 1B). Histological evidence of lung injury was manifested as significantly increased PVC and increased IAH. Higher LIS were observed in the dependent lung regions (dorsal) compared with non-dependent regions (ventral). In the




Figure 1. AHF model

Rats with AHF showed increased (A) ΔPaO_2 ; (B) Lung W/D ratio; (C) Nitrotyrosine (NO₂ tyrosine) content when compared with sham rats. L-NAME administration prior to induction of AHF (LN + AHF) prevented all indices of PE (PaO_2 , lung W/D). L-NAME (LN) bolus had no effect on the measured parameters. $n \geq 6$ /group; * $P < 0.05$ compared with Sham, ** $P < 0.01$ compared with Sham, *** $P < 0.001$ compared with Sham, ●●● $P < 0.001$ compared with AHF.

dorsal regions, total LIS for Sham was 0.17 ± 0.05 and for AHF rats = 0.38 ± 0.05 (Supplementary Figures S2 and S3).

Pressure-dependent hyperpermeability during AHF is not an inflammatory process

To determine if AHF and associated increase in P_{cp} caused an increase in leukocytes recruitment to the lungs that contributed to barrier failure, CD45 expression in lung sections, MPO activity in BAL and lung tissue were measured. As shown in Supplementary Figure S4A,B, lung MPO activity was not increased in the AHF BAL and lungs. In the same manner, no differences in CD45 staining was observed in lung sections from AHF lungs when compared with Sham lungs (Supplementary Figure S4D), ruling out increased white blood cell accumulation or activation as a cause for hyperpermeability. We also assessed the levels of interleukin 1- β (IL1- β), macrophage inflammatory protein 2 (MIP2), and tumor necrosis factor- α (TNF- α) by real-time PCR in lung lysates from sham and AHF rats. No differences in IL1- β , TNF- α , and MIP2 were found corroborating our previous data that this is not an inflammatory model of PE during AHF (Supplementary Figure S5).

Epithelial barrier is not damaged during AHF

To determine if epithelial barrier disruption contributed to the reduction in P_aO_2 and increase in lung W/D, albumin content of BAL fluid was measured. Acute increases in pulmonary vascular pressures were not associated with an increase in BAL albumin concentration (Supplementary Figure S4C) when compared with Sham BAL, indicating that the alveolar epithelial barrier remained intact.




Figure 2. Oxidative stress contributes to PE in AHF

Lungs collected from AHF rats showed (A) increased ROS levels when compared with lungs collected from sham rats. L-NAME administration before AHF (LN + AHF) prevented increase in lung ROS while L-NAME alone had no effect on ROS production in sham rats. (B) Apo attenuated the increase in lung W/D ratio in AHF rats but had no effect on control rat lung W/D. (C) Nox activity and (D,E) expression is not altered in AHF. L-NAME had no effect on Nox activity. $n \geq 4/\text{group}$; * $P < 0.05$ compared with Sham, ** $P < 0.01$ compared with Sham, ● $P < 0.05$ compared with AHF, ●● $P < 0.01$ compared with AHF.

NO in AHF

To determine if acute vascular pressure increased lung NO production, we indirectly assessed NO generation by quantitating nitration of tyrosine residues in lung lysates. As shown in Figure 1C, acute hypertension and associated increase in mPAP resulted in a 40% increase in nitro-tyrosine content, consistent with our previous findings [2] and indicates activation of endothelial mechanotransduction.

Inhibition of NOS prevents HF-dependent PE

To determine if pressure-dependent PE was caused by NO-mediated barrier failure, rats were treated with the NOS inhibitor, L-NAME, prior to the induction of AHF. L-NAME attenuated PE, returning P_aO_2 to Sham values (Figure 1A). In addition, L-NAME abolished the AHF-induced increase in lung W/D ratio (L-NAME + AHF wet/dry = 4.8 ± 0.41 compared with AHF = 6.4 ± 0.75) (Figure 1B). Pretreatment with L-NAME reduced histological evidence of lung injury during AHF. Histological LIS in the dorsal region were reduced in L-NAME + AHF to 0.381 ± 0.05 , compared with LIS = 0.287 ± 0.06 (in AHF); $P = 0.0098$ (Supplementary Figure S3).

Reactive oxygen species contribute to PE development in pressure-induced AHF

In order to evaluate if ROS contribute to PE during pressure-induced AHF, we first measured ROS concentrations in lung homogenates from Sham and AHF lungs. AHF lungs showed increased ROS concentration when compared with Sham lungs (299.7 ± 140.3 in AHF compared with 50.91 ± 35.47 in Sham; Figure 2A). The increase in ROS concentration was inhibited when rats were pretreated with L-NAME (L-NAME + AHF: 52.94 ± 45.32) (Figure 2A) suggesting eNOS uncoupling. L-NAME, alone, did not affect ROS concentration (111.7 ± 51.10 in L-NAME compared with 50.91 ± 35.47 in Sham). To investigate if ROS contribute to PE, we assessed lung W/D ratio in rats treated with apocynin prior to induction of AHF. Apo inhibited PE development in AHF (W/D = 4.80 ± 0.21 in Apo + AHF compared with 6.44 ± 0.75 in AHF). Rats that received either apocynin only or EtOH (vehicle used to dissolve Apo) did not show differences in lung W/D ratio (Apo = 5.02 ± 0.7 ; EtOH = 4.8 ± 0.36) when compared with sham rats (5.25 ± 0.59) (Figure 2B).




Figure 3. eNOS uncoupling during AHF

AHF results in (A) eNOS uncoupling, an effect inhibited by pretreatment of hypertensive rats with L-NAME (LN + AHF). L-NAME alone had no effect on eNOS monomer/dimer ratios. (B) Addition of BH4 or PEG-SOD to AHF lung lysates decreased ROS. (C) Treatment of rats with Folic Acid before induction of AHF resulted in protection from edema development. * $P < 0.05$ compared with Sham, •• $p < 0.01$ compared with AHF. $n \geq 3$ /group.

Uncoupled eNOS drives oxidative stress in pressure-dependent AHF

To evaluate the participation of Nox in ROS production during AHF we assessed Nox activity in isolated membrane fraction of lungs collected from Sham and AHF rats. Nox activity was not altered in AHF rats relative to Sham (Figure 2C). To confirm that Nox was not involved in ROS production in this model, we assessed the expression of Nox isoforms 1 and 2 in lung lysates from Sham and AHF rats. No changes were observed in NOX1 and NOX2 content in lung lysates collected from AHF rats when compared with Sham rats (Figure 2D,E).

As L-NAME decreased ROS levels during AHF, we then considered eNOS as a source for oxidative stress in AHF and assessed eNOS uncoupling by Western blot. AHF lungs showed eNOS uncoupling which was not observed in Sham lungs. Pretreatment of AHF rats with L-NAME attenuated eNOS uncoupling during AHF (Figure 3A) thus supporting the findings that L-NAME inhibited ROS production. Collectively, these results suggest that eNOS uncoupling occurs due to excessive eNOS stimulation and the lack of enzyme substrate.

Mechanism for eNOS uncoupling in AHF

We evaluated the hypothesis that depletion of BH4 was contributing to eNOS uncoupling. Exogenous BH4 was added to lung lysates *in vitro* and ROS production was assessed by lucigenin ECL. As shown in Figure 3B, supplementation of BH4 in lung lysates decreased ROS levels in AHF lungs (from 290.5 ± 55.41 in AHF to 125 ± 71.17 in AHF + BH4). The same effect was observed when PEG-SOD was added to the lysates (51.88 ± 24.13 in AHF + PEG SOD), indicating that superoxide anion was the ROS being measured. These results support the hypothesis that uncoupled eNOS was a source of ROS. To further investigate if uncoupled eNOS contributes to PE development *in vivo*, we treated rats with FA, known to recouple eNOS by promoting BH4 recycling, before inducing AHF and assessed lung W/D ratio. In the presence of FA (FA + AHF group), rats that underwent AHF did not develop PE as evidenced by similar lung W/D ratio when compared with sham rats (4.43 ± 0.78 in FA + AHF compared with 5.2 ± 0.59 in Sham), Figure 3C.




Figure 4. Caveolae and NOS isoforms during AHF

AHF did not result in (A) caveolae degradation, (B) changes in eNOS activity, (C) eNOS expression, or (D) iNOS content. $n \geq 4$ /group.

We then investigated if caveolae degradation was part of the mechanism(s) leading to eNOS uncoupling in AHF lungs. Western blot analysis of lung homogenates did not demonstrate caveolae degradation (Figure 4A), ruling out caveola deterioration as a pathway to eNOS uncoupling in pressure-dependent AHF. Additionally, we investigated eNOS activity in lung lysates from sham and AHF rats at the final time point of the experimental model (2 h). No significant changes in eNOS activity were observed in AHF lungs when compared with sham lungs (Figure 4B).

NOS isoforms

In order to determine if differences in NOS isoforms expression occurred during AHF, we assessed eNOS and iNOS expression in lung lysates. No differences in NOS isoform expression were seen between groups (Figures 4C,D), thus ruling out iNOS contribution to increase NO production during AHF.

Discussion

We used a hemodynamic model of HF that was induced by acute afterload mismatch, resulting in increased LVEDP, increased P_{pa} and subsequently, rapid PE [14–16]. This clinically relevant model allowed us to investigate the role of acute pressure-dependent endothelial mechanotransduction on vascular barrier failure as a mechanism of AHF-associated PE. Acute hypertension and the associated afterload mismatch resulted in: (i) a significant decrease in P_{aO_2} that occurred within 30 min, (ii) an increase in lung W/D ratio and, (iii) an increase in histological LIS. Inhibition of NOS during AHF prevented the reduction in P_{aO_2} , maintained normal lung W/D ratio, and reduced histological LIS. The major novel finding of the present study is that in the absence of NOS-dependent enhancement in endothelial permeability, acute increases in pulmonary vascular pressure alone are insufficient to cause significant PE. This represents a major paradigm shift in our understanding of the pathophysiology of hydrostatic PE.

These findings extend previous work on pressure-dependent endothelial mechanotransduction to a clinically relevant model of AHF. Kuebler et al. (2003) [17] were the first to demonstrate that acute increases in P_{pc} resulted in endothelial NO production. Previous work from our laboratory showed that pressure-dependent NOS activation, increased whole lung filtration coefficient (K_f) and endothelial hydraulic conductivity (L_p) [2,18].

Pathophysiology of hypertensive AHF

The increase in pulmonary vascular pressure during NE-infusion resulted from a combination of increased LV afterload and a translocation of blood volume from the peripheral circulation to the lungs [19]. The combined effect

of increased venous return and increased resistance to pulmonary outflow (increase in LVEDP) resulted in a significant increase in P_{pc} [20]. Because this is a model that combines two vasoactive effectors (pressure and NE) we also sought to identify the primary stimulus for endothelial hyperpermeability. In order to do so, we assessed the direct effects of NE on the pulmonary hemodynamics and edema development in the isolated perfused lung preparation. No changes in P_{pa} or in lung W/D ratio were found, corroborating previous findings by Krishnamoorthy et al. [21]. Given that HR did not increase during NE infusion, the increase in hydrostatic pressure appears the major stimulus for activation of NOS. Collectively, these results indicate that pressure, and not NE, is the main trigger for endothelial hyperpermeability

Pressure-dependent AHF is not an inflammatory process

Inflammation is a controversial process in HF progression [22]. Although many animal studies suggest that inflammation is a definitive component of AHF, clinical evidence indicates that is not always the case [23]. Particularly, in conditions of elevated pulmonary vascular pressure, localized inflammation may arise from increased leukocyte margination in the lungs [24,25]. In order to evaluate if leukocyte margination contributed to lung endothelial hyperpermeability in the current model of AHF, we assessed myeloperoxidase activation. Myeloperoxidase activity remained unaltered in AHF lungs and BAL. In the context of inflammation, iNOS could be contributing to hyperpermeability in AHF as it has been previously reported to participate in barrier dysfunction [26]. We validate that iNOS expression was not increased in AHF lungs relative to Sham ruling iNOS as a mediator in this model of AHF. Finally, immunohistochemistry for leukocytes failed to demonstrate an increase in immune cells in the lungs from AHF rats and measurement of IL1- β , TNF- α , and MIP2 indicated that production of these cytokines are not altered during AHF.

L-NAME protects against capillary stress failure

The protective effects of L-NAME on reducing histological indices of lung injury during acutely increased vascular pressure were most notable for the reduction in IAH. This was a novel and an unexpected finding and contributes to the hypothesis that capillary stress failure may be a regulated process [27]. In the dorsal sections of the lung, where hydrostatic pressure would be highest in a supine rat, L-NAME attenuated hypertension-induced IAH. Collectively, these results suggest that pressure-dependent increases in endothelial permeability, including capillary stress failure, occur as a part of NO-mediated process. Studies are underway to further characterize the role of pressure-dependent NO production in capillary stress failure.

Pressure-dependent mechanism(s) for NOS activation

Activation of NOS by mechanical forces is a hallmark of endothelial mechanotransduction [2–4,28,29] and the signaling pathways that lead to NOS activation include glyocalyx-dependent signaling [3,30], neutral sphingomyelinase (NSMase) [31], IK_{ATP} [32], and TRP channels [33,34]. We have recently demonstrated that inhibition of NSMase prevents pressure-dependent increase in the whole lung filtration coefficient [35]. It is likely therefore, that pressure-dependent activation of NSMase and subsequent release of ceramide is part of the pathway leading to NOS activation.

Reactive oxygen species in AHF

Little is known about the mechanisms that lead to pulmonary oxidative stress during AHF. We observed that ROS levels were increased in AHF lungs when compared with Sham lungs and we expected that Nox would play a role in ROS production as they are considered the main source for ROS in the vasculature [36]. Our results however indicated that this was not the case: we found no differences in Nox activity or isoforms expression (Nox1 and Nox2) in AHF lungs when compared with Sham lungs. The fact that we did not see increased leukocyte recruitment in BAL or MPO activity in lung tissue and BAL fluid corroborate our findings that Nox 2 is not involved in ROS production during AHF and support the rationale that macrophages are not a source for ROS during AHF.

The evidence clearly suggested that NOS is activated in this model, so we explored eNOS as a potential source for oxidative stress. ROS levels were decreased with pretreatment of AHF rats with L-NAME suggesting eNOS was uncoupled in AHF rat lungs. Immunoblots of lung lysates confirmed a higher eNOS monomer/dimer ratio in this model supporting the idea that uncoupled eNOS was the key element in pulmonary oxidative stress. Treatment of AHF rats with apocynin mitigated lung edema, confirming a role for ROS in barrier failure. We explored the mechanisms that result in eNOS uncoupling, first by evaluating substrate availability (e.g. BH4) and second by assessing




Figure 5. Schematic diagram

Schematic diagram illustrating the contribution of mechanotransduction to AHF: elevation of systemic blood pressure by sympathetic activation results in increases in pulmonary vascular pressure due to increased lung blood volume and increased LVEDP. The increase in P_{pa} activates mechanotransduction and leads to excessive eNOS activation that results in NO and superoxide anion ($O_2^{\bullet -}$) production in the lungs that promote vascular failure culminating in AHF and associated PE.

caveolae degradation. Supplementation of BH4 to lung lysates reduced ROS formation and confirmed that substrate depletion lead to eNOS uncoupling, thus supporting previous reports in the literature [37].

As eNOS is localized to the caveolae and caveolae dynamics is related to hyperpermeability in lung injury models [38,39], we investigated if caveolae degradation could be part of the mechanism leading to eNOS uncoupling [40]. No changes in caveolin-1 monomer/dimer ratio were found indicating that caveolae degradation is not a determinant for barrier failure in AHF. Our data support previous findings reporting uncoupled eNOS as a mediator of hyperpermeability [7] and as a mechanism in HF [37,41].

Novelty and clinical significance

This is the first report to demonstrate a role for pressure-dependent endothelial hyperpermeability in the pathogenesis of AHF using an intact animal model. The results indicate that pressure-dependent NOS activation contributes to PE development during AHF by increasing NO and ROS production. The observation that L-NAME attenuates PE without altering pulmonary hemodynamics indicate that the increase in pressure *per se* is insufficient to cause lung edema challenging the long-held notion that Starling forces alone can explain rapid and severe PE during AHF. To the contrary, our results suggest that rapid endothelial hyperpermeability, the end result of endothelial mechanotransduction is required in order for the prevailing hydrostatic pressure to cause clinically significant PE (Figure 5). This is a translational study that reveals NOS as a potential target for treating PE during AHF.

Perspectives

- The endothelial cell is an active participant in AHF.
- Endothelial mechanotransduction contributes to pathogenesis of AHF.
- Targetting endothelial hyperpermeability could be a novel approach for treating AHF.

Competing interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

This work was supported by the Department of Anesthesiology at University of Illinois at Chicago .

Author contribution

A.Z.C. wrote the manuscript and contributed to study conception, research design, data acquisition, analysis, and interpretation. A.I., R.R., J.S., and J.S. contributed to data acquisition. M.P. contributed to data acquisition and analysis. B.A.B. participated in writing the manuscript. R.O.D. wrote the manuscript and was responsible for study conception, research design, and supervised data acquisition and interpretation.

Abbreviations

AHF, acute heart failure; Apo, apocynin; BAL, bronchoalveolar lavage; BH4, tetrahydrobiopterin; eNOS, endothelial nitric oxide synthase; EtOH, ethanol; FA, folic acid; HF, heart failure; HR, heart rate; IAH, intra-alveolar hemorrhage; iNOS, inducible nitric oxide synthase; LIS, lung injury score; LVEDP, left ventricular end diastolic pressure; LVESP, left ventricular end systolic pressure; MAP, mean arterial pressure; MIP2, macrophage inflammatory protein 2; mPAP, mean pulmonary artery pressure; MPO, myeloperoxidase; NE, norepinephrine; NOS, nitric oxide synthase; Nox, NADPH oxidase; NSMase, neutral sphingomyelinase; PE, pulmonary edema; P_{pa}, pulmonary artery pressure; P_{pc}, pulmonary capillary pressure; PVC, perivascular cuffing; ROS, reactive oxygen species; TNF- α , tumor necrosis factor- α ; W/D, wet to dry.

References

- 1 Gandhi, S.K., Powers, J.C., Nomeir, A.M., Fowle, K., Kitzman, D.W., Rankin, K.M. et al. (2001) The pathogenesis of acute pulmonary edema associated with hypertension. *N. Engl. J. Med.* **344**, 17–22, <https://doi.org/10.1056/NEJM200101043440103>
- 2 Dull, R.O., Cluff, M., Kingston, J., Hill, D., Chen, H., Hoehne, S. et al. (2012) Lung heparan sulfates modulate K(fc) during increased vascular pressure: evidence for glyocalyx-mediated mechanotransduction. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **302**, L816–L828, <https://doi.org/10.1152/ajplung.00080.2011>
- 3 Dull, R.O., Mecham, I. and McJames, S. (2007) Heparan sulfates mediate pressure-induced increase in lung endothelial hydraulic conductivity via nitric oxide/reactive oxygen species. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **292**, L1452–L1458, <https://doi.org/10.1152/ajplung.00376.2006>
- 4 Kim, M.H., Harris, N.R. and Tarbell, J.M. (2005) Regulation of capillary hydraulic conductivity in response to an acute change in shear. *Am. J. Physiol. Heart Circ. Physiol.* **289**, H2126–H2135, <https://doi.org/10.1152/ajpheart.01270.2004>
- 5 Tilton, R.G., Chang, K.C., LeJeune, W.S., Stephan, C.C., Brock, T.A. and Williamson, J.R. (1999) Role for nitric oxide in the hyperpermeability and hemodynamic changes induced by intravenous VEGF. *Invest. Ophthalmol. Vis. Sci.* **40**, 689–696
- 6 Guequen, A., Carrasco, R., Zamorano, P., Rebolledo, L., Burboa, P., Sarmiento, J. et al. (2016) S-nitrosylation regulates VE-cadherin phosphorylation and internalization in microvascular permeability. *Am. J. Physiol. Heart Circ. Physiol.* **310**, H1039–H1044, <https://doi.org/10.1152/ajpheart.00063.2016>
- 7 Ulker, E., Parker, W.H., Raj, A., Qu, Z.C. and May, J.M. (2016) Ascorbic acid prevents VEGF-induced increases in endothelial barrier permeability. *Mol. Cell. Biochem.* **412**, 73–79, <https://doi.org/10.1007/s11010-015-2609-6>
- 8 Stevenson, L.W. and Perloff, J.K. (1989) The limited reliability of physical signs for estimating hemodynamics in chronic heart failure. *JAMA* **261**, 884–888, <https://doi.org/10.1001/jama.1989.03420060100040>
- 9 Melenovsky, V., Andersen, M.J., Andress, K., Reddy, Y.N. and Bortlaug, B.A. (2015) Lung congestion in chronic heart failure: haemodynamic, clinical, and prognostic implications. *Eur. J. Heart Fail.* **17**, 1161–1171, <https://doi.org/10.1002/ejhf.417>
- 10 Bommakanti, N., Isbatan, A., Bavishi, A., Dharmavaram, G., Chignalia, A.Z. and Dull, R.O. (2017) Hypercapnic acidosis attenuates pressure-dependent increase in whole-lung filtration coefficient (Kf). *Pulm Circ.* **7**, 719–726, <https://doi.org/10.1177/2045893217724414>
- 11 Chignalia, A.Z., Vogel, S.M., Reynolds, A.B., Mehta, D., Dull, R.O., Minshall, R.D. et al. (2015) p120-catenin expressed in alveolar type II cells is essential for the regulation of lung innate immune response. *Am. J. Pathol.* **185**, 1251–1263, <https://doi.org/10.1016/j.ajpath.2015.01.022>
- 12 Chignalia, A.Z., Oliveira, M.A., Debbas, V., Dull, R.O., Laurindo, F.R., Touyz, R.M. et al. (2015) Testosterone induces leucocyte migration by NADPH oxidase-driven ROS- and COX2-dependent mechanisms. *Clin. Sci. (Lond.)* **129**, 39–48, <https://doi.org/10.1042/CS20140548>
- 13 Benson, M.A., Batchelor, H., Chuaiphichai, S., Bailey, J., Zhu, H., Stuehr, D.J. et al. (2013) A pivotal role for tryptophan 447 in enzymatic coupling of human endothelial nitric oxide synthase (eNOS): effects on tetrahydrobiopterin-dependent catalysis and eNOS dimerization. *J. Biol. Chem.* **288**, 29836–29845, <https://doi.org/10.1074/jbc.M113.493023>
- 14 Rassler, B., Reissig, C., Briest, W., Tannapfel, A. and Zimmer, H.G. (2003) Pulmonary edema and pleural effusion in norepinephrine-stimulated rats—hemodynamic or inflammatory effect? *Mol. Cell Biochem.* **250**, 55–63, <https://doi.org/10.1023/A:1024942132705>
- 15 Rassler, B., Reissig, C., Briest, W., Tannapfel, A. and Zimmer, H.G. (2003) Catecholamine-induced pulmonary edema and pleural effusion in rats—alpha- and beta-adrenergic effects. *Respir. Physiol. Neurobiol.* **135**, 25–37, [https://doi.org/10.1016/S1569-9048\(03\)00062-4](https://doi.org/10.1016/S1569-9048(03)00062-4)
- 16 Viau, D.M., Sala-Mercado, J.A., Spranger, M.D., O’Leary, D.S. and Levy, P.D. (2015) The pathophysiology of hypertensive acute heart failure. *Heart* **101**, 1861–1867, <https://doi.org/10.1136/heartjnl-2015-307461>
- 17 Kuebler, W.M., Uhlig, U., Goldmann, T., Schael, G., Kerem, A., Exner, K. et al. (2003) Stretch activates nitric oxide production in pulmonary vascular endothelial cells *in situ*. *Am. J. Respir. Crit. Care Med.* **168**, 1391–1398, <https://doi.org/10.1164/rccm.200304-5620C>
- 18 Tarbell, J.M., Demaio, L. and Zaw, M.M. (1999) Effect of pressure on hydraulic conductivity of endothelial monolayers: role of endothelial cleft shear stress. *J. Appl. Physiol. (1985)* **87**, 261–268, <https://doi.org/10.1152/jappl.1999.87.1.261>

- 19 Jiang, C., Qian, H., Luo, S., Lin, J., Yu, J., Li, Y. et al. (2017) Vasopressors induce passive pulmonary hypertension by blood redistribution from systemic to pulmonary circulation. *Basic Res. Cardiol.* **112**, 21, <https://doi.org/10.1007/s00395-017-0611-8>
- 20 Lindsey, A.W., Banahan, B.F., Cannon, R.H. and Guyton, A.C. (1957) Pulmonary blood volume of the dog and its changes in acute heart failure. *Am. J. Physiol.* **190**, 45–48
- 21 Krishnamoorthy, V., Hiller, D.B., Ripper, R., Lin, B., Vogel, S.M., Feinstein, D.L. et al. (2012) Epinephrine induces rapid deterioration in pulmonary oxygen exchange in intact, anesthetized rats: a flow and pulmonary capillary pressure-dependent phenomenon. *Anesthesiology* **117**, 745–754, <https://doi.org/10.1097/ALN.0b013e31826a7da7>
- 22 Cocco, G., Jerie, P., Amiet, P. and Pandolfi, S. (2017) Inflammation in heart failure: known knowns and unknown unknowns. *Expert Opin. Pharmacother.* **18**, 1225–1233, <https://doi.org/10.1080/14656566.2017.1351948>
- 23 Dick, S.A. and Epelman, S. (2016) Chronic heart failure and inflammation: what do we really know? *Circ. Res.* **119**, 159–176, <https://doi.org/10.1161/CIRCRESAHA.116.308030>
- 24 Ichimura, H., Parthasarathi, K., Issekutz, A.C. and Bhattacharya, J. (2005) Pressure-induced leukocyte margination in lung postcapillary venules. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **289**, L407–L412, <https://doi.org/10.1152/ajplung.00048.2005>
- 25 Ichimura, H., Parthasarathi, K., Quadri, S., Issekutz, A.C. and Bhattacharya, J. (2003) Mechano-oxidative coupling by mitochondria induces proinflammatory responses in lung venular capillaries. *J. Clin. Invest.* **111**, 691–699, <https://doi.org/10.1172/JCI17271>
- 26 Jonkam, C.C., Bansal, K., Traber, D.L., Hamahata, A., Maybauer, M.O., Maybauer, D.M. et al. (2009) Pulmonary vascular permeability changes in an ovine model of methicillin-resistant *Staphylococcus aureus* sepsis. *Crit. Care* **13**, R19, <https://doi.org/10.1186/cc7720>
- 27 Bhattacharya, J. (2003) Pressure-induced capillary stress failure: is it regulated? *Am. J. Physiol. Lung Cell Mol. Physiol.* **284**, L701–2, <https://doi.org/10.1152/ajplung.00425.2002>
- 28 Koo, A., Nordsletten, D., Umeton, R., Yankama, B., Ayyadurai, S., Garcia-Cardena, G. et al. (2013) *In silico* modeling of shear-stress-induced nitric oxide production in endothelial cells through systems biology. *Biophys. J.* **104**, 2295–2306, <https://doi.org/10.1016/j.bpj.2013.03.052>
- 29 Kim, M.H., Harris, N.R. and Tarbell, J.M. (2005) Regulation of hydraulic conductivity in response to sustained changes in pressure. *Am. J. Physiol. Heart Circ. Physiol.* **289**, H2551–H2558, <https://doi.org/10.1152/ajpheart.00602.2005>
- 30 Yen, W., Cai, B., Yang, J., Zhang, L., Zeng, M., Tarbell, J.M. et al. (2015) Endothelial surface glycocalyx can regulate flow-induced nitric oxide production in microvessels *in vivo*. *PLoS ONE* **10**, e0117133, <https://doi.org/10.1371/journal.pone.0117133>
- 31 Czarny, M. and Schnitzer, J.E. (2004) Neutral sphingomyelinase inhibitor scyphostatin prevents and ceramide mimics mechanotransduction in vascular endothelium. *Am. J. Physiol. Heart Circ. Physiol.* **287**, H1344–H1352, <https://doi.org/10.1152/ajpheart.00222.2004>
- 32 Ahn, S.J., Fancher, I.S., Bian, J.T., Zhang, C.X., Schwab, S., Gaffin, R. et al. (2017) Inwardly rectifying K(+) channels are major contributors to flow-induced vasodilatation in resistance arteries. *J. Physiol.* **595**, 2339–2364, <https://doi.org/10.1113/JP273255>
- 33 Dragovich, M.A., Chester, D., Fu, B.M., Wu, C., Xu, Y., Goligorsky, M.S. et al. (2016) Mechanotransduction of the endothelial glycocalyx mediates nitric oxide production through activation of TRP channels. *Am. J. Physiol. Cell Physiol.* **311**, C846–C853, <https://doi.org/10.1152/ajpcell.00288.2015>
- 34 Yin, J., Hoffmann, J., Kaestle, S.M., Neye, N., Wang, L., Baeurle, J. et al. (2008) Negative-feedback loop attenuates hydrostatic lung edema via a cGMP-dependent regulation of transient receptor potential vanilloid 4. *Circ. Res.* **102**, 966–974, <https://doi.org/10.1161/CIRCRESAHA.107.168724>
- 35 Bommakanti, N., Isbatan, A., Bavishi, A., Dharmavaram, G., Chignalia, A. and Dull, R.O. (2017) Hypercapnic acidosis attenuates pressure-dependent increase in whole-lung filtration coefficient (Kf). *Pulm Circ.* **7**, 719–726, <https://doi.org/10.1177/2045893217724414>
- 36 Li, H., Horke, S. and Forstermann, U. (2014) Vascular oxidative stress, nitric oxide and atherosclerosis. *Atherosclerosis* **237**, 208–219, <https://doi.org/10.1016/j.atherosclerosis.2014.09.001>
- 37 Yamamoto, E., Hirata, Y., Tokitsu, T., Kusaka, H., Sakamoto, K., Yamamuro, M. et al. (2015) The pivotal role of eNOS uncoupling in vascular endothelial dysfunction in patients with heart failure with preserved ejection fraction. *Int. J. Cardiol.* **190**, 335–337, <https://doi.org/10.1016/j.ijcard.2015.04.162>
- 38 Maniatis, N.A., Kardara, M., Hecimovich, D., Letsiou, E., Castellon, M., Roussos, C. et al. (2012) Role of caveolin-1 expression in the pathogenesis of pulmonary edema in ventilator-induced lung injury. *Pulm. Circ.* **2**, 452–460, <https://doi.org/10.4103/2045-8932.105033>
- 39 Maniatis, N.A., Chernaya, O., Shinin, V. and Minshall, R.D. (2012) Caveolins and lung function. *Adv. Exp. Med. Biol.* **729**, 157–179, https://doi.org/10.1007/978-1-4614-1222-9_11
- 40 Cassuto, J., Dou, H., Czikora, I., Szabo, A., Patel, V.S., Kamath, V. et al. (2014) Peroxynitrite disrupts endothelial caveolae leading to eNOS uncoupling and diminished flow-mediated dilation in coronary arterioles of diabetic patients. *Diabetes* **63**, 1381–1393, <https://doi.org/10.2337/db13-0577>
- 41 Yamamoto, E., Kataoka, K., Shintaku, H., Yamashita, T., Tokutomi, Y., Dong, Y.F. et al. (2007) Novel mechanism and role of angiotensin II induced vascular endothelial injury in hypertensive diastolic heart failure. *Arterioscler. Thromb. Vasc. Biol.* **27**, 2569–2575, <https://doi.org/10.1161/ATVBAHA.107.153692>