



NOVEMBER 2019
SPECIAL ISSUE **2**

FEDERAL HEALTH REPORTING
JOINT SERVICE BY RKI AND DESTATIS

Journal of Health Monitoring

**30 years after the fall of the Berlin Wall:
Regional health differences in Germany**

Journal of Health Monitoring · 2019 4(S2)
DOI 10.25646/6077
Robert Koch Institute, Berlin

Thomas Lampert, Stephan Müters,
Benjamin Kuntz, Stefan Dahm,
Enno Nowossadeck

Robert Koch Institute, Berlin
Department of Epidemiology and
Health Monitoring

Submitted: 11.06.2019
Accepted: 28.08.2019
Published: 01.11.2019

30 years after the fall of the Berlin Wall: Regional health differences in Germany

Abstract

Following the fall of the Berlin Wall in November 1989, considerable effort was made to bring the living conditions and levels of social participation in the former East German federal states into line with the former West German federal states. As a result, differences in health between the East and the West diminished significantly, in many cases as early as the 1990s, examples being life expectancy and cardiovascular mortality. In regard to health behaviour, the overall tendency has also clearly been one of convergence. Thus, only very small differences can be observed today, for example in the use of tobacco or in the prevalence of obesity. Yet the results also highlight the insufficiency of regarding the remaining differences as a simple comparison between East and West. Instead, the focus should shift towards smaller-scale approaches that take regional differences in living conditions into account.

REGIONAL DIFFERENCES · EAST AND WEST GERMAN FEDERAL STATES · LIFE EXPECTANCY · HEALTH · RISK FACTORS

1. Introduction

After the fall of the Berlin Wall in November 1989, significant efforts were made to bring the living conditions and levels of social participation in the former East into line with the former West. Over the past 30 years, this has been achieved in many areas. Standard wages in East Germany, for example, are now 98% of the level in the West, and pensions in the East are set to achieve parity with pensions in the West by 2024 [1]. Differences in living standards between East and West, i.e. people's access to those consumer goods and durables they consider important, as well as their subjective satisfaction with living conditions, have also reduced significantly [2]. On the other hand, unemployment rates and the proportion of people affected by or at risk from

poverty continue to be significantly higher in the former East German federal states. However, this situation should not be ascribed to German Democratic Republic (GDR) history alone, but rather viewed against a backdrop of current economic, social and demographic developments.

It is clear that migration processes as an immanent element of social change vary significantly in the former East and West German federal states. Immigration from abroad has focused in particular on former West Germany. Moreover, many people have migrated and continue to migrate from East Germany to West Germany, which means that former East Germany has lost a significant number of people emigrating to former West Germany, predominantly young people and young, highly educated women (so-called age-selective migration) [3]. Thus, women leaving these

declining regions either became or might potentially become mothers, which has led to a spiral of emigration and declining birth rates [4]. Although there are indications that an increased number of people are now returning to the East [5], the majority of those returning are men.

The 2009 report, '20 Jahre nach dem Mauerfall: Wie hat sich die Gesundheit in Deutschland entwickelt?' strikingly demonstrates how important the living conditions and levels of social participation in former East and West Germany have been to people's health [6]. Drawing on a broad data basis, this report from the Robert Koch Institute, which was commissioned by the Federal Ministry of Health, takes stock of developments in health within East and West Germany. It covers an array of topics, from disease and mortality to health behaviour and its associated risk factors, and healthcare. As in the follow-up article published five years later [7], the initial report highlights the fact that many of the differences between health outcomes for the East and West have either diminished or no longer exist. Such is the case for average life expectancy, subjective health, as well as numerous chronic diseases and underlying risk factors. Moreover, the report clearly shows the inadequacy of an approach that focuses on the remaining differences between East and West. Instead, a smaller-scale regional analysis should be aimed for, one which takes regional differences in living conditions into account, for example in regard to economic power as well as employment and income opportunities [6-8].

In the present article in the Journal of Health Monitoring, we continue from previous reports, concerning ourselves on the 30th anniversary of the fall of the Berlin Wall with the question of whether the described developments

have continued, and if so, to what extent. Initially, the article examines average life expectancy and mortality (Chapter 3.1), before shifting focus to cardiovascular diseases, cancer and mental disorders (Chapter 3.2 to Chapter 3.4). In addition to subjective health (Chapter 3.5), tobacco use, obesity and physical inactivity (Chapters 3.6 to Chapter 3.8) are considered as important behavioural risk factors. To conclude, selected examples are used to illustrate the importance of a smaller-scale analysis of East-West differences that, where possible, takes the often substantial differences in people's living conditions and opportunities for social participation into account (Chapter 3.9).

2. Methodology

The findings presented in this article draw on a broad data basis. There are, however, very few studies and other data sources available that allow comparisons between the health of people in the former GDR and those in former West Germany. The current analysis, therefore, begins after the German reunification in 1990, when substantial efforts were made to improve the data basis for such comparisons. Between 1991 and 1992, for example, the Robert Koch Institute conducted a health survey in former East Germany that was broadly based on the health survey conducted in former West Germany between 1990 and 1991 [9, 10]. The first national health survey, the German National Health Interview and Examination Survey 1998, was conducted between 1997 and 1999 [11]. The present report also analyses a number of Robert Koch Institute surveys such as the German Health Interview and Examination Survey for Adults (DEGS1, 2008-2011) [10], as well as several waves

As early as the 1990s there was a significant decrease in differences in life expectancy between former East and West Germany.

of the German Health Update (GEDA) that were conducted in 2009, 2010, 2012 and 2014/2015 [12]. Findings on cardiovascular diseases are grounded in data on the lifetime prevalence (whether a person has ever contracted the disease in the course of their life) of significant cardiovascular disease (self-reported medical diagnosis of stroke, heart failure, infarction or any other coronary heart disease). GEDA 2014/2015-EHIS surveyed the prevalence of current depressive symptoms via the Patient Health Questionnaire (PHQ-8) [13].

Analyses of incidence case figures (number of new cases) or incidence rates (number of new cases per 100,000 inhabitants) draws on data from the German Centre for Cancer Registry Data at the Robert Koch Institute. Besides data from the Robert Koch Institute, this report also uses data from the Socio-Economic Panel (SOEP) of the German Institute for Economic Research [14] and the Microcensus of the Federal Statistical Office [15, 16]. Moreover, key insights are provided by further data sources from official statistics such as vital statistics [17], causes of death [18] and hospital diagnoses [19]. These findings are supplemented by results from individual health studies and/or surveys such as the survey of the German Olympics Sports Federation on membership numbers at sports clubs [20].

To compare disease and mortality rates of population groups with diverse age structures, age standardisations were calculated based on the old European standard population [21]. To analyse and evaluate regional health differences on a smaller scale than that of the federal state, these differences should be considered alongside differences in living conditions and opportunities for social participation. The

Robert Koch Institute has developed a corresponding index, which measures regional differences in socioeconomic deprivation (German Index of Socioeconomic Deprivation, GISD) based on a set of indicators that include unemployment, employment rate, net household income, debt ratio, tax revenue and the number of school leavers who leave without a certificate [22]. Multidimensional indices at the regional level are advantageous because they not only depict individual aspects but also the total socioeconomic advantages and disadvantages of specific socio-spatial areas.

3. Results

3.1 Life expectancy

Shortly after the German reunification, average life expectancy at birth was still significantly lower in the former East German states than in the West. Women in former East Germany had an average life expectancy at birth of 77.2 years, while in the West it was 79.5. For men, the difference was even greater, with 69.9 years compared to 73.1. In subsequent years, the East-West differences in average life expectancy at birth rapidly decreased. Parity for women was almost achieved in 2000, and since 2015/2017, the average life expectancy for women in former East Germany has even become slightly higher than in the former West German federal states [23, 24]. While male life expectancy has not achieved parity, the differences between the former East and West Germany have decreased significantly. As with women, this can be traced back to developments in the 1990s. The remaining disparity of slightly over a year less for men in former East Germany has not diminished any further (Table 1).

Table 1
Development of average life expectancy at birth (e_0) and further life expectancy at age 65 (e_{65}) in the former East and West German federal states by sex
Source: Vital statistics [17]

| | Women | | | | Men | | | |
|----------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | 1991/1993 | 1999/2001 | 2007/2009 | 2015/2017 | 1991/1993 | 1999/2001 | 2007/2009 | 2015/2017 |
| e_0 | | | | | | | | |
| Germany | 79.01 | 81.07 | 82.53 | 83.18 | 72.47 | 75.00 | 77.33 | 78.36 |
| East German federal states | 77.18 | 80.53 | 82.37 | 83.22 | 69.86 | 73.69 | 76.27 | 77.25 |
| West German federal states | 79.48 | 80.72 | 82.57 | 83.17 | 73.11 | 75.43 | 77.58 | 78.61 |
| Difference | -2.30 | -0.19 | -0.20 | 0.05 | -3.25 | -1.74 | -1.31 | -1.36 |
| e_{65} | | | | | | | | |
| Germany | 18.02 | 19.44 | 20.52 | 21.00 | 14.34 | 15.79 | 17.22 | 17.80 |
| East German federal states | 16.69 | 18.90 | 20.20 | 21.05 | 13.26 | 15.17 | 16.75 | 17.44 |
| West German federal states | 18.35 | 19.59 | 20.60 | 20.99 | 14.58 | 15.92 | 17.32 | 17.89 |
| Difference | -1.66 | -0.69 | -0.40 | 0.06 | -1.32 | -0.75 | -0.57 | -0.45 |

In the last 30 years, mortality rates for cardiovascular disease have reached a similar level in the former East German federal states to those of the former West German federal states.

A corresponding development can also be observed in regard to life expectancy at age 65. At the beginning of the 1990s, life expectancy at age 65 was one to two years lower for women and men from the former East German federal states compared to women and men from the former West. These rates have now more or less equalised; the life expectancy of women at age 65 is now marginally higher in former East Germany, while 65-year-old men from former East Germany still have a slightly lower life expectancy at age 65 than men from the former West.


3.2 Cardiovascular diseases

The prevalence of cardiovascular diseases in former East Germany is higher than in former West Germany. GEDA data from 2009, 2010 and 2012 indicates that the lifetime

prevalence of significant cardiovascular diseases (self-reported medical diagnosis of heart attack, stroke or heart failure) in the former East German federal states (including Berlin) was 13.0%, and therefore higher than in the former West German federal states (excluding Berlin) (11.7%) [25].

Mortality due to cardiovascular diseases is also elevated in the former East German federal states (Figure 1). While cardiovascular mortality rates had already begun to drop during the 1980s, convergence with levels encountered in the former West German federal states began after 1988 [24]. In 1990, age-standardised cardiovascular mortality in the former GDR was 1.52 times higher for women and 1.44 times higher for men compared to West Germany. This higher mortality then fell, and by 2016 it stood at 1.18 for women and 1.24 for men.

Figure 1
Development of cardiovascular mortality
(age-standardised) in the former East and West
German federal states 1990-2016 by sex
 Source: Cause of death statistics [18]



Eight largely avoidable risk factors (behavioural risk factors such as physical inactivity, drinking risky amounts of alcohol, smoking and low consumption of fruit and vegetables, as well as disease-related risk factors such as obesity, hypertension, diabetes and lipometabolic disorders) are behind over half of cardiovascular-related deaths [26]. The regional distribution of these risk factors in Germany was examined using data from the GEDA studies 2009, 2010 and 2012. Significant variations in prevalence among the federal states were found for all eight risk factors. Former East German federal states, with the exception of Berlin, presented the highest prevalence of physical inactivity, obesity, hypertension and diabetes for both sexes and risky alcohol consumption for men. The three city-states Berlin, Hamburg and Bremen ranked highly for smoking (Chapter 3.6). The prevalence of consuming less than one

portion of fruit, vegetable or juice per day was greatest in Saarland [27].

3.3 Cancer

The German Centre for Cancer Registry Data at the Robert Koch Institute estimated cancer incidence (without non-melanocytic skin cancers) in 2014 to be around 227,000 cases for women and 249,000 for men. Among women in 2014, the highest incidences were recorded for breast cancer (30.5% of new cases), bowel cancer (12.3%) and lung cancer (8.5%). For men, the highest incidences were found for prostate cancer (23.0% of new cases), lung cancer (13.9%) and bowel cancer (13.3%) [28]. Between 1970 and 2014, the annual number of new cancer incidences doubled. Since the risk of developing most types of cancer

increases with age, around half of this increase can be explained by demographic ageing, i.e. a shift in the age structure of the population towards a greater proportion of people in older age groups [29, 30]. Further factors, such as changes to behavioural risk factors, cancer-screening tests and improved diagnostic procedures, have also contributed to this increase [30].

Differences vary in overall cancer incidence rates (number of new cases (incidences) per 100,000 people) between the former East and West German states. Women in former East Germany present lower incidence rates than women in the West, while the opposite applies to men (Figure 2). At the same time, these differences have remained stable over the past years. Cancer mortality is




Figure 2
 Development of cancer incidence and mortality (ICD-10: C00-C96, excluding C44) per 100,000 inhabitants (age-standardised) in the former East and West German federal states 1999-2015 by sex
 Source: German Centre for Cancer Registry Data [28]

ICD-10=International Statistical Classification of Diseases and Related Health Problems, 10th revision

Depression is slightly less common in the former East than in the former West.

dropping, and for women, there are only minimal differences between the former East and West German federal states (Figure 2). For men, the differences are greater, to the detriment of those in former East German states, and these differences have not diminished in the period discussed here.

A different picture emerges for lung cancer. Within the trend towards an overall increase, lung cancer incidence rates and mortality for women are lower in the former East German federal states than in the former West German federal states. For men, the situation is different. Incidence rates and mortality in former East Germany are higher than in the West, while the overall trend is towards a decrease (Figure 3). The differences between women in the former East and West German states have increased, following a sharper increase in mortality in the former West German federal states than in the East. This is mainly owing to the proportion of female smokers, which was higher in the West for many years (Chapter 3.6). Differences in smoking rates between East and West have only converged in recent years [7].

Since the consequences of smoking often only manifest themselves as lung cancer many years later, around half of patients develop the disease after the age of 70, i.e. decades after they took up smoking. For this reason, the present lowered smoking rates will only be reflected in lower incidence and mortality after another two to three decades [31].

Breast cancer incidence rates in the former East and West German federal states continued to rise until 2008/2009, in particular after 2005. Since then, incidence rates have been decreasing (Figure 4) as a result of the

mammography screening programme established between 2005 and 2009 [32]. For women aged 50 to 69, access to the programme is free of charge. The programme helps diagnose tumours at an earlier stage. The earlier diagnosis leads initially to a rise in incidence rates followed by a drop [30]. During the entire observation period, the incidence rate of breast cancer is markedly higher in the former West German federal states than in the former East federal states.

Breast cancer mortality has been falling in both the East and the West German federal states since the 1990s (Figure 4), although mortality rates in the West are still roughly 20% higher than in the East. Higher birth rates, the lower age of first-time mothers, fewer women who opt not to have children and other lifestyle factors are assumed to be protective factors for women in the East German states [30, 33-35].

3.4 Mental health

Mental disorders are widespread and come with a high burden of disease [36]. Analysis of the additional mental health module from the DEGS1 study (DEGS1-MH) by the Robert Koch Institute shows that the general 18- to 79-year-old German population has a 27.7% 12-month prevalence for mental disorders. At the same time, the differences between the East and West German federal states are rather small. Whereas 36.6% of women in the East have been diagnosed with some kind of mental disorder, the figure for women in the West is 33.7%. The prevalence in men in the former East German federal states is 20.4%, compared to 23.0% in the former West German federal states [7, 37].





Figure 3
 Development of lung cancer incidence and mortality (ICD-10: C33-C34) per 100,000 inhabitants (age-standardised) in the former East and West German federal states 1999-2015 by sex
 Source: German Centre for Cancer Registry Data [28]

ICD-10=International Statistical Classification of Diseases and Related Health Problems, 10th revision

Figure 4
Development of breast cancer incidence and mortality (ICD-10: C50) per 100,000 inhabitants (age standardised) in the former East and West German federal states 1999-2015
 Source: German Centre for Cancer Registry Data [28]



ICD-10=International Statistical Classification of Diseases and Related Health Problems, 10th revision

According to GEDA 2014/2015-EHIS, the prevalence of current symptoms of depression in Germany is 10.1% [13]. It is slightly lower in East Germany (9.1%) compared to West Germany (10.3%). 10.8% of women from the East German federal states present with symptoms of depression, whereas for women in West Germany the figure is 11.7%. The prevalence in men is 7.3% and 8.9%, respectively. The 12-month prevalence of a self-reported medical diagnosis of depression in Germany, which GEDA 2014/15-EHIS also collected data on, is 8.1% [38], with East Germany at 6.6%, lower than West Germany at 8.3%. 8.8% of women from the East German federal states suffer from a medically diagnosed depression, compared to 9.8% in the West German federal states. Prevalence in men is 4.2% compared to 6.7%. Differentiated analyses

by federal state show that both depressive symptomatology and self-reported medical diagnoses of depression are more frequent in the West German federal states. One exception is Bavaria, where the prevalence is lower. Prevalence is higher in city-states than in non-city federal states [13].

Similar distributive patterns between the East and West German federal states are also reflected in health insurance billing data on depressive disorders [13, 38, 39]. Thus, when considering different age and sex structures, depression is more frequently diagnosed in the West German federal states than in the East [39]. Besides true differences in disease prevalence, possible explanations under discussion include disparities between health-care systems. Diagnoses should, for example, be made

by psychotherapists and psychiatrists. Particularly in the case of psychotherapists, there is a far greater concentration in the West German federal states than in the East, which means there are also better chances for diagnosis [39-41].

Hospital diagnosis statistics can provide further information on the treatment of mental disorders. A significant increase in the number of inpatient cases of mental and behavioural disorders was registered in the period from 2000 to 2011, a trend that has, however, not continued. This development can be seen in both the female and male population, and as much in the East as in the West German federal states. Since the turn of the century,

the differences between the East and West German federal states have not further increased.

It is a different story when considering specific groups of mental disorders. This is the case, for example, for the affective disorders (International Statistical Classification of Diseases and Related Health Problems, 10th revision, ICD-10: F30-F39), to which depression belongs, alongside neurotic disorders, stress disorders and somatoform disorders (ICD-10: F40-F48). Whereas inpatient case numbers in the diagnostic group of affective disorders are higher in the former West German federal states than in the East (Figure 5), these are higher in East Germany for neurotic disorders, stress-related disorders and somato-




Figure 5
Development of inpatient case numbers for affective disorders (ICD-10: F30-F39) per 100,000 inhabitants (age-standardised) in the former East and West German federal states 2000-2017 by sex

Source: Hospital diagnoses statistics [19]

ICD-10=International Statistical Classification of Diseases and Related Health Problems, 10th revision

form disorders (Figure 6). These differences between East and West German federal states became greater between 2000 and 2012 but have not diverged any further since.

3.5 Subjective health

Findings on the development of subjective health are also made possible by examining Socio-Economic Panel data. There are only slight differences between the East and West German federal states in regard to the proportion of 18- to 79-year-old women and men who describe their overall health as either good or very good. These differences were slight even at the beginning of the 1990s. During the past

20 years, the proportion of women and men who estimated their health as good or very good has not changed substantially. This applies to both the East and the West German federal states, as is confirmed by a differentiated analysis for the 18- to 39 and 40- to 54-year old groups. The 55- to 79-year-old group is the only one that shows an increase in the proportion of women and men with either good or very good subjective health. For the West German federal states, this trend is much stronger than for the East German federal states. As a result, for men in this age group in particular, the differences between the East and the West are now more pronounced than at the beginning of the 1990s (Figure 7).





Figure 6
Development of inpatient case numbers for neurotic, stress-related and somatoform disorders (ICD-10: F40-F48) per 100,000 inhabitants (age-standardised) in the former East and West German federal states 2000-2017 by sex

Source: Hospital diagnoses statistics [19]

ICD-10=International Statistical Classification of Diseases and Related Health Problems, 10th revision

Figure 7
Prevalence of 'good' or 'very good' self-reported health in the former East and West German federal states by sex
Source: SOEP 1994-2015 [14], own calculations




No differences in the prevalence of smoking or obesity persist between the former East and West.

3.6 Smoking

The development of tobacco use in the 25- to 69-year-old population in East and West Germany between 1990 and 2015 can be traced using data from the health surveys of the Robert Koch Institute (Figure 8). Shortly after the reunification, the proportion of male smokers was at 40.6% in the East German federal states, slightly higher than the 39.2% recorded for the West German federal states [42]. Male tobacco use has been in decline since the 2000s, while the slight difference between East and West has remained more or less stable. Overall, fewer

women smoke than men. At the beginning of the 1990s, 20.5% of women in the East were smokers, compared to a significantly higher 28.3% in the West German federal states. By the end of the 1990s, these figures for East and West German women had converged. Since the early 2000s, the percentage of female smokers in the East German federal states has remained relatively constant. In contrast, the number of female smokers in the West German federal states initially rose, before declining sharply after 2003. Most recently, the proportion of female smokers has even been slightly lower in the West than in the East (Figure 8).

Figure 8
Prevalence trends in smoking in the 25- to 69-year-old population in the former East and West German federal states by sex
Source: Health surveys of the Robert Koch Institute, updated according to Lampert 2010 [42]



The percentage of smokers over the age of 15 in individual federal states can be calculated using 2017 Microcensus data (Table 2) [15]. The highest rates are found among

| Federal state | Women % | Men % | Total % |
|-------------------------------|-------------|-------------|-------------|
| Baden-Wuerttemberg | 17.4 | 25.1 | 21.2 |
| Bavaria | 16.6 | 24.6 | 20.5 |
| Berlin | 21.3 | 29.9 | 25.5 |
| Brandenburg | 20.2 | 29.0 | 24.5 |
| Bremen | 24.2 | 30.9 | 27.4 |
| Hamburg | 18.9 | 27.8 | 23.2 |
| Hesse | 17.5 | 24.8 | 21.1 |
| Mecklenburg-Western Pomerania | 22.1 | 33.4 | 27.7 |
| Lower Saxony | 19.2 | 26.7 | 22.9 |
| North Rhine-Westphalia | 19.4 | 26.0 | 22.6 |
| Rhineland-Palatinate | 18.6 | 24.9 | 21.7 |
| Saarland | 17.7 | 23.5 | 20.6 |
| Saxony | 16.6 | 26.5 | 21.4 |
| Saxony-Anhalt | 20.0 | 29.8 | 24.8 |
| Schleswig-Holstein | 20.0 | 27.1 | 23.5 |
| Thuringia | 21.7 | 30.8 | 26.2 |
| Germany | 18.6 | 26.4 | 22.4 |

Table 2
Percentage of smokers aged over 15 by sex and federal state
Source: Microcensus 2017 [15]


men from Mecklenburg-Western Pomerania (33.4%). Rates in Thuringia, Saxony-Anhalt, and the city-states Berlin and Bremen are also relatively high (around 30%). Men from Saarland, Hesse, the Rhineland Palatinate and Bavaria smoke less often (<25%). Rates for women are highest in Bremen (24.2%), followed by Mecklenburg-Western Pomerania (22.1%) and Thuringia (21.7%). In a comparison of federal states, smoking rates for women from Bavaria and Saxony are the lowest (16.6%).

3.7 Obesity

During the initial years after German reunification, measured data indicated a significantly higher prevalence of obesity for both genders in the former East compared to the former West. Since then, prevalences in Germany have continued to rise [43], while the differences between East and West German federal states have gradually decreased [9]. The most recent available data (DEGS1) suggests that, in regard to the distribution of obesity, there are no longer any

Figure 9
Prevalence trends in obesity in 25- to 69-year-olds in the former East and West German federal states by sex

Source: Health surveys of the Robert Koch Institute, modified according to Finger et al. 2016 [9]



significant differences between the East and West for 25- to 69-year-old men. Small differences persist for women, with a higher prevalence in the East (Figure 9) [7], although it

| Federal state | Women % | Men % | Total % |
|-------------------------------|-------------|-------------|-------------|
| Baden-Wuerttemberg | 13.2 | 16.4 | 14.9 |
| Bavaria | 12.9 | 17.3 | 15.2 |
| Berlin | 12.2 | 13.8 | 13.0 |
| Brandenburg | 17.1 | 19.5 | 18.3 |
| Bremen | 14.6 | 18.1 | 17.8 |
| Hamburg | 10.0 | 12.8 | 12.2 |
| Hesse | 14.0 | 18.2 | 16.1 |
| Mecklenburg-Western Pomerania | 20.0 | 23.5 | 21.8 |
| Lower Saxony | 14.5 | 18.5 | 16.6 |
| North Rhine-Westphalia | 14.6 | 18.4 | 16.5 |
| Rhineland-Palatinate | 15.3 | 19.9 | 17.7 |
| Saarland | 11.8 | 19.6 | 16.3 |
| Saxony | 17.8 | 18.2 | 18.0 |
| Saxony-Anhalt | 19.8 | 21.7 | 20.8 |
| Schleswig-Holstein | 12.8 | 17.9 | 15.4 |
| Thuringia | 19.5 | 21.4 | 20.4 |
| Germany | 13.2 | 16.4 | 16.3 |

Table 3
Prevalence of obesity by sex and federal state
Source: Microcensus 2017 [16]

should be noted that the DEGS1 data was collected between 2008 and 2011. The next data set will be provided by the Health and Nutrition Survey in Germany (gern survey), which the Robert Koch Institute will jointly conduct with the Max Rubner Institute between 2020 and 2022.

Self-reported data on height and weight from the Microcensus can be used to analyse regional spread patterns. When doing so, it is important to note that prevalences based on self-reported data are usually far lower than those calculated using measurement data, since a certain proportion of participants tends to underestimate their weight and overestimate their height [44]. According to 2017 Microcensus data, the highest prevalence (totalling over 20%) are found in Mecklenburg-Western Pomerania, Saxony-Anhalt and Thuringia. Obesity levels are lowest in the city-states of Hamburg and Berlin (Table 3). An interpretation of these findings needs to take into account the – on average – younger age of the population in the city-states [16].


3.8 Physical inactivity

Data from the Robert Koch Institute health examination survey shows that, shortly after German reunification, levels of physical activity were lower in the East German federal states than in the West (Figure 10). In the 25- to 69-year-old age group, 49.2% of men in the East and 41.0% in the West stated that they did not do any sport. The corresponding figures for women were 58.7% in the East and 50.3% in the West German federal states. Towards the end of the 1990s, the differences between East and West had increased for men, whereas for women there were no longer any detectable differences. Most recently, levels of physical

Figure 10

Prevalence trends in physical inactivity in 25- to 69-year-olds in the former East and West German federal states by sex

Source: Health surveys of the Robert Koch Institute, modified according to Finger et al. 2016 [9]



People in the former East are less likely to belong to a sports club than people in the former West.

Physical inactivity have decreased significantly in both the East and the West German federal states. There are currently no differences between East and West either for women or for men.

The statistics of the German Olympics Sports Confederation are also informative [20]. They indicate that in 2018, a significantly smaller percentage of people in the East were

members of a sports club than in the West German federal states. This is apparent across all age groups regardless of gender (Figure 11). The highest levels of sports club membership can be found in the Saarland (37%), in Rhineland Palatinate (35%), as well as in Bavaria, Baden-Wuerttemberg and Hesse (all around 34%). In comparison, only 13% to 18% of the East German population are members of a sports club.

3.9 Differences on a smaller scale

In many cases, making distinctions between East and West or individual federal states is inadequate when analysing regional health differences. It should be noted that there can be substantial regional differences even within federal states in regard to a number of health indicators. For this reason, regional analyses on a smaller scale than the federal state level should be conducted whenever possible, for example in Germany's 96 planning regions (Raumordnungsregionen), 401 administrative rural and urban districts, and 4,504 collective municipalities.

Figure 11

Sports club members in relation to respective total populations in the former East and West German federal states

(as of 31 December 2017) by sex and age
Source: German Olympics Sports Confederation 2018 [20]




Figure 12 shows the differences between average life expectancy at birth and socioeconomic deprivation in rural and urban districts. In Saxony-Anhalt, Thuringia and Brandenburg, there is a relatively large number of rural and urban districts with a low average life expectancy, whereas

in Bavaria and Baden-Wuerttemberg, there are many rural and urban districts where life expectancy is high. It is clear that socioeconomic deprivation is highest in rural and urban districts with a low average life expectancy, and vice versa. The difference in average life expectancy between




Figure 12
Regional differences (rural and urban districts)
in relation to socioeconomic deprivation and
average life expectancy at birth

Source: Kroll et al. 2017 [22]; INKAR,
own calculations

1st quintile=lowest deprivation to 5th quintile=highest deprivation

rural and urban districts with the lowest and highest levels of socioeconomic deprivation is around five years.

For many other health indicators, too, a differentiated regional analysis demonstrably makes sense and can lead to different or additional findings. This can be seen in the case of sick days taken due to mental health issues. Data from company health insurance funds (Betriebskrankenkassen, BKK) members has shown that, in many rural and urban districts in Mecklenburg-Western Pomerania, Brandenburg and Saxony-Anhalt, the number of sick days due to mental health issues was far higher than the 2017 national average of 2.8 days. The same applies to some of the former West German federal states such as North Rhine-Westphalia, Lower Saxony and Saarland. On the other hand, the greatest deviation towards fewer sick days was found in Bavaria and Baden-Wuerttemberg, once again in rural and urban districts with a tendency toward lower levels of socioeconomic deprivation. However, there are also rural and urban districts in northern North Rhine-Westphalia, on Schleswig Holstein's North Sea coast and in parts of Saxony that register significantly fewer sick days than the national average. Overall, these results demonstrate that a smaller-scale analysis is helpful, for example for regional planning in regard to healthcare structures, prevention, and the promotion of health [45].

4. Discussion

The results presented here show that shortly after German reunification, the health situation in the former East and West German federal states differed in many aspects. 30 years of common development has, in many cases, led to

the convergence and in some cases to the equalisation of these differences. Remarkably, this equalisation was already taking place during the first 10 to 15 years after the reunification and can be ascribed to favourable developments that occurred faster in former East Germany than in former West Germany. Examples of such favourable developments include an increase in average life expectancy and a decrease in cardiovascular mortality. In some cases, equalisation actually occurred through negative trends, i.e. where the situation in the East was originally better but then became like the West. An example in this regard is the rise of tobacco use among women in former East Germany during the 1990s.

Data on the health situation of children and adolescents born after the reunification also indicate that the differences between the East and the West are no longer significant. The present article does not discuss these data; however, they were included in a report published by the Robert Koch Institute to commemorate the 20th anniversary of German reunification [8, 46]. According to this report, there have been no differences in the prevalence of allergic diseases and obesity amongst children and adolescents from the former East and West German federal states since the 2000s. There has also been a significant reduction in differences between East and West in regard to vaccination rates, which were initially higher in the former East German federal states, and in regard to participation rates for early detection examinations by children (U-Untersuchungen), which were more widely accepted in former West Germany. Based on these results, it can be assumed that differences between the East and the West will play an ever-diminishing role in future birth cohorts [8, 46].

The question of why the differences between East and West have diminished and why some regional disparities in health persist is not easy to answer. A significant contribution to the convergence of health opportunities in the East and West federal states was likely made by the increase in general living standards in former East Germany during the unification process, which was made possible by substantial investment and development aid. Contributing factors of note are economic restructuring, improvements in labour market conditions, improvements to urban development, housing conditions and environmental factors, and the rapid integration of the East into social security and healthcare systems.

Alongside these general processes, other developments likely played a role in diminishing (or maintaining) East-West differences. These can vary greatly depending on which health indicator is analysed. Any analysis of the continuing differences in sports club membership rates must allow for the fact that the role of club sport in mass sports was less important in the GDR than in former West Germany. The reason for this is that GDR sports clubs were more closely tied to promoting competitive sports. The higher child vaccination rates in the former East were probably also related to the compulsory GDR vaccination programme for many infectious diseases. Conversely, lower participation in early detection examinations in the former East can, to a certain degree, be explained by the fact that these programmes were first introduced in East Germany after the reunification.

In regard to health differences between federal states, and at the rural and urban district level, differences in living conditions and opportunities for social participation

are particularly important. The present article has referred to these using the German Index of Socioeconomic Deprivation, and by referring to individual indicators such as the at-risk-of-poverty rate, unemployment rate or gross domestic product. To an extent, the correlations can be traced to the socio-spatial segregation of people in different life situations. However, the results of multi-level analyses indicate that independent of individual socioeconomic status, the quality of the social environment has an effect on people's health prospects [22, 47].

The analysis of smaller-scale differences points to an increasing need, in the face of ever-diminishing differences between East and West, to apply regional criteria other than East and West or federal states. These could include smaller-scale approaches such as the NUTS2 regions, regional planning regions (Raumordnungsregionen) or district and municipality-based analytical approaches. As well as concepts orientated towards existing administrative boundaries, it could be useful to apply analysis that differentiates between urban and rural spaces, or, even more strongly differentiated, categorisations of district type or GPS-based health research.

To conclude, it is important to note that the development of both parts of Germany after the fall of the Berlin Wall did not take place in isolation but rather within the context of Europe. The breakdown of the socialist system in Europe caused similar upheavals in Poland, the Czech Republic, Hungary and other East European states. However, in comparison to East Germany, these countries did not become part of an economically stronger country and could not, therefore, count on associated investment and development aid. Against this historical backdrop,

the extent and pace of the positive developments in health and life expectancy in former East Germany stand out [48, 49].

Corresponding author

PD Dr Thomas Lampert
Robert Koch Institute
Department of Epidemiology and Health Monitoring
General-Pape-Str. 62–66
12101 Berlin, Germany
E-mail: LampertT@rki.de

Please cite this publication as

Lampert T, Müters S, Kuntz B, Dahm S, Nowossadeck E (2019)
30 years after the fall of the Berlin Wall:
Regional health differences in Germany.
Journal of Health Monitoring 4(S2): 2-23.
DOI 10.25646/6077

The German version of the article is available at:
www.rki.de/journalhealthmonitoring

Data protection and ethics

GEDA and DEGS₁ are subject to strict compliance with the data protection regulations of the Federal Data Protection Act and have been approved by the Federal Commissioner for Data Protection and Freedom of Information in Germany. Charité – Universitätsmedizin Berlin's ethics committee assessed the ethics of the DEGS₁ study and provided its approval (No.EA2/047/08). Participation in the studies GEDA and DEGS₁ was voluntary. The participants were informed about the aims and contents of the studies and about data protection. Informed consent was obtained in writing.

The data collected by the Socio-Economic Panel are subject to the provisions set out in the Federal Data Protection Act (BDSG). This means that the data collected through interviews are anonymised as part of the SOEP and individual respondents can no longer be identified.

Funding

GEDA and DEGS₁ were funded by the Federal Ministry of Health and the Robert Koch Institute.

Conflicts of interest

The authors declared no conflicts of interest.

References

1. Bundesministerium für Wirtschaft und Energie (Ed) (2019) Jahresbericht der Bundesregierung zum Stand der Deutschen Einheit 2018. BMWi, Berlin.
<https://www.bmwi.de/Redaktion/DE/Publikationen/Neue-Lae-nder/jahresbericht-zum-stand-der-deutschen-einheit-2018.html> (As at 18.04.2019)
2. Statistisches Bundesamt, Wissenschaftszentrum Berlin für Sozialforschung (Eds) (2018) Datenreport 2018. Ein Sozialbericht für die Bundesrepublik Deutschland. Bundeszentrale für politische Bildung, Bonn
3. Leibert T, Wiest K (2010) Ist die ländliche Peripherie männlich? In: Nationalatlas aktuell 4 (11.2010) 11 [24.11.2010]. Leibniz-Institut für Länderkunde, Leipzig
4. Nowossadeck E, Fiebig J (2017) „Älter, weniger, bunter“: der demografische Wandel in Deutschland. UMID: Umwelt und Mensch – Informationsdienst 2:62-72
5. Fuchs M, Nadler R, Roth D et al. (2017) Rückwanderung von Erwerbspersonen – aktuelle Deutschlandzahlen im regionalen Vergleich. In: Nationalatlas aktuell 11 (05.2017) 4 [22.05.2017]. Leibniz-Institut für Länderkunde, Leipzig.
6. Robert Koch Institute (Ed) (2009) 20 Jahre nach dem Fall der Mauer: Wie hat sich die Gesundheit in Deutschland entwickelt? Beiträge zur Gesundheitsberichterstattung des Bundes. RKI, Berlin.
<https://edoc.rki.de/handle/176904/3221> (As at 18.04.2019)

7. Prütz F, Rommel A, Kroll LE et al. (2014) 25 Years after the Fall of the Berlin Wall: Regional Differences in Health. GBE kompakt 5(3). RKI, Berlin.
<https://edoc.rki.de/handle/176904/3133> (As at 18.04.2019)
8. Lampert T, Ziese T, Kurth BM (2010) 20 Jahre Deutsche Einheit: Gesundheitliche Entwicklungen und Trends in den neuen und alten Bundesländern. Aus Politik und Zeitgeschichte 45:32-39
9. Finger JD, Busch MA, Du Y et al. (2016) Zeitliche Trends kardiometaboler Risikofaktoren bei Erwachsenen. Dtsch Arztebl International 113(42):712-719
10. Scheidt-Nave C, Kamtsiuris P, Gosswald A et al. (2012) German health interview and examination survey for adults (DEGS) - design, objectives and implementation of the first data collection wave. BMC Public Health 12:730
11. Bellach BM, Knopf H, Thefeld W (1998) Der Bundes-Gesundheitssurvey 1997/98. Gesundheitswesen 60 Suppl 2:S59-68
12. Lange C, Jentsch F, Allen J et al. (2015) Data Resource Profile: German Health Update (GEDA)—the health interview survey for adults in Germany. Int J Epidemiol 44(2):442-450
13. Bretschneider J, Kuhnert R, Hapke U (2017) Depressive symptoms among adults in Germany. Journal of Health Monitoring 2(3):77-83.
<https://edoc.rki.de/handle/176904/2826> (As at 18.04.2019)
14. Goebel J, Grabka MM, Liebig S et al. (2019) The German Socio-Economic Panel (SOEP). Journal of Economics and Statistics 239(2):345-360
15. Statistisches Bundesamt (Ed) (2018) Mikrozensus - Fragen zur Gesundheit - Rauchgewohnheiten der Bevölkerung 2017. Statistisches Bundesamt, Wiesbaden
16. Statistisches Bundesamt (Ed) (2018) Mikrozensus - Fragen zur Gesundheit - Körpermaße der Bevölkerung 2017. Statistisches Bundesamt, Wiesbaden
17. Statistisches Bundesamt (2019) Statistik der natürlichen Bevölkerungsbewegung.
www.gbe-bund.de (As at 18.04.2019)
18. Statistisches Bundesamt (2019) Todesursachenstatistik, Sterbefälle, Sterbeziffern (je 100.000 Einwohner, altersstandardisiert).
<http://www.gbe-bund.de> (As at 28.04.2019)
19. Statistisches Bundesamt (2019) Krankenhausstatistik – Diagnosedaten der Patienten und Patientinnen in Krankenhäusern.
<http://www.gbe-bund.de> (As at 24.5.2019)
20. Deutscher Olympischer Sportbund (Ed) (2018) Bestandserhebung 2018. DOSB, Frankfurt am Main
21. Robert Koch Institute (Ed) (2015) Gesundheit in Deutschland. Gesundheitsberichterstattung des Bundes. Gemeinsam getragen von RKI und Destatis. RKI, Berlin.
<https://edoc.rki.de/handle/176904/3248> (As at 24.5.2019)
22. Kroll LE, Schumann M, Hoebel J et al. (2017) Regional health differences – developing a socioeconomic deprivation index for Germany. Journal of Health Monitoring 2(2):98-114.
<https://edoc.rki.de/handle/176904/2657.2> (As at 18.04.2019)
23. Nowossadeck E, Von der Lippe E, Lampert T (2019) Developments in life expectancy in Germany. Current trends. Journal of Health Monitoring 4(1):38-45.
<https://edoc.rki.de/handle/176904/5914> (As at 05.04.2019)
24. Grigoriev P, Pechholdova M (2017) Health Convergence Between East and West Germany as Reflected in Long-Term Cause-Specific Mortality Trends: To What Extent was it Due to Reunification? Eur J Popul 33(5):701-731
25. Dornquast C, Kroll LE, Neuhauser HK et al. (2016) Regionale Unterschiede in der Prävalenz kardiovaskulärer Erkrankungen. Dtsch Arztebl International 113(42):704-711
26. World Health Organization (Ed) (2009) Global health risks. Mortality and burden of disease attributable to selected major risks. WHO, Geneva
27. Diederichs C, Neuhauser H, Kroll L et al. (2017) Regionale Unterschiede in der Prävalenz von kardiovaskulären Risikofaktoren bei Männern und Frauen in Deutschland. Bundesgesundheitsbl 60(2):151-162
28. Zentrum für Krebsregisterdaten im Robert Koch-Institut (2019) Datenbankabfrage zu aktuellen Krebsstatistiken für Deutschland.
www.krebsdaten.de/abfrage (As at 27.05.2019)
29. Nowossadeck E, Barnes B, Kowalski C (2018) Demografische Alterung und Krebs. Onkologie heute 3:48-50
30. Barnes B, Kraywinkel K, Nowossadeck E et al. (2016) Bericht zum Krebsgeschehen in Deutschland 2016. RKI, Berlin
31. Wienecke A, Kraywinkel K (2018) Tabakassoziierte Krebserkrankungen in Deutschland – Entwicklung der Inzidenz und Mortalität seit 1995. Bundesgesundheitsbl 61(11):1399-1406
32. Starker A, Buttman-Schweiger N, Krause L et al. (2018) Krebsfrüherkennungsuntersuchungen in Deutschland: Angebot und Inanspruchnahme. Bundesgesundheitsbl 61(12):1491-1499

33. Troisi R, Bjørge T, Gissler M et al. (2018) The role of pregnancy, perinatal factors and hormones in maternal cancer risk: a review of the evidence. *J Intern Med* 283(5):430-445
34. Nelson HD, Zakher B, Cantor A et al. (2012) Risk factors for breast cancer for women aged 40 to 49 years: a systematic review and meta-analysis. *Ann Intern Med* 156(9):635-648
35. Albrektsen G, Heuch I, Hansen S et al. (2005) Breast cancer risk by age at birth, time since birth and time intervals between births: exploring interaction effects. *Br J Cancer* 92(1):167-175
36. Plass D, Vos T, Hornberg C et al. (2014) Entwicklung der Krankheitslast in Deutschland. *Dtsch Arztebl International* 111(38):629-638
37. Jacobi F, Höfler M, Siegert J et al. (2014) Twelve-month prevalence, comorbidity and correlates of mental disorders in Germany: the Mental Health Module of the German Health Interview and Examination Survey for Adults (DEGS1-MH). *Int J Methods Psychiatr Res* 23(3):304-319
38. Thom J, Kuhnert R, Born S et al. (2017) 12-month prevalence of self-reported medical diagnoses of depression in Germany. *Journal of Health Monitoring* 2(3):68-76. <https://edoc.rki.de/handle/176904/2825> (As at 18.04.2019)
39. Bramesfeld A, Grobe T, Schwartz FW (2010) Prevalence of depression diagnosis and prescription of antidepressants in East and West Germany: an analysis of health insurance data. *Soc Psychiatry Psychiatr Epidemiol* 45(3):329-335
40. Melchior H, Schulz H, Härter M (2014) Stellenwert regionaler Variationen in der Prävalenz und Behandlung depressiver Erkrankungen und Implikationen für die Versorgungsforschung. *Bundesgesundheitsbl* 57(2):224-233
41. Rommel A, Bretschneider J, Kroll LE et al. (2017) The utilization of psychiatric and psychotherapeutic services in Germany – individual determinants and regional differences. *Journal of Health Monitoring* 2(4):3-22. <https://edoc.rki.de/handle/176904/2911.2> (As at 27.05.2019)
42. Lampert T (2010) Tabakkonsum, sportliche Inaktivität und Adipositas. Entwicklungen in den neuen und alten Bundesländern seit der Wiedervereinigung Deutschlands. In: Brähler E, Mohr I (Eds) 20 Jahre deutsche Einheit. Facetten einer geteilten Wirklichkeit. Psychosozial-Verlag, Gießen, P. 275-290
43. Mensink GBM, Schienkiewitz A, Haftenberger M et al. (2013) Übergewicht und Adipositas in Deutschland. Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsbl* 56(5-6):786-794
44. Mensink GBM, Lampert T, Bergmann E (2005) Übergewicht und Adipositas in Deutschland 1984-2003. *Bundesgesundheitsbl* 48(12):1348-1356
45. Knieps F, Pfaff H (Eds) (2018) BKK Gesundheitsreport 2018. Arbeit und Gesundheit Generation 50+. Zahlen, Daten, Fakten mit Gastbeiträgen aus Wissenschaft, Politik und Praxis. MWV Medizinisch Wissenschaftliche Verlagsgesellschaft, Berlin
46. Lampert T (2010) 20 Years of German Unity: Are there still differences between east and west as regards the health of children and adolescents? *GBE kompakt* 1(4). RKI, Berlin. <https://edoc.rki.de/handle/176904/3101> (As at 18.04.2019)
47. Dragano N, Bobak M, Wege N et al. (2007) Neighbourhood socioeconomic status and cardiovascular risk factors: a multilevel analysis of nine cities in the Czech Republic and Germany. *BMC Public Health* 7:255
48. Nolte E, Shkolnikov V, McKee M (2000) Changing mortality patterns in East and West Germany and Poland. I: Long term trends (1960–1997). *J Epidemiol Community Health* 54(12): 890-898
49. Nolte E, Scholz R, Shkolnikov V et al. (2002) The contribution of medical care to changing life expectancy in Germany and Poland. *Soc Sci Med* 55(11):1905-1921

Imprint

Journal of Health Monitoring

Publisher

Robert Koch Institute
Nordufer 20
13353 Berlin, Germany

Editors

Susanne Bartig, Johanna Gutsche, Dr Birte Hintzpeter,
Dr Franziska Prütz, Dr Martina Rabenberg, Dr Alexander Rommel,
Dr Livia Ryl, Dr Anke-Christine Saß, Stefanie Seeling,
Martin Thißen, Dr Thomas Ziese
Robert Koch Institute
Department of Epidemiology and Health Monitoring
Unit: Health Reporting
General-Pape-Str. 62–66
12101 Berlin, Germany
Phone: +49 (0)30-18 754-3400
E-mail: healthmonitoring@rki.de
www.rki.de/journalhealthmonitoring-en

Typesetting

Gisela Dugnus, Kerstin Möllerke, Alexander Krönke

Translation

Simon Phillips/Tim Jack

ISSN 2511-2708

Note

External contributions do not necessarily reflect the opinions of the Robert Koch Institute.



This work is licensed under a
Creative Commons Attribution 4.0
International License.



The Robert Koch Institute is a Federal Institute within
the portfolio of the German Federal Ministry of Health