

Molecular epidemiology of *Mycobacterium tuberculosis* isolates in Iran using spoligotyping

R. Ramazanzadeh^{1,2}, P. Shakib³, S. Rouhi^{4,5}, B. Mohammadi^{1,2}, P. Mohajeri⁶ and S. Borji⁷

1) Cellular and Molecular Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, 2) Department of Microbiology, Faculty of Medicine, Kurdistan University of Medical Sciences, Sanandaj, 3) Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, 4) Children Growth Research Center, Research Institute for Prevention of Non-Communicable Diseases, Qazvin University of Medical Sciences, Qazvin, 5) Clinical Research Development Unit, Kosar Hospital, Qazvin University of Medical Sciences, Qazvin, 6) Nosocomial Infections Research Center, Kermanshah University of Medical Sciences, and 7) Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran

Abstract

Spoligotyping can help assess the transmission of *Mycobacterium tuberculosis* strains. We aimed to study the genotyping of *M. tuberculosis* isolated from patients with tuberculosis from the west of Iran by spoligotyping. Forty-seven *M. tuberculosis* isolates were collected from the west of Iran. All samples were cultured on Löwenstein-Jensen medium incubated at 37°C for 8 weeks. Bacterial isolates were identified as *M. tuberculosis* using standard biochemical tests. Drug resistance patterns of *M. tuberculosis* to rifampicin and isoniazid were determined, and multidrug-resistant (MDR) strains were isolated. After DNA extraction, spoligotyping was performed. We found new spoligotypes 4162 and 4163, which correlated with atypical lineage. Atypical and unknown lineages also had correlations with the MDR tuberculosis rate (4%). The most prevalent spoligointernational types were orphan (34%), 2669 (23.4%) and 127 (14.8%) types. The most prevalent clades were Ural-2 (NEW-1) (25.53%) and atypical (23.40%) lineages. The predominant clade was Ural-2 (NEW-1) and an atypical lineage restricted to Iran. The rate of MDR was low. Knowledge of the circulating isolates in the west of Iran will help implement control programmes, so knowledge of the dynamic transmission of local isolates is crucial.

© 2020 Published by Elsevier Ltd.

Keywords: Epidemiology, Iran, Molecular, *Mycobacterium tuberculosis*, Spoligotyping

Original Submission: 13 May 2020; **Revised Submission:** 20 September 2020; **Accepted:** 21 September 2020

Article published online: 28 September 2020

Corresponding author: R. Ramazanzadeh, Cellular and Molecular Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran.
E-mail: atrop_t51@yahoo.com

Introduction

Tuberculosis (TB) is a classical infectious disease that threatens the world's public health; it is caused by bacteria of the *Mycobacterium tuberculosis* complex [1,2]. The World Health Organization reports 8.6 to 9.0 million occurrences of TB, which can cause 1.3 to 1.5 million deaths annually [3,4]. Risk factors include HIV infection, drug addiction, smoking and air pollution

as well as drug-resistant *M. tuberculosis* isolates that have high rates of TB incidence [5]. Combination therapy provided to treat TB includes the antibiotics rifampicin (RIF), isoniazid (INH), pyrazinamide and ethambutol or streptomycin [6]. Multidrug-resistant (MDR) TB strains are TB isolates with resistance to at least RIF and INH (the two most important anti-TB drugs available) with or without resistance to other drugs [7]. The mutations in individual drug target genes leads to the emergence of MDR-TB strains. Currently the rate of MDR-TB strains is increasing in different regions of the world. Further, MDR-TB is a major public health problem for TB control programmes in healthcare facilities [8].

In the 21st century, molecular genetics has greatly assisted the rapid diagnosis, fast initiation of therapy and evaluation of the source of TB [9]. Genotyping of TB strains includes mycobacterial interspersed repetitive unit-variable number of

tandem repeat (MIRU-VNTR), spacer oligonucleotide typing (spoligotyping) and IS6110 restriction fragment length polymorphism (IS6110-RFLP). Such genotyping helps TB control by permitting identification of unsuspected outbreaks and laboratory cross-contamination, and it helps distinguish exogenous reinfection from endogenous reactivation, thereby permitting understanding of the genetic diversity of strains [10–13]. Spoligotyping is a major genotyping assay which is PCR based and analyses the polymorphism in the spacer sequences, which are present in the direct repeat (DR) region of strains [14,15]. Spoligotyping is an easy, rapid, economical and highly reproducible assay for typing TB strains [13].

The aim of our study was to assess the molecular epidemiology of *M. tuberculosis* genotypes by spoligotype patterns from patients with TB from the west of Iran.

Materials and methods

Collection of *M. tuberculosis* isolates

In this cross-sectional study, 47 *M. tuberculosis* isolates were collected from Ardabil, Hamadan, Qazvin, Tabriz and Kurdistan provinces (west Iran). Demographic information and clinical information regarding age, sex, TB history and residential address were recorded.

All samples were cultured on Löwenstein-Jensen medium, incubated at 37°C and kept for 8 weeks [16]. Bacterial isolates were identified as *M. tuberculosis* using catalase, production of niacin and nitrate reduction.

Drug susceptibility testing and determining MDR-TB strains

Drug resistance patterns of *M. tuberculosis* to RIF and INH, as determined by culture on Löwenstein-Jensen medium and use of the proportions method, were analysed. The concentrations and proportions were 40 mg/L and 1.0% for RIF, and 0.2 mg/L and 1.0% for INH [17,18]. *M. tuberculosis* H37Rv strain was used for quality control testing in drug susceptibility assays. TB isolates that exhibited resistance to at least RIF and INH were defined as MDR strains [6,18].

DNA extraction

DNA extraction of *M. tuberculosis* isolates grown in Löwenstein-Jensen medium was performed in accordance with the manufacturer's instructions (Qiagen, Germantown, MD, USA). All the genomic DNA preparation materials were stored at –20°C until use. The quality and quantity of the extracted DNA were assessed by a NanoDrop machine (Thermo Fisher Scientific, Waltham, MA, USA). Distilled water was used as the negative control.

Spoligotyping

Spoligotyping was performed using amplification of direct repeat regions in the genome of *M. tuberculosis* complex by DRa and DRb primers and an available spoligotyping kit according to the manufacturer's protocol. Then the amplified products were hybridized on a membrane precoated with spacer oligos that characterized the spacer region of the identified sequence. After incubation with streptavidin–peroxidase and enhanced chemiluminescence detection, the presence of spacers was imaged on X-ray films as black squares [19]. *M. tuberculosis* H37Rv was used as the positive control (spoligotyping kit for detecting tuberculosis; Mapmygenome, Hyderabad, Andhra Pradesh, India).

Results

Spoligotype patterns of *M. tuberculosis* isolates are shown in Table 1. Orphans indicate spoligotyping patterns that are unique to the analysis and are not found in the SITVIT database (<http://www.pasteur-guadeloupe.fr:8081/SITVIT2/>). The frequency of the new spoligointernational types (SITs) 4162 and 4163 were 8.51% and 4.26% respectively. Orphan spoligotype patterns (34%) by definition were found only once and have not yet been described.

Table 2 lists the prevalence of lineages in this study. Ural-2 (NEW-1) had the highest rate (25.53%) of the lineage and of other clades, including atypical, unknown and Ural-2. The S lineage had the lowest rate.

Differences in the prevalence of drug resistance to RIF and INH are shown in Table 3. Atypical lineages had the highest proportion of members resistant to RIF and INH. Atypical and unknown lineages also had the highest correlation with the MDR-TB rate (4%).

Discussion

Molecular epidemiology may be applied for several aims, including comparing isolates worldwide [20]. TB control programmes depend on information about molecular epidemiology to obtain insight into TB's distribution and diversity in a specific area. In considering the SITVIT database, our study is the first to analyse spoligotyping in the west of Iran. Our study found the new spoligotypes 4162 and 4163 in this region. Orphan spoligotypes represent patterns that have been found only once and that have not been found in the past. They may indicate recent and/or sporadic TB infection in the studied area. For a profile to be considered a SIT, this profile must have been found at least twice in the present study. New SITs are created after a match

TABLE 1. Spoligotype patterns of *Mycobacterium tuberculosis* isolates

SIT	<i>M. tuberculosis</i> (%)	Total (%)
26	2.13	2.13
127	14.89	14.89
252	2.13	2.13
284	2.13	2.13
602	6.38	6.38
1965	2.13	2.13
2669	23.40	23.40
4162	8.51	8.51
4163	4.26	4.26
Orphan	34.04	34.04
Total	100	100

SIT, spoligointernational type.

with another orphan in the database, or if two or more strains belonging to the new pattern within this study were found. These SITs are from the Ural-2 (NEW-1) lineage. On the basis of the SITVIT database, this lineage is more prevalent in Iran. The most prevalent SITs were the orphan (34%), 2669 (23.4%) and 127 (14.8%) types. The orphan types mostly belong to the Ural-2 (NEW-1) lineage reported from Iran. The 2669 spoligotype belongs to atypical lineages mostly isolated from Saudi Arabia. The 127 spoligotype belongs to the Ural-2 lineage, which has mostly been isolated from Iran, Iraq and Saudi Arabia. Finding the most potent circulating clades is important in controlling TB and comparing them to other isolates.

Iran has a variety of geographical conditions; therefore, distribution of *M. tuberculosis* strains differs from state to state. Prevalence of lineages Ural-2 (NEW-1), atypical, Ural-2 and unknown were 25.53%, 23.4%, 14.89% and 14.89% respectively. As indicated earlier, the predominant lineage is reported from Iran. Therefore, most isolates are unique to our region. We are located on the border with Iraq, and because of ethnic group relationships, there is close communication between peoples; it was therefore obvious that the second prevalent lineage would be Ural-2.

The resistance rates to RIF and INH were as follows: atypical lineage, 8%; unknown lineage, 4%; and Ural-2, 4%. The rate of frequency of MDR strains was 4%, with strains belonging to atypical, Ural-2 and unknown lineages. The reported prevalence of drug resistance in Colombia is 2.38% [21]. The MDR-TB rate

TABLE 2. Prevalence of lineage in *Mycobacterium tuberculosis* isolates

Lineage	<i>M. tuberculosis</i> (%)	Total (%)
AFRI	6.38	6.38
Atypical	23.40	23.40
CAS1-Delhi	4.26	4.26
LAM9	2.13	2.13
S	2.13	2.13
TI	6.38	6.38
Unknown	14.89	14.89
Ural-2	14.89	14.89
Ural-2 (NEW-1)	25.53	25.53
Total	100	100

TABLE 3. Prevalence of lineage and multidrug resistance

Lineage	Rifampin (%)		Isoniazid (%)		MDR (%)	
	R	S	R	S	No	Yes
Atypical	8	36	8	36	40	4
CAS1-Delhi	0	8	0	8	8	0
LAM9	0	4	0	4	4	0
TI	0	4	0	4	4	0
Unknown	4	8	4	8	8	4
Ural-2	4	24	4	24	24	4
Total	16	84	16	84	88	12

MDR, multidrug resistant; R, resistant; S, susceptible.

is 11% in India [22]. In a previous study, the MDR rate was higher than in our study [23,24]. Therefore, TB control was successful in the west of Iran. Most resistant isolates are from the Beijing family. Interestingly, the Beijing genotype was not found in this study; most Beijing strains are prominent in populations of Afghani origin [24–27]. Because we do not have any Beijing types in this region, the rates of resistance are lower than in other parts of the world.

Conclusions

To our knowledge, ours is the first study to introduce the new spoligotypes 4162 and 4163 from the west of Iran. The predominant lineages were restricted to Iran. Also, the rate of MDR was low. A good control programme needs to understand the dynamic transmission of local isolates; therefore, we performed this study to identify circulating isolates in the west of Iran.

Conflicts of interest

None declared.

Acknowledgements

The DNA samples were collected as part of work on a microbiology thesis. Supported in part by the Kurdistan University of Medical Science and the Kermanshah University of Medical Sciences. The authors thank the health deputy of the Kurdistan University of Medical Science for preparation of the miniblitter. The authors also thank N. Rastogi and colleagues for SITVIT database registration.

References

[1] Walls T, Shingadia D. The epidemiology of tuberculosis in Europe. *Arch Dis Child* 2007;92:726–9.

- [2] Ravansalar H, Tadayon K, Mosavari N, Derakhshan M, Ghazvini K. Genetic diversity of *Mycobacterium tuberculosis* complex isolated from patients in the northeast of Iran by MIRU-VNTR and spoligotyping. *Jundishapur J Microbiol* 2017;10:e39568.
- [3] World Health Organization (WHO). Global tuberculosis report. Geneva: WHO; 2013. Available at: <https://apps.who.int/iris/handle/10665/91355>.
- [4] Zumla A, George A, Sharma V, Herbert RHN, Oxley A, Oliver M. The WHO 2014 global tuberculosis report—further to go. *Lancet Glob Health* 2015;3:e10–2.
- [5] Lönnroth K, Jaramillo E, Williams BG, Dye C, Raviglione M. Drivers of tuberculosis epidemics: the role of risk factors and social determinants. *Soc Sci Med* 2009;68:2240–6.
- [6] Ormerod LP. Multidrug-resistant tuberculosis (MDR-TB): epidemiology, prevention and treatment. *Br Med Bull* 2005;73:17–24.
- [7] World Health Organization (WHO). Anti-tuberculosis drug resistance in the world: third global report. Geneva: WHO; 2004. Available at: <https://apps.who.int/iris/handle/10665/43103>.
- [8] Ahmad S, Mokaddas E. Recent advances in the diagnosis and treatment of multidrug-resistant tuberculosis. *Respir Med* 2009;103:1777–90.
- [9] Jagielski T, Van Ingen J, Rastogi N, Dziadek J, Mazur PK, Bielecki J. Current methods in the molecular typing of *Mycobacterium tuberculosis* and other mycobacteria. *Biomed Res Int* 2014;2014:645802.
- [10] Reisig F, Kremer K, Amthor B, van Soolingen D, Haas WH. Fast ligation-mediated PCR, a fast and reliable method for IS6110-based typing of *Mycobacterium tuberculosis* complex. *J Clin Microbiol* 2005;43:5622–7.
- [11] Supply P, Lesjean S, Savine E, Kremer K, Van Soolingen D, Lochter C. Automated high-throughput genotyping for study of global epidemiology of *Mycobacterium tuberculosis* based on mycobacterial interspersed repetitive units. *J Clin Microbiol* 2001;39:3563–71.
- [12] Hermans P, Van Soolingen D, Bik E, De Haas P, Dale J, Van Embden J. Insertion element IS987 from *Mycobacterium bovis* BCG is located in a hot-spot integration region for insertion elements in *Mycobacterium tuberculosis* complex strains. *Infect Immun* 1991;59:2695–705.
- [13] Niemann S, Richter E, Rüsche-Gerdes S. Stability of *Mycobacterium tuberculosis* IS6110 restriction fragment length polymorphism patterns and spoligotypes determined by analyzing serial isolates from patients with drug-resistant tuberculosis. *J Clin Microbiol* 1999;37:409–12.
- [14] Brudey K, Driscoll JR, Rigouts L, Prodinger WM, Gori A, Al-Hajj SA, et al. *Mycobacterium tuberculosis* complex genetic diversity: mining the fourth international spoligotyping database (SpolDB4) for classification, population genetics and epidemiology. *BMC Microbiol* 2006;6:23.
- [15] Pang Y, Song Y, Xia H, Zhou Y, Zhao B, Zhao Y. Risk factors and clinical phenotypes of Beijing genotype strains in tuberculosis patients in China. *BMC Infect Dis* 2012;12:354.
- [16] Githui W, Meme H, Juma E, Kinyanjui P, Karimi F, Chakaya J, et al. Isolation of multidrug-resistant tuberculosis strains in patients from private and public health care facilities in Nairobi, Kenya. *Int J Tuberc Lung Dis* 2004;8:837–41.
- [17] Rieder HL, Chonde TM, Myking H, Urbanczik R, Laszlo A, Kim S, et al. The public health service national tuberculosis reference laboratory and the national laboratory network; minimum requirements, role and operation in a low-income country. Paris: International Union Against Tuberculosis and Lung Disease (IUATLD); 1998.
- [18] Farnia P, Varahram M, Doraghi M. Drug susceptibility test against *Mycobacterium tuberculosis*. Tehran: mycobacteriology research center (MRC). National Research Institute of Tuberculosis and Lung Disease (NRITLD); 2007.
- [19] Kamerbeek J, Schouls L, Kolk A, Van Agterveld M, Van Soolingen D, Kuijper S, et al. Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J Clin Microbiol* 1997;35:907–14.
- [20] Ramazanzadeh R, McNERNEY R. Variable number of tandem repeats (VNTR) and its application in bacterial epidemiology. *Pak J Biol Sci* 2007;10:2612–21.
- [21] Puerto G, Erazo L, Wintaco M, Castro C, Ribon W, Guerrero MI. *Mycobacterium tuberculosis* genotypes determined by spoligotyping to be circulating in Colombia between 1999 and 2012 and their possible associations with transmission and susceptibility to first-line drugs. *PLoS One* 2015;10:e0124308.
- [22] Gupta R, Amrathal RS, Prakash R, Jain S, Tiwari PK. Spoligotyping, phenotypic and genotypic characterization of *katG*, *rpoB* gene of *M. tuberculosis* isolates from Sahariya tribe of Madhya Pradesh India. *J Infect Publ Health* 2019;12:395–402.
- [23] Mohammadi B, Mohajeri P, Rouhi S, Ramazanzadeh R. The relationship between *embB306* and *embB406* mutations and ethambutol resistance in *Mycobacterium tuberculosis* isolated from patients in west of Iran. *Med J Islam Republic Iran* 2018;32:117.
- [24] Ramazanzadeh R, Farnia P, Amirmozafari N, Ghazi F, Ghadertotonchi Z, Kamran J, et al. Comparison between molecular epidemiology, geographical regions and drug resistance in *Mycobacterium tuberculosis* strains isolated from Iranian and Afghan patients. *Chemotherapy* 2006;52:316–20.
- [25] Ramazanzadeh R, Roshani D, Shakib P, Rouhi S. Prevalence and occurrence rate of *Mycobacterium tuberculosis* Haarlem family multi-drug resistant in the worldwide population: a systematic review and meta-analysis. *J Res Med Sci* 2015;20:78–88.
- [26] Ramazanzadeh R, Sayhemiri K. Prevalence of Beijing family in *Mycobacterium tuberculosis* in world population: systematic review and meta-analysis. *Int J Mycobacteriol* 2014;3:41–5.
- [27] Ramazanzadeh R, Farnia P, Amirmozafari N. Characterization of *Mycobacterium tuberculosis* complex isolated from Iranian and Afghani patients by spoligotyping method. *Braz J Microbiol* 2009;40:314–20.