

RESEARCH

Open Access



Circular RNA circ-DONSON facilitates gastric cancer growth and invasion via NURF complex dependent activation of transcription factor SOX4

Lixian Ding^{1,2†}, Yuying Zhao^{3†}, Shuwei Dang^{1,2†}, Yue Wang⁴, Xinglong Li^{1,2}, Xiaotong Yu^{1,2}, Zhongsheng Li^{1,2}, Jiufeng Wei^{1,2}, Ming Liu^{1,2} and Guodong Li^{1,2*} 

Abstract

Background: Circular RNAs (circRNAs) are a novel type of noncoding RNAs and play important roles in tumorigenesis, including gastric cancer (GC). However, the functions of most circRNAs remain poorly understood. In our study, we aimed to investigate the functions of a new circRNA circ-DONSON in GC progression.

Methods: The expression of circ-DONSON in gastric cancer tissues and adjacent normal tissues was analyzed by bioinformatics method, qRT-PCR, Northern blotting and in situ hybridization (ISH). The effects of circ-DONSON on GC cell proliferation, apoptosis, migration and invasion were measured by using CCK8, colony formation, EdU, immunofluorescence (IF), FACS and Transwell assays. qRT-PCR and Western blotting were utilized to validate how circ-DONSON regulates SOX4 expression. CHIP, DNA fluorescence in situ hybridization (DNA-FISH) and DNA accessibility assays were used to investigate how circ-DONSON regulates SOX4 transcription. The interaction between circ-DONSON and NURF complex was evaluated by mass spectrum, RNA immunoprecipitation (RIP), pulldown and EMSA assays. Xenograft mouse model was used to analyze the effect of circ-DONSON on GC growth in vivo.

Results: Elevated expression of circ-DONSON was observed in GC tissues and positively associated with advanced TNM stage and unfavorable prognosis. Silencing of circ-DONSON significantly suppressed the proliferation, migration and invasion of GC cells while promoting apoptosis. circ-DONSON was localized in the nucleus, recruited the NURF complex to SOX4 promoter and initiated its transcription. Silencing of the NURF complex subunit SNF2L, BPTF or RBBP4 similarly attenuated GC cell growth and increased apoptosis. circ-DONSON knockdown inhibited GC growth in vivo.

Conclusion: circ-DONSON promotes GC progression through recruiting the NURF complex to initiate SOX4 expression.

Keywords: circ-DONSON, Gastric cancer, NURF, SOX4, Progression

* Correspondence: liguodong@ems.hrbmu.edu.cn

[†]Lixian Ding, Yuying Zhao and Shuwei Dang contributed equally to this work.

¹Department of General Surgery, the Fourth Affiliated Hospital of Harbin Medical University, No. 37 Yiyuan Street, Nangang District, Harbin 150001, People's Republic of China

²Bio-Bank of Department of General Surgery, the Fourth Affiliated Hospital of Harbin Medical University, Harbin 150001, People's Republic of China

Full list of author information is available at the end of the article



Background

Gastric cancer (GC) is one of the most common cancers in gastrointestinal malignancies and remains the third leading cause of cancer-associated deaths around the world [1]. The development of GC is induced by several factors such as smoking [2], constant salty food intake [3] and genetic mutation [4]. As the developing of novel strategies for GC diagnosis and treatment, its incidence and mortality rates have been steadily decreased in the recent years [5]. However, the five-year overall survival rate of GC patients remains still lower than 29% because of tumor invasiveness and recurrence [6]. Hence, it is urgently required to investigate the molecular mechanisms of GC progression and develop more effective therapeutic methods.

Circular RNAs (circRNAs) are a recently discovered member of the noncoding RNA family and characterized by a covalently closed continuous loop and resistance to RNase R digestion [7, 8]. CircRNAs are highly stable and exist in various cell types. Increasing RNA-sequencing analyses have shown that circRNAs are highly expressed in tumor tissues, including GC tissues [9]. Emerging studies have indicated that circRNAs participate in tumorigenesis through regulating various biological processes, including proliferation, survival, invasion and differentiation [10–12]. For example, circPVT1 contributes to non-small cell lung cancer (NSCLC) cell growth and migration by inhibiting miR-125b to activate E2F2 expression [13]. Circ_0005230 is overexpressed in breast cancer and promotes tumor cell division and invasiveness through miR-618/CBX8 signaling [14]. CircMMP9 is upregulated by EIF4A3 in glioblastoma and contributes to tumor development by sponging miR-124 [15]. In GC, circ-SFMBT2 was found to initiate tumor growth [10]. These evidences demonstrate essential functions of circRNAs in cancer. However, there are still a large number of circRNAs in GC, whose roles are ill studied.

circ-DONSON (circbase ID: hsa_circ_0004339), derived from back-splicing of DONSON mRNA (from exon 3 to exon 8), is located on chromosome 21q22.11 and has 948 nucleotides in length. To our knowledge, the function of circ-DONSON has not been researched. In this study, we found that circ-DONSON was highly expressed in GC tissues and positively correlated with TNM stage and poor prognosis. circ-DONSON silencing suppressed GC cell proliferation, migration and invasion while promoting apoptosis in vitro. Moreover, circ-DONSON knockdown suppressed GC growth in vivo. Mechanistically, we demonstrated that circ-DONSON recruits the NURF complex to SOX4 promoter and initiates its transcription. In summary, circ-DONSON works as an oncogene in GC and might be a potential therapeutic target.

Methods

Human samples

A total of 142 GC tissues and paired adjacent normal tissues were obtained from the Fourth Affiliated Hospital of Harbin Medical University. The tissues were immediately stored in liquid nitrogen after surgery. These patients did not receive radiotherapy or chemotherapy prior to collection. This study was approved by the Ethics Committee of the Fourth Affiliated Hospital of Harbin Medical University. The informed consent was achieved from every patient.

Cell culture

GC cell lines (BGC-823, AGS, MGC-803, MKN74, HGC-27 and SGC-7901) and normal human gastric epithelial cell line GES-1 were bought from Shanghai Institutes for Biological Sciences, China. These cells were cultured with Dulbecco's Modified Eagle medium (DMEM) medium supplemented with 10% fetal bovine serum (FBS), 100 U/ml of penicillin, and 100 µg/ml of streptomycin (HyClone).

Lentivirus production and cell transfection

The lentivirus-containing short hairpin RNA (shRNA) targeting circ-DONSON, BPTF, SNF2L or RBBP4 was purchased from GenePharma (Shanghai, China). Both shRNAs were transfected into the GC cell lines using Lipofectamine 3000 (Invitrogen) according to the manufacturer's instructions. At 48 h post-transfection, the cells were selected with puromycin (2 µg/mL) for 2 weeks to construct stable cell lines. The transfection efficiency was verified by qRT-PCR.

Quantitative real-time PCR (qRT-PCR)

Total RNAs were isolated using TRIzol and inversely transcribed into cDNA using M-MLV and the SYBR Green Master Mix kit (Takara, Otsu, Japan). qPCR was completed as described previously [16].

Cell proliferation

Cell proliferation was determined using the Cell Counting Kit-8 (CCK8) assay according to a previous study [9].

Colony formation assay

BGC-823 and AGS cells were planted into the 6-well plates and incubated for 12 days at 37 °C. Then colonies were fixed and stained with 0.1% crystal violet. The colony numbers were counted finally.

5-Ethynyl-20-deoxyuridine (EdU) incorporation assay

The EdU assay was performed using a Cell-Light EdU DNA Cell Proliferation Kit (RiboBio, Shanghai, PR, China) according to the manufacturer's instructions.

Transwell assay

Transwell assays were used to detect cell migration and invasion and conducted as previously described [5].

RNA fluorescence in situ hybridization (RNA-FISH)

RNA-FISH was performed according to a previous study [5]. Briefly, the probes targeting the back-splicing site of circ-DONSON were used for this assay. The probe of circ-DONSON was marked with DIG-UTP (Roche, 11,209,256,910) for RNA labeling. Cells were first fixed with 4% paraformaldehyde for 10 min and then permeabilized in PBS with 0.5% Triton X-100 for 5 min. Next, the cells were hybridized with labeled FISH probe at 37 °C overnight. Afterwards, the cells were washed with 4× sodium citrate buffer containing 0.1% Tween-20 for 5 min and then washed with 1× SSC for 5 min. Finally, cells were stained with 4,6-diamidino-2-phenylindole for 10 min. The images were acquired using a fluorescence microscopy (Leica, SP8 laser confocal microscopy).

Animal assay

Four-week-old BABL/c female nude mice were purchased and maintained under specific pathogen-free conditions. For the in vivo tumor formation assay, BGC-823 cells (circ-DONSON silencing or control) were injected into the right flank of BABL/c nude mice (4 mice for each group). The tumor volume was measured every 5 days. 30 days after injection, the animals were sacrificed and the xenograft tumors were dissected and weighed. All animal studies were approved by the Animal Care and Use Committee of the Fourth Affiliated Hospital of Harbin Medical University.

Statistical analysis

Statistical analyses were carried out by using SPSS 20.0 (IBM, SPSS, Chicago, IL, USA) and GraphPad Prism. Student's t-test or ANOVA was used to assess the statistical significance for comparisons of two groups. The Pearson's correlation coefficient analysis was used to analyze the correlations. Overall survival (OS) and disease-free survival curves were analyzed using the Kaplan-Meier method and log-rank test. $P < 0.05$ was considered statistically significant.

Results


circ-DONSON is upregulated in GC tissues and positively correlated with poor prognosis

To identify important circRNAs involved in GC progression, we first analyzed overexpressed circRNAs in GC tissues compared to adjacent normal tissues according to online dataset (GSE83521). As shown, the circRNA circ-DONSON (probe ID: ASCRP003426) is the most upregulated circRNAs among all candidates (Fig. 1a, b).

circ-DONSON derives from back-splicing of DONSON mRNA (Additional file 1: Figure S1a). And its sequence is presented in Additional file 1: Table S1. Then qRT-PCR analysis was conducted to validate its expression. circ-DONSON expression was significantly upregulated in 142 GC tissues compared to their adjacent normal tissues (Fig. 1c). We then measured circ-DONSON expression through Northern blotting in 5 pairs of GC tissues and adjacent normal tissues. The results indicated that circ-DONSON levels were higher in tumor tissues (Fig. 1d), which was further confirmed by in situ hybridization (ISH) (Fig. 1e). Similarly, the expression of circ-DONSON was increased in GC cell lines compared to GES-1 cell line (Fig. 1f). Then we analyzed the correlation between circ-DONSON expression and clinical features. We found that circ-DONSON expression was positively correlated with TNM stage and lymphoid metastasis (Fig. 1g, h). Furthermore, higher expression of circ-DONSON in GC patients was correlated with lower overall survival rate and disease-free survival rate (Fig. 1i, j), indicating circ-DONSON might be a prognostic marker.

circ-DONSON silencing suppresses GC cell proliferation, migration and invasion, and induces apoptosis

We next investigated the roles of circ-DONSON in GC cell phenotypes. Because circ-DONSON level was relatively higher in BGC-823 and AGS cells (Fig. 1f), we performed following experiments using these two cells. Using two independent shRNAs targeting circ-DONSON, we effectively decreased its expression in BGC-823 and AGS cells (Fig. 2a). Through CCK8 assay, we found that circ-DONSON silencing significantly inhibited the proliferation of BGC-823 and AGS cells (Fig. 2b). EdU assay also illustrated that circ-DONSON knockdown reduced the incorporation of EdU (Fig. 2c). To further confirm it, we conducted colony formation assay, and found that the colony numbers were decreased after circ-DONSON silencing (Fig. 2d). Importantly, the Ki67 positive BGC-823 and AGS cells were reduced after circ-DONSON knockdown (Fig. 2e), supporting that circ-DONSON knockdown inhibited GC cell proliferation. Then we analyzed apoptosis, migration and invasion. Results demonstrated that circ-DONSON silencing induced more apoptosis while impairing the abilities of migration and invasion (Fig. 2f-h). Interestingly, the western blotting assay showed that loss of circ-DONSON increased the epithelial marker E-cadherin expression and decreased the mesenchymal marker N-cadherin expression in BGC-823 and AGS cells (Fig. 2i). To further rule out the effect of shRNA off-target, we overexpressed circ-DONSON by transfection with pcDNA3-circ-DONSON vector in BGC-823 and AGS cells (Additional file 1: Figure S1b). CCK8 and colony formation assays indicated that circ-DONSON overexpression promoted the proliferation




of BGC-823 and AGS cells (Additional file 1: Figure S1c, d). Furthermore, ectopic expression of circ-DONSON enhanced the migration and invasion of BGC-823 and AGS cells (Additional file 1: Figure S1e, f). Thus, these results demonstrated that circ-DONSON promotes GC growth and metastasis in vitro.

circ-DONSON regulates GC cell malignant behaviors through activating SOX4

Previous studies have reported that key signaling pathways, such as NOTCH signaling, Wnt signaling, NF- κ B signaling and Hedgehog signaling, play crucial roles in tumorigenesis [17, 18]. Thus, we speculated whether circ-DONSON regulates these signaling pathways. We measured the effects of circ-DONSON knockdown on the activation of these pathways by evaluating the expression of their target genes (HES6, HEY1, HES1, NRARP for NOTCH signaling; MYC, TIAM, KIAA,

CCND1, CCND2, SOX4, FN14, TCF1 for Wnt signaling; VEGF, BCL2L1, HIF1A, BIRC5, MMP2, TWIST1 for NF- κ B signaling; GLI1, PATCHED, GLI3 for Hedgehog signaling). Interestingly, we found that circ-DONSON silencing only significantly led to decreased expression of SOX4 (Fig. 3a, b), implying circ-DONSON might directly regulate SOX4 expression. Using circ-DONSON specific probes, we found that circ-DONSON could enrich on SOX4 promoter region ($-600 \sim -400$ bp from the transcription start site) (Fig. 3c). Pulldown assay also indicated that biotin labeled SOX4 promoter DNA precipitated circ-DONSON in cell lysates (Fig. 3d). Consistently, FISH assay showed that SOX4 promoter was co-localized with circ-DONSON in BGC-823 and AGS cells (Fig. 3e), suggesting circ-DONSON might regulate SOX4 transcription. As shown, we really found that circ-DONSON silencing suppressed the enrichment of transcriptional active marker H3K27ac on SOX4 promoter



(Fig. 3f). Moreover, after circ-DONSON knockdown, the SOX4 promoter was more resistant to DNaseI digestion (Fig. 3g), indicating circ-DONSON regulates SOX4 promoter accessibility. To further demonstrate that whether circ-DONSON-mediated SOX4 transcription promotes GC progression, we overexpressed SOX4. Through CCK8, EdU and colony formation assays, we found that SOX4 overexpression enhanced the proliferation of circ-DONSON-depleted BGC-823 and AGS cells (Fig. 3h-j). Moreover, restoration of SOX4 also reversed the effects of circ-DONSON silencing on apoptosis, migration and invasion (Fig. 3k-m). Taken together, circ-DONSON activates SOX4 transcription to promote GC progression.

circ-DONSON associates with the NURF complex by directly interacting with SNF2L subunit

Through FISH assay (Fig. 3e), we found circ-DONSON was mainly localized in the nucleus. We further validated it through qRT-PCR (Fig. 4a). To further investigate the molecular mechanism, we searched the potential protein that interacts with circ-DONSON. We performed pull-down assay and silver staining. We then chose the differential band in circ-DONSON lane for mass spectrum identification. SNF2L, an essential subunit of the NURF

complex, was identified (Fig. 4b). Through RIP and pull-down assay, we demonstrated their direct interaction (Fig. 4c, d). FISH assay also confirmed their colocalization in BGC-823 cells (Fig. 4e). Moreover, domain mapping assay indicated that the region 650–948 bp of circ-DONSON was essential for their interaction (Fig. 4f). EMSA assay further confirmed that SNF2L directly interacted with the region 650–948 bp of circ-DONSON (Fig. 4g). Finally, pull-down assay using circ-DONSON probes showed that circ-DONSON precipitated with SNF2L, BPTF and RBBP4 (three subunits of the NURF complex) in BGC-823 cells (Fig. 4h), indicating circ-DONSON associated with the NURF complex in GC.

circ-DONSON recruits the NURF complex to activate SOX4 transcription

The NURF complex is a chromatin remodeler and activates gene expression [19]. Therefore, we wondered whether the NURF complex participates in the regulation of SOX4 transcription. ChIP assay showed that SNF2L, BPTF and RBBP4 could enrich on the same region of SOX4 promoter as circ-DONSON (Fig. 5a). Notably, circ-DONSON silencing impaired the enrichment of SNF2L, BPTF and RBBP4 on SOX4 promoter (Fig. 5b). FISH assay also confirmed that circ-DONSON





Fig. 3 circ-DONSON regulates GC cell malignant behaviors through activating SOX4. **a** qRT-PCR analysis of indicated gene expression in BGC-823 and AGS cells after circ-DONSON depletion. **b** Western blotting result showed that circ-DONSON silencing suppressed SOX4 expression in BGC-823 and AGS cells. **c** ChIP assay was performed to measure the association of circ-DONSON with SOX4 promoter. **d** Pull-down assay showed that biotin-labeled SOX4 promoter region precipitated circ-DONSON in BGC-823 and AGS cell lysates. **e** DNA-FISH assay indicated the co-localization between circ-DONSON and SOX4 promoter in BGC-823 and AGS cells. Scale bar: 5 μ m. **f** ChIP assay showed that circ-DONSON silencing led to decreased enrichment of active marker H3K27ac on SOX4 promoter in BGC-823 and AGS cells. **g** SOX4 promoter was more resistant to DNaseI digestion after circ-DONSON knockdown. **h-j** CCK8, EdU and colony formation assays were performed to detect cell proliferation. **k** Restoration of SOX4 reduced the apoptosis of BGC-823 and AGS cells induced by circ-DONSON silencing. **l, m** Restoration of SOX4 rescued the abilities of migration and invasion in circ-DONSON knocked down BGC-823 and AGS cells. ** $P < 0.01$ and *** $P < 0.001$

knockdown abrogated the colocalization of SOX4 promoter with SNF2L (Fig. 5c). Interestingly, silencing of SNF2L, BPTF or RBBP4 also suppressed the enrichment of the active markers H3K27ac and H3K4me3 on SOX4 promoter (Fig. 5d, e), indicating that the NURF complex might regulate SOX4 transcription. We further performed luciferase reporter assay and demonstrated that overexpression of SNF2L, BPTF or RBBP4 promoted the luciferase activity of SOX4 promoter while circ-DONSON silencing abrogated it (Fig. 5f), demonstrating that the NURF complex promotes SOX4 transcription in a circ-DONSON-dependent manner. Really, silencing of SNF2L, BPTF or RBBP4 decreased the expression of

SOX4 in GC cells (Fig. 5g, h). Furthermore, we also observed that the expression of SNF2L was positively correlated with SOX4 in GC tissues (Fig. 5i). In summary, our data suggested that the circ-DONSON associated with the NURF complex to activate SOX4 transcription in GC.

The NURF complex modulates GC progression

Whether the NURF complex regulates GC progression has not been reported. Thus, we further explored the roles of the NURF complex on GC cells. Through TCGA database, we found that the expressions of BPTF and RBBP4 were significantly upregulated in GC tissues (Fig. 6a). qRT-PCR analysis also confirmed the upregulation of the



NURF complex in GC tissues and cell lines (Fig. 6b, c). Then we knocked down BPTF, RBBP4 or SNF2L and performed functional experiments. CCK8, colony formation and EdU assays showed that silencing of BPTF, RBBP4 or SNF2L significantly suppressed GC cell proliferation (Fig. 6d-f). FACS analysis and Transwell assay indicated that silencing of BPTF, RBBP4 or SNF2L induced apoptosis and inhibited cell migration and invasion (Fig. 6g-i). In conclusion, these findings suggested that the NURF complex also promotes GC progression.


Effects of circ-DONSON on GC growth in vivo

Then, we investigated the effect of circ-DONSON silencing on GC growth in vivo. BGC-823 cells with circ-DONSON knockdown or negative control were subcutaneously injected into the flank of nude mice.

Every 5 days, the tumor volumes were measured and after 30 days the tumor weights were determined. Results showed that circ-DONSON knockdown significantly decreased tumor volumes and weights (Fig. 7a-c). Furthermore, immunohistochemistry for SOX4 and Ki67 was conducted to detect SOX4 and Ki67 expression. Results showed that circ-DONSON silencing led to a substantial reduce of SOX4 and Ki67 protein levels (Fig. 7d, e), indicating circ-DONSON knockdown suppresses GC growth in vivo through SOX4.

Discussion

In this study, we investigated the functions of circ-DONSON in GC progression. circ-DONSON was highly expressed in GC tissues and cell lines. We also found that circ-DONSON overexpression predicted




advanced tumor stage, metastasis and poor prognosis. Moreover, we found that circ-DONSON silencing suppressed GC cell proliferation, migration and invasion while inducing apoptosis in vitro. Animal experiments indicated that circ-DONSON knockdown suppressed GC growth in vivo. We also found that circ-DONSON associated with the NURF complex through directly interacting with SNF2L. circ-DONSON recruited the NURF complex to SOX4 promoter and initiated its transcription. Collectively, our study demonstrated that circ-DONSON is a novel oncogenic circRNA through activation of SOX4 in GC.

In the recent years, large amounts of circRNAs are identified aberrantly expressed in tumor tissues, including GC. Emerging studies showed that circRNAs are important regulators for tumorigenesis by modulating malignant behaviors of tumor cells [11, 20]. For example,

circRNA hsa_circ_0000263 promotes cervical cancer progression via targeting miR-150-5p [21]. circRNA circ_0008450 is upregulated in hepatocellular carcinoma (HCC) and promotes proliferation and invasion of tumor cells [22]. In GC, only a few important circRNAs have been identified. hsa_circ_0000190 was identified as a diagnostic biomarker in GC [23]. Circular RNA_LARP4 was demonstrated to repress GC growth and invasion [5]. Additionally, Circ-SFMBT2 interacts with miR-182-5p to increase the growth of GC cells through upregulating CREB1 expression [10]. How circRNAs regulates GC development still remains ill understood. In our study, we screened a novel circRNA circ-DONSON. We demonstrated that circ-DONSON contributes to the malignant behaviors of GC cells.


To explore the molecular mechanism of circ-DONSON, we analyzed the downstream signaling pathway. We found



that circ-DONSON could promote SOX4 expression in GC cells. SOX4 is a key transcription factor involved in development of several cancers. For instance, activated SOX4 signaling was reported to promote breast cancer metastasis [24]. Upregulated SOX4 by UCA1 contributes to proliferation and invasion in renal cell carcinoma [25]. SOX4 is also found to promote GC progression [26, 27]. Consistent with above evidence, we also found that increased expression by circ-DONSON regulates the proliferation, migration, invasion and apoptosis of GC cells.

Recently, most studies about circRNAs demonstrated that circRNAs could work as competing endogenous RNAs (ceRNAs) to inhibit miRNAs and play functions [10, 13]. However, in our study, we found that

circ-DONSON was mainly located in the nucleus of GC cells, indicating circ-DONSON might be not a miRNA sponge. Up to date, how circRNAs exerts in the nucleus remains poorly investigated. In our study, we found that circ-DONSON directly deposited on the promoter of SOX4 and regulates its chromatin accessibility. Then through RNA pulldown and mass spectrum identification, we found that circ-DONSON interacted with SNF2L, an important subunit of the NURF complex [28]. We showed that circ-DONSON was associated with the NURF complex through directly interacting with SNF2L subunit in GC. Moreover, we demonstrated that the NURF complex was enriched on SOX4 promoter in a circ-DONSON-dependent manner. The



NURF complex is a critical chromatin remodeler and regulates gene expression [19, 29, 30]. Our results also indicated that NURF depletion suppressed SOX4 transcription and decreased SOX4 mRNA levels. Thus, our research revealed that circ-DONSON/NURF axis activates SOX4 signaling in GC.

Although the NURF complex has been reported to participate in some cancers, such as intestinal tumorigenesis [31] and HCC [32], whether it is involved in GC remains undefined. In our study, we found that its subunits RBBP4, BPTF and SNF2L were highly expressed in GC tissues. And knockdown of RBBP4, BPTF or SNF2L significantly suppressed GC cell proliferation, migration and invasion and promoted apoptosis. Thus, our results illustrated that the NURF complex contributes to GC development for the first time.

Conclusion

In summary, we identified a novel upregulated circRNA circ-DONSON that plays an oncogenic role in GC and associates with poor prognosis. Functional experiments demonstrated that circ-DONSON regulates GC cell proliferation, migration, invasion and apoptosis through the NURF complex-dependent activation of SOX4 signaling.

Additional file

Additional file 1: Figure S1. circ-DONSON overexpression promotes proliferation, migration and invasion of GC cells. a Diagram of back-splicing for circ-DONSON formation. b qRT-PCR analysis of circ-DONSON expression after transfection with pcDNA3-circ-DONSON or vector control. c CCK8 assay was used for proliferation evaluation. d Colony formation assay indicated that

circ-DONSON overexpression increased the colony numbers. e, f Transwell assays indicated that overexpression of circ-DONSON promoted migration and invasion of BGC-823 and AGS cells. $**P < 0.01$ and $***P < 0.001$. **Table S1.** Sequence of circ-DONSON. (DOCX 415 kb)

Abbreviations

ceRNA: Competing endogenous RNA; circRNA: circular RNA; FISH: Fluorescence in situ hybridization; GC: gastric cancer; IHC: Immunohistochemistry; ISH: In situ hybridization; qRT-PCR: Real-time quantitative polymerase chain reaction; RIP: RNA immunoprecipitation

Funding

This study was funded by Outstanding Youth Training Foundation of Academician Yu-Wei in Harbin Medical University (principal investigator Guodong Li) and Science Foundation for Key Project of the Fourth Affiliated Hospital of Harbin Medical University (Grant number: HYDSYJQ201602, principal investigator Guodong Li)

Availability of data and materials

All data are included in the manuscript.

Authors' contributions

GL, LD, YZ, SD and YW contributed to the design of the study. LD, YZ, SD, YW, XL, XY and ZL performed the experiments. JW and ML contributed to the material support of the study. LD, YZ, SD and GL analyzed the data and prepared all the figures and wrote the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The present study was approved by the Ethics Committee of the Fourth Affiliated Hospital of Harbin Medical University.

Consent for publication

Consent was achieved from all patients.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of General Surgery, the Fourth Affiliated Hospital of Harbin Medical University, No. 37 Yiyuan Street, Nangang District, Harbin 150001, People's Republic of China. ²Bio-Bank of Department of General Surgery, the Fourth Affiliated Hospital of Harbin Medical University, Harbin 150001, People's Republic of China. ³Department of Medical Oncology, The Fourth Affiliated Hospital of Harbin Medical University, Harbin 150001, People's Republic of China. ⁴Department of Pharmacology and Toxicology, Wright State University, Fairborn, OH 45435, USA.

Received: 11 January 2019 Accepted: 15 March 2019

Published online: 28 March 2019

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68:394–424.
- Vainio H, Heseltine E, Wilbourn J. Priorities for future IARC monographs on the evaluation of carcinogenic risks to humans. *Environ Health Perspect*. 1994;102:590–1.
- Ge S, Feng XH, Shen L, Wei ZY, Zhu QK, Sun J. Association between habitual dietary salt intake and risk of gastric Cancer: a systematic review of observational studies. *Gastroenterol Res Pract*. 2012;2012:808120.
- Funakoshi T, Miyamoto S, Kakiuchi N, Nikaido M, Setoyama T, Yokoyama A, Horimatsu T, Yamada A, Torishima M, Kosugi S, et al. Genetic analysis of a case of helicobacter pylori-uninfected intramucosal gastric cancer in a family with hereditary diffuse gastric cancer. *Gastric Cancer*. 2018. [Epub ahead of print]. <https://doi.org/10.1007/s10120-018-00912-w>
- Zhang J, Liu H, Hou L, Wang G, Zhang R, Huang Y, Chen X, Zhu J. Circular RNA_LARP4 inhibits cell proliferation and invasion of gastric cancer by sponging miR-424-5p and regulating LATS1 expression. *Mol Cancer*. 2017;16:151.
- Allemani C, Weir HK, Carreira H, Harewood R, Spika D, Wang XS, Bannon F, Ahn JV, Johnson CJ, Bonaventure A, et al. Global surveillance of cancer survival 1995-2009: analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet*. 2015;385:977–1010.
- Qu S, Zhong Y, Shang R, Zhang X, Song W, Kjemis J, Li H. The emerging landscape of circular RNA in life processes. *RNA Biol*. 2017;14:992–9.
- Qu S, Yang X, Li X, Wang J, Gao Y, Shang R, Sun W, Dou K, Li H. Circular RNA: a new star of noncoding RNAs. *Cancer Lett*. 2015;365:141–8.
- Ouyang Y, Li Y, Huang Y, Li X, Zhu Y, Long Y, Wang Y, Guo X, Gong K. CircRNA circPDS1 promotes the gastric cancer progression by sponging miR-186-5p and modulating NEK2. *J Cell Physiol*. 2019;234(7):10458–69.
- Sun HD, Xi PC, Sun ZQ, Wang Q, Zhu B, Zhou J, Jin H, Zheng WB, Tang WW, Cao HY, Cao XF. Circ-SFMBT2 promotes the proliferation of gastric cancer cells through sponging miR-182-5p to enhance CREB1 expression. *Cancer Manag Res*. 2018;10:5725–34.
- Li XN, Wang ZJ, Ye CX, Zhao BC, Li ZL, Yang Y. RNA sequencing reveals the expression profiles of circRNA and indicates that circDDX17 acts as a tumor suppressor in colorectal cancer. *J Exp Clin Cancer Res*. 2018;37:325.
- Li Y, Wan B, Liu L, Zhou L, Zeng Q. Circular RNA circMTO1 suppresses bladder cancer metastasis by sponging miR-221 and inhibiting epithelial-to-mesenchymal transition. *Biochem Biophys Res Commun*. 2019;508:991–6.
- Li X, Zhang Z, Jiang H, Li Q, Wang R, Pan H, Niu Y, Liu F, Gu H, Fan X, Gao J. Circular RNA circPVT1 promotes proliferation and invasion through sponging miR-125b and activating E2F2 signaling in non-small cell lung Cancer. *Cell Physiol Biochem*. 2018;51:2324–40.
- Xu Y, Yao Y, Leng K, Ji D, Qu L, Liu Y, Cui Y. Increased expression of circular RNA circ_0005230 indicates dismal prognosis in breast Cancer and regulates cell proliferation and invasion via miR-618/ CBX8 signal pathway. *Cell Physiol Biochem*. 2018;51:1710–22.
- Wang R, Zhang S, Chen X, Li N, Li J, Jia R, Pan Y, Liang H. EIF4A3-induced circular RNA MMP9 (circMMP9) acts as a sponge of miR-124 and promotes glioblastoma multiforme cell tumorigenesis. *Mol Cancer*. 2018;17:166.
- Wang Y, Cheng Q, Liu J, Dong M. Leukemia stem cell-released microvesicles promote the survival and migration of myeloid leukemia cells and these effects can be inhibited by MicroRNA34a overexpression. *Stem Cells Int*. 2016;2016:9313425.
- Huang G, Jiang H, Lin Y, Wu Y, Cai W, Shi B, Luo Y, Jian Z, Zhou X. IncAKHE enhances cell growth and migration in hepatocellular carcinoma via activation of NOTCH2 signaling. *Cell Death Dis*. 2018;9:487.
- Fu X, Zhu X, Qin F, Zhang Y, Lin J, Ding Y, Yang Z, Shang Y, Wang L, Zhang Q, Gao Q. Linc00210 drives Wnt/beta-catenin signaling activation and liver tumor progression through CTNNBIP1-dependent manner. *Mol Cancer*. 2018;17:73.
- Liu BY, Ye BQ, Yang LL, Zhu XX, Huang GL, Zhu PP, Du Y, Wu JY, Qin XW, Chen RS, et al. Long noncoding RNA lincKdm2b is required for ILC3 maintenance by initiation of Zfp292 expression. *Nat Immunol*. 2017;18:499–508.
- Xu Y, Yao Y, Gao P, Cui Y. Upregulated circular RNA circ_0030235 predicts unfavorable prognosis in pancreatic ductal adenocarcinoma and facilitates cell progression by sponging miR-1253 and miR-1294. *Biochem Biophys Res Commun*. 2019;509:138–42.
- Cai H, Zhang P, Xu M, Yan L, Liu N, Wu X. Circular RNA hsa_circ_0000263 participates in cervical cancer development by regulating target gene of miR-150-5p. *J Cell Physiol*. 2019;234(7):11391–400.
- Zhang J, Chang Y, Xu L, Qin L. Elevated expression of circular RNA circ_0008450 predicts dismal prognosis in hepatocellular carcinoma and regulates cell proliferation, apoptosis, and invasion via sponging miR-548p. *J Cell Biochem*. 2018. [Epub ahead of print]. <https://doi.org/10.1002/jcb.28224>
- Chen SJ, Li TW, Zhao QF, Xiao BX, Guo JM. Using circular RNA hsa_circ_0000190 as a new biomarker in the diagnosis of gastric cancer. *Clin Chim Acta*. 2017;466:167–71.
- Wang N, Liu W, Zheng Y, Wang S, Yang B, Li M, Song J, Zhang F, Zhang X, Wang Q, Wang Z. CXCL1 derived from tumor-associated macrophages promotes breast cancer metastasis via activating NF-kappaB/SOX4 signaling. *Cell Death Dis*. 2018;9:880.
- Liu Q, Li Y, Lv W, Zhang G, Tian X, Li X, Cheng H, Zhu C. UCA1 promotes cell proliferation and invasion and inhibits apoptosis through regulation of the miR129-SOX4 pathway in renal cell carcinoma. *Onco Targets Ther*. 2018;11:2475–87.
- Zou J, Xu Y. MicroRNA-140 inhibits cell proliferation in gastric Cancer cell line HGC-27 by suppressing SOX4. *Med Sci Monit*. 2016;22:2243–52.
- Liu E, Sun X, Li J, Zhang C. miR30a5p inhibits the proliferation, migration and invasion of melanoma cells by targeting SOX4. *Mol Med Rep*. 2018;18:2492–8.
- Eckey M, Kuphal S, Straub T, Rummele P, Kremmer E, Bosserhoff AK, Becker PB. Nucleosome remodeler SNF2L suppresses cell proliferation and migration and attenuates Wnt signaling. *Mol Cell Biol*. 2012;32:2359–71.
- Liu T, Han Z, Li H, Zhu Y, Sun Z, Zhu A. LncRNA DLEU1 contributes to colorectal cancer progression via activation of KPN3A. *Mol Cancer*. 2018;17:118.
- Li Y, Schulz VP, Deng C, Li G, Shen Y, Tusi BK, Ma G, Stees J, Qiu Y, Steiner LA, et al. Setd1a and NURF mediate chromatin dynamics and gene regulation during erythroid lineage commitment and differentiation. *Nucleic Acids Res*. 2016;44:7173–88.
- Zhu P, Wu J, Wang Y, Zhu X, Lu T, Liu B, He L, Ye B, Wang S, Meng S, et al. LncGata6 maintains stemness of intestinal stem cells and promotes intestinal tumorigenesis. *Nat Cell Biol*. 2018;20:1134–44.
- Zhao X, Zheng F, Li Y, Hao J, Tang Z, Tian C, Yang Q, Zhu T, Diao C, Zhang C, et al. BPTF promotes hepatocellular carcinoma growth by modulating hTERT signaling and cancer stem cell traits. *Redox Biol*. 2019;20:427–41.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

