

Open Access In Search of a Sensor: How Does CO₂ Regulate Alveolar Ion Transport?

Levels of carbon dioxide (CO₂), which is an end product of cellular energy metabolism, are well controlled by respiration. Therefore, abnormal alveolar ventilation strongly affects blood pH, in that hyperventilation and hypoventilation cause respiratory alkalosis and acidosis, respectively. Hypercapnia often occurs in patients on mechanical ventilation with elevated fractional oxygen content in the inspired air because of ventilation with low tidal volume designed to minimize ventilator-induced lung damage (1, 2), such as in acute respiratory distress syndrome. But hypercapnia severely impairs alveolar fluid clearance (1), causing or exaggerating alveolar edema in the already injured lung and resulting in tissue hypoxia, which also impairs alveolar fluid clearance and thus aggravates edema formation.

Alveolar fluid clearance is driven by vectorial Na transport mediated by apical epithelial Na channels (ENaC) and basolateral Na/K-ATPase (NKA). Both fluid reabsorption and Na transport are inhibited by hypercapnia, as shown in a series of publications by Sznajder and colleagues using various models that have unraveled bit by bit the signaling cascade (3–9). These publications show that hypercapnia, by transiently increasing the cytosolic Ca²⁺ concentration (Ca²⁺_i), initiates a cascade involving calcium-calmodulin-dependent kinases, mitogen-activated protein (MAP)-kinases, adenosine monophosphate (AMP)-activated kinase, protein kinase C (PKC)ζ, and c-Jun-terminal kinases (JNK) to cause internalization of the NKA and ENaC subunits and retention of the β1-NKA subunit in the endoplasmic reticulum (ER). It remains to be shown whether internalization of α/β-NKA and of α/β ENaC are caused by the same or by independent mechanisms.

In this issue of the *Journal*, Kryvenko and colleagues (pp. 615–629) (10) report that acute hypercapnia-exposed lung slices and A549 cells similarly attenuate the maturation of the β-NKA, a modulator of activity and supporter of membrane sorting of NKA (11). In both systems, hypercapnia caused retention of β-NKA in the ER by increasing the release of Ca²⁺ from the ER via 1,4,5-trisphosphate (IP₃) receptors. This activates IRE1-α to cause degradation of β-NKA. Genetic ablation or inhibition of IRE1-α prevented the hypercapnia effect, and induction of ER stress confirmed the role of endoplasmic-reticulum-associated degradation (ERAD) in degrading β-NKA during hypercapnia.

Although this extraordinary line of experiments clarifies many downstream signaling events leading to the internalization of Na⁺ transporters, the view on the CO₂ sensor and the initial signaling machinery remains blurred. Surprisingly, the most obvious mechanism, acidification, a potent regulator of both ENaC and NKA, seems not to play a significant role because CO₂ effects persist after clamping the extracellular pH and after inhibition (although incomplete) of carbonic anhydrases (CA) (3, 4). This ancient and

ubiquitously expressed enzyme catalyzes with high speed the interconversion between CO₂, H⁺, and bicarbonate and significantly contributes to lung CO₂ elimination (12). The elevation of bicarbonate in hypercapnia certainly affects the intracellular chloride concentration, which in turn might affect transepithelial transport of ions by altering the electrochemical gradients and might thus act independent of complex Ca²⁺ signaling. If it is not the CA mediating the CO₂ effects in alveolar epithelium in culture, in the *in vivo* situation, it might well play a role because of high activity of CA in erythrocytes (13).

At first view, one mechanism common to the internalization of α1- and β1-NKA subunits seems to be an increase in the intracellular Ca²⁺ concentration. However, the results are not consistent. BAPTA-AM, which is often used to scavenge intracellular Ca²⁺ and to prevent its signaling-induced concentration changes, fully prevented the increase in phospho-AMPK by hypercapnia (5). In the present paper, BAPTA-AM at normal CO₂ decreased β1-NKA in the ER and increased p-IRE1 levels, similar to hypercapnia. However, in presence of BAPTA-AM, hypercapnia still exerted its effects (see Figure 5 in Reference 10). Also, eliminating extracellular Ca²⁺ did not prevent hypercapnia-induced effects (see Figure E4 in Reference 10). Thapsigargin, which transiently increases Ca²⁺_i and prevents filling of intracellular stores by blocking SERCA, increased p-IRE1 and decreased β1-NKA in the ER (see Figure E1 in Reference 10) consistent with Ca²⁺_i being the relevant signal. The lack of effect of hypercapnia in the presence of BAPTA-AM plus thapsigargin was interpreted as indicative that a decrease in ER Ca²⁺ levels might be the relevant signal. These inconsistencies await clarification.

How should we proceed with shedding light on the signaling in hypercapnia? First, the CO₂ sensor and its immediate downstream signals need to be identified. Is there a target protein controlling Na²⁺ transport, whose function is altered directly by CO₂ binding (14) similar to changes in oxygen affinity of hemoglobin (15)? There is literature indicating CO₂-induced ATP-release. Extracellular ATP might cause an increase in Ca_i via purinergic receptors and Ca-induced Ca²⁺ release from intracellular stores modulates Ca-induced effects on ion transport in epithelial cells (16). CO₂-induced release of ATP via connexin hemichannels was found in medulla oblongata and HeLa cells independent of acidification (17), which might be caused by formation of carbamate (18).

Might comparison with below-ground-dwelling animals and hibernators provide insight into adjustments to hypercapnia? In fact, these species experience not only hypercapnia but also hypoxia, a situation very similar to patients with respiratory distress syndrome (ARDS) and acute lung injury (ALI). This points to another open and clinically very relevant question: What happens with ion transport in

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern (dgern@thoracic.org).

Supported by the German Center for Lung Research funded by the BMBF (82DZL004B1).

Originally Published in Press as DOI: 10.1165/rcmb.2021-0270ED on August 4, 2021

hypercapnia combined with hypoxia, which also inhibits alveolar reabsorption (19) by inhibiting ENaC and NKA (20)?

Is there a treatment to prevent the adverse effects of reduced alveolar fluid clearance in hypercapnia? Extracorporeal membrane oxygenation, which is used when CO₂ reaches dramatically high levels in ventilated patients, maintains both oxygen supply and removal of CO₂ and thus resolves the inhibition of reabsorption caused by hypercapnia. Considerably less invasive is the stimulation of Na⁺ reabsorption with β₂-adrenergic agents, which not only prevented but even reversed hypercapnic (5) and hypoxic inhibition (21) of alveolar reabsorption, but there might be contraindications for their use. Directly targeting the CO₂ sensor or early CO₂-specific signals might have the fewest unwanted effects. Intervention at levels of MAP kinases, AMPK, or at further distal sites might have too many side effects because of their involvement in multiple signaling events. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Emel Baloglu, M.D.
Department of Pharmacology
Acibadem Mehmet Ali Aydinlar University
School of Medicine
Istanbul, Turkey

Heimo Mairbäurl, Ph.D.
Translational Pneumology
University Hospital Heidelberg
Heidelberg, Germany
and
Translational Lung Research Center Heidelberg
German Center for Lung Research
Heidelberg, Germany

References

- Vadász I, Sznajder JI. Gas exchange disturbances regulate alveolar fluid clearance during acute lung injury. *Front Immunol* 2017;8:757.
- Bihari S, Thompson BT, Bersten AD. Do more injured lungs need more protection? Let's test it. *Am J Respir Crit Care Med* 2021;203:1334–1336.
- Chen J, Lecuona E, Briva A, Welch LC, Sznajder JI. Carbonic anhydrase II and alveolar fluid reabsorption during hypercapnia. *Am J Respir Cell Mol Biol* 2008;38:32–37.
- Briva A, Vadász I, Lecuona E, Welch LC, Chen J, Dada LA, et al. High CO₂ levels impair alveolar epithelial function independently of pH. *PLoS One* 2007;2:e1238.
- Vadász I, Dada LA, Briva A, Trejo HE, Welch LC, Chen J, et al. AMP-activated protein kinase regulates CO₂-induced alveolar epithelial dysfunction in rats and human cells by promoting Na,K-ATPase endocytosis. *J Clin Invest* 2008;118:752–762.
- Vadász I, Dada LA, Briva A, Helenius IT, Sharabi K, Welch LC, et al. Evolutionary conserved role of c-Jun-N-terminal kinase in CO₂-induced epithelial dysfunction. *PLoS One* 2012;7:e46696.
- Dada LA, Trejo Bittar HE, Welch LC, Vagin O, Deiss-Yehiely N, Kelly AM, et al. High CO₂ leads to Na,K-ATPase endocytosis via c-Jun amino-terminal kinase-induced LMO7b phosphorylation. *Mol Cell Biol* 2015;35:3962–3973.
- Gwoździńska P, Buchbinder BA, Mayer K, Herold S, Morty RE, Seeger W, et al. Hypercapnia impairs ENaC cell surface stability by promoting phosphorylation, polyubiquitination and endocytosis of β-ENaC in a human alveolar epithelial cell line. *Front Immunol* 2017;8:591.
- Kryvenko V, Wessendorf M, Morty RE, Herold S, Seeger W, Vagin O, et al. Hypercapnia impairs Na,K-ATPase function by inducing endoplasmic reticulum retention of the β-subunit of the enzyme in alveolar epithelial cells. *Int J Mol Sci* 2020;21:1467.
- Kryvenko V, Wessendorf M, Tello K, Herold S, Morty RE, Seeger W, et al. Hypercapnia induces inositol-requiring enzyme 1α-driven endoplasmic reticulum-associated degradation of the Na,K-ATPase β-subunit. *Am J Respir Cell Mol Biol* 2021;65:615–629.
- Clausen MV, Hilbers F, Poulsen H. The structure and function of the Na,K-ATPase isoforms in health and disease. *Front Physiol* 2017;8:371.
- Crandall ED, O'Brasky JE. Direct evidence of participation of rat lung carbonic anhydrase in CO₂ reactions. *J Clin Invest* 1978;62:618–622.
- Itani OA, Liu KZ, Cornish KL, Campbell JR, Thomas CP. Glucocorticoids stimulate human sgk1 gene expression by activation of a GRE in its 5'-flanking region. *Am J Physiol Endocrinol Metab* 2002;283:E971–E979.
- Linthwaite VL, Janus JM, Brown AP, Wong-Pascua D, O'Donoghue AC, Porter A, et al. The identification of carbon dioxide mediated protein post-translational modifications. *Nat Commun* 2018;9:3092.
- Mairbäurl H, Weber RE. Oxygen transport by hemoglobin. *Compr Physiol* 2012;2:1463–1489.
- Cuffe JE, Bielfeld-Ackermann A, Thomas J, Leipziger J, Korbmacher C. ATP stimulates Cl⁻ secretion and reduces amiloride-sensitive Na⁺ absorption in M-1 mouse cortical collecting duct cells. *J Physiol* 2000;524:77–90.
- Huckstepp RT, id Bihi R, Eason R, Spyer KM, Dicke N, Willecke K, et al. Connexin hemichannel-mediated CO₂-dependent release of ATP in the medulla oblongata contributes to central respiratory chemosensitivity. *J Physiol* 2010;588:3901–3920.
- Meigh L, Greenhalgh SA, Rodgers TL, Cann MJ, Roper DI, Dale N. CO₂ directly modulates connexin 26 by formation of carbamate bridges between subunits. *eLife* 2013;2:e01213.
- Vivona ML, Matthay M, Chabaud MB, Friedlander G, Clerici C. Hypoxia reduces alveolar epithelial sodium and fluid transport in rats: reversal by beta-adrenergic agonist treatment. *Am J Respir Cell Mol Biol* 2001;25:554–561.
- Mairbäurl H, Mayer K, Kim KJ, Borok Z, Bärtsch P, Crandall ED. Hypoxia decreases active Na transport across primary rat alveolar epithelial cell monolayers. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L659–L665.
- Baloglu E, Reingruber T, Bärtsch P, Mairbäurl H. β₂-adrenergics in hypoxia desensitize receptors but blunt inhibition of reabsorption in rat lungs. *Am J Respir Cell Mol Biol* 2011;45:1059–1068.