

Cite this: *Chem. Sci.*, 2018, 9, 380Received 15th September 2017
Accepted 23rd October 2017

DOI: 10.1039/c7sc04054g

rsc.li/chemical-science

Palladium-catalyzed regiodivergent hydroaminocarbonylation of alkenes to primary amides with ammonium chloride†

Bao Gao,^a Guoying Zhang,^a Xibing Zhou^a and Hanmin Huang^{id}*^{ab}

Palladium-catalyzed hydroaminocarbonylation of alkenes for the synthesis of primary amides has long been an elusive aim. Here, we report an efficient catalytic system which enables inexpensive NH₄Cl to be utilized as a practical alternative to gaseous ammonia for the palladium-catalyzed alkene-hydroaminocarbonylation reaction. Through appropriate choice of the palladium precursors and ligands, either branched or linear primary amides can be obtained in good yields with good to excellent regioselectivities. Primary mechanistic studies were conducted and disclosed that electrophilic acylpalladium species were capable of capturing the NH₂-moiety from ammonium salts to form amides in the presence of CO with NMP as a base.

Introduction

Ammonia is among the least expensive industrial chemicals, and has long been recognized as a cheap and ideal nitrogen-source for the synthesis of N-containing molecules *via* C–N bond formation reactions.¹ However, in spite of the utility and abundance of ammonia, it has not frequently been used in transition-metal catalyzed reactions, although notable exceptions include reductive amination,² hydroaminomethylation,³ hydroamination,⁴ allylic amination,⁵ aminocarbonylation,⁶ and coupling of ammonia with aryl halides.⁷ Besides the general technical complexities of using a gas, the potential reason for the scarcity of reports on exploiting ammonia in metal-catalyzed reactions may arise from its easy formation of unreactive Lewis acid-base adducts with transition-metals owing to its high basicity, small size and strong N–H bonds.⁸ An attractive and complementary approach would be to use ammonium salts as surrogates of ammonia gas for establishing transition-metal catalyzed reactions. In this respect, copper and palladium-catalyzed C–N bond formation reactions between aryl halides and ammonium salts have been established by Chang and Hartwig.⁹ However, super stoichiometric amounts of strong base were generally required in both of these systems to form the metal-amide species from the *in situ* released NH₃, leading to wasteful by-product generation. Therefore, the design of new and

efficient protocols which are capable of directly incorporating the nitrogen atom into the hydrocarbon from the ammonium salts in the absence of base is highly desirable.


Transition-metal-catalyzed hydroaminocarbonylation of alkenes constitutes one of the most economical and efficient methods for the construction of aliphatic amides from simple alkenes, CO and amines.¹⁰ Early studies in this field with Co, Ni, Fe and Ru-catalysts required harsh reaction conditions and suffered from poor chemoselectivity.¹¹ As a means of addressing the above synthetic problem, a palladium-catalyzed hydroaminocarbonylation reaction, initiated by palladium-hydride species, has been developed by Beller, Liu, our own group and others.^{12,13} While the utility of this process has been well demonstrated by the synthesis of various secondary and tertiary amides, it is surprising to note that the related reaction for the synthesis of aliphatic primary amides has never been realized, although they are more valuable building blocks in the synthesis of biologically active molecules and functional materials.¹⁴ One obvious reason for discouraging the study of this important unsolved synthetic problem is that the basic ammonia is thought to be necessarily utilized in the related hydroaminocarbonylation reaction, which is not only incompatible with the acids typically utilized for palladium-catalyzed hydroaminocarbonylation but also prone to forming unreactive Werner amine-complexes which deactivate the catalyst.⁸ For the past several years, our group has been focused on the development of palladium-catalyzed alkene hydroaminocarbonylation reactions.¹³ Crucial to the successful realization of these reactions is the formation of palladium-hydride species to initiate the catalysis. Consistent with this concept, we found that hydroxylamine hydrochloride (NH₂·OH·HCl) or alkylamine hydrochloride salts could act as co-catalysts or additives to generate palladium-hydride species *via* oxidative addition with Pd(0), leading to the establishment of efficient hydroaminocarbonylation reactions.¹³ In addition,

^aDepartment of Chemistry, Hefei National Laboratory for Physical Sciences at the Microscale, University of Science and Technology of China, Hefei, 230026, P. R. China. E-mail: hanmin@ustc.edu.cn

^bState Key Laboratory for Oxo Synthesis and Selective Oxidation, Lanzhou Institute of Chemical Physics, Chinese Academy of Sciences, Lanzhou 730000, P. R. China

† Electronic supplementary information (ESI) available. CCDC 1515391 (2e), 1515399 (3h), 1515405 (5i) and 1552470 (9). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7sc04054g





Scheme 1 Proposed hydroaminocarbonylation of alkenes to primary amides with NH_4Cl .

Beller and coworkers also demonstrated that the alkylamine hydrochloride salts could be directly used as a nitrogen source for palladium-catalyzed hydroaminocarbonylation.^{12e} Inspired by these results, we envisioned that the acidic ammonium salt would be a surrogate of ammonia for establishing an efficient catalytic hydroaminocarbonylation reaction to prepare primary amides. Herein, we report an efficient palladium-catalyzed regioselective hydroaminocarbonylation of alkenes to prepare aliphatic primary amides with NH_4Cl as the amino source. The regioselectivity for this reaction is accomplished by the appropriate choice of palladium-catalysts and reaction conditions, and both of the linear and branched primary amides can be constructed in high yields with good regioselectivities (Scheme 1).

Results and discussion

Initially, to test whether the key palladium-hydride species could be formed by the reaction of Pd(0) with an acidic ammonium salt, the reaction of Pd($t\text{-Bu}_3\text{P}$)₃ with NH_4Cl was investigated and monitored by NMR. After heating the mixture of Pd($t\text{-Bu}_3\text{P}$)₂ and NH_4Cl in NMP at 120 °C for 30 min, a new signal at -16.7 ppm appeared in the ¹H NMR spectra, which corresponded to the Pd-H resonance of HPd($t\text{-Bu}_3\text{P}$)₂Cl.^{13c,15} After prolonging the reaction time, the Pd-H resonance signal increased accordingly (see ESI†). These results convincingly supported our hypothesis that palladium-hydride species can be produced by the reaction of Pd($t\text{-Bu}_3\text{P}$)₂ and NH_4Cl .

The above results intrigued us to study the catalytic hydroaminocarbonylation of alkenes with NH_4Cl as the nitrogen source according to the proposed reaction pathway. On the basis of our previous studies of Pd-catalyzed hydroaminocarbonylation reactions,¹³ the present reaction was initially investigated with styrene (**1a**) and NH_4Cl in the presence of a catalytic amount of Pd($t\text{-Bu}_3\text{P}$)₂ under 30 atm of CO in NMP at 120 °C. We were pleased to find that the reaction did indeed proceed leading to the desired primary amides in 95% combined yield with excellent regioselectivity for exclusively giving the branched amide **2a** as the main product (Table 1, entry 1). Decreasing the pressure of CO to 10 atm led to a lower reactivity and selectivity (Table 1, entries 1–3). The simple Pd(PPh_3)₄ could also deliver the desired amide **2a** as the main product in excellent yield albeit with

Table 1 Optimization of the reaction conditions^a


| Entry | [Pd] | Ligand | CO (atm) | Yield (%) | 2a : 3a |
|-------------------|--|----------|----------|-----------|---------|
| 1 | Pd($t\text{-Bu}_3\text{P}$) ₂ | — | 30 | 95 | 96 : 4 |
| 2 | Pd($t\text{-Bu}_3\text{P}$) ₂ | — | 20 | 96 | 88 : 12 |
| 3 | Pd($t\text{-Bu}_3\text{P}$) ₂ | — | 10 | 82 | 85 : 15 |
| 4 | Pd(PPh_3) ₄ | — | 30 | 90 | 81 : 19 |
| 5 ^b | PdI ₂ | Xantphos | 30 | 86 | 20 : 80 |
| 6 ^b | PdI ₂ | DPEphos | 30 | 34 | 22 : 78 |
| 7 ^b | PdI ₂ | DPPF | 30 | 25 | 36 : 64 |
| 8 ^b | PdI ₂ | DPPH | 30 | 27 | 51 : 49 |
| 9 ^b | PdI ₂ | DPPEn | 30 | 23 | 21 : 79 |
| 10 ^b | PdI ₂ | DPPB | 30 | <5 | — |
| 11 ^b | PdI ₂ | DPPE | 30 | <5 | — |
| 12 ^b | PdI ₂ | DPPM | 30 | <5 | — |
| 13 ^{b,c} | PdI ₂ | Xantphos | 30 | 90 | 17 : 83 |
| 14 ^{b,c} | PdI ₂ | Xantphos | 20 | 65 | 17 : 83 |
| 15 ^{b,c} | PdI ₂ | Xantphos | 10 | 60 | 17 : 83 |
| 16 ^{b,c} | PdI ₂ | — | 30 | 20 | 32 : 68 |

^a Reaction conditions: **1a** (1.0 mmol), NH_4Cl (2.0 mmol), [Pd] (5 mol%), NMP (5 mL), 120 °C, and 24 h. The combined yield based on the alkene and the ratio (**2a** : **3a**) of the crude reaction mixture was determined by GC and GC-MS analysis using *n*-cetane as the internal standard. ^b [Pd] (2 mol%), ligand (2.5 mol%), NMP (3 mL), 120 °C, and 24 h. ^c 4 h.

a relatively lower regioselectivity. With Pd($t\text{-Bu}_3\text{P}$)₂ as the catalyst, the reaction time could be shortened to 15 h. Further evaluation of various ammonium salts revealed that the transformation was sensitive to the counter anion of the ammonium salts and found that NH_4Cl was a superb ammonium salt for this reaction (see ESI†). After identifying the optimized catalytic system for the synthesis of the branched amide **2a**, we next sought to determine whether the regioselectivity could be switched by changing the catalyst precursors and reaction conditions. This proved to be highly successful: using PdI₂ as a palladium source and Xantphos as the ligand, the regioselectivity was changed to the opposite direction, favoring the linear amide **3a** when the reaction was conducted at 120 °C under 30 atm of CO (Table 1, entry 5). Encouraged by this result, we then checked different bidentate phosphine ligands with distinct bite angles, such as DPEphos, DPPF, and DPPB, and found that Xantphos was the most efficient for giving the linear amide **3a** in terms of the reactivity and selectivity (Table 1, entries 6–12). With this optimal catalyst, we further screened other reaction parameters to maximize the efficiency of the reaction. This led to finding that the reaction time could be shortened to 4 hours (Table 1, entry 13). Reducing the CO-pressure resulted in a lower reactivity but the same regioselectivity was kept (Table 1, entries 13–15). Interestingly, while the reaction can still be carried out without Xantphos, a lower yield and regioselectivity were observed, revealing that the ligand is an important factor that contributes to a satisfactory result (Table 1, entry 16). In addition, we examined the effect of the solvent and observed that other solvents provided unsatisfactory results in terms of the reactivity and selectivity.



With the optimized reaction conditions in hand, we first examined the substrate scope for the synthesis of branched amides under the catalysis of $\text{Pd}(t\text{-Bu}_3\text{P})_2$, and the results are summarized in Table 2. A variety of styrenes bearing mono-substituents on the aryl ring were initially surveyed. Both electron-donating and -withdrawing groups are compatible with this transformation, and a series of functional groups, such as ether, halogen and ketone, are tolerated to give the desired products **2a–2j** in moderate to excellent yields with good to excellent regioselectivities. The structure of **2e** was confirmed by X-ray crystallographic analysis.¹⁶ For the styrenes with disubstituents on the aryl ring, including 3,4-dichlorostyrene and 2,6-dimethylstyrene, the reactions proceed smoothly to generate **2k** and **2l** in good yields with good selectivities. The reactions of 2-vinylnaphthalene and 1-vinylnaphthalene deliver **2m** and **2n** in 71% and 68% yields, respectively. In addition, the reaction of 9-vinylphenanthrene also provides product **2o** in 68% yield with good regioselectivity. Hydroaminocarbonylation

Table 2 Substrate scope for branched amide synthesis^a

^a Reaction conditions: **1** (1.0 mmol), NH_4Cl (2.0 mmol), NMP (5 mL), CO (30 atm), the isolated yield, and the ratio (B:L) of the crude reaction mixture determined by GC and GC-MS. ^b CO (50 atm) and 24 h.

of an estrone derivative alkene bearing a steroid scaffold was also conducted to produce the linear amide **2p** in 45% yield with a tolerating ketone group. This transformation indicated that the reaction system is amenable to functionalization and modification of complex alkenes bearing the skeleton of natural products, and shows high potential applications in biological evaluation. Unfortunately, aliphatic alkenes were less reactive and only 30% yield was obtained for allylbenzene when the reaction was conducted under 50 atm of CO for 24 h, which may be attributed to faster β -hydride elimination and slower CO insertion for aliphatic alkenes.

Inspired by the above results, we turned our attention to the investigation of the substrate scope for the synthesis of primary linear amides under the catalysis of $\text{PdL}_2/\text{Xantphos}$ in the presence of 30 atm CO pressure (Table 3). To our delight, a range of styrenes and vinylnaphthalenes are also suitable to react with NH_4Cl and CO under the optimized reaction

Table 3 Substrate scope for linear amide synthesis^a

Reaction scheme for Table 3: $\text{R-CH=CH}_2 + \text{NH}_4\text{Cl} + \text{CO} \xrightarrow[\text{NMP, 120 }^\circ\text{C, 4 h}]{\text{PdL}_2 \text{ (2 mol\%)}, \text{Xantphos (2.5 mol\%)}}$ $\text{R-CH}_2\text{-CH}_2\text{-C(=O)NH}_2$

| Entry | R | Product | Yield (%) | L : B |
|-------|---|-----------|-----------|---------|
| 1 | C_6H_5 | 3a | 72 | 83 : 17 |
| 2 | $4\text{-CH}_3\text{C}_6\text{H}_4$ | 3b | 73 | 80 : 20 |
| 3 | $3\text{-CH}_3\text{C}_6\text{H}_4$ | 3c | 69 | 82 : 18 |
| 4 | $2\text{-CH}_3\text{C}_6\text{H}_4$ | 3d | 86 | 94 : 6 |
| 5 | $4\text{-}t\text{-BuC}_6\text{H}_4$ | 3e | 77 | 83 : 17 |
| 6 | $4\text{-CH}_3\text{OC}_6\text{H}_4$ | 3f | 69 | 79 : 21 |
| 7 | $4\text{-FC}_6\text{H}_4$ | 3g | 67 | 82 : 18 |
| 8 | $4\text{-ClC}_6\text{H}_4$ | 3h | 67 | 82 : 18 |
| 9 | $2\text{-ClC}_6\text{H}_4$ | 3i | 77 | 95 : 5 |
| 10 | $4\text{-CH}_3\text{COC}_6\text{H}_4$ | 3j | 73 | 81 : 19 |
| 11 | $2,6\text{-(CH}_3)_2\text{C}_6\text{H}_3$ | 3k | 85 | 96 : 4 |
| 12 | $3,4\text{-Cl}_2\text{C}_6\text{H}_3$ | 3l | 69 | 85 : 15 |
| 13 | 2-Naphthyl | 3m | 78 | 81 : 19 |
| 14 | 1-Naphthyl | 3n | 82 | 96 : 4 |
| 15 | <i>t</i> -Bu | 3o | 74 | >99 : 1 |
| 16 | $\text{C}_6\text{H}_5\text{CH}_2$ | 3p | 54 | 79 : 21 |
| 17 | CyCH_2 | 3q | 56 | 91 : 9 |
| 18 | $\text{CH}_3(\text{CH}_2)_9$ | 3r | 59 | 89 : 11 |
| 19 | $\text{CH}_3(\text{CH}_2)_{11}$ | 3s | 56 | 88 : 12 |
| 20 | $\text{Cl}(\text{CH}_2)_4$ | 3t | 63 | 81 : 19 |
| 21 | $\text{Br}(\text{CH}_2)_5$ | 3u | 51 | 81 : 19 |
| 22 | $\text{MeOOC}(\text{CH}_2)_8$ | 3v | 55 | 87 : 13 |
| 23 | $\text{C}_6\text{H}_5\text{O}(\text{CH}_2)_2$ | 3w | 63 | 89 : 11 |

Additional products and yields (B:L):

- 3x**, 53%, (80:20)
- 3y**, 70%, (90:10)
- 3z**, 65%^b, (78:22)

^a Reaction conditions: **1** (1.0 mmol), NH_4Cl (2.0 mmol), PdL_2 (2 mol%), Xantphos (2.5 mol%), NMP (3 mL), CO (30 atm), 120 °C, and 4 h. The isolated yield based on the alkene and the ratio (L : B) of the crude reaction mixture was determined by GC and GC-MS. ^b $\text{Pd}(\text{COD})\text{Br}_2$ (5 mol%), Xantphos (6 mol%), NMP (5 mL), 120 °C, and 24 h.












conditions, and a series of functional groups, such as ethers, halides and ketones, can survive to give the desired linear amides **3a–3l** in 67–86% isolated yields with good regioselectivities. Notably, the regioselectivity is changed dramatically depending on the substitution. For instance, while the reaction took place with a high selectivity with substrates bearing a methyl group at the *ortho*-position (Table 3, entry 4), the selectivity was lower when it was located at the *para*- or *meta*-position (Table 3, entries 2 and 3). Similar excellent regioselectivities and good yields were observed for the reactions of 2-chlorostyrene, 2,6-dimethylstyrene and 1-vinylnaphthalene (Table 3, entries 9, 11 and 14). In addition to aryl alkenes, simple unactivated aliphatic alkenes are also suitable substrates for this reaction to give the corresponding linear amides in good yields and regioselectivities (Table 3, entries 15–19). Importantly, this reaction is not limited to simple aliphatic alkenes. For instance, functionalized aliphatic alkenes, bearing chlorine, bromine, ether, ester and phthalimide, can be successfully converted to the desired linear amides **3t–3y** in satisfactory yields and good regioselectivities. An estrone derivative was employed to deliver the linear amide **2p** in 65% yield under the catalysis of the Pd(COD)Br₂/Xantphos system. The solid structure of **3h** was further unambiguously confirmed by X-ray crystallographic analysis.¹⁶

To further examine and expand the scope of the reaction, the use of disubstituted alkenes was considered as they allow access to considerably more structurally diverse amides but pose a greater challenge. Indeed, the reaction of α -methyl styrene failed to yield the desired amide with a low conversion under the catalysis of the Pd(*t*-Bu₃P)₂ or PdI₂/Xantphos system. However, a slight change from PdI₂ to Pd(COD)Br₂ led to a successful hydroaminocarbonylation, allowing for a significant expansion of the scope (Table 4). Under these reaction conditions, the desired reaction was amenable to a broad range of α -methyl styrenes, 2-(prop-1-en-2-yl)naphthalene as well as 2-(prop-1-en-2-yl)thiophene, giving a range of primary amides in good to excellent yields with complete regioselectivities (Table 4, entries 1–9). Aside from simple α -methyl styrenes, many other α -substituted aromatic alkenes with different substituents at the α -position are also compatible with this novel reaction, generating the corresponding primary amides in 51–69% yields (**5j–5l**). The structure of the amide **5l** was also confirmed by single-crystal X-ray crystallographic analysis.¹⁶ In addition, cyclic alkenes, such as norbornene and cyclohexene, can be smoothly transferred to the corresponding primary amides **5m** and **5o** in 77% and 79% yield, respectively. Notably, the 1,5-cyclooctadiene is also readily converted to the desired primary amide **5n** in good yield with one double bond remaining. Interestingly, double carbonylation occurs when 3-bromocyclohex-1-ene is subjected to the present reaction conditions, providing the corresponding imide **5p** in 65% yield.

The hydroaminocarbonylation of simple ethylene to propionamide (**6**) with NH₄Cl on a 20 mmol scale was successfully realized in the presence of 0.01 mol% of Pd(*t*-Bu₃P)₂ (18% yield based on NH₄Cl, ton = 1800). In addition, with Pd(CH₃CN)₂Cl₂/Xantphos as the catalyst, the secondary amide *N*-propionylpropionamide (**7**) can be effectively assembled in 95% yield.

Table 4 Substrate scope for disubstituted alkenes^a


|  | | | | |
|--|--|---|-----------|--------------------------|
| Entry | R ¹ | R ² | Product | Yield (%) |
| 1 | C ₆ H ₅ | CH ₃ | 5a | 82 |
| 2 | 4-CH ₃ C ₆ H ₄ | CH ₃ | 5b | 71 |
| 3 | 4- <i>i</i> -BuC ₆ H ₄ | CH ₃ | 5c | 74 |
| 4 | 4-CH ₃ OC ₆ H ₄ | CH ₃ | 5d | 90 |
| 5 | 2-CH ₃ OC ₆ H ₄ | CH ₃ | 5e | 80 |
| 6 | 2-ClC ₆ H ₄ | CH ₃ | 5f | 71 |
| 7 | 3,5-(CF ₃) ₂ C ₆ H ₃ | CH ₃ | 5g | 73 |
| 8 | 2-Naphthyl | CH ₃ | 5h | 71 |
| 9 | 2-Thiophene | CH ₃ | 5i | 52 |
| 10 ^b | C ₆ H ₅ | <i>i</i> -Pr | 5j | 69 |
| 11 | C ₆ H ₅ | (CH ₂) ₇ CH ₃ | 5k | 50 |
| 12 ^b | C ₆ H ₅ | C ₆ H ₅ | 5l | 51 |
| 13 |  |  | 5m | 77(86 : 14) ^c |
| 14 |  |  | 5n | 62 |
| 15 |  |  | 5o | 79 |
| 16 |  |  | 5p | 65 |

^a Reaction condition: **4** (1.0 mmol), NH₄Cl (2.0 mmol), CO (30 atm), NMP (5 mL), 120 °C, and 24 h. The isolated yield. ^b 48 h. ^c dr of the crude reaction mixture determined by GC and GC-MS.


Furthermore, the ¹⁵N-labeled primary amide (**2m**) was obtained when ¹⁵NH₄Cl was utilized as the coupling partner, which provides a convenient method to incorporate ¹⁵N into amides. Finally, to demonstrate scalability, the reaction was performed on a preparative scale (15 mmol) with 1-(*tert*-butyl)-4-vinylbenzene (**1e**) as a substrate. The target reaction proceeded smoothly at a lower catalyst loading (0.1 mol%), affording the desired branched primary amide **2e** in 81% yield with excellent regioselectivity (Scheme 2, eqn (4)).¹⁷

To better understand the mechanism, a series of control experiments were performed. Initially, HPd(*t*-Bu₃P)₂Cl was prepared according to the reported method and utilized as a catalyst for the standard reaction.¹⁵ As expected, excellent reactivity and selectivity were observed for the desired reaction (Scheme 3, eqn (1)), suggesting the plausible intermediacy of palladium-hydride in the catalytic cycle. Moreover, the putative acylpalladium intermediate **9** was prepared *via* the reaction of 3-





Scheme 2 Synthetic applications.



Scheme 3 Preliminary mechanistic studies.

phenylpropanoyl chloride **8** with $\text{Pd}(\text{t-Bu}_3\text{P})_2$ (Scheme 3, eqn (2)). The structure of complex **9** was confirmed by X-ray crystallography.¹⁶ This complex adopts a monomeric T-shaped geometry at palladium, with the acyl ligand trans to the open coordination site, as has been reported for related aroylpalladium complexes.¹⁸ With the acylpalladium complex **9** in hand, we proceeded to execute a set of experiments to elucidate whether the acylpalladium **9** could be converted to an amide and how the NH_2 -moiety was installed into the desired amide from NH_4Cl . The treatment of acylpalladium **9** with two equivalents of NH_4Cl in the presence of CO in NMP at 120 °C afforded the linear amide **3a** and the branched amide **2a** in good yields (Scheme 3, eqn (3)). However, no desired amide was detected, but styrene was formed when the same reaction was conducted

in the absence of CO under other identical reaction conditions (Scheme 3, eqn (4)). These results indicate that the acylpalladium **9** can be converted to the desired amides and CO is essential for facilitating the process.^{18a} The formation of the branched amide **2a** from the linear acylpalladium **9** suggests that CO-deinsertion/insertion and β -hydride elimination/hydro-palladation occur and are reversible. However, the lower selectivity for getting the branched amide **2a** from the linear acylpalladium **9** indicated that the reverse reaction is slow under the reaction conditions, which was further supported by the deuterium studies, in which only partial D-scrambling was observed (Scheme 3, eqn (7) and (8)). As expected, the acylpalladium **9** can catalyze the present hydroaminocarbonylation to give the desired product in high yield under the standard conditions, suggesting the plausible intermediacy of the acylpalladium **9** in the catalytic cycle (Scheme 3, eqn (5)). Further control experiments disclosed that the desired amide **3a** could be readily produced by the reaction of the acid chloride **8** with NH_4Cl in NMP at 120 °C (Scheme 3, eqn (6)).¹⁹ This result together with previous reports¹⁸ that acid chloride could be generated by the reductive elimination of aroyl palladium complexes in the presence of CO suggested two possible pathways for the formation of amide from the acylpalladium **9**: (1) the NH_4Cl directly reacts with the acylpalladium **9** and (2) acid chloride is formed firstly from the acylpalladium **9**, which then reacts with NH_4Cl to generate the desired amide. In both pathways NMP may function as a base to capture the produced HCl (see ESI†).

To distinguish the two possibilities, kinetic reaction profiles for the two stoichiometric reactions of NH_4Cl with the acylpalladium complex (**9**) (eqn (3)) or the acid chloride **8** (eqn (6)) were investigated. The rate of reaction in eqn (3) is faster than that in eqn (6) (Fig. 1). In principle, if a reaction is composed of multiple steps, the observed global kinetic rate should be slower than the rate of each step, or at least equal to the slowest step if it exists. If the desired amide is generated from the acid chloride, eqn (6) should be part of the reaction pathway, and the




Fig. 1 Kinetic plots of the reaction in eqn (3) [acylpalladium **9** (0.2 mmol), NH_4Cl (0.4 mmol), CO (10 atm), and NMP (10 mL) at 120 °C] and the reaction in eqn (6) [**8** (0.2 mmol), NH_4Cl (0.4 mmol), and NMP (10 mL) at 120 °C].






Fig. 2 Plausible reaction mechanism.

observed rate of reaction in eqn (6) should be faster or at least equal to that in eqn (3). However, opposite results were gained, which excluded acid chloride as an intermediate in the reaction of eqn (3), and supported that the desired amide was most likely generated by the direct reaction of NH_4Cl with the acylpalladium complex in NMP.

While a precise reaction mechanism is not yet clear at the present stage, the most plausible mechanism in line with our experimental results and previous reports is illustrated in Fig. 2. First, an oxidative addition of NH_4Cl to $\text{Pd}(0)$ will produce the key palladium-hydride species. Reversible coordination and insertion of alkene into the palladium-hydride yield the Pd-alkyl intermediate **A** or **C**, which undergoes CO insertion to form **B** or **D**, respectively. Consequently, the resultant acylpalladium species **B** and **D** directly react with NH_4Cl in the presence of CO to generate the desired amides. The key palladium-hydride species is simultaneously generated to enter the next catalytic cycle and the released HCl might be trapped by NMP solvent (see ESI[†]) to finish the catalytic cycle. The regioselectivity was found to be highly influenced by the ligand used and the branched amide is exclusively formed for the aromatic alkenes with monodentate ligands. The phenyl group can stabilize the palladium intermediate **C** via delocalization to form an η^3 -benzyl palladium-complex.²⁰ *t*-Bu₃P appeared to be highly favorable to such stabilization due to its ability to create a coordinatively unsaturated palladium complex arising from its size and donor ability, thus facilitating the formation of **C** and/or the subsequent migratory insertion process to predominantly give the branched amide **2**. In contrast, bidentate Xantphos appeared to favor the linear palladium intermediate **A** likely due to the steric hindrance in the corresponding palladium catalyst created by the bidentate diphosphine ligand, consequently facilitating the formation of the linear amide.

Conclusions

In summary, we have successfully developed a practical palladium-catalyzed protocol for the synthesis of primary aliphatic amides via hydroaminocarbonylation of alkenes with NH_4Cl as a surrogate of ammonia in the presence of CO. A wide range of linear or branched primary amides have been obtained in high yields with good to excellent regioselectivities, which represents the first example of the direct conversion of NH_4Cl to

primary aliphatic amides in the absence of base. Apart from the synthetic value of this transformation, important mechanistic evidence regarding the origin of the palladium-hydride and the pathway for incorporation of an NH_2 -moiety from NH_4Cl into the amide was provided. These results suggest that the reaction proceeds through formation of palladium-hydride species generated by oxidative addition of NH_4Cl to $\text{Pd}(0)$ and the acylpalladium species is capable of directly capturing the NH_2 -moiety from the ammonium salt under the assistance of CO in NMP, and has promise as a valuable strategy for utilizing ammonium salts as practical alternatives to gaseous ammonia in a variety of C–N bond-forming manifolds.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (21672199 and 21702197), CAS Interdisciplinary Innovation Team, the Fundamental Research Funds for the Central Universities and the Anhui provincial Natural Science Foundation (1708085MB28).

Notes and references

- (a) J. R. J. LeBlanc, S. Madhavan, R. E. Porter and P. Kellogg, *Encyclopedia of Chemical Technology*, 2nd edn, Wiley, 1963, vol. 2; (b) Y. Aubin, C. Fischmeister, C. M. Thomas and J.-L. Renaud, *Chem. Soc. Rev.*, 2010, **39**, 4130; (c) J. L. Klinkenberg and J. F. Hartwig, *Angew. Chem., Int. Ed.*, 2011, **50**, 86; (d) J. Kim, H. J. Kim and S. Chang, *Eur. J. Org. Chem.*, 2013, 3201; (e) H. Kim and S. Chang, *Acc. Chem. Res.*, 2017, **50**, 482.
- (a) T. Gross, A. M. Seayad, M. Ahmad and M. Beller, *Org. Lett.*, 2002, **4**, 2055; (b) S. Ogo, K. Uehara, T. Abura and S. Fukuzumi, *J. Am. Chem. Soc.*, 2004, **126**, 3020.
- B. Zimmermann, J. Herwig and M. Beller, *Angew. Chem., Int. Ed.*, 1999, **38**, 2372.
- V. Lavallo, G. D. Frey, B. Donnadieu, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2008, **47**, 5224.
- (a) M. J. Pouy, A. Leitner, D. J. Weix, S. Ueno and J. F. Hartwig, *Org. Lett.*, 2007, **9**, 3949; (b) T. Nagano and S. Kobayashi, *J. Am. Chem. Soc.*, 2009, **131**, 4200.
- (a) X.-F. Wu, H. Neumann and M. Beller, *Chem.-Asian. J.*, 2010, **5**, 2168; (b) X.-F. Wu, J. Schranck, H. Neumann and M. Beller, *ChemCatChem*, 2012, **4**, 69; (c) X.-F. Wu, H. Neumann and M. Beller, *Chem.-Eur. J.*, 2012, **18**, 419; (d) K. M. Driller, H. Klein, R. Jackstell and M. Beller, *Angew. Chem., Int. Ed.*, 2009, **48**, 6041.
- (a) Q. Shen and J. F. Hartwig, *J. Am. Chem. Soc.*, 2006, **128**, 10028; (b) D. S. Surry and S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 10354; (c) T. Schulz, C. Torborg, S. Enthaler, B. Schäffner, A. Dumrath, A. Spannenberg, H. Neumann, A. Börner and M. Beller, *Chem.-Eur. J.*, 2009, **15**, 4528; (d)



- G. D. Vo and J. F. Hartwig, *J. Am. Chem. Soc.*, 2009, **131**, 11049.
- 8 (a) D. M. Roundhill, *Chem. Rev.*, 1992, **92**, 1; (b) J. Zhao, A. S. Goldman and J. F. Hartwig, *Science*, 2005, **307**, 1080.
- 9 (a) J. Kim and S. Chang, *Chem. Commun.*, 2008, 3052; (b) R. A. Green and J. F. Hartwig, *Org. Lett.*, 2014, **16**, 4388.
- 10 For recent reviews on the synthesis of amides *via* carbonylation, see: (a) X.-F. Wu, H. Neumann and M. Beller, *Chem. Soc. Rev.*, 2011, **40**, 4986; (b) X.-F. Wu, H. Neumann and M. Beller, *Chem. Rev.*, 2013, **113**, 1; (c) X.-F. Wu, X. Fang, L. Wu, R. Jackstell, H. Neumann and M. Beller, *Acc. Chem. Res.*, 2014, **47**, 1041.
- 11 (a) P. Pino and P. Paleari, *Gazz. Chim. Ital.*, 1951, **81**, 64; (b) P. Pino and R. Magri, *Chim. Ind.*, 1952, **34**, 511; (c) G. Natta, P. Pino and R. Ercoli, *J. Am. Chem. Soc.*, 1952, **74**, 4496; (d) W. Reppe and H. Main, *Chem. Abstr.*, 1953, **47**, 5428; (e) B. F. Crowe and O. C. Elmer, *Chem. Abstr.*, 1956, **50**, 16849; (f) T. J. Kealy and R. E. Benson, *J. Org. Chem.*, 1961, **26**, 3126; (g) A. Striegler and J. Weber, *J. Prakt. Chem.*, 1965, **29**, 281; (h) N. S. Imyanitov and D. M. Rudkvsku, *Chem. Abstr.*, 1967, **66**, 10530; (i) B. K. Nefedov, N. S. Sergeeva and Y. T. Eidus, *Chem. Abstr.*, 1975, **83**, 205731; (j) Y. Tsuji, T. Ohsumi, T. Kondo and Y. Watanabe, *J. Organomet. Chem.*, 1986, **309**, 333; (k) S. I. Lee, S. U. Son and Y. K. Chung, *Chem. Commun.*, 2002, 1310.
- 12 (a) X. Fang, R. Jackstell and M. Beller, *Angew. Chem., Int. Ed.*, 2013, **52**, 14089; (b) H. Liu, N. Yan and P. J. Dyson, *Chem. Commun.*, 2014, **50**, 7848; (c) X. Fang, H. Li, R. Jackstell and M. Beller, *J. Am. Chem. Soc.*, 2014, **136**, 16039; (d) H. Li, K. Dong, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2015, **54**, 10239; (e) J. Liu, H. Li, A. Spannenberg, R. Franke, R. Jackstell and M. Beller, *Angew. Chem., Int. Ed.*, 2016, **55**, 13544; (f) T. Xu, F. Sha and H. Alper, *J. Am. Chem. Soc.*, 2016, **138**, 6629.
- 13 (a) G. Zhang, B. Gao and H. Huang, *Angew. Chem., Int. Ed.*, 2015, **54**, 7657; (b) G. Zhang, X. Ji, H. Yu, L. Yang, P. Jiao and H. Huang, *Tetrahedron Lett.*, 2016, **57**, 383; (c) Y. Hu, Z. Shen and H. Huang, *ACS Catal.*, 2016, **6**, 6785.
- 14 (a) F. Matsuda, *Chem. Technol.*, 1977, **7**, 306; (b) C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, **40**, 3405.
- 15 I. D. Hills and G. C. Fu, *J. Am. Chem. Soc.*, 2004, **126**, 13178.
- 16 CCDC 1515391 (**2e**), 1515399 (**3h**), 1515405 (**5l**) and 1552470 (**9**) contain the supplementary crystallographic data for this paper.†
- 17 The utility of this catalytic reaction has been further demonstrated by streamlined synthesis of ibuprofen, nitriles and 3,4-dihydroquinolin-2(1*H*)-one from simple alkenes and NH₄Cl. See ESI† for details.
- 18 (a) J. S. Quesnel and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2013, **135**, 16841; (b) S. Korsager, R. H. Taaning and T. Skrydstrup, *J. Am. Chem. Soc.*, 2013, **135**, 2891; (c) J. Tjutris and B. A. Arndtsen, *J. Am. Chem. Soc.*, 2015, **137**, 12050.
- 19 No reaction occurred when the reaction was conducted in many other solvents such as toluene, CH₂Cl₂, THF, 1,4-dioxane and hexane, suggesting that basic NMP is essential for facilitating the reaction by capturing the released HCl (see ESI† for details).
- 20 (a) W. Ren, W. Chang, Y. Wang, J. Li and Y. Shi, *Org. Lett.*, 2015, **17**, 3544; (b) J. Li, W. Chang, W. Ren, J. Dai and Y. Shi, *Org. Lett.*, 2016, **18**, 5456.

