

# Correlation between Transabdominal Sonographic Prostate Volume and Anthropometric Parameters

Chibueze Okorie Udo<sup>1</sup>, Ebbi Donald Robinson<sup>1\*</sup>, Olukunmi Yetunde Ijeruh<sup>1</sup>, Nelson Chukwuemeka Nwankwo<sup>2</sup>

<sup>1</sup>Department of Radiology, Rivers State University Teaching Hospital, Port Harcourt, Rivers State, Nigeria, <sup>2</sup>Department of Radiology, University of Port Harcourt Teaching Hospital, Port Harcourt, Rivers State, Nigeria

## Abstract

**Background:** Prostate diseases commonly present with lower urinary tract symptoms (LUTS) resulting from prostatic enlargement. Prostate volume (PV) can be evaluated using transabdominal ultrasonography. Focus is currently on relative factors of prostatic enlargement which includes obesity and central adiposity. The aim of this study is to correlate transabdominal sonographic PV and anthropometric parameters in patients with LUTS in Port Harcourt. **Methods:** This was a prospective cross-sectional study carried out at the Radiology Department, Rivers State University Teaching Hospital, Port Harcourt, between September 2020 and January 2021. One hundred and twenty (120) males from 40 years and above who presented with LUTS were recruited. Transabdominal PV estimation was done and body mass index (BMI) as well as WC was assessed. Data were analyzed using a Statistical Package for Social Sciences; appropriate statistical tests were applied and  $P < 0.05$  was considered significant. **Results:** The mean PV was  $69.8 \pm 63.5 \text{ cm}^3$ , 79.2% of the subjects had enlarged prostate with volume  $\geq 30 \text{ cm}^3$ . PV was found to increase with age. The correlation between PV and anthropometric measures of obesity (BMI and WC) was statistically not significant. **Conclusion:** The work established that there is no correlation between PV and anthropometric measures of obesity – BMI and WC in negro population as opposed to nonblack population where there is correlation. Obesity may not be a considerable risk factor of prostatic enlargement in the studied population. Thus, anthropometrics may not be useful in predicting prostate size.

**Keywords:** Body mass index, International Prostate Symptom Score, obesity, prostate volume, waist circumference

## INTRODUCTION

Pathologies of the prostate gland which are mainly hyperplastic (benign prostatic hyperplasia), malignant (carcinoma of the prostate), and less commonly inflammatory (prostatitis) frequently give rise to increase in the organ size. Prostatic enlargement (prostatomegaly) leads to bladder outlet obstruction as a result of static compression as well as dynamic obstruction owing to the contraction of prostatic smooth muscles.<sup>[1]</sup> This could be asymptomatic in some cases; however, it commonly manifests as lower urinary tract symptoms (LUTS). PV assessment has become increasingly important because of its connections to disease progression, treatment response prediction, and therapeutic options.<sup>[2]</sup> Therefore, determinants of prostate volumes (PVs) will invariably influence management options, prognosis as well as preventive measures.

Radiologic evaluation of prostatic disease is commonly by ultrasonography, particularly in Nigeria as a developing country where availability of imaging equipment and cost of services are limiting factors.<sup>[3,4]</sup> Magnetic resonance imaging could be done, especially when malignant disease is suspected. Other modalities such as plain radiography, fluoroscopy, computed tomography as well as radionuclear studies can also image the gland. However, ultrasound remains the first-line modality which is safe, noninvasive, fast, cost-effective, and readily available imaging tool for the prostate gland. Transabdominal, transperineal, transurethral, and transrectal approaches are the common routes employed.<sup>[5]</sup>

Anthropometric measurements include body mass index (BMI), body circumferences, among others. The

**Address for correspondence:** Dr. Ebbi Donald Robinson,  
Department of Radiology, Rivers State University Teaching Hospital,  
Port Harcourt, Rivers State, Nigeria.  
E-mail: drebbirobinson@yahoo.co.uk

Received: 29-09-2021 Revised: 12-12-2021 Accepted: 22-12-2021 Available Online: 19-05-2022

### Access this article online

#### Quick Response Code:



**Website:**  
www.jmuonline.org

**DOI:**  
10.4103/jmu.jmu\_183\_21

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Udo CO, Robinson ED, Ijeruh OY, Nwankwo NC. Correlation between transabdominal sonographic prostate volume and anthropometric parameters. J Med Ultrasound 2022;30:261-5.

most common anthropometric measure of central obesity is waist circumference (WC), and hip circumference. The ethnic-specific definition of central obesity for African men by the International Diabetes Federation is  $WC \geq 94$  cm.<sup>[6,7]</sup> Central adiposity is a function of subcutaneous and visceral fats which are important components of metabolic syndrome; it may be considered “at-risk obesity.”<sup>[8]</sup>

Ultrasonography is the mainstay of PV assessment. Transabdominal and transrectal approaches are the common routes employed in imaging the prostate. The later, although it gives more accurate volumetric assessment, however, studies have shown no statistically significant difference between the transabdominal and transrectal sonographic PVs for clinical purposes.<sup>[9,10]</sup> In addition, patient’s discomfort, the need for rectal emptying, relatively more time consuming, and less cost-effectiveness relative to the former are the considerations in the choice of transabdominal approach for the study.

Currently, with emphasis on preventive than therapeutic medicine, more attention is given to determinants and risk factors of prostate enlargement with a view to identifying modifiable factors associated with prostate diseases.

Although studies have been carried out among Caucasians on the relationship between PV and obesity, limited data are available on researches done in sub-Saharan Africa. Particularly in Nigeria, few studies have been done in the area of the subject matter.<sup>[11,12]</sup> Thus, the purpose of this research project is to determine the association between PV and anthropometric measurements (particularly BMI and WC) and its relationship with symptom severity among men in South-South Nigeria.

## MATERIALS AND METHODS

This research was a prospective cross-sectional study, carried out at Rivers State University Teaching Hospital (RSUTH), Port Harcourt, from September 2020 to January 2021, among men above 40 years of age with complaints of LUTS, referred from the Urology Clinic to Radiology Department of RSUTH.

All the patients who presented for prostate ultrasound in the radiology department within the study period, who met the eligibility criteria, were recruited once they gave informed.

Informed consent was obtained. Then, the subject’s age and demographic and anthropometric parameters (height, weight, and WC) were taken. Transabdominal ultrasonography of the prostate gland was done and PV was measured. The information was appropriately recorded on the study data sheet. Participants were scanned with a moderately filled urinary bladder in a supine position using an ultrasound scanner fitted with an 3.5-MHz curvilinear transducer (Logiq F6, General Electric, USA, 2017). The patient was asked to lie supine in the ultrasound couch and his pelvic region was adequately exposed, then the area draped and cleaned. Coupling gel was applied to bridge acoustic impedance between the skin and probe surface.

PV was calculated from the dimensions obtained (in cm) using the default computer algorithm in the ultrasound machine based on the prostate ellipsoid formula<sup>[13]</sup> –  $PV \text{ (in cm}^3\text{)} = \text{length} \times \text{height} \times \text{width} \times 0.52$ , where the length, height, and width were the maximum cephalocaudal, anteroposterior, and transverse diameters, respectively, as shown in Figure 1.

## Anthropometric measurements

The subjects’ height, weight, and WC were obtained using standard anthropometric techniques as follows.

**Height:** With the subjects standing straight on the stadiometer base with barefoot, heels put together, while the buttocks and back make contact with the vertical rod, the heights were read. The measurements were taken in meters using a Z-16 Stadiometer (Wincom Company Limited, China, 2017).

**Weight:** Was also measured with the same Stadiometer, with the patient in the same position as above, having removed foot wares and heavy clothing. Readings were taken in kilograms.

**BMI** was calculated using the subjects’ height and weight measurements as  $BMI = \text{weight}/\text{height}^2 \text{ (kg/m}^2\text{)}$ .


**WC:** The subject in a standing position with the abdomen exposed, the lower rib margin and iliac crest were felt from behind, and measurements were taken in centimeters at the midpoint of the two bony landmarks at the end of expiration.

## Ethical consideration

Ethical approval was granted by the Ethical Committee of RSUTH (approval number: RSHMB/RSHREC/11.19/VOL 7/039).

## RESULTS

The mean age ( $\pm$ standard deviation [SD]) was  $65.1 \pm 9.6$  years, with an age range of 48–94 years [Table 1]. The lowest and highest age ranges of the participants were observed as 40–49 years ( $n = 6$ ; 5%) and 60–69 years ( $n = 50$ ;



**Figure 1:** Transabdominal sonograms of a participant showing grey-scale static images of the prostate gland – Longitudinal (a) and Transverse (b) sections with a prostatic volume of 98.90cm<sup>3</sup> NB: LONG: Longitudinal section, TRANS: Transverse section; \*: Urinary bladder

41.7%), respectively, as shown in Figure 2. The mean BMI was  $26.7 \pm 4.1$  kg/m<sup>2</sup>, while the mean WC was  $90.72 \pm 12.7$  cm ( $\pm$ SD). The mean PV measured among the participants was  $69.8 \pm 63.5$  cm<sup>3</sup> ( $\pm$ SD) [Table 1].

Table 2 shows that 41 (34.2%), 54 (45.0%), and 25 (28.8%) are of normal BMI, overweight, and obese, respectively, whereas 64 (53.3%) subjects are of normal group (WC <94 cm) and 56 (46.7%) in the central obesity group (WC  $\geq$ 94 cm).

As shown in Table 3, a total of 95 subjects (79.2%) had enlarged prostate of volume  $\geq 30$  cm<sup>3</sup>, while the higher proportion (66.7%) of the younger age group (40–49 years) had PV <30 cm<sup>3</sup>. PV was seen to vary among the various BMI (kg/m<sup>2</sup>) categories, with the mean volumes of 78.1 and 64.9 recorded in the normal and overweight (BMI 18.5–24.9 and 25.0–29.9), while 65.0 was noted in the obese (BMI  $\geq 30$ ) categories, respectively, as shown in Table 4. Table 3 also demonstrates that Fisher’s test between PV and BMI categories shows no significant association ( $P = 0.560$ ); the same also applies to the *t*-test between PV and WC categories ( $P = 0.064$ ).

The Pearson correlation of PV with BMI shows a very weak negative relationship which is not statistically significant ( $r = -0.120$ ,  $P = 0.192$ ), as illustrated in Table 5. So also, is PV with WC ( $r = -0.137$ ,  $P = 0.137$ ) which showed a very weak negative relationship that was not statistically significant [Table 5].

## DISCUSSION

In this cross-sectional study involving 120 men aged 40 years and above, the average age was 65.1 years. Majority of the patients were in the age range of 60–69 years (41.7%). The above findings are similar to the mean ages of 66.25 and 62.50 years reported locally by Udeh *et al.*<sup>[14]</sup> (Enugu) and Mohammed *et al.*<sup>[15]</sup> (Zaria), respectively, and 64.10 years by Rupam *et al.*<sup>[16]</sup> in India, with the highest frequencies (46%, 34.5%, and 42%, respectively) recorded within 60–69 years age range.

The mean transabdominal sonographic PV was  $69.84 \pm 63.5$  cm<sup>3</sup>. In a related study on transrectal PV correlation with prostate-specific antigen level in the same study setting,




Figure 2: Age distribution of participants

Robinson<sup>[17]</sup> reported a mean volume of  $66.13 \pm 30.43$  cm<sup>3</sup> among 143 symptomatic patients. The value is however lower than the mean  $83.8 \pm 37.7$  cm<sup>3</sup> and  $72.79 \pm 44.4$  cm<sup>3</sup> reported

Table 1: Demographic characteristics of the participants

Variables participants (n=120)	Mean $\pm$ SD	Median
Age (years)	65.1 $\pm$ 9.6	64.0
Height (m)	1.7 $\pm$ 0.05	1.7
Weight (kg)	72.7 $\pm$ 12.9	70.0
BMI (kg/m <sup>2</sup> )	26.7 $\pm$ 4.1	26.0
WC (cm)	90.7 $\pm$ 12.7	92.0
Prostate volume (cm <sup>3</sup> )	69.8 $\pm$ 63.5	51.6

BMI: Body mass index, WC: Waist circumference, SD: Standard deviation, IPSS: International Prostate Symptom Score

Table 2: Distribution of body mass index and waist circumference categories of the participants

Variables	Frequency (n=120) n (%)
BMI category (kg/m <sup>2</sup> )	
18.5-24.9	41 (34.2)
25.0-29.9	54 (45.0)
$\geq 30.0$	25 (28.8)
WC category (cm)	
<94	64 (53.3)
$\geq 94$	56 (46.7)

BMI: Body mass index, WC: Waist circumference

Table 3: Distribution of prostate volume with age group

Age (years)	Prostate volume cm <sup>3</sup>		Total, n (%)
	<30, n (%)	$\geq 30$ , n (%)	
40-49	4 (66.7)	2 (33.3)	6 (100.0)
50-59	4 (15.4)	22 (84.6)	26 (100.0)
60-69	8 (16.0)	42 (84.0)	50 (100.0)
$\geq 70$	9 (23.7)	29 (76.3)	38 (100.0)
Total	25 (20.8)	95 (79.2)	120 (100.0)

Table 4: Distribution of prostate volume and anthropometric (body mass index and waist circumference)

Variable	Frequency	Mean (cm <sup>3</sup> ) $\pm$ SD	F-test	P
BMI category (kg/m <sup>2</sup> )				
18.5-24.9	41	78.1 $\pm$ 83.5	0.582	0.560
25.0-29.9	54	64.9 $\pm$ 52.0		
$\geq 30$	25	65.0 $\pm$ 44.7		
Total	120	69.8 $\pm$ 63.5		
Waist circumference category (cm)				
<94	64	79.5 $\pm$ 79.1	1.875	0.064
$\geq 94$	56	58.8 $\pm$ 36.3		
Total	120	69.8 $\pm$ 63.5		

SD: Standard deviation

**Table 5: Prostate volume correlation with body mass index and waist circumference**

Variables	Correlation coefficient	P
Prostate volume versus BMI	-0.120	0.192
Prostate volume versus WC	-0.137	0.137

BMI: Body mass index, WC: Waist circumference

by Badmus *et al.*<sup>[11]</sup> in Ile-Ife and Udeh *et al.*<sup>[18]</sup> in Jos among 105 (mean age of 64.4 years) and 100 (mean age of 65.6 years) subjects, respectively. It is however higher than  $56.2 \pm 42.7 \text{ cm}^3$  recorded by Mohammed *et al.*<sup>[15]</sup> among 602 patients in Zaria with the age of 64.1 years. The differences may be attributed to numerical bias – more patients were recruited in this index study than was done in the work by Badmus *et al.* and that of Udeh *et al.*, while Mohammed *et al.* enrolled larger subjects in his study. The use of transrectal approach by Mohammed *et al.* compared to transabdominal volume estimation by the two studies as well as the index work may also account for the discrepancy.

The mean BMI recorded in this study is  $26.7 \pm 4.1 \text{ kg/m}^2$ . This is similar to the mean BMI of  $25.00 \pm 5.10$  reported by Robinson<sup>[17]</sup> among adult men aged 45–84 in the same facility, while Ejike<sup>[12]</sup> and Mubenga<sup>[19]</sup> recorded  $24.0 \pm 3.03$  and  $25.1 \pm 3.3$  in Eastern Nigeria and Congo Republic, respectively. The mean WC reported in this study was  $90.72 \pm 12.69 \text{ cm}$ , and it is similar to 88.4 cm accounted by Badmus *et al.*<sup>[11]</sup> in Ile-Ife in 2019, while  $94.6 \pm 10.3 \text{ cm}$  was recorded among Congolese.<sup>[20]</sup>

The association between prostate size and anthropometric measures of obesity has shown a wide range of variance from studies done both locally and in foreign settings. Different studies have shown significant correlation, either positive or negative, while others recorded no statistically significant association. This study found no significant correlation between PV and BMI ( $P = 0.192$ ) and also between PV and WC ( $P = 0.137$ ) respectively. The findings are in agreement with the two local studies done in Ife in 2013 and 2019 where PV was correlated with anthropometric in men presenting with LUTS. In the former, Badmus *et al.*<sup>[11]</sup> found no correlation between transabdominal PV and BMI among 105 men aged 40 and above who were managed for BPH ( $P = 0.840$ ). The latter study by Asaley *et al.*<sup>[21]</sup> among 90 men of similar age, correlation of transabdominal and transrectal PV with BMI ( $r = 0.156$ ,  $P = 0.144$ ) and PV with WC ( $r = -0.068$ ,  $P = 0.525$ ) was not significant in both cases. However, a positive but weak correlation was demonstrated between BMI and transrectal transitional zone volume ( $r = -0.230$ ,  $P = 0.029$ ).

Burke *et al.*<sup>[22]</sup> in an age-stratified random sample of 105 Caucasian males aged 43–88 years in Minnesota established that association between PV and anthropometric measures (BMI,  $P = 0.49$ , WC,  $P = 0.07$ ) was not significant. Similarly, the work done by Kim *et al.*<sup>[23]</sup> in South Korea found no correlation between PV and BMI ( $r = 0.164$ ,  $P < 0.001$ ) but

positive linear correlation between PV and WC ( $r = 0.217$ ,  $P < 0.01$ ). Conversely, a control study in Italy showed a moderate inverse relationship between BMI and BPH.<sup>[24]</sup> In the study, Zucchetto *et al.*<sup>[24]</sup> compared previous BMI of 1 year prior to histological diagnosis of BPH which was assessed by self-reported weight and height, with current PV in this case–control study. Thus, methodology may have accounted for the difference in findings.

On the other hand, most studies among nonblack men have shown a significant positive correlation between PV and anthropometric measures of obesity. In Pakistan, Raza *et al.*<sup>[25]</sup> found that transabdominal PV correlated positively with BMI and WC ( $P = 0.046$ ,  $P = 0.003$ , respectively). Jung *et al.*<sup>[26]</sup> and Fowke *et al.*<sup>[27]</sup> made similar findings in South Korea and the United States, respectively. In the same vein, Monowara *et al.*<sup>[28]</sup> reported a significant positive PV correlation coefficient of  $r = 0.352$  ( $P < 0.001$ ) with BMI. Furthermore, Li *et al.*<sup>[29]</sup> and Sokhal *et al.*<sup>[30]</sup> established similar relationship between PV and BMI among Chinese and Indian men (odds ratio = 1.772,  $P = 0.005$ ;  $r = 0.132$ ,  $P < 0.001$ ), respectively.

Most studies reported among the blacks, including this index study, however, demonstrate no significant relationship between obesity and prostate gland enlargement. This therefore suggests a strong influence of ethn racial factor in the determination of prostate size among adult males. Another possible explanation for the findings is the hypothesis of reduced testosterone levels in obese individuals as proposed by Zucchetto *et al.*<sup>[24]</sup> and La Vignera *et al.*<sup>[31]</sup>

## CONCLUSION

The work established that there is no correlation between PV and anthropometric measures of obesity – BMI and WC in black population as opposed to nonblack population where there is correlation. Obesity may not be a considerable risk factor of prostatic enlargement in the studied population. Thus, anthropometrics may not be useful in predicting prostate size, other risk factors aside obesity should be considered in the evaluation, prevention, and management of prostatic enlargement.

Furthermore, a longitudinal study design is recommended for further studies to address the limitations of this cross-sectional study in making an objective comparison of the variables over time. Finally, further studies on the scopes of this work are recommended in other geopolitical regions of Nigeria and beyond, including community-based surveys, to add to the existing body of knowledge.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

1. Foo KT. Pathophysiology of clinical benign prostatic hyperplasia. *Asian J Urol* 2017;4:152-7.
2. Fitzpatrick JM. The natural history of benign prostatic hyperplasia. *Br J Urol Int* 2006;97:3-6.
3. Ikpeme A, Ani N, Ago B, Effa E, Kosoko-Lasaki O, Ekpenyong A. The value of mobile ultrasound services in rural communities in south-south Nigeria. *Open Access Maced J Med Sci* 2017;5:1011-5.
4. Tyloch JF, Wiecek AP. The standards of an ultrasound examination of the prostate gland. Part 1. *J Ultrason* 2016;16:378-90.
5. Ebeye AO, Oyem JC, Iweariolor BE, Ubah SC. Ultrasonographic assessment of normal prostate volume and splenic length among Urhobo ethnic group in Delta State of Nigeria. *Ann Bioanthropol* 2016;4:101-4.
6. Poirier P. The many paradoxes of our modern world: Is there really an obesity paradox or is it only a matter of adiposity assessment? *Ann Intern Med* 2015;163:880-1.
7. Kim SY. The definition of obesity. *Korean J Fam Med* 2016;37:309.
8. Okafor CI, Raimi TH, Gezawa ID, Sabir AA, Enang O, Puepet F, *et al.* Performance of waist circumference and proposed cut-off levels for defining overweight and obesity in Nigerians. *Ann Afr Med* 2016;15:185-93.
9. Huang Foen Chung JW, de Vries SH, Raaijmakers R, Postma R, Bosch JL, van Mastrigt R. Prostate volume ultrasonography: The influence of transabdominal versus transrectal approach, device type and operator. *Eur Urol* 2004;46:352-6.
10. Ajayi I, Aremu A, Olajide A, Bello T, Olajide F, Adetiloye V. Correlation of transrectal and transabdominal ultrasound measurement of transition zone volume with post-operative enucleated adenoma volume in benign prostatic hypertrophy. *Pan Afr Med J* 2013;16:149.
11. Badmus TA, Asaleye CM, Badmus SA, Takure AO, Ibrahim MH, Arowola OA. Benign prostate hyperplasia: Average volume in southwestern Nigerians and correlation with anthropometrics. *Niger Postgrad Med J* 2013;20:52-6.
12. Chukwunonso E, Eze C. Prevalence of symptoms of benign prostatic hyperplasia in umudike and its relationship with measures of obesity. *Asian J Clin Nutr* 2015;7:1-8.
13. Aprikian S, Luz M, Brimo F, Scarlata E, Hamel L, Cury FL, *et al.* Improving ultrasound-based prostate volume estimation. *BMC Urol* 2019;19:68.
14. Udeh EI, Nnabugwu II, Ozoemena FO, Ugwumba FO, Aderibigbe AS, Ohayi SR, *et al.* Prostate-specific antigen density values among patients with symptomatic prostatic enlargement in Nigeria. *World J Surg Oncol* 2016;14:174.
15. Mohammed A, Bello A, Maitama HY, Ajibola HO, Lawal AT, Isah MJ, *et al.* Determination of prostatic volume and characteristics by transrectal ultrasound among patients with lower urinary tract symptoms in Zaria, Nigeria. *Arch Int Surg* 2014;4:31-5.
16. Rupam D, Bijoyananda D, Mustafa A, Rahman R. A study of relationship of prostate volume, prostate specific antigen and age in benign prostatic hyperplasia. *Int J Contemp Med Res* 2017;4:1582-6.
17. Robinson ED. Trans-rectal ultrasound prostate volume correlation with serum prostate specific antigen level in patients with prostatic enlargement in Port Harcourt. *Asian J Med Radiol Res* 2019;7:66-71.
18. Udeh EI, Dakum NK, Aderibigbe SA, Edeh JA. The utility of digital rectal examination in estimating prostate volume in a rural hospital setting. *Niger J Surg* 2015;21:111-4.
19. Bhindi B, Margel D, Trotter G, Hamilton RJ, Kulkarni GS, Hersey KM, *et al.* Obesity is associated with larger prostate volume but not with worse urinary symptoms: Analysis of a large multiethnic cohort. *Urology* 2014;83:81-7.
20. Mubenga LE, Chimanuka D, DeGroot P. Comparison of prostate size and anthropometric parameters between diabetic and non-diabetic Congolese patients who underwent transurethral prostate resection in the Democratic Republic of Congo. *Afr J Urol* 2019;25:2.
21. Asaleye CM, Omisore AD, Onigbinde SO, David RA. Obesity in benign prostatic enlargement: A cross-sectional study comparing sonographic and anthropometric indices of adiposity in a tertiary hospital in Southwestern Nigeria. *Niger J Clin Pract* 2019;22:1600-5.
22. Burke JP, Rhodes T, Jacobson DJ, McGree ME, Roberts RO, Girman CJ, *et al.* Association of anthropometric measures with the presence and progression of benign prostatic hyperplasia. *Am J Epidemiol* 2006;164:41-6.
23. Kim GW, Doo SW, Yang WJ, Song YS. Effects of obesity on prostate volume and lower urinary tract symptoms in Korean men. *Korean J Urol* 2010;51:344-7.
24. Zucchetto A, Tavani A, Dal Maso L, Gallus S, Negri E, Talamini R, *et al.* History of weight and obesity through life and risk of benign prostatic hyperplasia. *Int J Obes (Lond)* 2005;29:798-803.
25. Raza I, Nuzhat H, Pashmina G, Anis J, Nosheen Z, Naila Y. Determination of prostate gland volume by ultrasonography and its correlation with anthropometric measurements in a subset of Karachi population. *Br J Med Med Res* 2016;11:1-12.
26. Jung JH, Ahn SV, Song JM, Chang SJ, Kim KJ, Kwon SW, *et al.* Obesity as a risk factor for prostatic enlargement: A retrospective cohort study in Korea. *Int Neurourol J* 2016;20:321-8.
27. Fowke JH, Koyama T, Fadare O, Clark PE. Does inflammation mediate the obesity and BPH relationship? An epidemiologic analysis of body composition and inflammatory markers in blood, urine, and prostate tissue, and the relationship with prostate enlargement and lower urinary tract symptoms. *PLoS One* 2016;11:e0156918.
28. Monowara M, Ahmed AU, Mohiuddin AS, Taher MA, Nasrin Z, Hossain MM. Correlation between transabdominal sonographically measured prostate volume and anthropometric factors in normal healthy subjects. *Birdem Med J* 2012;2:29-32.
29. Li BH, Deng T, Huang Q, Zi H, Weng H, Zeng XT. Body mass index and risk of prostate volume, international prostate symptom score, maximum urinary flow rate, and post-void residual in benign prostatic hyperplasia patients. *Am J Mens Health* 2019;13:1557988319870382.
30. Sokhal AK, Jhanwar A, Sankhwar S, Singh K, Gupta AK. Does body mass index have an impact on prostate volume and serum prostate specific antigen? A prospective observational study in patients with lower urinary tract symptoms. *J Urol Nephrol* 2016;2:1-5.
31. La Vignera S, Condorelli RA, Russo GI, Morgia G, Calogero AE. Endocrine control of benign prostatic hyperplasia. *Andrology* 2016;4:404-11.