

Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects

André M. Eckardt

Department of Cranio-Maxillofacial Surgery, Faculty of Dentistry, Hannover Medical University, Hannover, Germany

Abstract (J Korean Assoc Oral Maxillofac Surg 2013;39:3-8)

Reconstruction of oral soft-tissue defects following resection of oral carcinomas can be achieved using various techniques including microsurgical tissue transfer. However, there seems to be a role for regional or local flaps. Small to medium-size defects can be functionally reconstructed with the platysma myocutaneous flap as an excellent choice particularly in medically compromised patients not being eligible for free tissue transfer. The present paper reviews the indication, surgical technique, and complications following reconstruction of defects of the oral cavity with the platysma myocutaneous flap.

Key words: Oral carcinoma, Functional reconstruction, Platysma myocutaneous flap, Head and neck neoplasms

[paper submitted 2012. 9. 5 / accepted 2012. 9. 9]

I. Introduction

Surgical resection of tumors of the oral cavity and oropharynx will result in severe functional impairment of the individual patient. Therefore adequate functional reconstruction of both hard and soft tissue defects plays a key role in the surgical treatment of head and neck tumors. Nowadays various surgical methods of reconstruction are available, including primary closure of mucosal defects, skin grafting, tongue flaps, pedicled myocutaneous island flaps, and free flaps. During the last three decades tremendous progress has been made in head and neck reconstructive surgery by introducing microvascular free tissue transfer from various donor sites for restoring extensive defects in the head and neck area^{1,2}. However, as most patients with head and neck tumours often have medical comorbidities a proper and critical preoperative assessment of the risk of postoperative complications is essential before such patients are selected for

extensive oncological and reconstructive surgery³⁻⁵. Therefore pedicled flaps might be indicated in specific cases resulting in comparable functional outcome⁶.

Pedicled myocutaneous flaps such pectoralis major flap, latissimus dorsi flap, trapezius flap also have a long tradition in head and neck reconstructive surgery, but they have been left for specific cases. Historically the use of the platysma myocutaneous flap can be traced back to 1887, when Robert Gersuny⁷, a surgeon from Austria described a reconstruction of a full-thickness cheek defect with a cervical skin/platysma flap which was rotated inward to provide a new lining for the buccal mucosa. This was probably the first description of the platysma myocutaneous flap used for head and neck reconstruction. However, it was not until 1978 that the platysma myocutaneous flap was introduced by Futrell et al.⁸ as an attractive reconstructive option with several advantages. Reviewing the literature shows large variety of indications to use the platysma myocutaneous flap for reconstruction of defects of the anterior floor of mouth, tongue, cheek, oropharynx, and facial skin⁹⁻³².

Detailed anatomic studies described the vascular anatomy with the submental artery as the predominant blood supply to the platysma muscle as well as to the skin paddle³³⁻³⁵. Apparently the issue of sacrificing the facial vessels is critical as far as flap survival is concerned, although there is some debate on this matter^{11,15,18}. McGuirt et al.¹¹ and Ruark et al.¹⁵ have stated that preservation of the facial artery is not a

André M. Eckardt

*Department of Cranio-Maxillofacial Surgery, Faculty of Dentistry, Hannover Medical University, Carl-Neuberg-Strasse 1, 30625 Hannover, Germany
TEL: +49-511-532-4879 FAX: +49-511-532-8879
E-mail: eckardt.andre@mh-hannover.de*

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2013 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.

prerequisite for survival of the platysma myocutaneous flap.

As the flap can be either used superiorly or posteriorly-based, it can provide an adequate amount of pliable soft tissue that is sufficient for reconstruction of small to medium-sized skin or mucosal defects of the facial skin, oral cavity, and pharynx³⁶. The platysma myocutaneous flap can be easily harvested and has low donor site morbidity, but has not gained widespread acceptance among head and neck surgeons because of problems with flap vascularity. There is still some reluctance to use the flap in view of reported flap-related complication rates in the range of 10-45%^{9,11,15,18,20,25,37}. Loss of skin paddle or formation of orocutaneous fistula have been described as main problem due to ischemic flap failure or venous congestion²⁰. Both arterial supply and venous drainage may be compromised during surgical dissection, as well as by previous neck dissection, scar formation, or previous radiotherapy¹⁸.

Although microvascular free tissue transfer using various donor sites is an established routine procedure for the majority of oncologic head and neck defects, the use of the platysma myocutaneous flap has been incorporated as a valuable alternative at our institution for selected indications. For medically compromised patients the platysma flap can be recommended as first choice with a predictable functional outcome^{12,38}. This report aims to illustrate clinical experience of intraoral soft tissue reconstruction with the superiorly pedicled platysma myocutaneous flap in a large number of oncologic patients and to critically review the existing literature on this topic.



Fig. 1. The intraoperative site with incision of the skin island during harvesting of a cranially pedicled platysma myocutaneous flap.

André M. Eckardt: Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013

II. Patients and Methods

From 2004 to 2012, 45 patients received a platysma myocutaneous island flap in a single stage procedure at the Department of Cranio-Maxillofacial Surgery, Hannover Medical University. The patient population consisted of 31 men and 14 women, with a mean age of 61.4 years (range 39 to 84 years). Except for one patient with chronic osteomyelitis and intraoral bone exposure, the majority of patients (n=41) were diagnosed with squamous cell carcinoma of the oral cavity. Three patients were diagnosed with carcinomas of minor salivary glands. All tumors were located in the oral cavity. Areas of resection included the anterior and lateral floor of mouth, tongue, and retromolar trigone. Primary tumor resection was combined with selective neck dissection level I-III ipsilateral or bilateral if indicated. Adjuvant treatment included postoperative radiotherapy in 3 patients (6.8%). No preoperative chemotherapy or chemoradiation was administered.

1. Surgical technique

Before flap harvesting the anticipated intraoral defect is measured. The outline of the platysma flap is initially made as an ellipse with the inferior margin low in the neck above the clavicle. Typically the dimensions of the skin island are 6 × 10 cm the maximum. If a standard apron-type flap is raised for a neck dissection, the inferior margin of that flap will be



Fig. 2. Following an apron-type incision a skin flap will be dissected up to the mandible leaving the platysma muscle with the skin island down.

André M. Eckardt: Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013




Fig. 3. Intraoral view of a cT2 carcinoma of the left lateral tongue border in a 75-year old woman (A). Intraoperative site following transoral resection of the carcinoma (B). A pedicled platysma myocutaneous flap is dissected (C) and rotated superiorly in to the intraoral defect of left tongue (D). Intraoperative site immediately after defect coverage of the left tongue border shows well vascularised pedicled platysma flap (E).

André M. Eckardt: *Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013*




Fig. 4. Cervical skin with acceptable scar of the lower neck one year after surgery.

André M. Eckardt: *Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013*

the inferior margin of the platysma flap.(Fig. 1) The inferior skin incision will be made through the platysma muscle, whereas the superior skin incision will pass only the skin. Subsequently a skin-only flap will be raised superior to the




Fig. 5. Intraoral situation one year after reconstruction using platysma myocutaneous flap with well vascularised skin island and unrestricted tongue mobility.

André M. Eckardt: *Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013*

skin paddle up to the mandible, leaving the platysma down. (Fig. 2) Whenever feasible the flap pedicle covered the full width of the platysma muscle at least 4 cm. The plane of elevation should include the sternocleidomastoid fascia, as well as the external jugular vein to provide adequate venous drainage. After tumor resection and neck dissection, the

platysma myocutaneous flap is rotated superiorly and 180 degrees into the intraoral defect and sutured into position. (Figs. 3. A-E) During rotation care must be taken to avoid excessive stretching, compression, or twisting of the flap pedicle. To avoid compression a sufficiently wide tunnel under the mandible should be dissected before the flap is rotated into the intraoral defect area. The neck incision is closed primarily. A long-term follow-up at 1 year shows an acceptable scar in the lower neck without any functional impairment.(Fig. 4) Intraorally the skin paddle has healed in without problems and allows for adequate tongue mobility.(Fig. 5)

III. Results

The patient population was evaluated for immediate and delayed postoperative complications, with follow-up ranging from 6 months to eight years. The tumors consisted of squamous cell carcinoma of the oral cavity in 88% of the cases (n=40) and the oropharynx in 2.3% (n=1). In 3 patients (6.8%) tumors were diagnosed as salivary gland carcinomas (adenoid-cystic carcinoma n=1, mucoepidermoid carcinoma n=2). Tumor size ranged from pT1 to pT4 (pT1: 26 (59%), pT2: 11 (25%), pT3: 1 (2.2%), and pT4: 3 (6.8%). There were 3 patients (6.8%) diagnosed with pTis. The nodal status was as follows: pN0 39 (88.6%), pN1 3 (6.8%), and pN2b 2 (4.5%). With regard to the anatomical site the floor of mouth was involved in 20 patients (44.4%), the cheek in 4 patients (8.9%), the tongue in 11 patients (24.4%), the alveolar crest in 8 patients (17.8%), and the oropharynx in 1 patient (2.2%). The size of the skin island varied in size from 3x5 cm to 5x7 cm. Flap harvesting was performed from one side of the neck in 43 patients (95.6%), a bilateral platysma flap was raised in 2 patients (4.4%). In all patients the reconstruction was performed as a single-stage procedure together with tumor resection using a transoral/transcervical approach. In 6 patients (13.3%) mandibular continuity resection with immediate alloplastic mandibular reconstruction using titanium 2.4 mm reconstruction plates was performed.

Selective neck dissection (level I-III) was performed in 44 patients (97.8%), as an ipsilateral procedure in 27 patients (61.3%) and as bilateral procedure in 14 patients (31.8%). The donor site was closed primarily and resulted in a good cosmetic result in the neck. Flap complications were noted in 4 patients (8.9%) with total flap loss in 3 patients (6.6%). Partial flap loss was managed by local measures, such as surgical wound debridement of necrotic tissue. A total of 4 patients (9%) developed tumor recurrences 6 months to

1 year after the primary resection. These patients received salvage surgery and adjuvant chemoradiation. However, 3 of these patients developed progressive disease after salvage treatment and died of their disease.

IV. Discussion

A major goal of oral and facial reconstruction after tumor resection is to restore adequate form and function thereby improving patients quality of life. During the last five decades important milestones in head and neck reconstructive surgery have been achieved including regional pedicled flaps and of course microvascular free tissue transfer. According to the literature pedicled flaps are mostly reserved for medium-sized defects or as a salvage option for tumor recurrences⁶. The platysma myocutaneous flap is a pedicled cervical flap initially described by the Austrian surgeon Gersuny⁷ in 1887; however this type of pedicled flap was then neglected for a long time before being introduced for intraoral soft tissue reconstruction by Futrell et al.⁸ in 1978. This flap is an attractive reconstructive option and offers several advantages: it is thin and pliable with no excessive bulk and easy to harvest. The near location to the oral cavity allows for reconstruction of various intraoral areas such as floor of mouth, tongue, pharyngeal wall^{9-12,14-17}. Despite these advantages the platysma myocutaneous flap has not gained widespread acceptance most probably due to a lack of its reliability and a high rate of complications (18-45%). The flap complications rate in our series was 8.9% with a total flap loss rate of 6.6% which is comparable to

Table 1. Complication rate of platysma myocutaneous flap reported in selected case studies

Author	No. of patients	Rate of flap loss (%)
Futrell et al. ⁸	14	7.1
Cannon et al. ¹⁰	22	16.7
Coleman et al. ⁹	24	29.6
Ozgen et al. ¹⁶	4	25
Ruark et al. ¹⁵	41	7.3
Esclamado et al. ²⁵	12	8.3
Ozcelik et al. ²⁷	20	0
Verschuur et al. ¹⁸	44	2.3
Koch ¹⁷	34	17.6
Peng et al. ³⁶	48	6.25
Grützenmacher et al. ²³	25	12
Pagani et al. ²¹	10	20
Puxeddu et al. ²²	11	0
Wang et al. ²⁴	10	0
Koch et al. ²⁰	70	10
Tosco et al. ¹⁹	91	13

André M. Eckardt: Platysma myocutaneous flap - its current role in reconstructive surgery of oral soft tissue defects. J Korean Assoc Oral Maxillofac Surg 2013

reported flap loss rates in larger case series. (Table 1) Szudek and Taylor³⁷ reported a flap complication rate of 33.7% in their meta-analysis including a total of 190 patients from 16 published reports. They further concluded that postoperative complications were not associated with age, sex, or preoperative radiotherapy, but were associated with recipient site and tumor stage³⁷. Verschuur et al.¹⁸ and Szudek and Taylor³⁷ reported a failure rate with total flap loss of 2.5-5.0%. A meticulous dissection technique with attention to the specific vascular supply of the platysma flap can avoid such complications as pointed out by Hurwitz et al.³³.

Relative contraindications to harvest the platysma flap are facial artery ligation, prior radiation or neck dissection. These factors are probably associated with impaired postoperative perfusion and were thought to be responsible for an increased flap failure rate. The blood supply of the platysma myocutaneous flap is extensive with the primary supply coming from the submental artery which branches from the facial artery. Additional supply comes from the transverse cervical, thyroid, occipital and posterior auricular vessels³⁹. The venous drainage is critical to this flap and, therefore, tension or compression should be avoided. The venous drainage pattern is vertical and the flap is more likely at risk due to venous congestion than arterial insufficiency³⁸. We agree with authors such as Ruark et al.¹⁵ and Verschuur et al.¹⁸ that the external jugular vein should be preserved and included in the flap pedicle whenever possible. Previous radiotherapy to the neck seems to be a relative contraindication for the platysma flap, although there is no conclusive evidence. In our series only 1 patient (2.2%) had previous radiation to the neck and developed no postoperative complications. Similarly Cannon et al.¹⁰ didn't observe any major complication in their series in which 5 patients received radiation with doses of 43 Gy or more.

There is controversy about the survival of the superiorly-based platysma flap when the facial artery has been ligated. Unfortunately not all studies provide details of the site of facial artery ligation or if was ligated at all. Coleman et al.⁹ emphasized that the platysma flap cannot be raised in standard radical neck dissection because the facial artery is transected which compromises flap survival. In contrast McGuirt et al.¹¹ found that only 1 in 20 patients with facial artery ligation had partial loss of the skin paddle. The flap did also well in patients in spite of preoperative radiation or previous neck dissection. McGuirt's clinical observations are supported by the detailed anatomical studies performed by Hurwitz et al.³³ who came to the conclusion that an intact

facial artery is not crucial for survival of the platysma flap. Interestingly, some studies have emphasized that the success also depends on careful preoperative patient selection and careful surgical technique^{12,15,17,18}. In their review Szudek and Taylor³⁷ reported a success rate of 95% in 190 patients. This is comparable to success rates of free tissue transfer being in the range of 91-99%¹⁻³. Intraoperative situations such as an extended radical neck dissection with multiple vessel ligations must be included in the surgical planning^{11,18,25}. We would agree with Tosco et al¹⁹. that the main contraindications to use the platysma myocutaneous flap are previous surgical scarring or previous neck radiation therapy, the need of tissue bulk in the defect zone, defect size larger than 7×10 cm, and when there is radiologic suspicion or evidence of neck metastases with extracapsular spread.

V. Conclusion

Although free tissue transfer has gained widespread acceptance as a standard technique for restoring extensive oncological defects in the head and neck region, pedicled regional flaps such as the platysma myocutaneous flap should not be neglected since they are a valuable alternative for smaller to medium size defects and particularly useful in medically compromised patients. Watchful observation of indications and contraindication together with careful patient selection are crucial factors to be considered in the surgical planning process. Taking the mentioned contraindications and limitations into account, the platysma myocutaneous flap is an elegant and attractive reconstructive option for a variety of intraoral locations and should be included in the repertoire of any cranio-maxillofacial surgeon involved in the surgical treatment of oral and oropharyngeal malignancies.

References

1. Nakatsuka T, Harii K, Asato H, Takushima A, Ebihara S, Kimata Y, et al. Analytic review of 2372 free flap transfers for head and neck reconstruction following cancer resection. *J Reconstr Microsurg* 2003;19:363-8.
2. Eckardt A, Meyer A, Laas U, Hausamen JE. Reconstruction of defects in the head and neck with free flaps: 20 years experience. *Brit J Oral Maxillofac Surg* 2007;45:11-5.
3. Singh B, Cordeiro PG, Santamaria E, Shaha AR, Pfister DG, Shah JP. Factors associated with complications in microvascular reconstruction of head and neck defects. *Plast Reconstr Surg* 1999; 103:403-11.
4. Genden EM, Rinaldo A, Suárez C, Wei WI, Bradley PJ, Ferlito A. Complications of free flap transfers for head and neck reconstruction following cancer resection. *Oral Oncol* 2004;40:979-84.

5. Shestak KC, Jones NF, Wu W, Johnson JT, Myers EN. Effect of advanced age and medical disease on the outcome of microvascular reconstruction for head and neck defects. *Head Neck* 1992;14:14-8.
6. Schusterman MA, Kroll SS, Weber RS, Byers RM, Guillaumondegui O, Goepfert H. Intraoral soft tissue reconstruction after cancer ablation: a comparison of the pectoralis major flap and the free radial forearm flap. *Am J Surg* 1991;162:397-9.
7. Gersuny R. Plastischer ersatz der wangenschleimhaut. *Zentralbl Chir* 1887;14:706.
8. Futrell JW, Johns ME, Edgerton MT, Cantrell RW, Fitz-Hugh GS. Platysma myocutaneous flap for intraoral reconstruction. *Am J Surg* 1978;136:504-7.
9. Coleman JJ 3rd, Jurkiewicz MJ, Nahai F, Mathes SJ. The platysma musculocutaneous flap: experience with 24 cases. *Plast Reconstr Surg* 1983;72:315-23.
10. Cannon CR, Johns ME, Atkins JP Jr, Keane WM, Cantrell RW. Reconstruction of the oral cavity using the platysma myocutaneous flap. *Arch Otolaryngol* 1982;108:491-4.
11. McGuirt WF, Matthews BL, Brody JA, May JS. Platysma myocutaneous flap: caveats reexamined. *Laryngoscope* 1991;101:1238-44.
12. Vriens JP. A reappraisal of the platysma myocutaneous island flap. A follow-up study of the functional outcome after reconstruction of the floor of the mouth. *Int J Oral Maxillofac Surg* 1995;24:212-5.
13. Manni JJ, Bruaset I. Reconstruction of the anterior oral cavity using the platysma myocutaneous island flap. *Laryngoscope* 1986;96:564-7.
14. Posnick JC, McCraw JB. Reconstruction of the chin-lower cheek complex with a platysma myocutaneous flap. *J Oral Maxillofac Surg* 1988;46:152-5.
15. Ruark DS, McClairn WC Jr, Schlehaider UK, Abdel-Misih RZ. Head and neck reconstruction using the platysma myocutaneous flap. *Am J Surg* 1993;165:713-8.
16. Ozgen K, Berberoglu U, Altinok M, Ozdemir E, Celen O. Reconstruction of full-thickness cheek defects using platysma myocutaneous flaps. *Eur J Plast Surg* 1993;16:208-11.
17. Koch WM. The platysma myocutaneous flap: underused alternative for head and neck reconstruction. *Laryngoscope* 2002;112:1204-8.
18. Verschuur HP, Dassonville O, Santini J, Vallicioni J, Poissonnet G, Laudoyer Y, et al. Complications of the myocutaneous platysma flap in intraoral reconstruction. *Head Neck* 1998;20:623-9.
19. Tosco P, Garzino-Demo P, Ramieri G, Tanteri G, Pecorari G, Caldarelli C, et al. The platysma myocutaneous flap (PMF) for head and neck reconstruction: a retrospective and multicentric analysis of 91 T1-T2 patients. *J Craniomaxillofac Surg* 2012;40:e415-8.
20. Koch M, Künzel J, Mantsopoulos K, Zenk J, Iro H. Defect closure after oral and pharyngeal tumor resection with the superiorly pedicled myocutaneous platysma flap: indications, technique, and complications. *Eur Arch Otorhinolaryngol* 2012;269:2111-9.
21. Pagani D, Capaccio P, Fontanella W, Sambataro G, Broich G, Pignataro L. Vertical platysma myocutaneous flap reconstruction for T2-staged oral carcinoma. *Anticancer Res* 2007;27:2961-4.
22. Puxeddu R, Dennis S, Ferrelli C, Caldera S, Brennan PA. Platysma myocutaneous flap for reconstruction of skin defects in the head and neck. *Br J Oral Maxillofac Surg* 2008;46:383-6.
23. Grützenmacher S, Steinmeier E, Hosemann W. The use of the platysma myocutaneous flap for reconstruction in the head-neck area -- a retrospective study. *Laryngorhinotologie* 2005;84:733-7.
24. Wang KH, Hsu EK, Shemen LJ. Platysma myocutaneous flap for oral cavity reconstruction. *Ear Nose Throat J* 2010;89:276-9.
25. Esclamado RM, Burkey BB, Carroll WR, Bradford CR. The platysma myocutaneous flap. Indications and caveats. *Arch Otolaryngol Head Neck Surg* 1994;120:32-5.
26. Su T, Zhao YF, Liu B, Hu YP, Zhang WF. Clinical review of three types of platysma myocutaneous flap. *Int J Oral Maxillofac Surg* 2006;35:1011-5.
27. Özçelik T, Aksoy S, Gökler A. Platysma myocutaneous flap: use for intraoral reconstruction. *Otolaryngol Head Neck Surg* 1997;116:493-6.
28. Lazaridis N, Dimitrakopoulos I, Zouloumis L. The superiorly based platysma flap for oral reconstruction in conjunction with neck dissection: a case series. *J Oral Maxillofac Surg* 2007;65:895-900.
29. Baur DA, Helman JI. The posteriorly based platysma flap in oral and facial reconstruction: a case series. *J Oral Maxillofac Surg* 2002;60:1147-50.
30. Ariyan S. The transverse platysma myocutaneous flap for head and neck reconstruction: an update. *Plast Reconstr Surg* 2003;111:378-80.
31. Handa Y, Kitajima T, Takagi N, Yasuoka T, Naitoh K, Tatematsu N, et al. The platysma myocutaneous flap for oral reconstruction. Experience with MacFee's cervical incision. *J Craniomaxillofac Surg* 1989;17:318-22.
32. Mazzola RF, Benazzo M. Platysma flap for oral reconstruction. *Clin Plast Surg* 2001;28:411-9.
33. Hurwitz DJ, Rabson JA, Futrell JW. The anatomic basis for the platysma skin flap. *Plast Reconstr Surg* 1983;72:302-14.
34. Uehara M, Helman JI, Lillie JH, Brooks SL. Blood supply to the platysma muscle flap: an anatomic study with clinical correlation. *J Oral Maxillofac Surg* 2001;59:642-6.
35. Kocer U, Ozdemir R, Ulusoy MG, Uysal A, Sungur N, Sahin B, et al. Anatomy of the platysma muscle and the evaluation of it for the reconstruction of facial defects. *J Craniofac Surg* 2005;16:463-70.
36. Peng LW, Zhang WF, Zhao JH, He SG, Zhao YF. Two designs of platysma myocutaneous flap for reconstruction of oral and facial defects following cancer surgery. *Int J Oral Maxillofac Surg* 2005;34:507-13.
37. Szudek J, Taylor SM. Systematic review of the platysma myocutaneous flap for head and neck reconstruction. *Arch Otolaryngol Head Neck Surg* 2007;133:655-61.
38. Berenholz L, Kessler A, Segal S. Platysma myocutaneous flap for intraoral reconstruction: an option in the compromised patient. *Int J Oral Maxillofac Surg* 1999;28:285-7.
39. Rabson JA, Hurwitz DJ, Futrell JW. The cutaneous blood supply of the neck: relevance to incision planning and surgical reconstruction. *Br J Plast Surg* 1985;38:208-19.