

Research Article

The Correlation between the Triglyceride-Glucose Index and Coagulation Markers in Patients with Recent Acute Myocardial Infarction

Daniel Košuta ^{1,2}, Marko Novakovič^{1,2}, Mojca Božič Mijovski ^{2,3} and Borut Jug ^{1,2}

¹Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

²Department of Vascular Diseases, University Medical Centre Ljubljana, Ljubljana, Slovenia

³Faculty of Pharmacy, University of Ljubljana, Ljubljana, Slovenia

Correspondence should be addressed to Daniel Košuta; daniel.kosuta@kclj.si

Received 27 June 2022; Revised 30 September 2022; Accepted 5 October 2022; Published 20 October 2022

Academic Editor: Serena Del Turco

Copyright © 2022 Daniel Košuta et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Metabolic abnormalities and hypercoagulability seem to have an important predictive role in patients with coronary artery disease (CAD). The triglyceride-glucose (TyG) index has emerged as a good marker for metabolic syndrome with predictive value for cardiovascular events. Overall haemostatic potential (OHP) is a reliable global haemostatic assay to identify hypercoagulability in CAD patients. The aim of our study was therefore to evaluate a possible correlation between the TyG index and haemostatic derangements in patients with CAD. **Methods.** Consecutive patients referred for the first follow-up visit after acute myocardial infarction between December 1, 2018, and March 31, 2020, and did not meet exclusion criteria were included. We determined OHP, overall coagulation potential (OCP), overall fibrinolytic potential (OFP), fibrinogen, D-dimer, and von Willebrand factor from peripheral blood samples. The TyG index was calculated with the previously described and validated formula. Linear regression models were constructed for the multivariate analysis. **Results.** A total of 117 patients (mean age 56 ± 10 years, 20% women) were included. A correlation was found between TyG index and OCP ($r = 0.229, p = 0.026$), TyG index and OHP ($r = 0.202, p = 0.050$), and TyG index and fibrinogen ($r = 0.271, p = 0.005$). In the multivariate model which accounted for sex, age, and BMI, the correlation between TyG index and OCP (R^2 0.108; ANOVA for regression $p = 0.035$; beta 2.08 [0.79-4.01], $p = 0.042$) and between TyG index and fibrinogen (R^2 0.11; ANOVA for regression $p = 0.015$; beta 0.35 [0.08-0.62], $p = 0.012$) emerged as statistically significant. **Conclusion.** The TyG index, a marker of metabolic syndrome, has a strong correlation with a hypercoagulability state in CAD, as determined by the OCP and higher fibrinogen levels. Our findings suggest that metabolic syndrome may be an important driver of atherothrombotic risk in patients with CAD.

1. Introduction

Coronary artery disease (CAD) is a form of atherosclerotic vascular disease associated with significant morbidity and mortality despite mounting efforts of prevention, early diagnosis, and aggressive management [1]. CAD is characterized by progressive atherosclerotic plaque build-up, which is prone to rupture, and atherothrombotic coronary artery occlusion, yielding myocardial infarction. On the one hand, atherosclerosis progression is associated with cardiovascular risk factors. While some risk factors, such as smoking, have been stagnating or declining in

western populations, others—especially cardiometabolic risk factors, such as obesity, diabetes, and dyslipidaemia—have been on the rise [2, 3]. On the other hand, atherothrombotic events are associated with a procoagulant state [4], which may discern stable chronic CAD from unstable acute coronary syndromes.

Obesity and insulin resistance are notable hallmarks of cardiometabolic risk, suggesting underlying derangements in glucose and lipid metabolism. However, recent studies suggest that cardiometabolic abnormalities can be detected in nonobese individuals as well [5, 6], and identification of

cardiometabolic risk beyond apparent clinical characteristics—possibly through biomarkers—is being pursued. One such biomarker is the triglyceride-glucose (TyG) index, which has been proposed as a marker of metabolic syndrome and has been associated with carotid atherosclerosis, coronary artery calcification, and higher risk of cardiovascular disease [7–9]. Moreover, recent studies have identified the TyG index as an independent predictor of coronary calcification progression, of clinical outcomes in patients with stable CAD and premature CAD, and of cardiovascular outcomes in acute coronary syndromes [7, 10–13].

In addition to risk factors-driven coronary atherosclerosis progression, haemostatic derangements play an important role in the development of myocardial infarction. CAD has been associated with a procoagulant state with mounting evidence of high haemostatic factor activity—such as increased fibrinogen levels, a procoagulant platelet response, and thrombogenicity [4, 14–17]. Procoagulant activity is associated with both the severity of chronic CAD and the likelihood of its progression to myocardial infarction [4, 18]. Since coagulation is a complex and multifactorial process, global haemostatic essays—as opposed to the measurement of individual procoagulant and fibrinolytic factors—have recently gained popularity [19]. Overall haemostatic potential (OHP) is a global haemostasis essay [20, 21], which has been validated in healthy individual as well as various groups of patients [22–24]. In CAD patients, OHP detected a hypercoagulability state, driven primarily by impaired fibrinolysis [23, 25].

Clinical features of the metabolic syndrome are associated with a prothrombotic state. Several coagulation markers are increased, and fibrinolysis is impaired in patients with metabolic syndrome, which may contribute to the adverse atherothrombotic pathophysiology and high-risk features of CAD in patients with dysmetabolic profiles [26, 27]. In the present study, we sought to appraise a possible association between the TyG index, a marker of the metabolic syndrome beyond traditional clinical features, and the OHP, an essay for haemostatic derangements, in patients with CAD.

2. Methods

2.1. Study Design and Population. Consecutive patients referred for the first follow-up visit after acute myocardial infarction between December 1, 2018, and March 31, 2019, were screened for inclusion. Inclusion criteria were 18–70 years of age and recent myocardial infarction (within 90 days from inclusion). Patients with advanced heart failure (New York Heart Association classes III and IV), patients on anticoagulation treatment, and pregnant women were excluded. The study was conducted in accordance with the Declaration of Helsinki (1964) and approved by the National Medical Ethics Committee (approval number 0120-77/2020/3).

After signing informed consent, patients were included in the study and had a fasting (>12 hours after the last meal) blood sample drawn from the cubital vein; blood was collected into 4.5 mL 0.109 M sodium citrate tubes. Citrated blood was then centrifuged for 20 minutes at 2000 × g; the extracted plasma was then stored at –70°C for further analysis.

We collected baseline data, cardiovascular risk factors, and laboratory tests (glucose (mg/dL), total cholesterol (mg/dL), HDL cholesterol (mg/dL), LDL cholesterol (mg/dL), triglycerides (mg/dL), fibrinogen (g/L), D-dimer (μg/L), and von Willibrand factor (%)). The researcher who collected the aforementioned data was blinded to the coagulation test results.

Overall haemostatic potential (OHP, abs-sum), overall coagulation potential (OCP, abs-sum), and overall fibrinolytic potential (OFP, %) were determined as previously described [20, 23] using bovine thrombin (Sigma Chemical Company, St. Louis, USA) and recombinant tissue-type plasminogen activator (Actilyse 0.1 mL/mL, Boehringer Ingelheim, Germany) by absorbance measurements at 405 nm in 1-minute intervals for 40 minutes. Areas under the curve were constructed for OHP and OCP with the obtained measurements; OFP was calculated as the difference between the two aforementioned areas: $OFP = [(OHP - OCP)/OCP] \times 100 (\%)$.

The TyG index was calculated with the previously described and validated formula: $TyG\ index = \ln [triglycerides\ (mg/dL) \times plasma\ glucose\ (mg/dL)/2]$ [11].

2.2. Statistical Analysis. Baseline characteristics are expressed as the mean (±standard deviation) for normally distributed continuous variables, as median (interquartile range) for nonnormally distributed continuous variables, and as frequency (%) for categorical variables. Between-group differences were assessed by the *t*-test for normally distributed variables and by the Mann–Whitney *U* test for nonnormally distributed variables; proportions were compared using the χ^2 test. Correlations were explored with Pearson's tests. Linear regression models were constructed for the multivariate analysis; results are expressed as beta with corresponding 95% confidence intervals (CI). A 2-tailed $p < 0.05$ was considered significant. Statistical analysis was carried out using SPSS Statistics version 23 (SPSS Inc, Chicago, USA).

3. Results

A total of 143 patients were assessed for eligibility; 26 patients were excluded (20 did not meet the inclusion criteria and 6 declined to participate), 117 patients after a recent myocardial infarction were included, and no participant had a previous history of arterial or venous thrombotic events. Baseline characteristics and risk factor presence are reported in Table 1. Medical therapy, prescribed at the time of the event, is listed in Table 1. The median triglyceride value was 115 mg/dL (IQR 88), median glucose value was 100 mg/dL (IQR 14), and median TyG index value was 8.68 (IQR 0.5). We divided participants according to the median value of the TyG index and compared the two groups: BMI and total cholesterol were found to be significantly increased and HDL cholesterol to be significantly decreased in the group with the TyG index above median (Tables 1 and 2). A comparison between the lowest and highest tertile of the TyG index was performed (Table 3); a significant difference was found in diabetes prevalence and a trend to significance for BMI. Among coagulation markers, OCP and OHP were found to be significantly higher in patients with the TyG index above median (Table 2).

TABLE 1: Baseline demographic and risk factors.

	All (<i>n</i> = 117)	TyG below median (<i>n</i> = 58)	TyG above median (<i>n</i> = 59)	<i>p</i> value
Women*	23 (19.7)	12 (20.7)	11 (18.9)	0.808
Age (years)**	56 (10.3)	57.6 (10.8)	55.1 (10.0)	0.223
BMI (kg/m ²)**	28.8 (4.6)	28.1 (4.0)	29.9 (4.8)	0.013
Event				
STEMI*	71 (60.7)	35 (60.3)	36 (61.0)	0.554
NSTE-ACS*	46 (39.3)	23 (39.7)	23 (39.0)	
Arterial hypertension*	86 (73.5)	44 (75.8)	42 (71.2)	0.495
Diabetes mellitus*	10 (8.5)	4 (6.9)	6 (10.2)	0.093
Dyslipidaemia*	75 (64.1)	37 (63.8)	38 (64.4)	0.423
Family history*	46 (39.3)	24 (41.4)	22 (37.3)	0.695
Smoker*	54 (46.2)	25 (43.1)	29 (49.2)	0.181
Antiplatelet drugs				
ASA*	116 (99.1)	58 (100)	58 (98.3)	
Clopidogrel*	12 (10.3)	7 (12.1)	5 (8.5)	0.940
Prasugrel*	25 (21.4)	12 (20.7)	13 (22.0)	0.824
Ticagrelor*	73 (62.4)	36 (62.1)	37 (62.7)	
ACE inhibitor*	91 (77.8)	47 (81.0)	44 (74.6)	0.615
Beta-blocker*	95 (81.2)	46 (79.3)	49 (83.1)	0.580
Statins*	117 (100)	58 (100)	59 (100)	
Rosuvastatin*	104 (88.9)	53 (91.4)	51 (86.4)	0.761
Atorvastatin*	13 (11.1)	5 (8.6)	8 (13.6)	

ASA: acetylsalicylic acid; BMI: body mass index; NSTE-ACS: non-ST-elevation acute coronary syndrome; SD: standard deviation; STEMI: ST-elevation myocardial infarction; TyG: triglyceride-glucose index; *: results displayed as number (%); **: results displayed as mean (standard deviation).

TABLE 2: Clinical, biochemical, and coagulation markers.

	All (<i>n</i> = 117)	TyG below median (<i>n</i> = 58)	TyG above median (<i>n</i> = 59)	<i>p</i> value
Systolic blood pressure (mmHg)*	124.1 (7.1)	122.8 (15.8)	125.7 (17.6)	0.387 ^a
Diastolic blood pressure (mmHg)*	78.8 (10.3)	77.4 (10.7)	80.5 (10.2)	0.132 ^a
Total cholesterol (mg/dL)**	123.7 (108.3, 150.8)	119.9 (104.4, 143.1)	127.6 (112.1, 158.6)	0.045 ^b
HDL cholesterol (mg/dL)**	42.5 (34.8, 50.3)	42.5 (38.7, 54.1)	38.7 (34.8, 46.4)	<0.001 ^b
LDL cholesterol (mg/dL)**	58 (42.5, 73.5)	54.1 (46.4, 69.6)	58 (42.5, 81.2)	0.617 ^b
OCP, abs-sum**	22.6 (20.4, 26.6)	22.4 (20.3, 25.5)	24.3 (20.6, 29.2)	0.040 ^b
OHP, abs-sum**	8.0 (6.5, 9.8)	7.5 (5.6, 9.1)	8.3 (6.9, 9.8)	0.041 ^b
OFF (%)**	66.0 (60.0, 71.0)	66.0 (60, 73)	66 (60.3, 69)	0.517 ^b
Fibrinogen (g/L)**	3.3 (2.9, 3.9)	3.2 (2.9, 3.9)	3.6 (3.0, 4.0)	0.092 ^b
D-dimer (μg/L)**	304.5 (197.0, 478.0)	293.0 (215.5, 404.5)	333.0 (195.0, 572.5)	0.414 ^b
von Willebrand factor (%)**	150.5 (120.5, 192.5)	141.0 (115.0, 180.0)	169.0 (124.5, 205.0)	0.146 ^b
Time from AMI to visit (days)**	58.0 (47.5, 79.5)	58.0 (49.0, 82.0)	56.0 (47.0, 79.0)	0.959 ^b

AMI: acute myocardial infarction; OCP: overall coagulation potential; OFF: overall fibrinolytic potential; OHP: overall haemostatic potential; TyG: triglyceride-glucose index; *: results displayed as mean (SD); **: results displayed as median (25th, 75th percentile). ^a*t*-test; ^bMann-Whitney *U* test.

A correlation analysis using Pearson's test showed a correlation between TyG index and total cholesterol ($r = 0.382, p < 0.001$), TyG index and HDL cholesterol ($r = -0.364, p < 0.001$), TyG index and LDL cholesterol ($r = 0.199, p = 0.038$), TyG index

and OCP ($r = 0.229, p = 0.026$), TyG index and OHP ($r = 0.202, p = 0.049$), and TyG index and fibrinogen ($r = 0.271, p = 0.005$). No correlation was found between TyG index and OFF ($r = -0.035, p = 0.738$), TyG index and D-

TABLE 3: Baseline demographic and risk factor comparison between lowest and highest tertile of TyG index.

	Lowest tertile TyG index ($n = 37$)	Highest tertile TyG index ($n = 36$)	p value
Women*	9 (24)	8 (22)	0.309
Age (year)**	57.2 (10.7)	54.9 (9.9)	0.339
BMI (kg/m^2)**	28.1 (4.5)	30.4 (5.5)	0.057
Event			
STEMI*	21 (56.8)	24 (66.7)	0.665
NSTE-ACS*	16 (43.2)	12 (33.3)	
Arterial hypertension*	29 (78.4)	27 (75.0)	0.936
Diabetes mellitus*	1 (2.7)	8 (22.2)	0.004
Dyslipidaemia*	23 (62.2)	25 (69.4)	0.793
Family history*	17 (45.9)	42 (38.5)	0.401
Smoker*	15 (40.5)	19 (52.8)	0.564

BMI: body mass index; NSTE-ACS: non-ST-elevation acute coronary syndrome; SD: standard deviation; STEMI: ST-elevation myocardial infarction; TyG: triglyceride-glucose index; *: results displayed as number (%); **: results displayed as mean (standard deviation).

dimer ($r = 0.077$, $p = 0.435$), and TyG index and von Willebrand factor ($r = 0.019$, $p = 0.846$).

In the multivariate model which accounted for sex, age, and BMI, the correlation between TyG index and total cholesterol (R^2 0.184; ANOVA for regression $p < 0.001$; beta 0.362 [0.117–0.329], $p < 0.001$), between TyG index and HDL cholesterol (R^2 0.146; ANOVA for regression $p < 0.001$; beta -0.357 [-0.878–-0.238], $p = 0.001$), between TyG index and OCP (R^2 0.108; ANOVA for regression $p = 0.035$; beta 2.08 [0.79–4.01], $p = 0.042$), and between TyG index and fibrinogen (R^2 0.11; ANOVA for regression $p = 0.015$; beta 0.35 [0.08–0.62], $p = 0.012$) emerged as statistically significant (Figure 1).

To further explore the association between coagulation biomarkers and TyG index, a ROC analysis was performed. Both OCP and fibrinogen showed a modest predictivity for the TyG index (ROC area 0.60, 95% CI 0.480–0.713 and ROC area 0.59, 95% CI 0.484–0.701, respectively).

4. Discussion

In the present study, we have shown a significant association between the TyG index (a biomarker of cardiometabolic risk) and haemostatic abnormalities in patients after myocardial infarction. The finding suggests a possible link between cardiometabolic and coagulation derangements beyond traditional clinical features of the metabolic syndrome. To the best of our knowledge, ours is the first study to appraise a strong positive correlation between a procoagulant state (OCP and fibrinogen levels) and the TyG index, suggesting a pathophysiological interplay between cardiometabolic abnormalities and atherothrombotic potential in patients after a recent coronary event.

The metabolic syndrome is a known predictor of cardiovascular events in patients with established CAD [28–30]. Recent research has identified the TyG index as a reliable marker of metabolic syndrome and a strong predictor of CAD, premature CAD, and adverse events in CAD patients [10–13, 28]. In our study, values of the TyG index were comparable to those from previous studies in patients with CAD,

wherein TyG index values were also associated with an increased event risk [11]. Our data complete these observations by proposing a pathophysiological link between the TyG index and haemostatic derangements; as the majority of CAD events represent thrombotic complications of atherosclerosis, an association between cardiometabolic and haemostatic derangements provides an important further insight in the unfavourable atherothrombotic prognostic role of the metabolic syndrome.

Previous studies have shown that insulin resistance and metabolic syndrome may confer a procoagulant state through several mechanisms. Adipose tissue may impair fibrinolysis through direct production of coagulation factors (such as PAI-1 and thrombin activatable fibrinolysis inhibition), dysmetabolism-associated nonalcohol liver steatosis may alter the hepatic synthesis of haemostatic factors, and adipokines may affect platelet function; insulin resistance-associated low-grade inflammation and endothelial dysfunction may modulate the genetic expression, activity, and interactions of haemostatic factors [27, 31–33]. In our study, we detected a strong correlation between known markers of metabolic disorder and TyG index and the association of the TyG index with a procoagulant state. The traditional assessment of a procoagulant state relies on measuring individual coagulant factors but may fail to capture the complexity of the overall haemostatic processes, and even in our study, D-dimer and von Willebrand factor did not show a significant correlation with the TyG index, but a strong correlation, independent of BMI, was found for fibrinogen, a known marker of coagulation. The finding supports coagulation activation in patients with metabolic syndrome and CAD. Global haemostatic assays for net yields of coagulation and fibrinolysis have recently been proposed as more suitable indicators for overall haemostasis appraisal [19]. They have been extensively validated in diverse populations, including patients with CAD [20, 24, 25, 34]. In our study, OCP levels—indicating hypercoagulability—were comparable to other studies of patients with CAD [23, 25] and significantly increased in patients with an elevated TyG index. Hence, our results corroborate that CAD confers a




FIGURE 1: (a) Correlation between triglyceride-glucose index and overall coagulation potential; (b) correlation between triglyceride-glucose index and fibrinogen. TyG: triglyceride-glucose index; OCP: overall coagulation potential.

procoagulant state, which is especially pronounced in the CAD subpopulation with metabolic syndrome. Moreover, the association between TyG index and procoagulation was found to be independent of BMI. While clinical features of the metabolic syndrome (i.e., obesity, blood pressure, and metabolic dyslipidaemia) represent pivotal characteristics for the diagnosis and management of the condition, preclinical cardiometabolic biomarkers—such as the TyG index—may provide an independent, additional tool for early and meaningful detection of dysmetabolic derangements.

Hence, our results suggest that the TyG index is an independent marker of both cardiometabolic and haemostatic derangements. In this respect, an independent association of the TyG index and a procoagulant state in patients after a recent myocardial infarction may confer an important starting point for future risk assessment strategies for patients with CAD.

Limitations of our study primarily pertain to its realisation in a single centre and to its cross-sectional observational design and the study of surrogate biomarkers. Firstly, the single-centre nature of our study is somewhat counterbalanced by the fact that it was carried out at a national referral centre gathering a wide and varied patient population. In fact, the characteristics and risk factor presence in our participant group are similar to data from the literature, and hence, our results can be fairly generalized to the population of patients with CAD after a recent myocardial infarction. Secondly, the cross-sectional design can carry some measurement bias. Our measurements were in line with those reported in the literature for participants with CAD. Thirdly, we were able to include in the analysis only BMI as a marker of excess body fat. Including waist circumference or ectopic fat deposition would have added a deeper layer of understanding about the interplay between various components of the metabolic syndrome and coagulation markers. Fourthly, ours was a study to find an association; while it provides interesting and potentially important insights into the pathophysiological interplay between cardiometabolism and haemostasis in CAD, further prospective cohort studies are needed to establish a potential clinical relevance of our findings.

5. Conclusions

In conclusion, the TyG index represents an important and potentially relevant pathophysiological link between cardiometabolic and haemostatic derangements in patients with CAD. The TyG index is a reliable marker of metabolic syn-

drome and has a strong correlation with a procoagulant state. Our findings suggest that metabolic syndrome may be an important driver of atherothrombotic risk in patients with CAD.

Data Availability

The data used to support the findings of this study are available upon request to the corresponding author.

Disclosure

The authors alone are responsible for the content and writing of the paper.

Conflicts of Interest

The authors declare no conflict of interest.

Acknowledgments

We thank all the participants in this study, all nurses, and laboratory and administrative staff for generously helping us in this research. This research was cofunded by the University Medical Centre Ljubljana's intern tertiary grant and by the Slovenian Research Agency (research project grant V3-2104 "Secondary prevention and quality of care after myocardial infarction in Slovenia").

References

- [1] G. A. Roth, G. A. Mensah, C. O. Johnson et al., "Global burden of cardiovascular diseases and risk factors, 1990-2019: update from the GBD 2019 study," *Journal of the American College of Cardiology*, vol. 76, no. 25, pp. 2982–3021, 2020.
- [2] M. G. Saklayen, "The global epidemic of the metabolic syndrome," *Current Hypertension Reports*, vol. 20, no. 2, p. 12, 2018.
- [3] M. Ng, T. Fleming, M. Robinson et al., "Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013," *The Lancet*, vol. 384, no. 9945, pp. 766–781, 2014.
- [4] L. Pasalic, E. Wing-Lun, J. K. Lau et al., "Novel assay demonstrates that coronary artery disease patients have heightened

- procoagulant platelet response,” *Journal of Thrombosis and Haemostasis*, vol. 16, no. 6, pp. 1198–1210, 2018.
- [5] F. Guo and W. T. Garvey, “Cardiometabolic disease risk in metabolically healthy and unhealthy obesity: stability of metabolic health status in adults,” *Obesity*, vol. 24, no. 2, pp. 516–525, 2016.
 - [6] D. G. Carey, A. B. Jenkins, L. V. Campbell, J. Freund, and D. J. Chisholm, “Abdominal fat and insulin resistance in normal and overweight women: direct measurements reveal a strong relationship in subjects at both low and high risk of NIDDM,” *Diabetes*, vol. 45, no. 5, pp. 633–638, 1996.
 - [7] K. Park, C. W. Ahn, S. B. Lee et al., “Elevated TyG index predicts progression of coronary artery calcification,” *Diabetes Care*, vol. 42, no. 8, pp. 1569–1573, 2019.
 - [8] L. Sánchez-Íñigo, D. Navarro-González, A. Fernández-Montero, J. Pastrana-Delgado, and J. A. Martínez, “The TyG index may predict the development of cardiovascular events,” *European Journal of Clinical Investigation*, vol. 46, no. 2, pp. 189–197, 2016.
 - [9] C. Irace, C. Carallo, F. B. Scavelli et al., “Markers of insulin resistance and carotid atherosclerosis. A comparison of the homeostasis model assessment and triglyceride glucose index,” *International Journal of Clinical Practice*, vol. 67, no. 7, pp. 665–672, 2013.
 - [10] Q. Mao, D. Zhou, Y. Li, Y. Wang, S. C. Xu, and X. H. Zhao, “The triglyceride-glucose index predicts coronary artery disease severity and cardiovascular outcomes in patients with non-ST-segment elevation acute coronary syndrome,” *Disease Markers*, vol. 2019, Article ID 6891537, 11 pages, 2019.
 - [11] J. L. Jin, Y. X. Cao, L. G. Wu et al., “Triglyceride glucose index for predicting cardiovascular outcomes in patients with coronary artery disease,” *Journal of Thoracic Disease*, vol. 10, no. 11, pp. 6137–6146, 2018.
 - [12] Z. Wu, L. Liu, W. Wang et al., “Triglyceride-glucose index in the prediction of adverse cardiovascular events in patients with premature coronary artery disease: a retrospective cohort study,” *Cardiovascular Diabetology*, vol. 21, no. 1, p. 142, 2022.
 - [13] Z. Wu, H. Cui, W. Li et al., “Comparison of three non-insulin-based insulin resistance indexes in predicting the presence and severity of coronary artery disease,” *Frontiers in Cardiovascular Medicine*, vol. 9, article 918359, 2022.
 - [14] L. F. Hong, X. L. Li, S. H. Luo et al., “Association of fibrinogen with severity of stable coronary artery disease in patients with type 2 diabetic mellitus,” *Disease Markers*, vol. 2014, Article ID 485687, 8 pages, 2014.
 - [15] G. Ndrepepa, S. Braun, L. King et al., “Relation of fibrinogen level with cardiovascular events in patients with coronary artery disease,” *The American Journal of Cardiology*, vol. 111, no. 6, pp. 804–810, 2013.
 - [16] G. De Luca, M. Verdoia, E. Cassetti et al., “High fibrinogen level is an independent predictor of presence and extent of coronary artery disease among Italian population,” *Journal of Thrombosis and Thrombolysis*, vol. 31, no. 4, pp. 458–463, 2011.
 - [17] R. Chaudhary, A. Sukhi, R. Chaudhary et al., “Gender differences in thrombogenicity among patients with angina and non-obstructive coronary artery disease,” *Journal of Thrombosis and Thrombolysis*, vol. 48, no. 3, pp. 373–381, 2019.
 - [18] T. Tsujimoto and H. Kajio, “Thrombotic/thrombolytic balance as a cardiac treatment determinant in patients with diabetes mellitus and coronary artery disease,” *Journal of the American Heart Association*, vol. 8, no. 2, p. e011207, 2019.
 - [19] N. Tynngård, T. L. Lindahl, and S. Ramström, “Assays of different aspects of haemostasis - what do they measure?,” *Thrombosis Journal*, vol. 13, no. 1, p. 8, 2015.
 - [20] A. Antovic, “The overall hemostasis potential: a laboratory tool for the investigation of global hemostasis,” *Seminars in Thrombosis and Hemostasis*, vol. 36, no. 7, pp. 772–779, 2010.
 - [21] A. Antovic, “Screening haemostasis – looking for global assays: the overall haemostasis potential (ohp) method – a possible tool for laboratory investigation of global haemostasis in both hypo- and hypercoagulable conditions,” *Current Vascular Pharmacology*, vol. 6, no. 3, pp. 173–185, 2008.
 - [22] S. Anžej, M. Božič, A. Antovič et al., “Evidence of hypercoagulability and inflammation in young patients long after acute cerebral ischaemia,” *Thrombosis Research*, vol. 120, no. 1, pp. 39–46, 2007.
 - [23] S. He, A. Antovic, and M. Blombäck, “A simple and rapid laboratory method for determination of haemostasis potential in plasma: II. Modifications for use in routine laboratories and research work,” *Thrombosis Research*, vol. 103, no. 5, pp. 355–361, 2001.
 - [24] M. Rakusa, M. Jensterle, M. Božič-Mijovski, and A. Janez, “Increased coagulation and decreased fibrinolysis as measured with overall hemostatic potential are dependent on BMI and not associated with PCOS,” *Metabolic Syndrome and Related Disorders*, vol. 15, no. 4, pp. 194–198, 2017.
 - [25] C. J. Reddel, J. L. Curnow, J. Voitl et al., “Detection of hypofibrinolysis in stable coronary artery disease using the overall haemostatic potential assay,” *Thrombosis Research*, vol. 131, no. 5, pp. 457–462, 2013.
 - [26] M. Nieuwdorp, E. S. G. Stroes, J. C. M. Meijers, and H. Büller, “Hypercoagulability in the metabolic syndrome,” *Current Opinion in Pharmacology*, vol. 5, no. 2, pp. 155–159, 2005.
 - [27] S. S. Anand, Q. Yi, H. Gerstein et al., “Relationship of metabolic syndrome and fibrinolytic dysfunction to cardiovascular disease,” *Circulation*, vol. 108, no. 4, pp. 420–425, 2003.
 - [28] D. Y. Lee, E. S. Lee, J. H. Kim et al., “Predictive value of triglyceride glucose index for the risk of incident diabetes: a 4-year retrospective longitudinal study,” *PloS One*, vol. 11, no. 9, p. e0163465, 2016.
 - [29] K. Aung, C. Lorenzo, M. A. Hinojosa, and S. M. Haffner, “Risk of developing diabetes and cardiovascular disease in metabolically unhealthy normal-weight and metabolically healthy obese individuals,” *The Journal of Clinical Endocrinology and Metabolism*, vol. 99, no. 2, pp. 462–468, 2014.
 - [30] J. Arnlöv, E. Ingelsson, J. Sundström, and L. Lind, “Impact of body mass index and the metabolic syndrome on the risk of cardiovascular disease and death in middle-aged men,” *Circulation*, vol. 121, no. 2, pp. 230–236, 2010.
 - [31] Y. Aso, “Plasminogen activator inhibitor (PAI)-1 in vascular inflammation and thrombosis,” *Frontiers in Bioscience-Landmark*, vol. 12, no. 8, pp. 2957–2966, 2007.
 - [32] I. Mertens and L. F. Van Gaal, “Obesity, haemostasis and the fibrinolytic system,” *Obesity Reviews*, vol. 3, no. 2, pp. 85–101, 2002.
 - [33] D. Vykoukal and M. G. Davies, “Vascular biology of metabolic syndrome,” *Journal of Vascular Surgery*, vol. 54, no. 3, pp. 819–831, 2011.
 - [34] S. He, K. Bremme, and M. Blombäck, “A laboratory method for determination of overall haemostatic potential in plasma. I. Method design and preliminary results,” *Thrombosis Research*, vol. 96, no. 2, pp. 145–156, 1999.