



Scientific Comment

Does gender influence reference values of platelet indices? ☆



Magnun Nueldo Nunes dos Santos  *

Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

Platelets, the smallest cellular component of blood, are anucleated, originate from the cytoplasmic fragmentation of the megakaryocyte in the bone marrow, and play a central role in hemostasis.¹ Based on data, automated hematological analyzers are essential in clinical laboratories, especially for the diagnosis and follow-up of patients with disorders of hemostasis.²

Significant technological development in automated blood cell counters has been observed in recent years, providing excellent analytical performance and robust information. However, specific knowledge becomes necessary in order to elucidate the appropriate clinical applications of these data.³ Some of these analyzers provide data on platelet indices, such as mean platelet volume (MPV), platelet distribution width (PDW), percentage of large platelets (P-LCR), and the percentage of blood volume occupied by platelets (plateletcrit – PCT),⁴ which are valuable in a number of clinical, hematological or non-hematological conditions.^{5–7}

Platelet indices have been investigated in cardiovascular diseases, cerebrovascular diseases, immune thrombocytopenia, diabetes mellitus, among others. Some studies indicate that these indices can be considered as markers of platelet activation in inflammatory and thrombotic conditions.^{5–9} In addition, the use of these indices as markers of the quality of platelet concentrates in transfusion therapy has been highlighted.¹⁰ Despite the accessibility and scientific interest in investigating the association of these data with various clinical conditions, platelet indices are not routinely used in

clinical-laboratory practice. Perhaps, the underutilization of these indices may be due to limited knowledge regarding their reference values.⁵

The purpose of laboratory tests may be diagnostic, prognostic, or for monitoring of treatment. To achieve this purpose, it is necessary that some aspects of each laboratory test be considered. An important aspect is the limit of significance that is the determination of its reference range or interval.¹¹

The reference interval of laboratory tests is of fundamental relevance for the interpretation of laboratory results and provision of useful clinical information in order to aid proper medical decision. Thus, reliability of the reference interval may play a crucial role in the clinical interpretation of results.¹²

Several laboratory parameters have specific reference values for men and women due to differences in anatomical structure, muscle mass or hormonal profile.¹³ Specifically for the determination of platelet indices, there are contradictory reports about these differences.^{5,13,14}

In this volume of the Brazilian Journal of Hematology and Hemotherapy, Ali et al. analyzed the reference ranges of MPV, PDW, P-LCR and PCT according to gender, of 2356 apparently healthy individuals from a population of the United Kingdom.¹⁴ In the study, individuals were found to have statistically significant differences in the reference range of MPV, P-LCR and PCT for men and women. However, no differences were identified in the reference range of PDW.

DOI of original article: <https://doi.org/10.1016/j.htct.2018.09.005>.

☆ See paper by Ali U et al. on pages [XX-XX].

* Correspondence to: Departamento de Patologia Clínica, Faculdade de Ciências Médicas, Universidade de Campinas (UNICAMP), Campinas, SP, Brazil.

E-mails: magnun@fcm.unicamp.br, magnun.nunes@gmail.com

<https://doi.org/10.1016/j.htct.2019.04.001>

2531-1379/© 2019 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

An important aspect was the sample size analyzed in the mentioned study and, as highlighted by Ali et al., the results obtained confirmed reports in the literature.¹⁴ However, when compared to other studies, the findings were discordant, and support the relevance of establishing reference intervals for each region and specific population as recommended by the Clinical and Laboratory Standards Institute (CLSI).¹⁵ More importantly, this work reinforces the idea of having a local reference schedule with regards to gender, thus, permitting a more accurate assessment of the quality and information of the results of clinical laboratory tests.

Conflicts of interest

The author declares no conflicts of interest.

REFERENCES

- Gibbins JM. Platelet adhesion signalling and the regulation of thrombus formation. *J Cell Sci.* 2004;117 Pt 16:3415–25.
- Wada A, Takagi Y, Kono M, Morikawa T. Accuracy of a new platelet count system (PLT-F) depends on the staining property of its reagents. *PLOS ONE.* 2015;10(10):e0141311.
- Buttarelo M, Plebani M. Automated blood cell counts: state of the art. *Am J Clin Pathol.* 2008;130(1):104–16.
- Kim HK, Kim JE, Ham CK, Lee DS, Park S, Cho HI. Prognostic value of platelet indices as determined by ADVIA 120 in patients suspected of having disseminated intravascular coagulation. *Int J Lab Hematol.* 2008;30(2):117–23.
- Maluf CB, Barreto SM, Vidigal PG. Standardization and reference intervals of platelet volume indices: insight from the Brazilian longitudinal study of adult health (ELSA-BRASIL). *Platelets.* 2015;26(5):413–20.
- Nkambule BB, Davison GM, Ipp H. The evaluation of platelet indices and markers of inflammation, coagulation and disease progression in treatment-naïve, asymptomatic HIV-infected individuals. *Int J Lab Hematol.* 2015;37(4):450–8.
- Bora K, Jitani AK, Raphael V, Ruram AA, Borah P, Khonglah Y. Association between lipid profile and platelet indices: the importance of considering the influence of lipid profile while evaluating the clinical utility of platelet indices. *Int J Lab Hematol.* 2016;38(4):e80–3.
- Kaito K, Otsubo H, Usui N, Yoshida M, Tanno J, Kurihara E, et al. Platelet size deviation width, platelet large cell ratio, and mean platelet volume have sufficient sensitivity and specificity in the diagnosis of immune thrombocytopenia. *Br J Haematol.* 2005;128(5):698–702.
- Venkatesh V, Kumar R, Varma DK, Bhatia P, Yadav J, Dayal D. Changes in platelet morphology indices in relation to duration of disease and glycemic control in children with type 1 diabetes mellitus. *J Diabetes Complicat.* 2018;32(9):833–8.
- Singh H, Chaudhary R, Ray V. Platelet indices as quality markers of platelet concentrates during storage. *Clin Lab Haematol.* 2003;25(5):307–10.
- Slhessarenko N, Andriolo A. The importance of determining reference intervals for Laboratory Medicine. *J Bras Patol Med Lab.* 2016;52(2):68–9.
- Bimerew LG, Demie T, Eskinder K, Getachew A, Bekele S, Cheneke W, et al. Reference intervals for hematology test parameters from apparently healthy individuals in southwest Ethiopia. *SAGE Open Med.* 2018;6, 2050312118807626.
- Butkiewicz AM, Kemona H, Dymicka-Piekarska V, Matowicka-Karna J, Radziwon P, Lipska A. Platelet count, mean platelet volume and thrombocytopenic indices in healthy women and men. *Thromb Res.* 2006;118(2):199–204.
- Ali U, Gibbs R, Knight G, Tsitsikas D. Sex-divided reference intervals for mean platelet volume, platelet large cell ratio and plateletcrit using the Sysmex XN-10 automated hematology analyzer in a UK population. *Hematol Transfus Cell Ther.* 2019;41(2):153–7.
- CLSI. Defining, establishing, and verifying reference intervals in the Clinical Laboratory; approved guideline. CLSI document EP28-A3c. 3rd ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2008.