

# Potentiation of TRPM7 Inward Currents by Protons

Jianmin Jiang, Mingjiang Li, and Lixia Yue

Center for Cardiology and Cardiovascular Biology, Department of Cell Biology, University of Connecticut Health Center, Farmington, CT 06032

TRPM7 is unique in being both an ion channel and a protein kinase. It conducts a large outward current at +100 mV but a small inward current at voltages ranging from  $-100$  to  $-40$  mV under physiological ionic conditions. Here we show that the small inward current of TRPM7 was dramatically enhanced by a decrease in extracellular pH, with an  $\sim 10$ -fold increase at pH 4.0 and 1–2-fold increase at pH 6.0. Several lines of evidence suggest that protons enhance TRPM7 inward currents by competing with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding sites, thereby releasing blockade of divalent cations on inward monovalent currents. First, extracellular protons significantly increased monovalent cation permeability. Second, higher proton concentrations were required to induce 50% of maximal increase in TRPM7 currents when the external  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations were increased. Third, the apparent affinity for  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  was significantly diminished at elevated external  $\text{H}^+$  concentrations. Fourth, the anomalous-mole fraction behavior of  $\text{H}^+$  permeation further suggests that protons compete with divalent cations for binding sites in the TRPM7 pore. Taken together, it appears that at physiological pH (7.4),  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  bind to TRPM7 and inhibit the monovalent cationic currents; whereas at high  $\text{H}^+$  concentrations, the affinity of TRPM7 for  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  is decreased, thereby allowing monovalent cations to pass through TRPM7. Furthermore, we showed that the endogenous TRPM7-like current, which is known as  $\text{Mg}^{2+}$ -inhibitible cation current (MIC) or Mg nucleotide–regulated metal ion current (MagNuM) in rat basophilic leukemia (RBL) cells was also significantly potentiated by acidic pH, suggesting that MIC/MagNuM is encoded by TRPM7. The pH sensitivity represents a novel feature of TRPM7 and implies that TRPM7 may play a role under acidic pathological conditions.

## INTRODUCTION

TRPM7 is a ubiquitously distributed ion channel that belongs to the long or melastatin-related transient receptor potential (TRPM) ion channel subfamily (Harteneck et al., 2000; Montell, 2001; Clapham, 2003; Fleig and Penner, 2004). It is unique in being both an ion channel and a protein kinase. Although the physiological functions of the kinase are not well understood, recent studies have suggested that TRPM7 plays important roles in cellular  $\text{Mg}^{2+}$  homeostasis (Schmitz et al., 2003), anoxic neuronal cell death (Aarts et al., 2003), cell proliferation and viability (Nadler et al., 2001; Hanano et al., 2004), and diseases caused by abnormal magnesium absorption (Schlingmann et al., 2002; Walder et al., 2002; Chubanov et al., 2004).

TRPM7 produces pronounced outward currents at nonphysiological voltages ranging from +50 to +100 mV and small inward currents at negative potentials between  $-100$  to  $-40$  mV when expressed heterologously in mammalian cells (Nadler et al., 2001; Runnels et al., 2001; Monteilh-Zoller et al., 2003; Schmitz et al., 2003). Unlike some other TRP channels that are gated or potentiated by activation of the PLC pathway (Clapham, 2003), TRPM7 is inhibited by depletion of  $\text{PIP}_2$

mediated by PLC activation (Runnels et al., 2002; Aarts et al., 2003). The basal activity of TRPM7 is regulated by millimolar levels of intracellular MgATP and  $\text{Mg}^{2+}$ , so that TRPM7 is activated by depletion of intracellular MgATP and  $\text{Mg}^{2+}$ , and is inhibited by high concentrations of MgATP and  $\text{Mg}^{2+}$  with an  $\text{IC}_{50}$  of  $\sim 0.6$  mM (Nadler et al., 2001). The mechanism by which  $\text{Mg}^{2+}$  inhibits TRPM7, however, is not yet entirely clear (Nadler et al., 2001; Hermosura et al., 2002; Prakriya and Lewis, 2002; Runnels et al., 2002; Kerschbaum et al., 2003; Kozak and Cahalan, 2003; Monteilh-Zoller et al., 2003; Schmitz et al., 2003). Other divalent cations have also been reported to inhibit TRPM7 (Kozak and Cahalan, 2003).

Although inactivation of TRPM7 has been extensively investigated, the activation mechanism of TRPM7 under physiological conditions remains unknown. Intracellular  $\text{Mg}^{2+}$  levels (0.5–1 mM) under physiological conditions can inactivate  $>50\%$  of TRPM7 channel activities (Nadler et al., 2001; Kozak and Cahalan, 2003). Thus, the inward current amplitude, which is usually 1/30 to 1/10 of the outward current amplitude measured at +100 mV (Nadler et al., 2001; Runnels et al., 2001; Schmitz et al., 2003), may be very small under

J. Jiang and M. Li contributed equally to this work.

Correspondence to Lixia Yue: lyue@uchc.edu

M. Li's present address is Renmin Hospital of Wuhan University, People's Republic of China.

Abbreviations used in this paper: DVF, divalent-free solution; MagNuM, Mg nucleotide–regulated metal ion current; MIC,  $\text{Mg}^{2+}$ -inhibitible cation; RBL, rat basophilic leukemia; TRPM, melastatin-related transient receptor potential.

physiological internal  $Mg^{2+}$  levels in the native cells. Given the potential important physiological functions (Nadler et al., 2001; Runnels et al., 2002; Schlingmann et al., 2002; Walder et al., 2002; Aarts et al., 2003; Ryazanova et al., 2004; Schmitz et al., 2003; Chubanov et al., 2004; Hanano et al., 2004), it is likely that TRPM7 inward current may be potentiated by physiological or pathological stimuli. A recent study showed that TRPM7 is up- and down-regulated in a cAMP- and PKA-dependent manner (Takezawa et al., 2004), with the changes assessed by outward current amplitude.  $H_2O_2$  was reported to increase TRPM7 inward currents by one- to twofold after prolonged (30–50 min) incubation (Aarts et al., 2003). However, the mechanism by which  $H_2O_2$  regulates TRPM7 remains unclear.

In the present study, we demonstrate that protons markedly potentiate TRPM7 inward currents. Lowering extracellular pH increases the inward current by  $\sim 10$ -fold, whereas the outward current is only changed by  $< 30\%$ . This transforms the normal outward rectification of TRPM7 to rectification in both the inward and outward directions. Further, our data suggest that protons enhance TRPM7 inward currents by competing with external  $Ca^{2+}$  and  $Mg^{2+}$  for binding sites in the TRPM7 pore. Therefore, the effect of acidic pH on TRPM7 is more pronounced when extracellular  $Ca^{2+}$  concentration is decreased. Importantly, we show that the well-characterized endogenous TRPM7-like current MIC ( $Mg^{2+}$ -inhibitable cation)/MagNum ( $Mg$  nucleotide-regulated metal ion) in rat basophilic leukemia (RBL) cells is similarly potentiated by a decrease in extracellular pH. As high proton concentrations (pH  $< 6$ ) can be generated during various forms of injury, including infection, inflammation, and ischemia (Jacobus et al., 1977; Stevens et al., 1991; Steen et al., 1992), the significant increase in TRPM7 inward currents by protons suggests that TRPM7 may play a role under acidic pathological conditions.

## MATERIALS AND METHODS

### Cell Culture

HEK-293 cells stably transfected with a FLAG-tagged murine TRPM7 in pCDNA4/TO vector were provided by A. Scharenberg (University of Washington, Seattle, WA). Cells were grown with DMEM medium supplemented with 10% FBS, 100 U/ml penicillin and 100 mg/ml streptomycin, blasticidin (5  $\mu$ g/ml), and zeocin (0.4 mg/ml). Expression of TRPM7 was induced 12–24 h before experiments by adding 1  $\mu$ g/ml tetracycline to the culture medium (Nadler et al., 2001; Schmitz et al., 2003). Unless otherwise stated, experiments were conducted using TRPM7 expressing HEK-293 cells after tetracycline induction.

RBL-2H3 cells were provided by D. Clapham (Harvard Medical School, Boston, MA). Cells were cultured in DMEM supplemented with 10% FBS and 100 U/ml penicillin and 100 mg/ml streptomycin. For electrophysiological experiments, cells were plated onto glass coverslips and used 12 h thereafter.


### Electrophysiology

Whole-cell currents were recorded using an Axopatch 200B (Axon Instruments, Inc.) amplifier. Data were digitized at 20 kHz and low-pass filtered at 2 kHz. pClamp9 software was used for data acquisition and analysis. Patch electrodes were pulled from borosilicate glass and fire polished to a resistance of  $\sim 3$  M $\Omega$  when filled with internal solutions. Series resistance ( $R_s$ ) was compensated up to 90% to reduce series resistance errors to  $< 5$  mV. Cells in which  $R_s$  was  $> 10$  M $\Omega$  were discarded (Yue et al., 2002). All experiments were conducted at  $22 \pm 2^\circ C$ .

Voltage stimuli lasting 250 ms were delivered at 1- to 5-s intervals, with either voltage ramps or voltage steps ranging from  $-120$  to  $+100$  mV. Unless otherwise stated, 3–5 min were allowed to let TRPM7 current develop and reach a steady state after break-in. A fast perfusion system was used to exchange extracellular solutions. A complete solution exchange can be achieved in  $\sim 1$ –3 s.

The internal pipette solution (P1) for whole cell current recordings in HEK-293 cells stably expressing TRPM7 contained (in mM) 145 Cs-methanesulfonate, 8 NaCl, 10 EGTA, and 10 HEPES, pH adjusted to 7.2 with CsOH. In some experiments,  $Mg^{2+}$  was added to the pipette solution and the free  $Mg^{2+}$  concentration was titrated to 3 mM (calculated with the MaxChelator software, available at <http://www.stanford.edu/~cpatton/webmaxc.htm>). For current recordings in RBL cells, pipette solution (P2) contained (in mM) 145 Cs-methanesulfonate, 8 NaCl, 1 EGTA, 0.084  $Ca^{2+}$ , 1  $MgCl_2$ , 5 ATP- $Na_2$ , 10 HEPES, pH adjusted to 7.2 with CsOH. Free  $Ca^{2+}$  and  $Mg^{2+}$  concentrations were estimated at  $\sim 15$  nM and 26  $\mu$ M, respectively (MaxChelator). In experiments designed to diminish outward currents, pipette solution (P3) contained (in mM) 120 NMDG, 108 glutamic acid, 10 HEPES, 10 EGTA, 10 CsCl, pH adjusted 7.2 with NMDG.

The standard extracellular Tyrode's solution contained (in mM) 140 NaCl, 5 KCl, 2  $CaCl_2$ , 20 HEPES, and 10 glucose, pH adjusted to 7.4 (NaOH). HEPES (20 mM) was used in the solutions at pH 7.0 and 7.4, and was replaced by 10 mM HEPES and 10 mM MES for the solutions at pH  $\leq 6$  (Jordt et al., 2000; Askwith et al., 2004; Yermolaieva et al., 2004). Divalent-free solution (DVF) contained (in mM) 145 NaCl, 20 HEPES, 5 EGTA, 2 EDTA and 10 glucose, with estimated free  $[Ca^{2+}] < 1$  nM at pH 7.4 and free  $[Mg^{2+}] \approx 10$  nM at pH 7.4 (MaxChelator). HEPES (20 mM) was replaced by 10 mM HEPES and 10 mM MES in DVF solutions at pH 4.0, and the estimated free  $[Ca^{2+}]$  was 7.7  $\mu$ M and free  $[Mg^{2+}]$  was 9.9  $\mu$ M in DVF at pH 4.0 (MaxChelator). Appropriate  $Ca^{2+}$  or  $Mg^{2+}$  was added to the DVF at pH 7.4 to prepare solutions containing  $\leq 10$   $\mu$ M  $Mg^{2+}$  or  $Ca^{2+}$  (Fig. 6). Solutions containing 1, 2, and 10 mM  $Mg^{2+}$  or  $Ca^{2+}$  at both pH 4.0 and 7.4 were prepared by omitting EDTA and EGTA in the DVF solution, and by adding the appropriate concentrations of  $Mg^{2+}$  or  $Ca^{2+}$ . Isotonic  $Ca^{2+}$  or  $Mg^{2+}$  solution contains 120 mM  $Ca^{2+}$  or  $Mg^{2+}$ , 10 mM HEPES, 10 mM glucose, with pH adjusted to pH 7.4 or pH 4.0. Anomalous mole fraction behavior of  $Ca^{2+}$  permeation (Fig. 3 F) was evaluated in a series of external solutions, including isotonic  $Ca^{2+}$  (120 mM), 10 mM  $Ca^{2+}$ , 2 mM  $Ca^{2+}$ , 1 mM  $Ca^{2+}$ , 100  $\mu$ M  $Ca^{2+}$ , and nominally  $Ca^{2+}$ -free solution in which free  $Ca^{2+}$  concentration was estimated at 10  $\mu$ M (Vennekens et al., 2000; Yue et al., 2001). The solutions containing 100  $\mu$ M to 10 mM  $Ca^{2+}$  were prepared from normal Tyrode's solution by adding appropriate concentration of  $Ca^{2+}$ , with reductions in  $Na^+$  concentration when necessary to keep the constant osmolarity. The same method was used to prepare a series of solutions containing various  $Mg^{2+}$  concentrations for anomalous mole fraction behavior of  $Mg^{2+}$  permeation experiment shown in Fig. 3 F. Cells were usually exposed to acidic solutions for  $\sim 30$  s to avoid desensitization unless otherwise stated. Current amplitude was measured at  $-120$  or  $+100$  mV. Amiloride was added to the perfusate as indicated in the text. All the chemicals for electrophysiological experiments were from Sigma-Aldrich.



**Figure 1.** Potentiation of TRPM7 inward currents by external protons in HEK-293 cells stably expressing TRPM7. (A) Representative TRPM7 currents evoked by voltage ramps ranging from  $-120$  to  $+100$  mV in the external Tyrode's solutions at pH 7.4 and pH 4.0. (B) Normalized outward ( $+100$  mV) and inward currents ( $-120$  mV) at pH 4.0 and pH 7.4. The acidic external solution (pH 4.0) was applied after the current amplitude reached a steady-state ( $\sim 5$  min after break-in). Repetitive application of acidic solution produced a similar increase in inward and outward currents. (C and D) TRPM7 currents elicited by voltage steps ranging from  $-120$  to  $+100$  mV with an increment of  $20$  mV at pH 7.4 (C) and pH 4.0 (D). Only inward current was significantly increased at pH 4.0 (D). (E) Currents recorded at various time points in a cell dialyzed with the pipette solution containing  $3$  mM free  $Mg^{2+}$ . a, immediately after formation of whole-cell configuration; b, before the external solution was changed from pH 7.4 to pH 4.0; c, the first time application of external solution at pH 4.0 induced an increase in inward current; d, the fifth time application

## Data Analysis

Pooled data are presented as mean  $\pm$  SEM. Concentration-response curves were fitted by an equation of the form:  $E = E_{max} \{1/[1 + (EC_{50}/C)^n]\}$ , where  $E$  is the effect at concentration  $C$ ,  $E_{max}$  is maximal effect,  $EC_{50}$  is the concentration for half-maximal effect and  $n$  is the Hill coefficient (Yue et al., 2000).  $EC_{50}$  is replaced by  $IC_{50}$  if the effect is an inhibitory effect. Statistical comparisons were made using two-way analysis of variance (ANOVA) and two-tailed  $t$  test with Bonferroni correction;  $P < 0.05$  indicated statistical significance.

## RESULTS


### Potentiation of TRPM7 Inward Currents by Protons

TRPM7 currents were elicited by voltage ramps or voltage steps ranging from  $-120$  to  $+100$  mV from a holding potential of  $0$  mV (Runnels et al., 2001). As previously reported, TRPM7 produced large outward currents and small inward currents (Fig. 1 A, red trace) (Nadler et al., 2001; Runnels et al., 2001; Schmitz et al., 2003). After break-in,  $3$ – $5$  min was allowed to let TRPM7 current amplitude reach a steady state (see MATERIALS AND METHODS) before changing external solutions. As the measurable outward current occurs at nonphysiological range, we investigated whether the TRPM7 inward current can be potentiated by pathological stimuli. As shown in Fig. 1 A, the inward current of TRPM7 was dramatically increased by a decrease in the extracellular pH to 4.0. In the same cell as shown in Fig. 1 A, a similar degree of increase in inward current was observed upon a second application of the external solution at pH 4.0, indicating that the effect of pH 4.0 on TRPM7 was reversible and reproducible (Fig. 1 B). While the inward currents were increased by  $\sim 10$ -fold (at  $-120$  mV) at pH 4.0, the outward currents measured at  $+100$  mV were only mildly changed, showing an  $\sim 30\%$  increase (Fig. 1 B, top). The large increase in inward current compared with the small change in outward current induced by acidic pH was also evident in the currents elicited by voltage steps ranging from  $-120$  to  $+100$  mV (Fig. 1, C and D).

of external solution at pH 4.0 could not induce any change because TRPM7 was completely blocked by intracellular  $Mg^{2+}$ . (F) In the same cell as shown in E, continuous changes in inward (top) and outward (middle) currents measured at  $+100$  mV and  $-120$  mV were plotted as a function of time. The inward currents labeled 1, 2, 3, 4, 5, and 6 (middle) represent the time points when the cell was exposed to the acidic external solution (pH 4.0). (Bottom) Normalized inward current at pH 4.0 (filled circle, red) superimposed with the normalized outward current at pH 7.4 (filled triangle, black). The green line obtained by fitting the normalized outward current represents currents decay. (G) TRPM7 currents recorded at pH 7.4 and pH 4.0 with and without  $200$   $\mu$ M amiloride (Ami), respectively. (H) Time-dependent changes of the inward current induced by pH 4.0 in the presence and absence of  $200$   $\mu$ M amiloride (Ami).


To test whether the changes elicited by low pH were mediated by TRPM7, we studied the effects of acidic pH solution on control cells. In HEK-293 cells without tetracycline induction of TRPM7 expression, small endogenous TRPM7-like currents were recorded. The endogenous inward currents were also enhanced by  $\sim 10$ -fold at pH 4.0 ( $n = 6$ ; unpublished data). Although the outwardly rectifying I-V curve suggests that the endogenous current in the HEK-293 cells is TRPM7-like current, we cannot exclude the possibility that the current may be from some leaky expression. To further confirm that the pH 4.0-elicited increases in inward current are through TRPM7, we did the following experiments. As TRPM7 is ubiquitously expressed in various cell types (Nadler et al., 2001; Hermosura et al., 2002; Kozak et al., 2002; Prakriya and Lewis, 2002; Runnels et al., 2002; Aarts et al., 2003; Jiang et al., 2003; Kozak and Cahalan, 2003), including the HEK-293 cells used to create the stable TRPM7 expressing cell lines (Nadler et al., 2001; Takezawa et al., 2004), it is difficult to find a cell type that does not have endogenous TRPM7 expression. Therefore, we used a pipette solution containing 3 mM free  $Mg^{2+}$  to suppress TRPM7 currents and then measured the response of the cells to acidic pH. TRPM7 currents were apparent immediately after formation of the whole-cell configuration (Fig. 1 E, a and b), but then the current amplitude gradually decreased (Fig. 1 F). Application of external solution at pH 4.0 induced dramatic increases in inward current with a small increase in outward currents early after forming the whole cell configuration (Fig. 1 E, c, and Fig. 1 F, 1(c)). Before the current was completely inhibited by intracellular  $Mg^{2+}$ , exposing the cell to the external solution at pH 4.0 for a second, third, and fourth time repeatedly induced significant increases in TRPM7 inward currents (Fig. 1 F, 1, 2, 3, and 4). After TRPM7 was totally blocked by intracellular  $Mg^{2+}$ , pH 4.0 failed to induce any change in current amplitude (Fig. 1 E, d, and Fig 1 F, 5 and 6). Similar results were observed in five other cells, suggesting that pH 4.0-elicited increases in current amplitude are through TRPM7. In addition, the normalized inward current at pH 4.0 plotted as a function of time superimposed with that of normalized outward current at pH 7.4, indicating that the inward current at pH 4.0 and the outward current at pH 7.4 decay at the same rate when the pipette solution contains 3 mM free  $Mg^{2+}$  (Fig. 1 F, bottom), further indicating that the pH 4.0-induced increase in inward current is mediated by TRPM7.

To rule out the possibility that proton-activated Na channels (Waldmann and Lazdunski, 1998) were involved in the increased inward current elicited by low pH, amiloride (200  $\mu M$ ) was added to the external solution. No significant difference was observed in inward currents elicited by pH 4.0 in the presence and absence of 200  $\mu M$  amiloride (Fig. 1, G and H), sug-



**Figure 2.** Concentration-dependent effects of protons on TRPM7 currents. (A–E) Potentiation of TRPM7 inward currents by protons at pH 7.0 (A), pH 6.0 (B), pH 5.0 (C), pH 4.0 (D), and pH 3.0 (E). Representative traces were elicited by voltage ramps ranging from  $-120$  to  $+100$  mV in the external solutions with NaCl replaced by  $NaSO_3CH_3$ . The y axis in each panel was scaled to illustrate the changes in inward currents. (F) Concentration-dependent changes in inward current amplitude measured at  $-120$  mV from the recordings elicited by voltage ramps at indicated pH. The dashed line represents zero current levels. (G) The increase in current amplitude at the indicated pH was normalized to the value at pH 3.0. Best fit of the normalized data yielded an  $EC_{50} = 4.5 \pm 0.5$  pH unit (mean  $\pm$  SEM,  $n = 8$ ; Hill coefficient was 1).

gesting that the pH 4.0-induced inward currents were not due to acid-sensitive channels of the degenerin family (Waldmann and Lazdunski, 1998). Similarly, contamination by a proton-activated  $Cl^-$  current (Cherny et al., 1997) was excluded because acidic pH-induced increases in TRPM7 currents were not affected by replacement of NaCl with  $NaSO_3CH_3$  in the external solution (see Fig. 2). All the above-mentioned results suggest that the marked increases in inward currents evoked by acidic pH in TRPM7-overexpressing cells were mediated by TRPM7 channels.



**Figure 3.** Monovalent permeability of TRPM7 at pH 7.4. (A and B) Original recordings of TRPM7 currents obtained using P3 pipette solution in the following external solutions at pH 7.4 (mM): a, 2 Ca<sup>2+</sup>/NMDG; b, isotonic NMDG; c, 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/NMDG; d, 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 Na<sup>+</sup>; e, 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 K<sup>+</sup>; f, 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 Cs<sup>+</sup>; g, isotonic Mg<sup>2+</sup> (120 mM); h, isotonic Ca<sup>2+</sup> (120 mM). The insets show an expanded section of the graph to illustrate changes in reversal potential in different solutions. (C) Inward current amplitude in a representative cell in the indicated external solutions. (D) Changes in reversal potential obtained by subtracting reversal potential in isotonic NMDG from those in test solutions (mean ± SEM, *n* = 6). \*\*, *P* < 0.01 in comparison with the value obtained in 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/NMDG. (E) Averaged TRPM7 current amplitude in various test solutions. Note that the current amplitude in 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 Na<sup>+</sup>, 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 K<sup>+</sup>, and 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/150 Cs<sup>+</sup> is significantly larger than that in 2 Ca<sup>2+</sup>/1 Mg<sup>2+</sup>/NMDG. (F) Ca<sup>2+</sup>- and Mg<sup>2+</sup>-dependent anomalous mole-fraction effects. In the left panel, changes in inward current measured at -120 mV were normalized in reference to the value at 10 mM extracellular Ca<sup>2+</sup> (extracellular solutions were nominally Mg<sup>2+</sup> free). In the right


panel, changes in inward current measured at -120 mV were normalized in reference to the value at 2 mM extracellular Mg<sup>2+</sup> (extracellular solutions were nominally Ca<sup>2+</sup> free). Note that the current amplitude is minimal in the external solutions with 10 mM Ca<sup>2+</sup> (left) or 2 mM Mg<sup>2+</sup> (right) (mean ± SEM, *n* = 6).

We next studied concentration-dependent effects of protons on TRPM7. A small increase in the inward current was seen at pH 7.0 (Fig. 2 A); the increase was significant at pH 6.0 and reached a maximum at pH 3.0 (Fig. 2, B–E). The concentration-dependent increase of TRPM7 inward currents from the same cell is shown in Fig. 2 F. At pH 3.0, the average inward current amplitude measured at -120 mV was increased by 10.2 ± 1.3-fold compared with the current amplitude at pH 7.4. The pH required for inducing half-maximal increase in inward current was 4.5 (Fig. 2 G). Changes in

the outward current (measured at +100 mV) were much smaller than those of the inward current. For example, 13.1 ± 1.1%, 23.1 ± 2.5%, 23.3 ± 3.1%, and 24.8 ± 3.4% increases were observed at pH 7.0, 6.0, 5.0, and 4.0 (*P* < 0.05), respectively; whereas a 26.5 ± 4.0% (*P* < 0.05) decrease was seen at pH 3.0.

#### Protons Increase Monovalent Cation Permeability through TRPM7


We previously reported that TRPM7 is a nonselective cation channel that is permeable to both monovalent



**Figure 4.** Protons enhance TRPM7 inward currents by increasing monovalent cation permeability. (A, C, and E) Representative recordings of TRPM7 in 0, 10, 30, 100, and 150 mM Na<sup>+</sup> (A), K<sup>+</sup> (C), and Cs<sup>+</sup> (E) in the presence of 10 mM Ca<sup>2+</sup> at pH 4.0, respectively. The external solution containing 10 mM Ca<sup>2+</sup>/150 mM NMDG (pH 4.0) was used as a control solution. When monovalent ion concentrations were increased, equal molar NMDG was omitted from the solution to keep the osmolality constant. Note the concentration-dependent increase of TRPM7 conductance in A, C, and E. The insets show expanded section of the graphs to illustrate changes in reversal potential in various monovalent ion concentrations. (B, D, and F) Average changes in reversal potential (top) and inward current density (bottom) measured at -120 mV in the external solutions with various concentrations of monovalent cations. The changes in reversal potential were obtained by subtracting reversal potential in the control solution (10 mM Ca<sup>2+</sup>/NMDG) from those in test solutions with various monovalent cation concentrations (mean  $\pm$  SEM;  $n = 6$  [B];  $n = 7$  [D];  $n = 9$  [F]). \*,  $P < 0.05$ ; \*\*,  $P < 0.01$  in comparison with  $E_{rev}$  in 10 mM Ca<sup>2+</sup>/NMDG.

and divalent cations (Runnels et al., 2001). Monovalent permeability in the presence of divalent cations was also reported for the MIC channel in Jurkat cells (Prakriya and Lewis, 2002). Other studies suggested that TRPM7 is selective for divalent ions, and the inward current was carried exclusively by divalent cations in external solutions containing 10 mM Ca<sup>2+</sup> and 2 mM Mg<sup>2+</sup> (Nadler et al., 2001). To test whether monovalent cations contribute to TRPM7 inward currents under physiological Ca<sup>2+</sup> (2 mM) and Mg<sup>2+</sup> (1 mM) concentrations, we compared the TRPM7 current amplitude and reversal potentials in extracellular solutions con-

taining different monovalent cations, with those in the isotonic divalent cation solutions (Fig. 3 B). The P3 pipette solution containing lowered Cs<sup>+</sup> concentration (10 mM, see MATERIALS AND METHODS) was used to minimize outward current amplitude (Fig. 3). In the presence of 2 mM Ca<sup>2+</sup> and 1 mM Mg<sup>2+</sup>, changing the external solution from nonpermeable NMDG (Jiang et al., 2003; Kozak and Cahalan, 2003) to the solutions containing 150 mM Na<sup>+</sup>, K<sup>+</sup>, or Cs<sup>+</sup> significantly increased inward current amplitude and shifted reversal potentials (Fig. 3, A, C, D, and E), indicating that under physiological divalent cation concentrations,



**Figure 5.** Acidic pH does not increase TRPM7 divalent conductance. Currents were elicited by voltage ramps ( $-120$  to  $+100$  mV) with P1 pipette solution. (A) Divalent current amplitude measured at  $-120$  mV in the external solutions containing  $2$  mM  $\text{Ca}^{2+}$ / $1$  mM  $\text{Mg}^{2+}$ /NMDG at pH  $7.4$  and pH  $4.0$ , respectively. Dash line represents zero current level. (B) Averaged divalent current density of TRPM7 measured at  $-120$  mV in the external solutions as shown in A (mean  $\pm$  SEM,  $n = 8$ ). Current density at pH  $4.0$  was significantly smaller than that at pH  $7.4$ . (C) Current amplitude of TRPM7 measured at  $-120$  mV in the isotonic  $\text{Mg}^{2+}$  and isotonic  $\text{Ca}^{2+}$  solutions at pH  $7.4$  and pH  $4.0$ , respectively. Dashed lines represent zero current level. (D) Mean current density at  $-120$  mV in the isotonic  $\text{Mg}^{2+}$  or  $\text{Ca}^{2+}$  solutions at pH  $7.4$  and pH  $4.0$ , respectively (mean  $\pm$  SEM,  $n = 6$ ).

monovalent cationic currents contribute to TRPM7 inward currents. We further studied the relative effects of divalent cations on TRPM7 monovalent currents. Fig. 3 F shows anomalous mole fraction behavior of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  permeation (Fig. 3 F). The smallest current amplitude was observed in the external solutions containing  $10$  mM  $\text{Ca}^{2+}$  or  $2$  mM  $\text{Mg}^{2+}$ , respectively.


We proceeded to study the monovalent conductance under acidic conditions by using external solutions containing different concentrations of  $\text{Na}^+$ ,  $\text{K}^+$ , or  $\text{Cs}^+$  in the presence of  $10$  mM  $\text{Ca}^{2+}$  (Fig. 4). The P3 pipette solution was used in this experiment. Whole-cell configuration was established in the  $2$  mM  $\text{Ca}^{2+}$  Tyrode's solution at pH  $7.4$ , and cells were then exposed to  $0$ ,  $10$ ,  $30$ ,  $100$ , and  $150$  mM  $\text{Na}^+$ ,  $\text{K}^+$ , or  $\text{Cs}^+$  solutions in the presence of  $10$  mM  $\text{Ca}^{2+}$  at pH  $4.0$  for  $30$ – $60$  s. Concentration-dependent increases in current amplitude and shift of reversal potentials at high monovalent cation concentrations at pH  $4.0$  were observed (Fig. 4, A, C, and E, also see insets). These results (Fig. 4) indicate that the proton-evoked increases in TRPM7 inward currents were attributed to the increased  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cs}^+$  conductance.

To ensure that protons enhance monovalent conductance without increasing divalent conductance, we evaluated the effects of acidic pH on divalent currents. At physiological concentrations of  $\text{Ca}^{2+}$  ( $2$  mM) and  $\text{Mg}^{2+}$  ( $1$  mM) in the presence of NMDG solution, protons significantly decreased current amplitude measured at  $-120$  mV (Fig. 5, A and B). This result suggests that protons may compete with divalent cations for binding sites in the pore, so that low concentration of divalent cations were outcompeted by protons, therefore the di-

valent current amplitude at pH  $4.0$  was smaller than that at pH  $7.4$ . In agreement with this notion, we found that the isotonic  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  currents at pH  $4.0$  were not significantly different from those at pH  $7.4$  (Fig. 5, C and D). Thus, the acidic external solution-induced increase in inward currents was mediated by increasing monovalent cation permeability through TRPM7.

#### Protons Increase TRPM7 Currents by Competing with $\text{Ca}^{2+}$ and $\text{Mg}^{2+}$ for Binding Sites

To investigate the mechanism by which protons potentiate TRPM7 monovalent inward currents, we studied the effects of protons on TRPM7 currents in the presence of various external  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  concentrations. At pH  $4.0$ , protons induced a maximal increase in inward current in the external solution containing  $0.5$  mM  $\text{Ca}^{2+}$ , but only  $\sim 30\%$  of maximal response in the external solution containing  $10$  mM  $\text{Ca}^{2+}$  (Fig. 6 A). The  $\text{EC}_{50}$  for protons was changed from pH  $5.1$  to  $4.5$  and  $3.4$  when the external  $\text{Ca}^{2+}$  concentration was increased from  $0.5$  mM to  $2$  and  $10$  mM (Fig. 6 B), respectively. Similarly, pH  $3.5$  elicited a maximal increase in inward current in the external solution containing  $1$  mM  $\text{Mg}^{2+}$ , but only  $\sim 50\%$  maximal increase in TRPM7 inward current in the external solution containing  $10$  mM  $\text{Mg}^{2+}$ . The  $\text{EC}_{50}$  was changed from pH  $4.6$  to pH  $3.6$  when the external  $\text{Mg}^{2+}$  concentration was increased from  $1$  to  $10$  mM (Fig. 6, C and D). The  $1.1$  and  $1.7$  pH unit change in  $\text{EC}_{50}$  when external  $\text{Ca}^{2+}$  was changed from  $0.5$  to  $2$  and  $10$  mM and  $1$  pH unit change in  $\text{EC}_{50}$  when external  $\text{Mg}^{2+}$  was increased from  $1$  to  $10$  mM indicate that increasing external  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  concentrations decreases the affinity of TRPM7 for protons, and suggests




**Figure 6.** Protons increase TRPM7 inward currents by competing with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding sites. Current recorded in 2 mM  $\text{Ca}^{2+}$  Tyrode's at pH 7.4 (red) is shown in A, C, E, and G as a reference. (A) Representative traces of TRPM7 currents recorded in 0.5 and 10 mM external  $\text{Ca}^{2+}$  at pH 4.0. (B) Normalized increase in TRPM7 inward currents at indicated pH in the external solutions containing 0.5 and 10 mM  $\text{Ca}^{2+}$ . The dotted lines represent the results obtained in 2 mM external  $\text{Ca}^{2+}$  as shown in Fig 2. The pH values for 50% maximal response were  $5.1 \pm 0.2$ ,  $4.5 \pm 0.5$ , and  $3.4 \pm 0.1$  in 0.5, 2, and 10 mM external  $\text{Ca}^{2+}$  (mean  $\pm$  SEM,  $n = 6, 8,$  and  $6$ ), respectively. The Hill coefficient factors were 1.0 for the dose-response curves in B, D, F, and H. (C) Representative TRPM7 currents elicited by voltage ramps in 1 and 10 mM external  $\text{Mg}^{2+}$  at indicated pH values. (D) Normalized increase in TRPM7 inward current amplitude at indicated pH in the external solutions containing various  $\text{Mg}^{2+}$  concentrations. The pH values for 50% maximal response were  $4.6 \pm 0.5$  and  $3.6 \pm 0.2$  in 1 and 10 mM external  $\text{Mg}^{2+}$ , respectively (mean  $\pm$  SEM,  $n = 6$  for each group). (E) Representative currents elicited by voltage ramps in 100  $\mu\text{M}$  external  $\text{Ca}^{2+}$  at pH 7.4 and pH 4.0. (F) Normalized current amplitude at pH 4.0 and pH 7.4 was plotted as a function of external  $\text{Ca}^{2+}$  concentrations. The  $\text{Ca}^{2+}$  concentration required to block 50% of the monovalent currents was  $47.1 \pm 7.2 \mu\text{M}$  at pH 7.4 and  $5.6 \pm 0.7 \text{ mM}$  at pH 4.0 ( $n = 6$ ). (G) Original recordings of TRPM7 in 9.9 and 10  $\mu\text{M}$  external  $\text{Mg}^{2+}$  at indicated pH values. (H) Normalized current plotted as a function of  $\text{Mg}^{2+}$  concentrations at pH 7.4 and pH 4.0. The  $\text{IC}_{50}$  was  $5.9 \pm 1.2 \text{ mM}$  at pH 4.0, and  $5.4 \pm 0.5 \mu\text{M}$  at pH 7.4 (mean  $\pm$  SEM,  $n = 6$ ), respectively.

that protons compete with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for the binding sites in the TRPM7 pore.

We observed that protons produced different effects on outward currents under different external divalent cation conditions. For example, compared with the currents at pH 7.4 in 2 mM external  $\text{Ca}^{2+}$ , the outward current amplitude at pH 4.0 measured at +100 mV was changed by  $+41.1 \pm 5.4\%$  ( $n = 6$ ),  $+276.7 \pm 35.3\%$

( $n = 6$ ),  $-22.0 \pm 2.2\%$  ( $n = 6$ ),  $+24.8 \pm 3.4\%$  ( $n = 8$ ), and  $+26.2 \pm 8.9\%$  ( $n = 6$ ) in the external solutions containing 1 mM  $\text{Mg}^{2+}$ , 10 mM  $\text{Mg}^{2+}$ , 0.5 mM  $\text{Ca}^{2+}$ , 2 mM  $\text{Ca}^{2+}$ , and 10 mM  $\text{Ca}^{2+}$ , respectively (see Fig. 6, A and C, but the entire outward currents were not shown). The increase in outward current by low pH was larger in the external solutions with higher divalent concentrations; and the effect of protons on outward





**Figure 7.** Currents carried by protons through TRPM7 channels. (A) Original recordings in isotonic NMDG solution at pH 7.4 and pH 4.0 using P3 pipette solution (note the inward current induced at pH 4.0, red), and in 10  $\text{Ca}^{2+}$ /NMDG solution at pH 4.0 (green) in a HEK-293 cell overexpressing TRPM7. The proton current was largely blocked by 10  $\text{mM}$   $\text{Ca}^{2+}$ . (B) Averaged proton current at various test pH. Protons carry measurable currents at pH 5.0, and pronounced currents at pH 4.0 and 3.0 were observed (mean  $\pm$  SEM,  $n = 6$ ). \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ . (C and D) Anomalous mole fraction behavior of  $\text{H}^+$  permeation at pH 4.0. Current amplitude measured at  $-120$  mV at indicated  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  concentration was normalized in reference to the current amplitude at 10  $\mu\text{M}$  extracellular  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  (mean  $\pm$  SEM,  $n = 7$ ).

TRPM7 current was more pronounced in  $\text{Mg}^{2+}$ -containing than in  $\text{Ca}^{2+}$ -containing external solutions. This is presumably due to the fact that, at normal pH (7.4),  $\text{Mg}^{2+}$  exhibits a stronger block on TRPM7 outward current than that of  $\text{Ca}^{2+}$  (Kerschbaum et al., 2003; Monteilh-Zoller et al., 2003). Therefore, when the  $\text{Mg}^{2+}$  block is removed by protons, larger changes in TRPM7 outward current are observed. These results are in agreement with the notion that protons compete with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding sites in the TRPM7 pore.

If protons and divalent cations compete for binding sites, and the binding of protons to the external sites of the channel pore allows monovalent ions to pass through TRPM7, one would expect that the inhibitory effects of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  on monovalent currents should be influenced by proton concentrations. We therefore studied the inhibitory effects of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  on TRPM7 monovalent currents at pH 7.4 and pH 4.0. At pH 4.0, the free  $\text{Ca}^{2+}$  concentration was 7.7  $\mu\text{M}$  and the free  $\text{Mg}^{2+}$  concentration was 9.9  $\mu\text{M}$ , although the same concentrations of EDTA and EGTA can de-


crease free  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations to  $<1$  or 10 nM at pH 7.4 (see MATERIALS AND METHODS). Fig. 6 E shows representative recordings of TRPM7 in the presence of 100  $\mu\text{M}$   $\text{Ca}^{2+}$  at pH 4.0 and 7.4, respectively.  $\text{IC}_{50}$  was 47.1  $\mu\text{M}$   $\text{Ca}^{2+}$  at pH 7.4 and 5.6  $\text{mM}$   $\text{Ca}^{2+}$  at pH 4.0. Similarly, 10  $\mu\text{M}$   $\text{Mg}^{2+}$  produced more inhibition on TRPM7 at pH 7.4 than that at pH 4.0 (Fig. 6 G). The  $\text{IC}_{50}$  of  $\text{Mg}^{2+}$  at pH 4.0 (5.9  $\text{mM}$ ) is about 1,000-fold different from that at pH 7.4 (5.4  $\mu\text{M}$ ) (Fig. 6 H). The significantly decreased  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  affinities (Fig. 6, F and H) to TRPM7 channels in the acidic external solutions further indicate that protons compete with divalent  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding sites, thereby allowing monovalent cations to pass through TRPM7 channels.

#### Anomalous-mole Fraction Behavior of $\text{H}^+$ Permeation

As protons compete with divalent cations for binding sites in the channel pore, we tested whether protons pass through TRPM7 channels at low external divalent cation concentrations. No current was observed in the isotonic NMDG solution at pH 7.4, 7.0, or 6.0 (Fig. 7, A and B). However, inward currents were observed in isotonic NMDG solutions at pH 5.0, 4.0, and 3.0. Since NMDG is nonpermeant (Jiang et al., 2003; Kozak and Cahalan, 2003), it seems that the inward current is carried by protons. This notion is supported by the fact that current amplitude increased with increasing proton concentrations (Fig. 7 B). A high intracellular free  $\text{Mg}^{2+}$  concentration (3  $\text{mM}$ ), which inactivates TRPM7 channels, prevented the development of proton currents in TRPM7-expressing cells (unpublished data), suggesting that the proton-carried current was mediated by TRPM7 channels. Fig. 7 (C and D) shows anomalous mole fraction behavior of  $\text{H}^+$  permeation. The largest current amplitude was observed at 10  $\mu\text{M}$  extracellular  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$ , whereas the smallest current amplitude occurred at 10  $\text{mM}$  extracellular  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$ . These anomalous-mole fraction effects further indicate that protons compete with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding sites in the external pore of TRPM7, consistent with the results shown in Fig. 6.

#### Effects of Protons on the Endogenous TRPM7-like Current MIC/MagNuM in RBL Cells

Endogenous TRPM7-like currents MIC/MagNuM have been identified in a variety of cells (Nadler et al., 2001; Hermosura et al., 2002; Kozak et al., 2002; Prakriya and Lewis, 2002; Runnels et al., 2002; Aarts et al., 2003; Jiang et al., 2003). Since MIC/MagNuM in RBL cells have been well characterized (Hermosura et al., 2002; Kozak et al., 2002; Kozak and Cahalan, 2003, 2004), we chose to use RBL cells to study whether protons regulate endogenous MIC/MagNuM channels. As high concentration of intracellular  $\text{Ca}^{2+}$  blocks TRPM7 (Monteilh-Zoller et al., 2003), a pipette solution containing 15 nM



**Figure 8.** Effects of protons on MIC/MagNuM in RBL cells. Experiments were performed in external solutions with NaCl replaced by  $\text{NaSO}_3\text{CH}_3$ . (A) Representative MIC currents elicited by voltage ramps ranging from  $-120$  to  $+100$  mV in the external solutions at pH 7.4 (red) and 4.0 (black) in a RBL cell. (B) Concentration-dependent effects of protons on MIC inward currents in a representative RBL cell. The y axis was scaled to illustrate inward currents. (C) Inward current amplitude measured at  $-120$  mV at different test pH. External solution was changed to Tyrode's solution at pH 7.4 after each application of low pH external solution. (D) Concentration-dependent effects of protons on MIC currents. Best fit of the normalized change yielded an  $\text{EC}_{50} = 4.2 \pm 0.4$  (mean  $\pm$  SEM,  $n = 6$ ; Hill coefficient factor was 1).

$\text{Ca}^{2+}$  with weak buffering (1 mM EGTA) was used in this experiment. A low intracellular  $\text{Ca}^{2+}$  buffering condition (1 mM EGTA) was reported to be able to deactivate  $I_{\text{CRAC}}$  after its activation (Zweifach and Lewis, 1995) and was used for recording of TRPM7-like MIC/MagNuM currents in Jurkat cells (Prakriya and Lewis, 2002). We also included 5 mM CsCl in the external solution to eliminate the endogenous potassium current. NaCl was replaced by  $\text{NaSO}_3\text{CH}_3$  in the Tyrode's solution at pH 4.0 to prevent contamination from proton-activated  $\text{Cl}^-$  currents. After break-in, a voltage ramp protocol was used to monitor current for 10 min to ensure the complete deactivation of  $I_{\text{CRAC}}$  (Prakriya and Lewis, 2002) and full activation of TRPM7 (Kozak et al., 2002). Similar to TRPM7 currents in the heterologous expression system, the MIC/MagNuM currents in RBL cells were increased dramatically when the pH of the external Tyrode's solution changed from 7.4 to 4.0 (Fig. 8 A). The increase elicited by low pH was not affected by adding 200  $\mu\text{M}$  amiloride (unpublished data), suggesting that the increase in MIC/MagNuM current was not due to proton-activated  $\text{Na}^+$  channels. A best fit of normalized concentration-dependent increases in inward current (Fig. 8, B and C) yielded an  $\text{EC}_{50}$  of pH  $4.2 \pm 0.4$  (Fig. 8 D), which is similar to the pH value required to induce 50% of maximal response in the heterologously expressed TRPM7 currents (see Fig. 2 G). The similar response of TRPM7 and MIC/MagNuM to acidic pH provides further evidence that MIC/MagNuM is encoded by TRPM7.

## DISCUSSION

We have shown that low extracellular pH significantly enhanced TRPM7 inward current by increasing the

monovalent cation permeability. The mechanism by which protons increase monovalent currents is likely to be through competition with divalent cations for binding sites in the external pore of TRPM7, thereby removing the divalent cation block on the monovalent currents. The pH sensitivity of TRPM7 and native MIC/MagNuM channels suggests that TRPM7 may play an important role under acidic pathological conditions.

### TRPM7 Is an Acid-sensitive Ion Channel

Previous studies have shown that TRPM7 exhibits many unique features (Nadler et al., 2001; Runnels et al., 2001; Montell-Zoller et al., 2003; Schmitz et al., 2003), including the observation that it conducts small inward current at physiological voltages ( $-100$  to  $-40$  mV) and large outward current at  $+100$  mV, producing the characteristic outwardly rectifying I-V. The present study extends our understanding about TRPM7 by showing that TRPM7 is also a pH-sensitive ion channel. We show that a decrease in external pH evokes a marked increase in TRPM7 inward currents, with an  $\sim 10$ -fold increase at pH 4.0 and  $\sim 1$ -2-fold increase at pH 6.0 in the presence of a physiological external  $\text{Ca}^{2+}$  concentration (2 mM). The dramatic increase in TRPM7 inward current by pH 4.0 transforms the normally outwardly rectifying I-V curve to a double-rectifying I-V shape. We found that the increase in TRPM7 current by protons was more pronounced when the external  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  concentrations were decreased. Although the physiological function of TRPM7 is not completely understood, it has been shown that TRPM7 plays an important role in neuronal cell death caused by anoxia (Aarts et al., 2003). Given that native TRPM7 is only active to a small degree in the presence of physiological intracellular  $\text{Mg}^{2+}$ , an approximately one-

twofold increase in TRPM7 inward current at pH 6.0 suggests that TRPM7 may play an important role under acidic pathological conditions in which extracellular pH may decrease to pH 6.0 (Jacobus et al., 1977; Stevens et al., 1991; Steen et al., 1992).

Potential of TRPM7 inward currents by external protons is a novel feature of TRPM7. A previous study showed that a decrease in intracellular pH ( $pH_i$ ) inhibits monovalent  $Na^+$  currents through MIC channels in Jurkat T lymphocytes (Kerschbaum and Cahalan, 1998) and RBL cells (Braun et al., 2001). It is proposed that inhibition of MIC or TRPM7 by internal  $Mg^{2+}$ , other polyvalent cations, and  $H^+$  represents a general electrostatic cationic screening process (Kozak and Cahalan, 2003, 2004). We did not study how intracellular low pH affects TRPM7 in the present study, because the intracellular pH should not be changed due to the high concentration of HEPES buffering under our experimental conditions. The marked increase in TRPM7 inward current by external acidic pH shown in the present study and the inhibitory effects on TRPM7 outward currents by  $pH_i$  as previously reported (Kerschbaum and Cahalan, 1998; Braun et al., 2001; Kozak and Cahalan, 2003, 2004) indicate that TRPM7 is an acid-sensitive ion channel.

#### Potential Mechanism by which Protons Potentiate TRPM7 Inward Currents

We showed that both  $Ca^{2+}$  and  $Mg^{2+}$  exhibit anomalous mole-fraction effects at normal physiological pH. The apparent affinity of TRPM7 is  $47.1 \mu M$  for  $Ca^{2+}$  and  $5.4 \mu M$  for  $Mg^{2+}$ , similar to the previously reported  $Ca^{2+}$  ( $20 \mu M$ , at  $-120 mV$ ) (Fomina et al., 2000) and  $Mg^{2+}$  affinity ( $3 \mu M$ , at  $-120 mV$ ) to native MIC/MagNum channels (Kerschbaum et al., 2003). Under normal physiological  $Ca^{2+}$  ( $2 mM$ ) and  $Mg^{2+}$  ( $0.7$ – $1.1 mM$ ) concentrations (Konrad et al., 2004), we showed that monovalent cations contribute to the inward currents of TRPM7 (Fig. 3), and the contribution of monovalent currents becomes more pronounced under acidic conditions (pH 4.0, Fig. 4).

Several lines of evidence shown in the present study indicate that external protons increase TRPM7 inward currents by competing with divalent cations for binding sites in the TRPM7 pore, thereby enhancing monovalent cation permeability. First, there was a concentration-dependent increase in monovalent cation conductance and reversal potential for  $Na^+$ ,  $K^+$ , and  $Cs^+$  at pH 4.0, indicating that the enhanced inward TRPM7 currents resulted from an increased monovalent cation permeability. Second, the half-maximal pH was changed toward acidic pH direction by 0.6 and 1.7 pH units when the extracellular  $Ca^{2+}$  was increased from 0.5 to 2 and 10 mM, respectively; similarly, an increase of external  $Mg^{2+}$  concentration from 1 to 10 mM

shifted the half-maximal pH toward acidic pH direction by 1.0 pH unit (Fig. 6). The decreased proton affinity to TRPM7 at higher extracellular divalent concentrations indicate that protons compete with  $Ca^{2+}$  and  $Mg^{2+}$  for the same binding sites, therefore, at higher concentrations of divalent cations, more protons are required to induce 50% of maximal response. Third, high proton concentrations significantly decreased the affinity of  $Ca^{2+}$  and  $Mg^{2+}$  to TRPM7. The  $Ca^{2+}$  affinity to TRPM7 was decreased by  $\sim 100$ -fold and the  $Mg^{2+}$  affinity was decreased by  $\sim 1,000$ -fold when the external pH was changed from 7.4 to 4.0. Fourth, anomalous mole fraction permeation of protons indicate that protons compete with  $Ca^{2+}$  and  $Mg^{2+}$  for the binding sites in the external pore of TRPM7 (Fig. 7). Taken together, it seems that at physiological pH, TRPM7 only permeates a small inward current due to the  $Ca^{2+}$  and  $Mg^{2+}$  block on the monovalent current; whereas at acidic pH,  $Ca^{2+}$  and  $Mg^{2+}$  are outcompeted by protons, which relieves the block on the monovalent current and elicits a large inward current carried by monovalent cations.

Where are the binding sites for divalent cations and protons? A high affinity site for binding  $Mg^{2+}$  within the electric field and two low-affinity sites have been proposed for MIC channel by Kerschbaum et al. (2003) using the Eyring rate model. Kerschbaum et al. also proposed that the internal  $Mg^{2+}$  inhibits MIC in a voltage-independent manner, suggesting that internal  $Mg^{2+}$  is unable to access the pore from the inside (Kerschbaum et al., 2003). Our results show that external protons elicit a marked increase in TRPM7 currents at hyperpolarized potentials but only a small increase at positive potentials. These voltage-dependent effects suggest that protons can access the TRPM7 pore and compete with  $Mg^{2+}$  and  $Ca^{2+}$  for binding. The anomalous mole fraction behavior of  $Ca^{2+}$ ,  $Mg^{2+}$  (Fig. 3 F), and  $H^+$  permeation (Fig. 7, C and D) indicates that they can bind to the TRPM7 pore, and compete for binding sites within the pore (Fig. 6).

Proton competition for  $Ca^{2+}$  binding sites and the consequent channel opening has been proposed as the gating mechanism for ASIC3 (Immke and McCleskey, 2001, 2003), a proton-activated  $Na^+$  channel of the degenerin family (Waldmann and Lazdunski, 1998; Immke and McCleskey, 2003). Immke and McCleskey (2003) showed that the  $Ca^{2+}$  affinity was changed from  $K_d = 12 \mu M$  at pH 7.4 to  $K_d = 100 \mu M$  at pH 7.0, such that  $Ca^{2+}$  is released from a binding site and  $Na^+$  can pass through the channel at millimolar  $Ca^{2+}$  concentrations. The authors predict that, like the  $Ca^{2+}$  chelator EGTA (with four titratable acid groups), the titratable  $Ca^{2+}$  binding site of ASIC3 is able to bind four protons (Hill coefficient is 4). Our data suggest that protons enhance TRPM7 current by a similar competing mecha-

nism, such that binding of protons to TRPM7 relieves the blockade of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , thereby allowing  $\text{Na}^+$  to pass through TRPM7. Since the dose–response curves can be best fitted by sigmoidal dose–response equation (see MATERIALS AND METHODS) with the Hill coefficient factor of 1, we assume that there is probably one binding site. We do not know if the binding site is a specific amino acid residue or a site formed by several amino acid residues. It was reported that protonation of voltage-gated  $\text{Ca}^{2+}$  channels requires multiple carboxylates of the glutamic acid (Glu) residues to form a single high affinity site (Chen and Tsien, 1997). The Glu residues are also important for proton regulation on TRPV1 (Tominaga et al., 1998; Jordt et al., 2000; Ryu et al., 2003) and TRPV5 (Yeh et al., 2003). There are seven negatively charged Glu and Aspartic acid (Asp) residues between transmembrane domain 5 (TM5) and TM6 of TRPM7, which may be involved in  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , or  $\text{H}^+$  binding. The half-maximum pH (pH 4.5) for TRPM7 is also close to the pKa of free Glu ( $\sim$ pH 4.0) and Asp ( $\sim$ pH 3.8). Thus, it is possible that Glu or Asp in the TRPM7 pore may serve as the binding sites for external  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , and are also able to bind to protons, so that monovalent cations can readily pass through when the binding sites are occupied by protons. Alternatively, instead of competitive binding, protons may titrate away the block of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  on monovalent current by causing conformational changes. Further studies are required to elucidate the proton binding sites and detailed mechanisms by which protons increase TRPM7 inward monovalent currents.

#### Potential Significance

It has been suggested that TRPM7 plays an important physiological role in  $\text{Mg}^{2+}$  homeostasis, neuronal cell death, and cell viability (Nadler et al., 2001; Schlingmann et al., 2002; Walder et al., 2002; Aarts et al., 2003; Schmitz et al., 2003; Chubanov et al., 2004). We demonstrated here that acidic pH 6.0 increases TRPM7 inward current by approximately one- to twofold. Such an acidic condition (pH < 6) can occur during various forms of tissue injury (Jacobus et al., 1977; Stevens et al., 1991; Steen et al., 1992), or during repetitive nerve activities, ischemia, and seizures (Siesjo, 1988; Chesler and Kaila, 1992), suggesting that TRPM7 may play a role under acidic pathological conditions. However, without knowing the real physiological functions of TRPM7, it is difficult to predict a potential role of TRPM7 under acidic conditions. We have investigated if there are other factors that may change the pH sensitivity of TRPM7 closer to physiological pH. With 20  $\mu\text{M}$   $\text{PIP}_2$  in the pipette solution, the  $\text{EC}_{50}$  was pH  $4.7 \pm 0.7$  (mean  $\pm$  SEM,  $n = 6$ , Hill coefficient 1.0; unpublished data), which is not significantly different from the  $\text{EC}_{50}$

obtained without  $\text{PIP}_2$  in the pipette solution (Fig. 2). In addition, with normal physiological intracellular  $\text{Mg}^{2+}$  concentrations in a native system, the MIC/MagNum channel is only active to a small degree. Further studies are required to elucidate the physiological and pathological roles of TRPM7 under normal and acidic conditions.

Like TRPM7, the native MIC/MagNum current in RBL cells showed a similar acidic potentiation to that seen for TRPM7 currents in the heterologous expression system (Fig. 8), indicating that MIC/MagNum is encoded by TRPM7. A recent study by Gwanyanya et al. (2004) showed that MIC in cardiac myocytes and RBL cells were inhibited by acidic pH. It is not clear why there is a discrepancy between our and their results. One difference is that they evaluated pH effects on MIC in divalent-free solutions, whereas we used physiological external solutions containing divalent cations in the present study.

Protons have been reported to regulate channel activities in different channel superfamilies (Hille, 2003; Holzer, 2003), including TRP channel superfamily. It has been shown that TRPV1 and TRPV4 are enhanced by low pH, whereas TRPV5 (Yeh et al., 2003) and TRPM5 (Liu et al., 2005) are inhibited by protons. We show here that protons dramatically potentiate TRPM7 inward current by competing with divalent cations for binding sites. The pH sensitivity is a novel feature of TRPM7, and the results in the present study not only provide a clue as to the potential functions of TRPM7 in vivo, but also may help to identify the amino acid residues that are important in the ion selectivity of TRPM7 channels.

#### Potential Limitations

Proton permeation has been observed for voltage-gated  $\text{Na}^+$  channels when the external solution is free of  $\text{Na}^+$  (Mozhayeva and Naumov, 1983; Hille, 2003). We showed anomalous mole fraction behavior of  $\text{H}^+$  permeation in the external solutions free of permeant monovalent cations (Fig. 7). In 2 mM  $\text{Ca}^{2+}$ /NMDG/pH 4.0 solution, the proton carried current is 4.4 pA/pF, corresponding to 17% of the current amplitude obtained in 10  $\mu\text{M}$   $\text{Ca}^{2+}$ /NMDG/pH 4.0 (Fig. 7 C). This proton current amplitude is similar to the value shown in Fig. 5 B (6.5 pA/pF, pH 4.0), indicating that the current at pH 4.0 (Fig. 5 B) is mainly carried by protons, and consistent with the notion that protons compete with divalent cations and therefore almost blocked all the divalent cation current under the conditions shown in Fig. 5 (A and B).

It is possible that in the normal Tyrode's solution at pH 4.0, protons may pass through TRPM7 along with  $\text{Na}^+$  or  $\text{K}^+$ . If this were the case, proton-carried current may have contributed to the inward monovalent cur-

rents at pH 4.0. However, given that the current amplitude (6.5 pA/pF, Fig. 5 B) in NMDG solution at pH 4.0 is only ~2.5% of the inward current amplitude (247 pA/pF, Fig. 2) obtained in normal Tyrode's solutions at pH 4.0, contribution of proton-carried inward currents in the normal Tyrode's solutions at pH 4.0 should be <2.5%, and should not contaminate the experimental results.

At pH 3.0, the I-V curve of TRPM7 elicited by voltage ramps seems different from those at pH >3.0 (Fig. 2 E), and the TRPM7 outward currents were inhibited by pH 3.0. In addition, a strong inactivation or desensitization was observed (Fig. 2 F) when cells were continuously exposed to the external solution at pH 3.0. The following potential mechanisms may account for the above observations. First, TRPM7 is desensitized at pH 3.0; second, protons may pass through TRPM7, resulting in low intracellular pH, which inhibits TRPM7 outward current (Kozak and Cahalan, 2004); and third, protons may exhibit complex effects on TRPM7. Further studies are required to reveal the detailed mechanisms.

### Conclusion

In conclusion, we have demonstrated that acidic pH significantly increases TRPM7 inward monovalent currents by competing with divalent cations for binding sites. The pH sensitivity represents a novel feature of TRPM7. We showed that MIC/MagNum currents were similarly potentiated by protons, suggesting that MIC/MagNum is encoded by TRPM7. The large TRPM7 inward current elicited by low pH suggests that TRPM7 may play a role under acidic pathological conditions. Further studies are required to elucidate the mechanism by which protons potentiate TRPM7, as well as the potential significance of TRPM7 under acidic pathological conditions.

We thank Dr. Andrew Scharenberg for providing the HEK-293 cells stably expressing TRPM7. We thank Dr. Alan Fein for constructive suggestions and comments. And we'd like to thank David Clapham, Bruce Liang, Kim Dodge, Haoxin Xu, Loren Rannels, Laurinda Jaffe, and Dejian Ren for helpful comments.

This work was supported in part by an award from the American Heart Association and by a National Institutes of Health grant R01 HL078960-01A1 to L. Yue.

Olaf S. Andersen served as editor.

Submitted: 22 September 2004

Accepted: 10 June 2005

### REFERENCES

Aarts, M., K. Iihara, W.L. Wei, Z.G. Xiong, M. Arundine, W. Cerwinski, J.F. MacDonald, and M. Tymianski. 2003. A key role for TRPM7 channels in anoxic neuronal death. *Cell*. 115:863–877.

Askwith, C.C., J.A. Wemmie, M.P. Price, T. Rokhlina, and M.J. Welsh. 2004. Acid-sensing ion channel 2 (ASIC2) modulates ASIC1 H<sup>+</sup>-activated currents in hippocampal neurons. *J. Biol.*

*Chem.* 279:18296–18305.

Braun, F.J., L.M. Broad, D.L. Armstrong, and J.W. Putney Jr. 2001. Stable activation of single Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> channels in divalent cation-free solutions. *J. Biol. Chem.* 276:1063–1070.

Chen, X.-H., and R.W. Tsien. 1997. Aspartate substitutions establish the concerted action of P-region glutamates in repeats I and III in forming the protonation site of L-type Ca<sup>2+</sup> channels. *J. Biol. Chem.* 272:30002–30008.

Cherny, V.V., L.M. Henderson, and T.E. DeCoursey. 1997. Proton and chloride currents in Chinese hamster ovary cells. *Membr. Cell Biol.* 11:337–347.

Chesler, M., and K. Kaila. 1992. Modulation of pH by neuronal activity. *Trends Neurosci.* 15:396–402.

Chubanov, V., S. Waldegger, M. Mederos y Schnitzler, H. Vitzthum, M.C. Sassen, H.W. Seyberth, M. Konrad, and T. Gudermann. 2004. Disruption of TRPM6/TRPM7 complex formation by a mutation in the TRPM6 gene causes hypomagnesemia with secondary hypocalcemia. *Proc. Natl. Acad. Sci. USA.* 101:2894–2899.

Clapham, D.E. 2003. TRP channels as cellular sensors. *Nature.* 426: 517–524.

Fleig, A., and R. Penner. 2004. Emerging roles of TRPM channels. *Novartis Found. Symp.* 258:248–258; discussion 258–266.

Fomina, A.F., C.M. Fanger, J.A. Kozak, and M.D. Cahalan. 2000. Single channel properties and regulated expression of Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> (CRAC) channels in human T cells. *J. Cell Biol.* 150:1435–1444.

Gwanyanya, A., B. Amuzescu, S.I. Zakharov, R. Macianskiene, K.R. Sipido, V.M. Bolotina, J. Vereecke, K. Mubagwa. 2004. Magnesium-inhibited, TRPM6/7-like channel in cardiac myocytes: permeation of divalent cations and pH-mediated regulation. *J. Physiol.* 559:761–776.

Hanano, T., Y. Hara, J. Shi, H. Morita, C. Umebayashi, E. Mori, H. Sumimoto, Y. Ito, Y. Mori, and R. Inoue. 2004. Involvement of TRPM7 in cell growth as a spontaneously activated Ca<sup>2+</sup> entry pathway in human retinoblastoma cells. *J. Pharmacol. Sci.* 95:403–419.

Harteneck, C., T.D. Plant, and G. Schultz. 2000. From worm to man: three subfamilies of TRP channels. *Trends Neurosci.* 23:159–166.

Hermosura, M.C., M.K. Monteilh-Zoller, A.M. Scharenberg, R. Penner, and A. Fleig. 2002. Dissociation of the store-operated calcium current I(CRAC) and the Mg-nucleotide-regulated metal ion current MagNum. *J. Physiol.* 539:445–458.

Hille, B. 2003. *Ion Channels of Excitable Membranes*. Third edition. Sinauer Associates, Inc. Sunderland, MA. 471–502.

Holzer, P. 2003. Acid-sensitive ion channels in gastrointestinal function. *Curr. Opin. Pharmacol.* 3:618–625.

Immke, D.C., and E.W. McCleskey. 2001. Lactate enhances the acid-sensing Na<sup>+</sup> channel on ischemia-sensing neurons. *Nat. Neurosci.* 4:869–870.

Immke, D.C., and E.W. McCleskey. 2003. Protons open acid-sensing ion channels by catalyzing relief of Ca<sup>2+</sup> blockade. *Neuron.* 37:75–84.

Jacobus, W.E., G.J. Taylor, D.P. Hollis, and R.L. Nunnally. 1977. Phosphorus nuclear magnetic resonance of perfused working rat hearts. *Nature.* 265:756–758.

Jiang, X., E.W. Newell, and L.C. Schlichter. 2003. Regulation of a TRPM7-like current in rat brain microglia. *J. Biol. Chem.* 278: 42867–42876.

Jordt, S.E., M. Tominaga, and D. Julius. 2000. Acid potentiation of the capsaicin receptor determined by a key extracellular site. *Proc. Natl. Acad. Sci. USA.* 97:8134–8139.

Kerschbaum, H.H., and M.D. Cahalan. 1998. Monovalent permeability, rectification, and ionic block of store-operated calcium channels in Jurkat T lymphocytes. *J. Gen. Physiol.* 111:521–537.

- Kerschbaum, H.H., J.A. Kozak, and M.D. Cahalan. 2003. Polyvalent cations as permeant probes of MIC and TRPM7 pores. *Biophys. J.* 84:2293–2305.
- Konrad, M., K.P. Schlingmann, and T. Gudermann. 2004. Insights into the molecular nature of magnesium homeostasis. *Am. J. Physiol. Renal Physiol.* 286:F599–F605.
- Kozak, J.A., and M.D. Cahalan. 2003. MIC channels are inhibited by internal divalent cations but not ATP. *Biophys. J.* 84:922–927.
- Kozak, J.A., and M.D. Cahalan. 2004. Regulation of MIC and TRPM7 channels by internal polyvalent cations. *Biophys. J.* Abstract.
- Kozak, J.A., H.H. Kerschbaum, and M.D. Cahalan. 2002. Distinct properties of CRAC and MIC channels in RBL cells. *J. Gen. Physiol.* 120:221–235.
- Liu, D., Z. Zhang, and E.R. Liman. 2005. Extracellular acid block and acid-enhanced inactivation of the Ca<sup>2+</sup>-activated cation channel TRPM5 involve residues in the S3-S4 and S5-S6 extracellular domains. *J. Biol. Chem.* 280:20691–20699.
- Monteilh-Zoller, M.K., M.C. Hermosura, M.J. Nadler, A.M. Scharenberg, R. Penner, and A. Fleig. 2003. TRPM7 provides an ion channel mechanism for cellular entry of trace metal ions. *J. Gen. Physiol.* 121:49–60.
- Montell, C. 2001. Physiology, phylogeny, and functions of the TRP superfamily of cation channels. *Sci STKE.* 2001:RE1.
- Mozhayeva, G.N., and A.P. Naumov. 1983. The permeability of sodium channels to hydrogen ions in nerve fibres. *Pflugers Arch.* 396:163–173.
- Nadler, M.J., M.C. Hermosura, K. Inabe, A.L. Perraud, Q. Zhu, A.J. Stokes, T. Kurosaki, J.P. Kinet, R. Penner, A.M. Scharenberg, and A. Fleig. 2001. LTRPC7 is a Mg.ATP-regulated divalent cation channel required for cell viability. *Nature.* 411:590–595.
- Prakriya, M., and R.S. Lewis. 2002. Separation and characterization of currents through store-operated CRAC channels and Mg<sup>2+</sup>-inhibited cation (MIC) channels. *J. Gen. Physiol.* 119:487–507.
- Runnels, L.W., L. Yue, and D.E. Clapham. 2001. TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science.* 291:1043–1047.
- Runnels, L.W., L. Yue, and D.E. Clapham. 2002. The TRPM7 channel is inactivated by PIP(2) hydrolysis. *Nat. Cell Biol.* 4:329–336.
- Ryazanova, L.V., M.V. Dorovkov, A. Ansari, and A.G. Ryazanov. 2004. Characterization of the protein kinase activity of TRPM7/ChaK1, a protein kinase fused to TRP ion channel. *J. Biol. Chem.* 279:3708–3716.
- Ryu, S., B. Liu, and F. Qin. 2003. Low pH potentiates both capsaicin binding and channel gating of VR1 receptors. *J. Gen. Physiol.* 122:45–61.
- Schlingmann, K.P., S. Weber, M. Peters, L. Niemann Nejsum, H. Vitzthum, K. Klingel, M. Kratz, E. Haddad, E. Ristoff, D. Dinour, et al. 2002. Hypomagnesemia with secondary hypocalcemia is caused by mutations in TRPM6, a new member of the TRPM gene family. *Nat. Genet.* 31:166–170.
- Schmitz, C., A.L. Perraud, C.O. Johnson, K. Inabe, M.K. Smith, R. Penner, T. Kurosaki, A. Fleig, and A.M. Scharenberg. 2003. Regulation of vertebrate cellular Mg<sup>2+</sup> homeostasis by TRPM7. *Cell.* 114:191–200.
- Siesjo, B.K. 1988. Acidosis and ischemic brain damage. *Neurochem. Pathol.* 9:31–88.
- Steen, K.H., P.W. Reeh, F. Anton, and H.O. Handwerker. 1992. Protons selectively induce lasting excitation and sensitization to mechanical stimulation of nociceptors in rat skin, in vitro. *J. Neurosci.* 12:86–95.
- Stevens, C.R., R.B. Williams, A.J. Farrell, and D.R. Blake. 1991. Hypoxia and inflammatory synovitis: observations and speculation. *Ann. Rheum. Dis.* 50:124–132.
- Takezawa, R., C. Schmitz, P. Demeuse, A.M. Scharenberg, R. Penner, and A. Fleig. 2004. Receptor-mediated regulation of the TRPM7 channel through its endogenous protein kinase domain. *Proc. Natl. Acad. Sci. USA.* 101:6009–6014.
- Tominaga, M., M.J. Caterina, A.B. Malmberg, T.A. Rosen, H. Gilbert, K. Skinner, B.E. Raumann, A.I. Basbaum, and D. Julius. 1998. The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron.* 21:531–543.
- Vennekens, R., J.G.J. Hoenderop, J. Prenen, M. Stuver, P.H.G.M. Willems, G. Droogmans, B. Nilius, and R.J.M. Bindels. 2000. Permeation and gating properties of the novel epithelial Ca<sup>2+</sup> channel. *J. Biol. Chem.* 275:3963–3969.
- Walder, R.Y., D. Landau, P. Meyer, H. Shalev, M. Tsolia, Z. Borochowitz, M.B. Boettger, G.E. Beck, R.K. Englehardt, R. Carmi, and V.C. Sheffield. 2002. Mutation of TRPM6 causes familial hypomagnesemia with secondary hypocalcemia. *Nat. Genet.* 31:171–174.
- Waldmann, R., and M. Lazdunski. 1998. H<sup>+</sup>-gated cation channels: neuronal acid sensors in the NaC/DEG family of ion channels. *Curr. Opin. Neurobiol.* 8:418–424.
- Yeh, B.-I., T.-J. Sun, J.-Z. Lee, H.-H. Chen, and C.-L. Huang. 2003. Mechanism and molecular determinant for regulation of rabbit transient receptor potential type 5 (TRPV5) channel by extracellular pH. *J. Biol. Chem.* 278:51044–51052.
- Yermolaieva, O., A.S. Leonard, M.K. Schnizler, F.M. Abboud, and M.J. Welsh. 2004. Extracellular acidosis increases neuronal cell calcium by activating acid-sensing ion channel 1a. *Proc. Natl. Acad. Sci. USA.* 101:6752–6757.
- Yue, L., J.L. Feng, Z. Wang, and S. Nattel. 2000. Effects of ambasilide, quinidine, flecainide and verapamil on ultra-rapid delayed rectifier potassium currents in canine atrial myocytes. *Cardiovasc. Res.* 46:151–161.
- Yue, L., B. Navarro, D. Ren, A. Ramos, and D. Clapham. 2002. The cation selectivity filter of the bacterial sodium channel, NaCh-Bac. *J. Gen. Physiol.* 160:845–853.
- Yue, L., J.B. Peng, M.A. Hediger, and D.E. Clapham. 2001. CaT1 manifests the pore properties of the calcium-release-activated calcium channel. *Nature.* 410:705–709.
- Zweifach, A., and R.S. Lewis. 1995. Slow calcium-dependent inactivation of depletion-activated calcium current. Store-dependent and -independent mechanisms. *J. Biol. Chem.* 270:14445–14451.