

ORIGINAL ARTICLE

Impaired procedural memory in narcolepsy type 1

Amanda Asp¹ | Frida Lund¹ | Christian Benedict²  | Pontus Wasling^{1,3} 

¹Department of Clinical Neuroscience, Institute of Neuroscience and Physiology, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

²Molecular Neuropharmacology (Sleep Science Lab), Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden

³Department of Neurology, Sahlgrenska University Hospital, Gothenburg, Sweden

Correspondence

Pontus Wasling, Department of Clinical Neuroscience, Institute of Neuroscience and Physiology, Sahlgrenska Academy at Gothenburg University, Blå Stråket 7, PO Box 430, SE-40530 Gothenburg, Sweden. Email: pontus.wasling@gu.se

Funding information

Neuro Association, grant F2020-0121

Objectives: Sleep enhances the consolidation of memories. Here, we investigated whether sleep-dependent memory consolidation differs between healthy subjects and narcolepsy type 1 (NT1) patients.

Material and Methods: We recruited 18 patients with NT1 and 24 healthy controls. The consolidation of spatial (declarative memory; 2-dimensional object location) and procedural (non-declarative memory; finger sequence tapping) memories was examined across one night of at-home sleep. Sleep was measured by an ambulatory sleep recording device.

Results: The overnight gain in the number of correctly recalled sequences in the finger-tapping test was smaller for NT1 patients than healthy subjects (+8.1% vs. +23.8% from pre-sleep learning to post-sleep recall, $p = .035$). No significant group differences were found for the overnight consolidation of spatial memory. Compared to healthy subjects, the sleep of NT1 patients was significantly more fragmented and shallow. However, no significant correlations were found between sleep parameters and overnight performance changes on the memory tests in the whole group.

Conclusion: The sleep-dependent consolidation of procedural but not spatial memories may be impaired among patients with NT1. Therefore, future studies are warranted to examine whether sleep improvement, for example, using sodium oxybate, can aid the sleep-dependent formation of procedural memories among NT1 patients.

1 | INTRODUCTION

Nocturnal sleep represents a time of the day where the consolidation of newly encoded procedural and explicit memories is facilitated.¹ However, under conditions of disturbed sleep, the consolidation of newly acquired memories during sleep appears dysfunctional. For example, a study involving young adults demonstrated that the consolidation of verbal memories was impaired when sleep in the post-learning night was fragmented.² Furthermore, reduced time in slow-wave sleep, a non-rapid eye movement (REM) sleep stage, has been linked to a less pronounced consolidation of declarative information.³ Sleep deprivation reduces success rate on declarative and procedural memory tasks,^{4,5} and a small study showed that

sleep-associated consolidation of procedural memories might be impaired among patients with primary insomnia.⁶

Narcolepsy is a chronic sleep disorder characterized by excessive daytime sleepiness (EDS) and disturbed nocturnal sleep. A major sleep characteristic of narcolepsy is sleep-onset REM periods (SOREMPs); however, non-REM sleep is also affected. Patients with narcolepsy often spend more time in lighter and less time in deeper non-REM sleep stages.⁷⁻⁹ Frequent nocturnal awakenings from sleep are also highly prevalent in narcolepsy.^{8,10,11} Thus, narcolepsy may interfere with sleep-dependent memory consolidation. In line with this assumption, one study found that the consolidation of procedural memories across one night of sleep was less pronounced among patients with narcolepsy than healthy controls.¹² However,

Amanda Asp and Frida Lund contributed equally to this study.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Acta Neurologica Scandinavica* published by John Wiley & Sons Ltd.

a recent review concluded that studies on memory and learning show largely unimpaired functions in patients with narcolepsy.¹³ In contrast, a common complaint among patients is their perceived impairment in memory functions. Whether the sleep-dependent consolidation of explicit memories is altered by narcolepsy has not been examined yet.

Therefore, in the present study, we compared the overnight consolidation of newly acquired procedural skills and spatial information (a type of explicit memory) between healthy controls and patients suffering from narcolepsy type 1 (NT1). Given their disturbed sleep, we hypothesized that the sleep-dependent consolidation of procedural and spatial memories would be less pronounced among NT1 patients than healthy subjects

2 | METHODS

2.1 | NT1 patients

Eighteen patients (age range: 19–32; 56% female) fulfilling the criteria for NT1 diagnosis according to The International Classification of Sleep Disorders 3rd edition (ICSD-3)¹⁴ participated in the study. All patients had documented descriptions of cataplexy, with a sudden loss of muscle tone in the neck, face and/or arms and legs related to the onset of an emotional trigger, such as laughter, anger, and/or surprise. In addition, EDS was present in all subjects.

As shown by a multiple sleep latency test (MSLT), 11 patients had a sleep latency ≤ 8 min and ≥ 2 SOREMPs. The remaining seven NT1 patients fulfilled one of these two MSLT criteria. Lumbar puncture performed in seven patients revealed cerebrospinal fluid (CSF) orexin-A levels below 133 pg/ml in six patients, that is, the diagnostic criteria for NT1.¹⁵ The cutoff level was set according to the ICSD-3 criteria to one-third of the normal reference level (>400 pg/ml) for the in-house method presently used in the study. Three of the subjects not fulfilling both MSLT criteria had low orexin-A levels in their CSF. Out of the four remaining subjects, three had a sleep latency >8 min and one had 0 SOREMPs, but all four had given credible descriptions of cataplexy and EDS. Seven of the narcolepsy subjects took one wake-promoting medication, and three subjects took two wake-promoting drugs. Seven were medicated with antidepressants (selective serotonin/noradrenalin uptake inhibitors) to alleviate cataplexy. Eight of the narcoleptic subjects were ordinarily medicated

with sodium oxybate but withheld from taking that drug during the study night. One participant with narcolepsy did not take any medication.

2.2 | Healthy subjects

Twenty-four healthy young adults (age range: 20–26 years; 46% female) were recruited from the local community in Gothenburg (Sweden). Inclusion criteria comprised age between 18 and 35 years, general good health status, and normal sleep habits (sleep duration between 7–9 h/night and self-reported sleep-onset time between 9.30 p.m. and 01.00 a.m.). Subjects were not invited to participate in the study when they reported a sleep disorder or the use of sleep medication. Three participants took prescription drugs (insulin, mood stabilizers, and hormonal treatment); the remaining subjects were free of medication.

2.3 | Study protocol

This prospective case-control study was conducted between February and May 2021 and was approved by the Ethical Review Board in Uppsala, Sweden, (#2020-03328). All subjects gave written informed consent to the study that conformed to the Declaration of Helsinki. The study was performed in at-home settings, and in each study night, subjects were told to go to bed when they felt sufficiently sleepy and go up the following day at their usual wake-up time.

Before the experimental night, each participant took part in an adaption night, familiarizing them with wearing the sleep device (Dreem 2 SAS). This device is worn as a headband, including six dry electroencephalogram (EEG)-sensors, a pulse oximeter, and a three-dimensional accelerometer. It provides the sleep-related physiological signals of heart rate, respiratory rate, body movements, and body position. In addition, a deep learning algorithm enables automatic sleep stage analysis.¹⁶ Before the onset of the adaption night, subjects also filled in the Epworth Sleepiness Scale (ESS),¹⁷ the Hospital Anxiety and Depression Scale (HADS),¹⁸ and their BMI was measured. A sum over the cutoff value 11 was used to define depression and anxiety in the HADS-Depression and HADS-Anxiety subscales, respectively.¹⁸




FIGURE 1 Experimental design. During the learning session in the evening, all subjects performed the finger-tapping test (FTT) and the 2D memory game (2DMG), and within 3 h, they went to sleep. The test session was set within 1 h after wake-up, where they performed the test versions of two tasks

During the experimental night, all participants underwent first the finger tapping and then the 2D-object location task (for a description, please see below). The learning session (45 min) was scheduled 3h before each participant's usual bedtime. Re-testing (15 min) was scheduled the following day within 1h after habitual awakening (Figure 1) and before the intake of wake-promoting medication. Explicit (or declarative) and implicit (or non-declarative) memory formation constitute the two main forms of long-term memories in man.¹⁹ One memory test for each memory domain was chosen in the present study.

2.4 | Finger-tapping test

The computerized finger-tapping test has previously been used to examine the consolidation of procedural memory across sleep.²⁰ Non-declarative procedural learning such as acquisition of a new motor skill is thought to be mediated by cortical-striatal-cerebellar based circuits.²¹

In this task, participants used their non-dominant hand to type a 5-digit target sequence displayed on a computer screen (e.g., 4-1-3-2-4) as fast and accurately as possible. Learning consisted of 12 blocks of 30 s each (with 30-s breaks between two consecutive blocks).

At re-testing (i.e., after sleep), participants had three 30-s blocks to tap the target sequence. The overnight gain was defined by dividing the increased number of correctly tapped sequences at re-testing by the number of correctly tapped sequences of the final three learning trials, times 100.

2.5 | Two-dimensional (2-D) memory task

The 2D-object location task resembles the board game Memory. The task reliably measures the sleep-dependent consolidation of hippocampus-dependent memories.²² Participants were shown 15

card pairs of pictures showing animals or everyday objects on a computer screen (5*6 matrix). After two repetitions, the immediate recall session started to ensure sufficient encoding. They were instructed to use the mouse to find the second card after the first card of a pair had been uncovered. After each choice, the correct location of the second card was presented to stimulate re-encoding. The pre-sleep learning session was over when at least nine card pair locations were correctly learned.

At re-testing (i.e., after sleep), participants were again instructed to recall as many card pair locations as possible on the computer screen. The overnight consolidation success was calculated by dividing the number of correctly located card pairs at re-testing by the number of correctly recalled card pair locations at learning times 100.

2.6 | Data analysis

Data were analyzed using GraphPad Prism Software version 8.4.2 for Mac (GraphPad Software). Data are reported as mean \pm standard deviation (SD), unless otherwise stated. Group differences in demographic data, questionnaire scores, dependent variables in memory testing (i.e., trials until criterion, learning, recall, and overnight change in memory performance), and sleep variables were analyzed with the Mann-Whitney *U*-test, as most distributions were skewed according to the Shapiro-Wilk test. Chi-square exact test was used for between-group comparisons of dichotomous variables. The Wilcoxon signed-rank test was applied in cases of paired non-parametric analysis. We investigated possible interactions of the within-subjects factor session (learning/test) with the between-subjects factor health status (healthy control/NT1) with repeated measures analyses of variances (ANOVA) and Sidak's multiple comparison test was used post hoc. To analyze associations between results from the memory tasks, demographic data, questionnaire scores, and sleep variables, bivariate associations were performed. All comparisons were two-tailed, and significance was set at the .05 level.

	Healthy controls(n = 24)	Narcolepsy(n = 18)	Statistics ^a
Age (years)	23.1 \pm 3.0	25.3 \pm 3.0	<i>p</i> = .0028
Sex (% women)	45.8	55.5	<i>p</i> = .53
BMI (kg/m ²)	22.9 \pm 3.3	23.2 \pm 4.2	<i>p</i> = .83
HADS-D	2.6 \pm 2.5	6.2 \pm 3.2	<i>p</i> < .0002
% with depression (n)	0 (0)	16.7 (3)	<i>p</i> = .038
HADS-A	5.3 \pm 3.0	8.7 \pm 3.7	<i>p</i> = .0039
% with anxiety (n)	8.3 (2)	27.8 (5)	<i>p</i> = .09
ESS	6.3 \pm 3.6	15.4 \pm 3.3	<i>p</i> < .0001

TABLE 1 Group characteristics

Note. All values are expressed as mean \pm standard deviation.

Abbreviations: BMI, body mass index; ESS, Epworth sleepiness scale; HADS-A, hospital anxiety and depression scale - anxiety; HADS-D, hospital anxiety and depression scale - depression.

^aBolded analyses = statistically significant (*p* < .05) with Mann-Whitney *U* test in all comparisons, except χ^2 for gender and proportion of subjects with depression and anxiety.

3 | RESULTS

3.1 | Group characteristics

The majority of participants in both groups were in their twenties; however, NT1 patients were on average slightly older than healthy subjects. No significant group differences in gender ratio or BMI were observed. As expected, NT1 patients scored significantly higher on ESS compared to the controls. Further characteristics can be found in [Table 1](#).

Compared to healthy subjects, NT1 patients were more awake after onset of sleep, woke up more often during the night, had a lower sleep efficiency, a shorter sleep time, spent more time in sleep stage N2 and less time in both N3 and REM during the post-learning night. More sleep characteristics can be found in [Table 2](#).

3.2 | Finger sequence tapping task

The mean number of correctly recalled sequences during the last three learning blocks and the three re-testing blocks did not significantly differ between healthy subjects and NT1 patients ([Table 3](#)). After on night of sleep, the number of correctly recalled sequences increased in healthy controls, but not in NT1 (from 17.5 ± 4.8 to 21.0 ± 4.7 in HC, $p = .037$; from 17.9 ± 4.7 to 19.0 ± 4.1 in NT1, $p = .42$; [Figure 2A](#)). When calculating the overnight performance change, we found that healthy subjects showed a performance gain (percentual change in accuracy) of about 23.8%, whereas NT patients only increased their performance by about 8.1% between the learning and re-testing session ([Figure 2B](#); $p = .035$ for NT1 group vs. healthy group). In line with this result pattern, a repeated-measures ANOVA utilizing *TIME* as within-subjects factor (i.e., learning vs. re-testing),

HEALTH as between-subjects factor (i.e., healthy control vs. NT1 patients), and the number of correctly recalled sequences as dependent variable revealed significant effects of *TIME* ($F[1,42] = 28.11$, $p < .0001$) and *TIME*HEALTH* ($F[1, 42] = 7.835$, $p = .0078$) but not *HEALTH* ($F[1, 42] = 0.39$, $p = .54$).

3.3 | 2D memory task

As shown in [Table 3](#), the number of trials required to remember at least nine card pairs and the number of card pairs recalled during the learning session for the 2D memory task did not differ between healthy subjects and subjects with NT1. No overnight performance change was observed in healthy controls, whereas a decline in performance was observed in NT1 subjects (-6.4 ± 3.6 in HC, $p = .13$; 15.8 ± 4.1 in NT1, $p = .002$; [Figure 3A](#)). However, the overnight change in the number of correctly recalled card pairs did not differ between the NT1 and healthy control groups ([Figure 3B](#)). A repeated-measures ANOVA revealed a main effect of *TIME* ($F[1, 42] = 15.97$, $p = .0003$). In contrast, neither *HEALTH* ($F[1, 42] = 0.37$, $p = 0.38$) nor the interaction *TIME*HEALTH* reached significance ($F[1, 42] = 2.78$, $p = .10$).

3.4 | Correlation between sleep parameters and overnight consolidation scores

We next explored whether the finger sequence tapping performance or the 2D memory task, that is, relative performance change from pre-sleep to post-sleep test, was related to any of the sleep parameters in the entire group. None of the correlational analyses reached significance.

TABLE 2 Sleep variables in subjects with narcolepsy and healthy controls

	Healthy controls(n = 24)	Narcolepsy type 1(n = 18)	Statistics ^a
TST (min)	431.8±37.8	355.8±94.9	p = .0012
SL (min)	23.8±15.6	19.3±1.5	p = .14
WASO (min)	13.2±2.1	55.0±62.1	p < .0001
SE (%)	91.7±4.4	82.5±16.2	p = .033
N1 (% of TST)	0.1±0.2	0.4±0.8	p = .087
N2 (% of TST)	45.1±7.1	61.7±15.5	p < .0001
N3 (% of TST)	24.5±6.3	18.0±8.8	p = .012
REM (% of TST)	30.3±7.9	19.9±11.3	p = .013
NOW (n)	4.1±2.4	8.3±3.4	p < .0001
HR	55.7±6.5	61.1±6.4	p = .012
RR	15.0±1.6	14.8±1.6	p = .89

Note. All values are presented as mean ± standard deviation.

Abbreviations: HR, heart rate; N1, sleep stage 1; N2, sleep stage 2; N3, sleep stage 3; NOW, number of wakes; REM, rapid eye movement sleep; RR, respiratory rate; SE, sleep efficiency; SL, sleep-onset latency; TST, total sleep time; WASO, wake after sleep onset.

^aBolded analyses = statistically significant ($p < .05$) with Mann-Whitney *U* test.

	Healthy controls	Narcolepsy	Statistics ^a
Finger-tapping task			
No. of correctly tapped sequences			
Learning	17.5 ± 4.8	17.9 ± 4.7	$p = .82$
Test	21.0 ± 4.7	19.0 ± 4.1	$p = .36$
Overnight change (sequences)	3.5 ± 2.6	1.1 ± 2.9	$p = .009$
Overnight change (%)	23.8 ± 26.0	8.1 ± 17.6	$p = .035$
2D Memory task			
Trials to the 60% criterion	2.6 ± 2.0	2.9 ± 1.6	$p = .25$
No. of correctly recalled card pairs			
Learning	11.0 ± 1.4	11.1 ± 2.0	$p = .87$
Test	10.3 ± 2.2	9.2 ± 2.2	$p = .20$
Overnight change (card pairs)	-0.7 ± 2.0	-1.8 ± 2.0	$p = .15$
Overnight change (%)	-6.4 ± 17.7	-15.8 ± 17.2	$p = .15$

Note. All values are presented as mean or mean ± standard deviation.

^aBolded analyses = statistically significant ($p < .05$) with Mann-Whitney U test.




FIGURE 2 Overnight change in the finger-tapping test. (A) The number of correctly tapped sequences for the final three learning trials (10, 11, 12) and test performance, defined as the average number of correct sequences on the three test trials (1, 2, 3), for healthy controls and the narcolepsy subjects. (B) Relative change of accuracy in finger-tapping test performance. Abbreviations: HC, healthy controls; NT1, narcolepsy type 1. Means ± SD are shown

Additional whole-group analyses did not reveal significant correlations of age, depression prevalence, and EDS with overnight memory changes.

3.5 | Sensitivity analysis

Sensitivity analyses were performed to exclude that medical treatment affected the results. Thus, we performed two repeated measures ANOVA on the number of correctly recalled sequences in the finger-tapping task with subjects removed from the group of healthy controls and NT1, respectively. Also without the four subjects in the healthy control group who had a medical treatment (lithium and quetiapine, insulin, somatropin and contraception pills), a significant effect was found for $TIME$ ($F(1, 38) = 26.77, p < .0001$) and $TIME*HEALTH$ ($F(1, 38) = 7.831, p = .0082$) but not $HEALTH$ ($F(1, 38) = 0.72, p = .40$). When we removed subjects in the NT1 group from the analysis, who normally were medicated with sodium oxybate, but refrained from it during the test night, significant effect was still


found for $TIME$ ($F(1, 34) = 22.43, p < .0001$) and $TIME*HEALTH$ ($F(1, 34) = 6.31, p = .017$) but not $HEALTH$ ($F(1, 34) = 0.29, p = .59$).

4 | DISCUSSION

The present results suggest that the sleep-dependent consolidation of procedural but not spatial memory may be impaired among patients with NT1. Although patients with NT1 exhibited significantly worse sleep, we could not find any correlation between the measured sleep variables and memory consolidation. Importantly, although NT1 patients were slightly older, more often depressed, and anxious than healthy controls, none of these variables correlated with overnight memory consolidation.

In previous studies examining procedural memory consolidation in patients with narcolepsy, Mazzetti and coauthors investigated the consolidation of procedural motor skills using the finger-tapping test.^{12,23} They could not observe a significant interaction in their factor analysis (health status and time) in speed (number of

FIGURE 3 Overnight change in the two-dimensional (2-D) memory game. (A) The number of correctly recalled card pairs at learning trials and test, for healthy controls and the narcolepsy subjects. (B) Relative change in 2D memory task performance. Abbreviations: HC, healthy controls; NT1, narcolepsy type 1. Means \pm SD are shown



sequences) or accuracy (number of correct sequences).¹² A post hoc analysis showed an improved overnight speed in the finger-tapping task in controls but not in NT1. In the other study comparing children and adults with narcolepsy, subjects had a learning session in the morning and a retrieval session in the afternoon with one or more free naps allowed.²³ In adults, no improvement was found in finger-tapping task results during the day until the first retrieval session in the late afternoon. When subjects were tested after 24 h, an overnight increase in speed but not accuracy (number of correct sequences) was found. In another study in subjects with narcolepsy, the consolidation of procedural visual discrimination skills using the texture discrimination task was used, but no significant differences were observed.²⁴ These studies used an experimental design where the learning session was set in the morning Day 1, followed by one or two test sessions Day 1 or Day 2. In two of the studies, the test session was performed after sleep in the morning Day 2.^{12,24} In one study, two test sessions were used, one in the evening Day 1 and one in the morning Day 2.²³ Based on studies showing that wakefulness may affect the consolidation of procedural memory,²⁵ the timing of the learning session may explain the different findings compared to our study. Brawn et al.²⁶ showed that finger-tapping performance deteriorates during 1 day of wakefulness and is restored after one night of sleep in healthy subjects. This finding in their study contrasted with the results obtained when the training session was set in the evening, where performance remained stable across a night of sleep. Thus, the conditions in our and previous studies might not be comparable. This difference in experimental setup could explain why we found a group difference in the overnight performance gain of correctly tapped sequences, which was not seen in previous studies.

We performed analyses of possible correlations between change in performance in the two memory tests and the sleep variables, but could not find any correlation. Studies have shown that sleep deprivation reduces the success rate on declarative and procedural memory task.^{4,5} The lower total sleep time (TST) in narcolepsy subjects compared to healthy controls in the present study points toward narcoleptic sleep deprivation. In line with this, we observed less time spent in slow-wave sleep (SWS) and more time in N2, which is in line with published data from studies performed

in sleep laboratories.⁷ SWS is closely associated with hippocampal reactivation and redistribution of newly encoded memories, leading to memory consolidation.^{27,28} Learning of motor skills has also been associated with the proportion of N2.²⁹ Still, no correlation was found between the variables of sleep or other group characteristics of the participants for either of the sleep variable, in accordance with other previous studies.^{12,25,30} The doubtful identification of sleep stage N1 (see below), however, makes this conclusion somewhat uncertain.

Another aspect to consider is the matter of psychiatric comorbidity. A higher prevalence of depression and anxiety in narcolepsy subjects than healthy controls were observed in this study, using the HADS. This overrepresentation of psychiatric comorbidity in narcoleptic patients aligns with previous studies.^{31,32} A study using the same sequential finger-tapping test as in this study showed that patients with depression failed to show any overnight improvement in performance. In contrast, healthy controls improved by ~18%.³³ There was, however, no correlation between the scoring on HADS and memory performance in this study.

As to the 2D memory game, the task relies on temporal lobe structure, including the hippocampus, and has been used to measure overnight declarative memory consolidation.^{22,34} We found a higher degree of forgetting in NT1 subjects than in healthy controls, suggesting that declarative memories are more fragile in NT1. However, in the end, the groups did not significantly differ in their overnight consolidation of spatial memory. As was observed with the finger-tapping task, no correlation was found with any sleep variable.

4.1 | Limitations

In a study of this limited size, we could observe heterogeneity in the severity of NT1 symptoms, the number of medicines used, and the level of psychiatric comorbidity. We acknowledge that the use of antidepressant medication in some NT1 subjects could for example affect sleep quality and thereby influence the results. NT1 subjects were also, on average, 2 years older than the control subjects. Another factor that should be considered when interpreting the results is that the study was performed in at-home settings, and

the environment was not controlled in the same strict manner as possible in a sleep laboratory. A reduced degree of vigilance after wake-up in the morning could possibly also affect the result in NT1 subjects, as compared to healthy controls.

The study was performed using a portable device with a limited amount of validation¹⁶ and only data from healthy individuals. The received output lacked sleep variables such as sleep fragmentation, sleep stage shift index, and arousal index. It also does not report indices of sleep-disordered breathing disorders. The possible correlation between learning and any of these variables could therefore not be determined in this study.

The healthy controls in this study slept close to the approximated standard reported for healthy individuals. We observed that the healthy participants sleep slightly less in N1 and N2 but marginally more in the N3 and REM stages than in other studies.^{35,36} In addition, the NT1 patients in our study slept close to or almost the same amount of time in the different sleep stages as the narcolepsy subjects in other studies.^{7-9,37} One caveat is, however, that the automatic sleep detection largely lacked identification of N1 sleep, as seen in a recent study with the Dreem headband device,³⁸ possibly affecting the distribution of other sleep stages.

When directly comparing the healthy controls and narcolepsy subjects, there are some differences as compared to previous studies. In the present study, the narcolepsy patients slept significantly longer in N2 and shorter in N3 and REM than healthy controls. In contrast, no significant differences were found in time spent in N1, even though the sleep staging of N1 may be inaccurate. This result pattern is different from earlier studies where narcolepsy patients usually sleep significantly longer in N1 and significantly shorter in N3,⁷⁻⁹ whereas N2 and REM were equal in amount. However, a shorter time in N2 and longer time in REM sleep has also been observed in NT1 as compared to healthy control.⁸

5 | CONCLUSION

Our findings suggest that the sleep-dependent consolidation of procedural memories may be impaired among patients with NT1. Therefore, future studies are warranted to examine whether sleep improvement, for example, using sodium oxybate, can aid the sleep-dependent formation of procedural memories among NT1 patients. Although polysomnography is the gold standard method for sleep monitoring in NT1 patients, our data suggest that the Dreem headband used herein may represent a cheap and easily applicable alternative to track sleep in patients with NT1. However, the limitations of the device must be taken into consideration.

ACKNOWLEDGMENTS

We thank Elisa de Mello e Souza Meth and Shervin Bukhari for technical assistance.

CONFLICT OF INTEREST

The authors report there are no competing interests to declare.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/ane.13651>.

DATA AVAILABILITY STATEMENT

Data available upon request.

ORCID

Christian Benedict  <https://orcid.org/0000-0002-8911-4068>

Pontus Wasling  <https://orcid.org/0000-0002-9106-9032>

REFERENCES

- Diekelmann S, Born J. The memory function of sleep. *Nat Rev Neurosci*. 2010;11(2):114-126.
- Ficca G, Lombardo P, Rossi L, Salzarulo P. Morning recall of verbal material depends on prior sleep organization. *Behav Brain Res*. 2000;112(1-2):159-163.
- Backhaus J, Born J, Hoeckesfeld R, Fokuhl S, Hohagen F, Junghanns K. Midlife decline in declarative memory consolidation is correlated with a decline in slow wave sleep. *Learn Mem*. 2007;14(5):336-341.
- Stepan ME, Fenn KM, Altmann EM. Effects of sleep deprivation on procedural errors. *J Exp Psychol Gen*. 2019;148(10):1828-1833.
- Cousins JN, Sasmita K, Chee MWL. Memory encoding is impaired after multiple nights of partial sleep restriction. *J Sleep Res*. 2018;27(1):138-145.
- Nissen C, Kloepper C, Nofzinger EA, Feige B, Voderholzer U, Riemann D. Impaired sleep-related memory consolidation in primary insomnia—a pilot study. *Sleep*. 2006;29(8):1068-1073.
- Rogers AE, Aldrich MS, Caruso CC. Patterns of sleep and wakefulness in treated narcoleptic subjects. *Sleep*. 1994;17(7):590-597.
- Ferri R, Miano S, Bruni O, et al. NREM sleep alterations in narcolepsy/cataplexy. *Clin Neurophysiol*. 2005;116(11):2675-2684.
- Mukai J, Uchida S, Miyazaki S, Nishihara K, Honda Y. Spectral analysis of all-night human sleep EEG in narcoleptic patients and normal subjects. *J Sleep Res*. 2003;12(1):63-71.
- Bixler EO, Kales A, Vela-Bueno A, Drozdiak RA, Jacoby JA, Manfredi RL. Narcolepsy/cataplexy. III: nocturnal sleep and wakefulness patterns. *Int J Neurosci*. 1986;29(3-4):305-316.
- Sorensen GL, Knudsen S, Jennum P. Sleep transitions in hypocretin-deficient narcolepsy. *Sleep*. 2013;36(8):1173-1177.
- Mazzetti M, Plazzi G, Campi C, et al. Sleep-dependent consolidation of motor skills in patients with narcolepsy-cataplexy. *Arch Ital Biol*. 2012;150(2-3):185-193.
- Filardi M, D'Anselmo A, Agnoli S, et al. Cognitive dysfunction in central disorders of hypersomnolence: a systematic review. *Sleep Med Rev*. 2021;59:101510.
- American Academy of Sleep Medicine (AASM). In: Darien I, ed. *International Classification of Sleep Disorders (ICSD)*. 3rd ed. American Academy of Sleep Medicine; 2014.
- Portelius E, Soininen H, Andreasson U, et al. Exploring Alzheimer molecular pathology in Down's syndrome cerebrospinal fluid. *Neurodegener Dis*. 2014;14(2):98-106.
- Arnal PJ, Thorey V, Debellemanni E, et al. The Dreem headband compared to polysomnography for electroencephalographic signal acquisition and sleep staging. *Sleep*. 2020;43(11):zsaa097.
- Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep*. 1991;14(6):540-545.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67(6):361-370.
- Graf P, Schacter DL. Implicit and explicit memory for new associations in normal and amnesic subjects. *J Exp Psychol Learn Mem Cogn*. 1985;11(3):501-518.

20. Cedernaes J, Sand F, Liethof L, et al. Learning and sleep-dependent consolidation of spatial and procedural memories are unaltered in young men under a fixed short sleep schedule. *Neurobiol Learn Mem.* 2016;131:87-94.
21. Squire LR. Memory systems of the brain: a brief history and current perspective. *Neurobiol Learn Mem.* 2004;82(3):171-177.
22. Cedernaes J, Rangtell FH, Axelsson EK, et al. Short sleep makes declarative memories vulnerable to stress in humans. *Sleep.* 2015;38(12):1861-1868.
23. Mazzetti M, Bellucci C, Cipolli C, et al. Age-related differences in sleep-dependent consolidation of motor skills in patients with narcolepsy type 1. *Sleep Med.* 2016;24:80-86.
24. Cipolli C, Campana G, Campi C, et al. Sleep and time course of consolidation of visual discrimination skills in patients with narcolepsy-cataplexy. *J Sleep Res.* 2009;18(2):209-220.
25. Nettersheim A, Hallschmid M, Born J, Diekelmann S. The role of sleep in motor sequence consolidation: stabilization rather than enhancement. *J Neurosci.* 2015;35(17):6696-6702.
26. Brawn TP, Fenn KM, Nusbaum HC, Margoliash D. Consolidating the effects of waking and sleep on motor-sequence learning. *J Neurosci.* 2010;30(42):13977-13982.
27. Huber R, Ghilardi MF, Massimini M, Tononi G. Local sleep and learning. *Nature.* 2004;430(6995):78-81.
28. Sirota A, Csicsvari J, Buhl D, Buzsaki G. Communication between neocortex and hippocampus during sleep in rodents. *Proc Natl Acad Sci USA.* 2003;100(4):2065-2069.
29. Nishida M, Walker MP. Daytime naps, motor memory consolidation and regionally specific sleep spindles. *PLoS One.* 2007;2(4):e341.
30. Rangtell FH, Karamchedu S, Andersson P, et al. Learning performance is linked to procedural memory consolidation across both sleep and wakefulness. *Sci Rep.* 2017;7(1):10234.
31. Wasling HB, Bornstein A, Wasling P. Quality of life and procrastination in post-H1N1 narcolepsy, sporadic narcolepsy and idiopathic hypersomnia, a Swedish cross-sectional study. *Sleep Med.* 2020;76:104-112.
32. Cohen A, Mandrekar J, St Louis EK, Silber MH, Kotagal S. Comorbidities in a community sample of narcolepsy. *Sleep Med.* 2018;43:14-18.
33. Dresler M, Kluge M, Genzel L, Schussler P, Steiger A. Impaired off-line memory consolidation in depression. *Eur Neuropsychopharmacol.* 2010;20(8):553-561.
34. Rasch B, Buchel C, Gais S, Born J. Odor cues during slow-wave sleep prompt declarative memory consolidation. *Science.* 2007;315(5817):1426-1429.
35. Mitterling T, Hogg B, Schonwald SV, et al. Sleep and respiration in 100 healthy Caucasian sleepers - a polysomnographic study according to American Academy of sleep medicine standards. *Sleep.* 2015;38(6):867-875.
36. Hertenstein E, Gabryelska A, Spiegelhalter K, et al. Reference data for polysomnography-measured and subjective sleep in healthy adults. *J Clin Sleep Med.* 2018;14(4):523-532.
37. Zorick F, Roehrs T, Wittig R, Lamphere J, Sicklesteel J, Roth T. Sleep-wake abnormalities in narcolepsy. *Sleep.* 1986;9(1 Pt 2):189-193.
38. Pepin JL, Bailly S, Mordret E, et al. Greatest changes in objective sleep architecture during COVID-19 lockdown in night owls with increased REM sleep. *Sleep.* 2021;44(9):zsab075.

How to cite this article: Asp, A., Lund, F., Benedict, C. & Wasling, P. (2022). Impaired procedural memory in narcolepsy type 1. *Acta Neurologica Scandinavica*, 146, 186–193. <https://doi.org/10.1111/ane.13651>